

A New Structural Model for Multidimensional Default Risk

Chenchen Zhou

The Florida State University

czhou@math.fsu.com


April 12, 2017

Overview

- 1 Motivation
- 2 A New Structural Model
- 3 Multidimensional Default Risk
- 4 Closed Form

Structural versus Reduced Form Models

- Structural
 - Black and Scholes in 1973, Merton 1974, Black and Cox(1976),etc
- Reduced Form
 - Jarrow and Turnbull in 1995



Structural versus Reduced Form Models

- Structural or Reduced Form?
- default time: predictable vs inaccessible
- Robert A.Jarrow and Philip Protter: complete information vs incomplete information

Structural versus Reduced Form Models

Structural Model with Partial Information

Let X_t be the cash balances of the firm, $g(t) := \sup\{s \leq t : X_s = 0\}$ denote the last time (before t) that cash balances hit zero. Let

$$\tau_\alpha := \inf\{t > 0 : t - g(t) \geq \frac{\alpha^2}{2}, \text{ where } X_s < 0 \text{ for } s \in (g(t-), t)\}$$

τ_α is the first time that the firm's cash balances have continued to be negative for at least $\frac{\alpha^2}{2}$ units of time. The default time is defined as

$$\tau := \inf\{t > \tau_\alpha : X_t = 2X_{\tau_\alpha}\}$$

Reduced Form Model

Model Setup

Suppose τ is the default time, $H_t = I_{\tau \leq t}$, and $\mathcal{H}_t = \sigma(H_s : s \leq t)$ a filtration denotes the default time information. Denote $F(t)$ the survival probability, i.e. $F(t) = P(\tau > t)$.

Definition

Hazard Function The function $\Gamma: R^+ \rightarrow R^+$ given by

$$\Gamma(t) = -\log(F(t))$$

is called the hazard function. If F is absolutely continuous, define the intensity function

$$\lambda(t) = \Gamma'(t)$$

$$F(t) = e^{-\int_0^t \lambda(s) ds}$$

- advantage: explicit formulas for survival probabilities

$$P(t, T) = 1_{\tau > t} E^Q[e^{-\int_t^T \lambda(X_s) ds} | \mathcal{F}_t]$$

- interest rate model
- disadvantage: lack economic interpretation

Definition

(*New structural Model: one dimension*) Let S be the stock price of a firm which is described by an exponential Lévy process . We define the time of default as the first time the log-return of S_t jumps below a level $a(t) < 0$.

$$\tau = \inf\{t > 0 : \log S_t / S_{t-} \leq a(t)\}$$

We call $a(t)$ the default level of the firm, $a(t)$ could be stochastic.

- stock price proxy for firm's value
- exponential lévy process to describe stock price
- jump of log return

Lévy Process

Definition

(Lévy Process) A process $X = X_t : t \geq 0$ defined on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ is said to be a Lévy process if it possesses the following properties:

- (i) The paths of X are \mathbb{P} -almost surely right continuous with left limits.
- (ii) $\mathbb{P}(X_0 = 0) = 1$.
- (iii) For $0 \leq s \leq t$, $X_t - X_s$ is equal in distribution to X_{t-s} .
- (iv) For $0 \leq s \leq t$, $X_t - X_s$ is independent of $X_u : u \leq s$.




Figure: A sample path of a Variance Gamma process.

Lévy Process: random measure

Definition

(*Random measure*) Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and (S, \mathcal{S}) a measurable space. The function $M: \Omega \times \mathcal{S} \rightarrow [0, \infty]$ is called a random measure if

- (i) $\forall A \in \mathcal{S}, M(., A)$ is a random variable on $(\Omega, \mathcal{F}, \mathbb{P})$.
- (ii) $\forall \omega \in \Omega, M(\omega, .)$ is a measure on (S, \mathcal{S})

Definition

(*Poisson random measure*) Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and (S, \mathcal{S}, η) a measurable space. A random measure X is a Poisson random measure on \mathcal{S} with intensity η if:

- (i) $\forall A \in \mathcal{S}, X(A)$ is a Poisson random variable on $(\Omega, \mathcal{F}, \mathbb{P})$ with parameter $\eta(A)$.
- (ii) $\forall \{A_i\}_{i=1}^n \in \mathcal{S}$ such that $A_i \cap A_j = \emptyset$ if $i \neq j$, the r.v. $X(A_i)$ and $X(A_j)$ are independent.

Definition

(*Temporal Poisson random measure-TPRM*) Let $(\Omega, \mathcal{F}, \mathbb{P}, (\mathcal{F}_t)_{t \geq 0})$ be a filtered probability space, (E, \mathcal{E}, π) a measure space, and Leb denote Lebesgue measure on $[0, \infty)$. A temporal Poisson random measure (TPRM) X on E with intensity π is an \mathcal{F} -adapted Poisson random measure on $([0, \infty) \times E, \mathcal{B}[0, \infty) \otimes \mathcal{E})$ with intensity $\text{Leb} \times \pi$. For $t \geq 0, A \in \mathcal{E}$ we write $X_t(A) = X([0, t] \times A)$.

With this TPRM, we can define the integral with this measure. For $t \geq 0, A \in \mathcal{E}$, we can write the measure in an integral form

$$X(\omega, [0, t] \times A) = \int_0^t \int_A X(\omega, ds \times dx) \quad (1)$$

Further, we consider integral with measurable function x ,

$$\int_0^t \int_A x X(\omega, ds \times dx) \quad (2)$$

Definition

(Tail Process) Let Y be a Lévy process with Levy measure λ . The tail process Y^a of Y at the level a is the counting process defined by

$$Y_t^a = \#\{0 \leq s \leq t : \Delta Y_s(\omega) \leq a\}, \omega \in \Omega, t \geq 0$$

Important Fact

The tail process Y^a of Y is a Poisson process with mean $\Lambda(a)$ and the survival time is an exponential random variable with parameter $\Lambda(a)$. where $\Lambda(x) = \int_{-\infty}^x \lambda(d\omega)$ is the tail integral of the process Y .

Then the survival probability of the firm up to time $t > 0$ is given by

$$P(\tau > t) = E[e^{-\int_0^t \Lambda(a_u) du}]$$

Further more, for almost every $t \geq 0$, the local default rate LDR_t exists and is given by

$$LDR_t \equiv \lim_{h \downarrow 0} \frac{\mathbb{P}(\tau \leq t + h | \tau > t)}{h} = E(\Lambda(a_t) | \tau > t)$$

Theorem

Let N and M be two Poisson processes on $(\Omega, \mathcal{F}, P, (\mathcal{F}_t)_{t \geq 0})$ with parameters λ and μ . Suppose (N, M) is a two dimensional Poisson process. Then there exists three independent adapted Poisson processes L_1, L_2, L_{12} , with respective parameters $\lambda - \rho$, $\mu - \rho$, ρ , such that,

$$N_t^1 = L_1 + L_{12}$$

$$M_t^2 = L_2 + L_{12}$$

Multidimensional Lévy process

$Y = (Y_t^1, Y_t^2, \dots, Y_t^d)$ is a d dimensional Lévy process, then there exists a vector $\vec{b} \in \mathbb{R}^d$, a unique Σ , and a Temporal Poisson random measure (TPRM) X on $E = \mathbb{R}^d \setminus \{0\}$ with intensity π such that

$$Y_t(\omega) = -\vec{b}t + \Sigma W_t + \int_0^t \int_{|x| \geq 1} x X(ds \times dx) \\ + \lim_{\epsilon \downarrow 0} \int_0^t \int_{\epsilon < |x| < 1} x \{X(dx \times dx) - ds\pi(dx)\} \quad (3)$$

Apply to Tail Processes

$$N_t := (X_1^{a_1(t)}, X_2^{a_2(t)}, X_3^{a_3(t)})$$

where $X_i^{a_i(t)}$ is the tail process of X_i at level $a_i(t)$, $i = 1, 2, 3$. there exists 7 independent adapted Poisson processes L_l

$$N_t^1 = L_1 + L_{12} + L_{13} + L_{123}$$

$$N_t^2 = L_2 + L_{12} + L_{23} + L_{123}$$

$$N_t^3 = L_3 + L_{13} + L_{23} + L_{123}$$

$$\begin{aligned} & \mathbb{P}(\tau_u^1 > x_1, \tau_u^2 > x_2, \tau_u^3 > x_3) \\ &= E[e^{-\int_u^{u+x_1} \Lambda_1(s) ds} e^{-\int_u^{u+x_2} \Lambda_2(s) ds} e^{-\int_u^{u+x_3} \Lambda_3(s) ds} \\ & e^{\int_u^{u+\min(x_1, x_2)} \Lambda_{12}(s) ds} e^{\int_u^{u+\min(x_1, x_3)} \Lambda_{13}(s) ds} e^{\int_u^{u+\min(x_2, x_3)} \Lambda_{23}(s) ds} \\ & e^{-\int_u^{u+\min(x_1, x_2, x_3)} \Lambda_{123}(s) ds} | \mathcal{F}_u] \end{aligned}$$

Idea of poof

$$N_t^1 = \int_0^t \int_{A_1} X(ds \times d\vec{x})$$

$$A_1 = B_1 \cup B_{12} \cup B_{13} \cup B_{123}$$

$$\begin{aligned} N_t^1 &= \int_0^t \int_{B_1} X(ds \times d\vec{x}) + \int_0^t \int_{B_{12}} X(ds \times d\vec{x}) + \int_0^t \int_{B_{13}} X(ds \times d\vec{x}) \\ &\quad + \int_0^t \int_{B_{123}} X(ds \times d\vec{x}) \\ &= L_1 + L_{12} + L_{13} + L_{123} \end{aligned}$$

In summary,

$$N_t^1, N_t^2, N_t^3$$

represented by 7 independent Poisson processes.

Multidimensional Model

Suppose there is a portfolio consisted of d obligors. Consider a d -dimensional stock price vector $\vec{S}_t = (S_t^1, S_t^2, \dots, S_t^d)$ where S_t^i denotes i^{th} firm's stock price. Let τ_i be the default time of name i , τ_i is defined as the first time log-return of S_t^i jumps below a level $a^i(t) < 0$.

$$\tau^i = \inf\{t > 0 : \log S_t^i / S_{t-}^i \leq a^i(t)\}$$

$a^i(t)$ is called default level, $a^i(t)$ could be stochastic.

Assume the price vector \vec{S}_t follows an exponential Lévy process, i.e.

$$S_t^j = S_0^j \exp\{rt + Y_t^j + t\psi^j(-i)\}$$

where S_0^j is the initial stock price, r is interest rate, $Y = (Y_t^1, Y_t^2, \dots, Y_t^d)$ is a d dimensional Lévy process defined on \mathbb{R}^n , ψ^j is the characteristic exponent of Y_1^j , $j = 1, 2, \dots, n$.

If the default level $\vec{a} = (a_1(t), a_2(t), a_3(t))$ is stochastic, then

$$\mathbb{P}(\tau_u^1 > x_1, \tau_u^2 > x_2, \tau_u^3 > x_3) = E[e^{-\int_u^{u+\max(x_1, x_2, x_3)} \vec{1}^T \Lambda^* \vec{1} ds} | \mathcal{F}_u]$$

$\Lambda^* =$

$$\begin{pmatrix} \Lambda_1 1_{s < \min(x_1)} & -\Lambda_{12} 1_{s < \min(x_1, x_2)} & -\Lambda_{13} 1_{s < \min(x_1, x_3)} \\ 0 & \Lambda_2 1_{s < \min(x_2)} & -\Lambda_{23} 1_{s < \min(x_2, x_3)} \\ \Lambda_{123} 1_{s < \min(x_1, x_2, x_3)} & 0 & \Lambda_3 1_{s < \min(x_3)} \end{pmatrix}$$

Multidimensional Risk

- default dependency
- dependency measure
 - correlation coefficient, rank correlation, coefficients of tail dependence.
 - copula function

Definition

(*Copula Functions*) U is a uniform random variable if it has a uniform distribution on the interval $[0,1]$. For d uniform random variables U_1, U_2, \dots, U_d , the joint distribution function C , defined as

$$C(u_1, u_2, \dots, u_d) = P(U_1 \leq u_1, U_2 \leq u_2, \dots, U_d \leq u_d)$$

is called a copula function

Theorem

Sklar's theorem Let F be a joint distribution function with margins F_1, F_2, \dots, F_d , then there exists a copula C such that for all $(x_1, x_2, \dots, x_d) \in R^d$,

$$F(x_1, x_2, \dots, x_d) = C(F_1(x_1), F_2(x_2), \dots, F_d(x_d))$$

Conversely, if C is a copula function and F_1, F_2, \dots, F_d are distributions, then the function F defined by above formula is a joint distribution function with margins F_1, F_2, \dots, F_d .

- Lévy copulas parallels the notion of a copula on the level of Lévy measures.
- Lévy copulas are used to characterize the dependence among components of multidimensional Lévy processes.
- A version of Sklar's theorem: the law of a general multivariate Lévy process is obtained by combining arbitrary univariate Lévy processes with an arbitrary Lévy copula.
- Copulas allow to separate the dependence structure of a random vector from its univariate margins.

Multidimensional Risk

- Assume that the 2 stock prices $S = (S_t^1, S_t^2)$ of 2 firms respectively can be written in terms of a 2 dimensional Lévy process $Y = (Y_t^1, Y_t^2)$
- Default Levels: $\vec{a} = (a_1(t), a_2(t))$
- Joint Survival Probability

$$\mathbb{P}(\tau^1 > t_1, \tau^2 > t_2)$$



- tail integrals

$$\Lambda_{12}(x_1, x_2) := \pi((-\infty, x_1) \times (-\infty, x_2))$$


$$\Lambda_1(x) := \mu_1((-\infty, x))$$

$$\Lambda_2(x) := \mu_2((-\infty, x))$$

Simulation of Two Dimensional Default



Simulation of Two Dimensional Default



survival probability by Lévy copula

$$\mathbb{P}(\tau_u^1 > x_1, \tau_u^2 > x_2, \tau_u^3 > x_3) = E[e^{-\int_u^{u+\max(x_1, x_2, x_3)} \vec{1}^T \Lambda^* \vec{1} ds} | \mathcal{F}_u]$$

where $\Lambda^* =$

$$\begin{pmatrix} \Lambda_1 1_{s < \min(x_1)} & -F^{12}(-\Lambda_1, -\Lambda_2) 1_{s < \min(x_1, x_2)} & -F^{13}(-\Lambda_1, -\Lambda_3) 1_{s < \min(x_1, x_3)} \\ 0 & \Lambda_2 1_{s < \min(x_2)} & -F^{23}(-\Lambda_2, -\Lambda_3) 1_{s < \min(x_2, x_3)} \\ -F^{123}(-\Lambda_1, -\Lambda_2, -\Lambda_3) 1_{s < \min(x_1, x_2, x_3)} & 0 & \Lambda_3 1_{s < \min(x_3)} \end{pmatrix}$$

Conclusion: Two threshold

reduced form model and structural model

Suppose there are d firms. For each obligor $1 \leq i \leq d$, we define

- 1 The default intensity $\lambda^i(t)$: a deterministic function. We usually assume it to be a step function.
- 2 The survival function $S^i(t)$:

$$S^i(t) := \exp\left(-\int_0^t \lambda^i(u) du\right).$$

- 3 The default trigger variables U_i : uniform random variables on $[0,1]$. The d -dimensional vector $U = (U_1, U_2, \dots, U_d)$ is distributed according to the d -dimensional copula C .
- 4 The time of default τ_i of obligor i , where $i = 1, \dots, d$,

$$\tau_i := \inf\{t : S^i(t) \leq 1 - U_i\}.$$

Basket CDS Pricing formula

$$m_t = (1 - R) \frac{\int_t^T E^Q(e^{-\int_t^u r_s + \Lambda(\vec{a}(s)) ds} \cdot (\Lambda(\vec{a}(u))) | \mathcal{H}_t) du}{\int_t^T E^Q(e^{-\int_t^u r_s + \Lambda(\vec{a}(s)) ds} | \mathcal{H}_t) du}$$

$$\begin{aligned} \Lambda(\vec{a}(s)) &= \Lambda_1(a_1) + \Lambda_2(a_2) + \Lambda_3(a_3) \\ &\quad - F_{12}(\bar{\Lambda}(a_1), \bar{\Lambda}(a_2)) - F_{13}(\bar{\Lambda}(a_1), \bar{\Lambda}(a_3)) - F_{23}(\bar{\Lambda}(a_2), \bar{\Lambda}(a_3)) \\ &\quad + F_{123}(\bar{\Lambda}(a_1), \bar{\Lambda}(a_2), \bar{\Lambda}(a_3)) \end{aligned}$$

$$\bar{\Lambda}(x_1, x_2, x_3) = \prod sgn(x_1, x_2, x_3) \lambda \left(\prod_{i=1}^3 \mathcal{I}(x_i) \right)$$

$$E(\exp(-\int_0^T (r_t + \lambda_t)dt))$$

and

$$E(\lambda_t \exp(-\int_0^t (r_u + \lambda_u)du))$$

Interest rate and Default rate

$$\lambda_t = k_t r_t + b_t,$$

where λ_t is the default rate and r_t is interest rate, k_t and b_t are deterministic functions.

One Factor Model

$$dr(t) = k[\theta - r(t)]dt + \sigma dW(t), r(0) = r_0$$

$$P(t, T) = A(t, T)e^{-B(t, T)r(t)},$$

where

$$A(t, T) = \exp\left\{\left(\theta - \frac{\sigma^2}{2k^2}\right)[B(t, T) - T + t] - \frac{\sigma^2}{4k}B(t, T)^2\right\}$$

$$B(t, T) = \frac{1}{k}[1 - e^{-k(T-t)}]$$

Closed Form: One dimensional case

$$m_t = (1 - R) \cdot \frac{f_1}{f_2} \quad (4)$$

$$\begin{aligned} f_1 = & \int_t^T \exp\left\{-r_t \int_t^u (a_s + 1)e^{k(t-s)} ds + \int_t^u (-k\theta_s \int_s^u (a_s + 1)e^{k(s-m)} dm \right. \\ & \left. + \frac{1}{2}\sigma^2 \left(\int_s^u (a_s + 1)e^{k(s-m)} dm\right)^2 - b_s\right) ds\} du \end{aligned} \quad (5)$$

$$\begin{aligned} f_2 = & \int_t^T \exp\left\{-r_t \int_t^u (a_s + 1)e^{k(t-s)} ds + \int_t^u (-k\theta_s \int_s^u (a_s + 1)e^{k(s-m)} dm \right. \\ & \left. + \frac{1}{2}\sigma^2 \left(\int_s^u (a_s + 1)e^{k(s-m)} dm\right)^2 - b_s\right) ds - \ln(a_u r_u + b_u)\} du \end{aligned} \quad (6)$$

Closed Form: Model from Structural Perspective

Interest rate and Default level

$$a_t = k_t r_t + b_t,$$

where a_t is the default level and r_t is interest rate, k_t and b_t are deterministic functions.

Lévy measure

Assume the stock price follows an exponential compound poisson process. We are only interested in jump component

$$\sum_{i=1}^{N_t} U_i$$

where N_t is a Poisson process with intensity λ , and U_i is uniform random on $[-\lambda, 0)$, thus the Lévy measure, which is product of jump arrival rate λ and $\mu(F)$, where F is some jump size set. In this case, the Lévy measure is $\lambda \frac{1}{\lambda} 1_{[-\lambda, 0)}$. So the tail integral would be

$$\Lambda(a_t) = \begin{cases} 0 & a_t < -\lambda \\ a_t + \lambda & -\lambda \leq a_t < 0 \end{cases} \quad (7)$$

Definition

(Clayton Levy Copula, two dimensional case). A Clayton Levy copula is a function $F(u,v)$ with parameter θ, η defined as

$$(|u|^{-\theta} + |v|^{-\theta})^{-\frac{1}{\theta}} (\eta 1_{uv \geq 0} - (1 - \eta) 1_{uv < 0}) \quad (8)$$

where $\theta > 0$ controls the dependence of absolute value of jumps for two levy processes, and $0 \leq \eta \leq 1$ controls measures the correlation between two levy processes. The closer η to 1, the more two levy processes are related.

To get explicit formula, I select $\theta = 1$. So the Clayton copula becomes

$$\frac{|uv|}{|u| + |v|} \eta \quad (9)$$

We assume that

$$|u| = p_k |v| \quad (10)$$

$$|u| \eta \quad (11)$$

$$\begin{aligned}
m_t/(1-R) &= \frac{\int_t^T E^Q(e^{-\int_t^u r_s + 2(a_s + \lambda) - (a_s + \lambda)\eta ds} \cdot (2(a_u + \lambda) - (a_u + \lambda)\eta) | \mathcal{H}_t)}{\int_t^T E^Q(e^{-\int_t^u r_s + 2(a_s + \lambda) - (a_s + \lambda)\eta ds} | \mathcal{H}_t) du} \\
&\frac{E^Q(e^{-\int_t^u r_s + (2-\eta)a_s + 2\lambda - \eta\lambda ds} \cdot ((2-\eta)a_u + 2\lambda - \eta\lambda) | \mathcal{H}_t)}{E^Q(e^{-\int_t^u r_s + (2-\eta)a_s + 2\lambda - \eta\lambda ds} | \mathcal{H}_t)} \\
&\frac{E^Q(e^{-\int_t^u r_s + (2-\eta)(k_t r_s + b_s) + 2\lambda - \eta\lambda ds} \cdot ((2-\eta)(k_u r_u + b_u) + 2\lambda - \eta\lambda) | \mathcal{H}_t)}{E^Q(e^{-\int_t^u r_s + (2-\eta)(k_t r_s + b_s) + 2\lambda - \eta\lambda ds} | \mathcal{H}_t)} \\
&\frac{E^Q(e^{-\int_t^u (1+(2-\eta)k_t)r_s + (2-\eta)b_s + 2\lambda - \eta\lambda ds} \cdot ((2-\eta)k_u r_u + (2-\eta)b_u + 2\lambda - \eta\lambda) | \mathcal{H}_t)}{E^Q(e^{-\int_t^u (1+(2-\eta)k_t)r_s + (2-\eta)b_s + 2\lambda - \eta\lambda ds} | \mathcal{H}_t)}
\end{aligned} \tag{12}$$

Two Factor Model

$$r(t) = x(t) + y(t) + \varphi(t), r(0) = r_0$$

$$dx(t) = -ax(t)dt + \sigma dW_1(t), x(0) = 0,$$

$$dy(t) = -by(t)dt + \eta dW_2(t), y(0) = 0,$$

where (W_1, W_2) is a two-dimensional Brownian motion with instantaneous correlation ρ as from

$$dW_1(t)dW_2(t) = \rho dt,$$

References I



Pierre Garreau and Alec Kercheval.

A structural jump threshold framework for credit risk.
2014.



Peter Tankov.

Simulation and option pricing in lévy copula models.
Mathematical Modelling of Financial Derivatives, IMA Volumes in Mathematics and Applications, Springer, 2006.



R Jarrow and Philip Protter.

Structural versus reduced form models: a new information based perspective.
Journal of Investment management, 2(2):1–10, 2004.



Sudheer Chava and Robert A Jarrow.

Bankruptcy prediction with industry effects.
Review of Finance, 8(4):537–569, 2004.

References II



Umut Cetin, Robert A Jarrow, Philip Protter, and Yildiray Yildirim.
Modeling credit risk with partial information.
Annals of Applied Probability, forthcoming, 2004.

Thank You!