

Exp 6: Surface Analysis

BY YUEJIAN MO

April 20, 2018

Introduction

Here, we will study the surface of Human hemoglobin, including hydrophobicity, conservation and electrostatic potential.

APBS, the Adaptive Poisson-Boltzmann Solver, is a [freely](#) available macromolecular electrostatics calculation program released under the [GPL](#). It is a cost-effective but uncompromised alternative to [GRASP](#), and it can be used within PyMOL. PyMOL can display the results of the calculations as an electrostatic potential molecular surface.

Methods

1. Load human hemoglobin structure (PDB id: 4HHB) in PyMOL

```
PyMOL> fetch 4HHB
```

2. Create new objects containing beta-subunit.

```
PyMOL> select beta, chain B
```

3. Separate human hemoglobin and hemo group into two objects

4. Generate the surfaces to display the two different properties of the protein

- Hydrophobicity

```
PyMOL> set surface_color, white, beta,
```

```
PyMOL> show surface, beta
```

```
PyMOL> select hydrophobicity, resn Ala+Val+Ile+Leu+Met+Phe+Tyr+Trp in  
beta
```

```
PyMOL> remove backbone in hydrophobicity
```

```
PyMOL> set surface_color, blue, hydrophobicity
```

```
PyMOL> show surface, hydrophobicity
```

```
PyMOL> set transparency, 0.2
```

- Conservation

I aligned protein 4HHB-B, 4BJA-A, 4MPM-A, 4MPM-B using structural information by tcoffee. The identical residues were marked as red, and highly similar residues were marked as raspberry, and similar residues were marked as warmpick.

5. Generate the electrostatic potential surfaces for the tetrameric hemoglobin

Action → generate → vacuum electrostatics → protein contact potential (local)

- 6.

```

select hydrophobicity, resn Ala+Val+Ile+Leu+Met+PheTyr+Trp
set surface_color, green, phobic
set transparency, 0.2

```

Results

- 1.
- 2.
- 3.
4. Hydrophobicity

Save the result in Hydrophobicity.pse since 003. The hole which heme insert is hydrophobicity, in order to keep heme inside hemoglobin.(Figure 1) Many sunken surface is hydrophobic(Figure 2), which suggests show the folding occur to decrease system energy. However, here are raised surface is hydrophobic.(Figure 3) The hydrophobic environment between subunit interact edge may explain this exception.




Figure 1. Heme insert to a hydrophobic hole




Figure 2. Much sunken surface is hydrophobic




Figure 3. Subunit interact surface is more hydrophobic

5. Convserision

I find following :

- i. The hemo binding sites are identical convservation.
- ii. Most interaction edge is similar residues.
- iii. At non subunit interaction surface, most loops and trun are conservation




Figure 4. The Hemo binding site is identical convservation




Figure 5. Most interaction edge is similar residues




Figure 6. At no interaction surface,most loops and turn are conservation

6. Electrostatic in vacuum

Because 2,3-BPG is negative charge, I find a positive charged surface between two β subunits.(Figure 4)(Reference 4)(electrostatic.pse sence 3). I find that four Heme interactional surface on hemoglobin are positive charged (electrostatic.pse sence 4). However, Heme doesn't show much negative features.




Figure 7. The positive charged pocket for 2,3-BPG binding. Bottom black circle is a PO_4^{4-} , and top black circle is responding positive surface to hold PO_4^{4-} . (Two β subunits are colored by cyans).




Figure 8. Here are positive charged surface on hemoglobin around a Heme (Black circle).




Figure 9. Heme b group (Ref 2)

7. Electrostatic potential surface using APBS program

Conclusions

References

1. https://pymolwiki.org/index.php/Property_Selectors
2. <https://en.wikipedia.org/wiki/Heme>
3. https://en.wikipedia.org/wiki/2,3-Bisphosphoglyceric_acid
4. http://cbc.chem.arizona.edu/classes/bioc462/462a/NOTES/hemoglobin/hemoglobin_function.htm
5. <https://www.rcsb.org/structure/4HHB>
6. <https://pymolwiki.org/index.php/APBS>