


# REGULATION BY POST- TRANSLATIONAL MODIFICATION

---


**Dr. Zhiyi Wei**  
SUSTC

# Post-translational modification




# Addition of chemical groups



- Phosphorylation
- Methylation
- Acetylation
- Nitrosylation
- Hydroxylation
- ...



# Amino acid specificity in modification




# Methylation




# Regulation by methylation

- A common modification of proteins in eukaryotic nuclei
  - Histone
    - Heterogeneous nuclear ribonucleoprotein (hnRNP)
      - Arginine methylation
- Methyltransferase (“writer”)
  - Using S-adenosylmethionine as the methyl donor
- Demethylase (“eraser”)
- Effects by methylation on Lys/Arg
  - No changes on the overall charge
  - Altering the steric interactions the group can make
  - Eliminating possible hydrogen-bond donors
- Methylation on Lys/Arg creating a new and highly specific protein binding site and altering the protein-protein interaction

# Methyl-lysine readers



# N-acetylation




- Acetyltransferases
  - Using an acetyl group from acetyl-CoA
  - Amino terminus of protein backbone can be modified by sequence specific N-acetyltransferases
  - Lysine sidechain of **histone** can be modified by other specific acetyltransferases
- Deacetylase




# Regulation by N-acetylation

- Histone methylation may induce either an active or an inactive state of chromatin
  - Depending on the position and the nature of the methyl group
- Histone acetylation is always associated with an active state of chromatin
  - Promoted by chromatin-remodeling enzymes recruited to the DNA by proteins containing bromodomains that specifically recognize acetylated lysines
- Unlike methylation, acetylation changes the charge on Lys
- Similar with methylation, acetylation on Lys creating a unique chemical group for protein-protein interaction


# Acetyl-lysine readers






AF9\_YEATS-H3K9ac




BRD4\_Bromo1-H3K14ac



DPF3b\_PHD12-H3K14ac



# Histone is highly modified by methylation and acetylation



# Histone modification and epigenetics




# Cell Signaling Technology


**Epigenetics Regulation & Associated Product Offerings**

# Addition of complex molecules

- Proteins can be modified by other large biomolecules
- By glycan
  - Glycosylation
- By lipid (lipidation)
  - Myristoylation
  - Palmitoylation
  - Prenylation
  - ...
- By nucleotide
  - ADP-ribosylation
  - ...



# Glycosylation




- Glycosylation
  - Attaching carbohydrate chains (oligosaccharides) to asparagine, serine and threonine residues on protein surfaces
  - The most complex and diverse post-translational modification
  - N-linked (Asn) or O-linked (Ser/Thr) types



# Regulation by glycosylation

- Functions for sugars attached to proteins
  - Providing recognition sites that tag glycoproteins for recognition by other proteins
  - Shielding large areas of the protein surface, therefore providing protection from proteases and nonspecific protein–protein interactions
  - In some cases, oligosaccharides increase the solubility of nascent glycoproteins and prevent their aggregation




- Protecting vulnerable regions of the molecule from proteolysis
- Providing binding sites for bacteria, thereby protecting the mucosal surface from infection

IgA protects mucosal surfaces from pathogens





# Oligosaccharide processing in cell

- Almost all secreted and membrane-associated proteins of eukaryotic cells are glycosylated
- Oligosaccharide processing on newly synthesized glycoprotein




## Diverse and complex structures of oligosaccharides on glycoproteins



The structure of Glc3Man9GlcNAc2

# Lipid modification (lipidation)





- **Protein targeting** by lipid modification
  - Covalent attachment of lipids targets proteins to membranes and other proteins



# Four types of lipid modifications

- Classification of lipid modification according to the identity of the attached lipid


- Myristoylation
- Palmitoylation
- Prenylation

Lipid Modifications		Signals	Enzymes
Prenylation	<b>Farnesylation</b> 	—CaaX	Farnesyltransferase Geranylgeranyltransferase I
	<b>Geranylgeranylation</b> 	—CC or —CXC (Rab proteins only)	Geranylgeranyltransferase II
N-myristoylation		MGxxS—	N-myristoyltransferase
Palmitoylation		Poorly defined	Palmitoyltransferase

- The modification by a glycosylphosphatidylinositol (GPI) anchor
  - Attached to the C-terminus of a protein

# GPI anchoring

- A way for protein of glycoprotein transport by vesicle (ER-Golgi)
- GPI modification is reversible
- PLC cleavage is a releasing mechanism



The protein is connected through an amide linkage to a phosphoethanolamine molecule (E-P) that is attached to a core tetrasaccharide composed of three mannose sugars (M) and a single glucosamine sugar (G). The tetrasaccharide is in turn attached to phosphatidylinositol (I)

# GTPase and lipid modification


- Some GTPases are reversibly associated with internal membranes of the cell via lipid modification
- Rab GTPases
  - A large family of small GTPases, regulating membrane traffic
    - vesicle formation, movement, tethering, and membrane fusion
  - Directing transport vesicles between the membrane-bounded compartments of the cell

**Cargo selection/  
budding/coat formation**

**Transport**

**Tethering**

**Fusion**



BIO446 Protein Structure and Function

