# Exp 6: Surface Analysis

BY YUEJIAN MO April 14, 2018

### 1 Introduction

Here, we will study the surface of Huaman hemoglobin, including hydrophobicity, conservation and electrostatic potential.

#### 2 Methods

- 1. Load human hemoglobin structure (PDB id: 4HHB) in PyMOL PyMOL> fetch 4HHB
- 2. Create new objecs containing beta-subunit.

```
PyMOL> select beta, chain B
```

- 3. Separate human hemoglobin and hemo group into two objects
- 4. Generate the surfaces to display the two different properties of the protein
  - Hydrophobicity

```
PyMOL> set surface_color, white, beta,
PyMOL> show surface, beta

PyMOL> select hydrophobicity, resn Ala+Val+IIe+Leu+Met+Phe+Tyr+Trp in beta

PyMOL> remove backbone in hydrophobicity

PyMOL> set surface_color, blue, hydrophobicity

PyMOL> show surface, hydrophobicity

PyMOL> set transparency, 0.2
```

- Conservation
- 5. Generate the electronstatic potential surfaces for the tetrameric hemoglobin

6.

```
select hydrophobicity, resn Ala+Val+lle+Leu+Met+PheTyr+Trp set surface_color, green, phobic set transparency, 0.2
```

#### 3 Results

1.

2.

3.

#### 4. Hydrophobicity

Save the result in Hydrophobicity.pse sence 003. The hole which heme insert is hydrophobicity, in oder to keep heme inside hemoglobin.(Figure 1) Many sunkens surface is hydrophobic(Figure 2), which suggestes show the folding occuar to decrease system energy. However, here are raised surface is hydrophobic.(Figure 3) The hydrophobic environment between subunit interacte edge may explain this exception.




Figure 1. Hem insert to a hydrophobic hole

 ${\bf Figure \ 2.} \ \ {\bf Much \ sunken \ surface \ is \ hydrophobic }$ 




Figure 3. Subunit interacte surface is more hydrophobic

5.

## 4 Conclusions

## 5 References

- $\bullet \quad https://pymolwiki.org/index.php/Property\_Selectors$
- https://en.wikipedia.org/wiki/Heme