



# DISCRETE MATHEMATICS FOR COMPUTER SCIENCE

Dr. QI WANG

Department of Computer Science and Engineering

Office: Room903, Nanshan iPark A7 Building

Email: [wangqi@sustc.edu.cn](mailto:wangqi@sustc.edu.cn)

Please submit your  
assignments before class!



# The Birthday Paradox

- Suppose that 25 students are in a room. What is the probability that at least two of them share a birthday?



# The Birthday Paradox

- Suppose that 25 students are in a room. What is the probability that at least two of them share a birthday?

It's greater than  $1/2$ ! (only need 23)



# The Birthday Paradox

- Suppose that 25 students are in a room. What is the probability that at least two of them share a birthday?

It's greater than  $1/2$ ! (only need 23)

$A_n$  – “there are  $n$  students in a room and at least two of them share a birthday.”



# The Birthday Paradox

- Suppose that 25 students are in a room. What is the probability that at least two of them share a birthday?

It's greater than  $1/2$ ! (only need 23)

$A_n$  – “there are  $n$  students in a room and at least two of them share a birthday.”

We may assume that a year has 365 days and there are no twins in the room.



# The Birthday Paradox

- Suppose that 25 students are in a room. What is the probability that at least two of them share a birthday?

It's greater than  $1/2$ ! (only need 23)

$A_n$  – “there are  $n$  students in a room and at least two of them share a birthday.”

We may assume that a year has 365 days and there are no twins in the room.

This will be very similar to the analysis of hashing  $n$  keys into a table of size 365.



# The Birthday Paradox

- $A_n$  – “there are  $n$  students in a room and at least two of them share a birthday.”

Sample space:  $|S| = 365^n$





# The Birthday Paradox

- $A_n$  – “there are  $n$  students in a room and at least two of them share a birthday.”

Sample space:  $|S| = 365^n$

$B_n$  – “there are  $n$  students in a room and none of them share a birthday.”



# The Birthday Paradox

- $A_n$  – “there are  $n$  students in a room and at least two of them share a birthday.”

Sample space:  $|S| = 365^n$

$B_n$  – “there are  $n$  students in a room and none of them share a birthday.”

$$\#B_n = 365 \times 364 \times \cdots \times (365 - (n - 1))$$



# The Birthday Paradox


- $A_n$  – “there are  $n$  students in a room and at least two of them share a birthday.”

Sample space:  $|S| = 365^n$

$B_n$  – “there are  $n$  students in a room and none of them share a birthday.”


$$\#B_n = 365 \times 364 \times \cdots \times (365 - (n - 1))$$

$$\#A_n + \#B_n = 365^n$$



# The Birthday Paradox

$n$	$A_n$	$B_n$	$n$	$A_n$	$B_n$
1	0.00000000	1.00000000	16	0.28360400	0.71639599
2	0.00273972	0.99726027	17	0.31500766	0.68499233
3	0.00820416	0.99179583	18	0.34691141	0.65308858
4	0.01635591	0.98364408	19	0.37911852	0.62088147
5	0.02713557	0.97286442	20	0.41143838	0.58856161
6	0.04046248	0.95953751	21	0.44368833	0.55631166
7	0.05623570	0.94376429	22	0.47569530	0.52430469
8	0.07433529	0.92566470	23	0.50729723	0.49270276
9	0.09462383	0.90537616	24	0.53834425	0.46165574
10	0.11694817	0.88305182	25	0.56869970	0.43130029
11	0.14114137	0.85885862	26	0.59824082	0.40175917
12	0.16702478	0.83297521	27	0.62685928	0.37314071
13	0.19441027	0.80558972	28	0.65446147	0.34553852
14	0.22310251	0.77689748	29	0.68096853	0.31903146
15	0.25290131	0.74709868	30	0.70631624	0.29368375



# Euclidean Algorithm

- The Euclidean algorithm in pseudocode

## ALGORITHM 1 The Euclidean Algorithm.

```
procedure gcd(a, b: positive integers)
  x := a
  y := b
  while y ≠ 0
    r := x mod y
    x := y
    y := r
  return x{gcd(a, b) is x}
```

The number of **divisions** required to find  $\text{gcd}(a, b)$  is  $O(\log b)$ , where  $a \geq b$ . (this will be proved later.)



# Euclidean Algorithm

- The Euclidean algorithm in pseudocode

## ALGORITHM 1 The Euclidean Algorithm.

```
procedure gcd( $a, b$ : positive integers)
 $x := a$ 
 $y := b$ 
while  $y \neq 0$ 
     $r := x \bmod y$ 
     $x := y$ 
     $y := r$ 
return  $x$  {gcd( $a, b$ ) is  $x$ }
```

The number of **divisions** required to find  $\text{gcd}(a, b)$  is  $O(\log b)$ , where  $a \geq b$ . (this will be proved later.)

**Why ?**



# Euclidean Algorithm

- Key steps in the Euclidean algorithm

$$r_0 = r_1 q_1 + r_2 \quad 0 \leq r_2 < r_1,$$

$$r_1 = r_2 q_2 + r_3 \quad 0 \leq r_3 < r_2,$$

.

.

.

$$r_{n-2} = r_{n-1} q_{n-1} + r_n \quad 0 \leq r_n < r_{n-1},$$

$$r_{n-1} = r_n q_n .$$

# Euclidean Algorithm

- Key steps in the Euclidean algorithm

$$\begin{aligned}r_0 &= r_1 q_1 + r_2 & 0 \leq r_2 < r_1, \\r_1 &= r_2 q_2 + r_3 & 0 \leq r_3 < r_2, \\&\cdot \\&\cdot \\&\cdot \\r_{n-2} &= r_{n-1} q_{n-1} + r_n & 0 \leq r_n < r_{n-1}, \\r_{n-1} &= r_n q_n.\end{aligned}$$

**Observation:**

$$r_{i+2} = r_i \bmod r_{i+1}$$



# Euclidean Algorithm

- Key steps in the Euclidean algorithm

$$\begin{aligned}r_0 &= r_1 q_1 + r_2 & 0 \leq r_2 < r_1, \\r_1 &= r_2 q_2 + r_3 & 0 \leq r_3 < r_2, \\&\cdot \\&\cdot \\&\cdot \\r_{n-2} &= r_{n-1} q_{n-1} + r_n & 0 \leq r_n < r_{n-1}, \\r_{n-1} &= r_n q_n.\end{aligned}$$

**Observation:**

$$r_{i+2} = r_i \bmod r_{i+1}$$

We claim that  $r_{i+2} < \frac{1}{2} r_i$

# Euclidean Algorithm

- Key steps in the Euclidean algorithm

$$\begin{aligned}r_0 &= r_1 q_1 + r_2 & 0 \leq r_2 < r_1, \\r_1 &= r_2 q_2 + r_3 & 0 \leq r_3 < r_2, \\&\vdots \\&\vdots \\&\vdots \\r_{n-2} &= r_{n-1} q_{n-1} + r_n & 0 \leq r_n < r_{n-1}, \\r_{n-1} &= r_n q_n.\end{aligned}$$

**Observation:**

$$r_{i+2} = r_i \bmod r_{i+1}$$

We claim that  $r_{i+2} < \frac{1}{2} r_i$

Case (i):  $r_{i+1} \leq \frac{1}{2} r_i$ :  $r_{i+2} < r_{i+1} \leq \frac{1}{2} r_i$ .

Case (ii):  $r_{i+1} > \frac{1}{2} r_i$ :  $r_{i+2} = r_i \bmod r_{i+1} = r_i - r_{i+1} < \frac{1}{2} r_i$ .

# Euclidean Algorithm

## ■ Key steps in the Euclidean algorithm

$$r_0 = r_1 q_1 + r_2 \quad 0 \leq r_2 < r_1,$$

$$r_1 = r_2 q_2 + r_3 \quad 0 \leq r_3 < r_2,$$

.

.

.

$$r_{n-2} = r_{n-1} q_{n-1} + r_n \quad 0 \leq r_n < r_{n-1},$$

$$r_{n-1} = r_n q_n.$$

See [Theorem 1 p. 347].

## Observation:

$$r_{i+2} = r_i \bmod r_{i+1}$$

We claim that  $r_{i+2} < \frac{1}{2} r_i$

Case (i):  $r_{i+1} \leq \frac{1}{2} r_i$ :  $r_{i+2} < r_{i+1} \leq \frac{1}{2} r_i$ .

Case (ii):  $r_{i+1} > \frac{1}{2} r_i$ :  $r_{i+2} = r_i \bmod r_{i+1} = r_i - r_{i+1} < \frac{1}{2} r_i$ .

# Solving Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree  $k$*  with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where  $c_1, c_2, \dots, c_k$  are real numbers, and  $c_k \neq 0$ .



# Solving Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree  $k$*  with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where  $c_1, c_2, \dots, c_k$  are real numbers, and  $c_k \neq 0$ .

- ◇ **linear**: it is a linear combination of previous terms



# Solving Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree  $k$*  with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where  $c_1, c_2, \dots, c_k$  are real numbers, and  $c_k \neq 0$ .

- ◇ **linear**: it is a linear combination of previous terms
- ◇ **homogeneous**: all terms are multiples of  $a_j$ 's



# Solving Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree  $k$*  with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where  $c_1, c_2, \dots, c_k$  are real numbers, and  $c_k \neq 0$ .

- ◇ **linear**: it is a linear combination of previous terms
- ◇ **homogeneous**: all terms are multiples of  $a_j$ 's
- ◇ **degree  $k$** :  $a_n$  is expressed by the previous  $k$  terms



# Solving Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree  $k$*  with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where  $c_1, c_2, \dots, c_k$  are real numbers, and  $c_k \neq 0$ .

- ◇ **linear**: it is a linear combination of previous terms
- ◇ **homogeneous**: all terms are multiples of  $a_j$ 's
- ◇ **degree  $k$** :  $a_n$  is expressed by the previous  $k$  terms
- ◇ **constant coefficients**: coefficients are constants





# Solving Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree  $k$*  with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where  $c_1, c_2, \dots, c_k$  are real numbers, and  $c_k \neq 0$ .

- ◇ **linear**: it is a linear combination of previous terms
- ◇ **homogeneous**: all terms are multiples of  $a_j$ 's
- ◇ **degree  $k$** :  $a_n$  is expressed by the previous  $k$  terms
- ◇ **constant coefficients**: coefficients are constants

By induction, such a recurrence relation is **uniquely** determined by this recurrence relation, and  **$k$  initial conditions**  $a_0, a_1, \dots, a_{k-1}$ .



# Examples of Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree  $k$*  with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where  $c_1, c_2, \dots, c_k$  are real numbers, and  $c_k \neq 0$ .

## Examples


$$P_n = (1.11)P_{n-1}$$

$$f_n = f_{n-1} + f_{n-2}$$

$$a_n = a_{n-1} + a_{n-2}^2$$

$$H_n = 2H_{n-1} + 1$$

$$B_n = nB_{n-1}$$



# Examples of Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree  $k$*  with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where  $c_1, c_2, \dots, c_k$  are real numbers, and  $c_k \neq 0$ .

## Examples


$$P_n = (1.11)P_{n-1} \quad \text{linear homogeneous recurrence relation of degree 1}$$

$$f_n = f_{n-1} + f_{n-2}$$

$$a_n = a_{n-1} + a_{n-2}^2$$

$$H_n = 2H_{n-1} + 1$$

$$B_n = nB_{n-1}$$



# Examples of Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree  $k$*  with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where  $c_1, c_2, \dots, c_k$  are real numbers, and  $c_k \neq 0$ .

## Examples


$P_n = (1.11)P_{n-1}$       linear homogeneous recurrence relation of degree 1

$f_n = f_{n-1} + f_{n-2}$       linear homogeneous recurrence relation of degree 2

$$a_n = a_{n-1} + a_{n-2}^2$$

$$H_n = 2H_{n-1} + 1$$

$$B_n = nB_{n-1}$$



# Examples of Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree  $k$*  with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where  $c_1, c_2, \dots, c_k$  are real numbers, and  $c_k \neq 0$ .

## Examples


$P_n = (1.11)P_{n-1}$       linear homogeneous recurrence relation of degree 1

$f_n = f_{n-1} + f_{n-2}$       linear homogeneous recurrence relation of degree 2

$a_n = a_{n-1} + a_{n-2}^2$       NOT linear

$H_n = 2H_{n-1} + 1$

$B_n = nB_{n-1}$



# Examples of Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree  $k$*  with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where  $c_1, c_2, \dots, c_k$  are real numbers, and  $c_k \neq 0$ .

## Examples


$P_n = (1.11)P_{n-1}$       linear homogeneous recurrence relation of degree 1

$f_n = f_{n-1} + f_{n-2}$       linear homogeneous recurrence relation of degree 2

$a_n = a_{n-1} + a_{n-2}^2$       NOT linear

$H_n = 2H_{n-1} + 1$       NOT homogeneous

$B_n = nB_{n-1}$



# Examples of Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree  $k$*  with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where  $c_1, c_2, \dots, c_k$  are real numbers, and  $c_k \neq 0$ .

## Examples


$P_n = (1.11)P_{n-1}$       linear homogeneous recurrence relation of degree 1

$f_n = f_{n-1} + f_{n-2}$       linear homogeneous recurrence relation of degree 2

$a_n = a_{n-1} + a_{n-2}^2$       NOT linear

$H_n = 2H_{n-1} + 1$       NOT homogeneous

$B_n = nB_{n-1}$       coefficients are not constants



# Solving Linear Recurrence Relations

- **Example** Consider the recurrence relation

$$a_n = 2a_{n-1} - a_{n-2},$$

Which of the following are solutions?

◇  $a_n = 3n$ :

◇  $a_n = 2^n$ :

◇  $a_n = 5$ :





# Solving Linear Recurrence Relations

- **Example** Consider the recurrence relation

$$a_n = 2a_{n-1} - a_{n-2},$$

Which of the following are solutions?

◇  $a_n = 3n$ : YES

◇  $a_n = 2^n$ : NO

◇  $a_n = 5$ : YES



# Solving Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree  $k$*  with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where  $c_1, c_2, \dots, c_k$  are real numbers, and  $c_k \neq 0$ .



# Solving Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree  $k$*  with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where  $c_1, c_2, \dots, c_k$  are real numbers, and  $c_k \neq 0$ .

**Basic idea:** Look for solutions of the form  $a_n = r^n$ , where  $r$  is a constant.



# Solving Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree  $k$*  with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$


where  $c_1, c_2, \dots, c_k$  are real numbers, and  $c_k \neq 0$ .

**Basic idea:** Look for solutions of the form  $a_n = r^n$ , where  $r$  is a constant.

◇ Bring  $a_n = r^n$  back to the recurrence relation:

i.e.,

$$r^n = c_1 r^{n-1} + c_2 r^{n-2} + \cdots + c_k r^{n-k},$$
$$r^{n-k} (r^k - c_1 r^{k-1} - \cdots - c_k) = 0$$



# Solving Linear Recurrence Relations

- **Definition** A *linear homogeneous relation of degree  $k$*  with *constant coefficients* is a recurrence relation of the form

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k},$$

where  $c_1, c_2, \dots, c_k$  are real numbers, and  $c_k \neq 0$ .

**Basic idea:** Look for solutions of the form  $a_n = r^n$ , where  $r$  is a constant.


- ◇ Bring  $a_n = r^n$  back to the recurrence relation:

i.e.,

$$r^n = c_1 r^{n-1} + c_2 r^{n-2} + \cdots + c_k r^{n-k},$$
$$r^{n-k}(r^k - c_1 r^{k-1} - \cdots - c_k) = 0$$

- ◇ The solutions to the *characteristic equation* can yield an explicit formula for the sequence.

$$(r^k - c_1 r^{k-1} - \cdots - c_k) = 0$$



# Recall: Problem IV

## ■ Fibonacci number

$$F_0 = 0, F_1 = 1, F_n = F_{n-1} + F_{n-2} \text{ for } n \geq 2$$



# Recall: Problem IV

## ■ Fibonacci number

$$F_0 = 0, F_1 = 1, F_n = F_{n-1} + F_{n-2} \text{ for } n \geq 2$$

◇ What is the closed-form expression of  $F_n$ ?



# Recall: Problem IV

## ■ Fibonacci number

$$F_0 = 0, F_1 = 1, F_n = F_{n-1} + F_{n-2} \text{ for } n \geq 2$$

◇ What is the closed-form expression of  $F_n$ ?

Consider  $x^n = x^{n-1} + x^{n-2}$ , with  $x \neq 0$ . There are two different roots

$$\phi = \frac{1 + \sqrt{5}}{2}, \quad \psi = \frac{1 - \sqrt{5}}{2}$$

Then  $F_n$  can be the form of  $a\phi^n + b\psi^n$ . By  $F_0 = 0$  and  $F_1 = 1$ , we have  $a + b = 0$  and  $\phi a + \psi b = 1$ , leading to  $a = \frac{1}{\sqrt{5}}$ ,  $b = -a$ . Therefore,

$$F_n = \frac{\phi^n - \psi^n}{\sqrt{5}}$$





# Solving Linear Recurrence Relations of degree 2

- Consider an arbitrary linear homogeneous relation of degree 2 with constant coefficients:

$$a_n = c_1 a_{n-1} + c_2 a_{n-2}.$$



# Solving Linear Recurrence Relations of degree 2

- Consider an arbitrary linear homogeneous relation of degree 2 with constant coefficients:

$$a_n = c_1 a_{n-1} + c_2 a_{n-2}.$$

The characteristic equation (CE) is:

$$r^2 - c_1 r - c_2 = 0.$$



# Solving Linear Recurrence Relations of degree 2

- Consider an arbitrary linear homogeneous relation of degree 2 with constant coefficients:

$$a_n = c_1 a_{n-1} + c_2 a_{n-2}.$$

The characteristic equation (CE) is:

$$r^2 - c_1 r - c_2 = 0.$$

**Theorem** If this CE has 2 roots  $r_1 \neq r_2$ , then the sequence  $\{a_n\}$  is a solution of the recurrence relation **if and only if**  $a_n = \alpha_1 r_1^n + \alpha_2 r_2^n$  for  $n \geq 0$  and constants  $\alpha_1, \alpha_2$ .



# Solving Linear Recurrence Relations of degree 2

- Consider an arbitrary linear homogeneous relation of degree 2 with constant coefficients:

$$a_n = c_1 a_{n-1} + c_2 a_{n-2}.$$

The characteristic equation (CE) is:

$$r^2 - c_1 r - c_2 = 0.$$

**Theorem** If this CE has 2 roots  $r_1 \neq r_2$ , then the sequence  $\{a_n\}$  is a solution of the recurrence relation **if and only if**  $a_n = \alpha_1 r_1^n + \alpha_2 r_2^n$  for  $n \geq 0$  and constants  $\alpha_1, \alpha_2$ .

**Proof?**



# Solving Linear Recurrence Relations of degree 2

- **Example 1**  $a_n = a_{n-1} + 2a_{n-2}$ , with  $a_0 = 2$ ,  $a_1 = 7$



# Solving Linear Recurrence Relations of degree 2

- **Example 1**  $a_n = a_{n-1} + 2a_{n-2}$ , with  $a_0 = 2$ ,  $a_1 = 7$

The *characteristic equation* is

$$r^2 - r - 2 = 0.$$



# Solving Linear Recurrence Relations of degree 2

- **Example 1**  $a_n = a_{n-1} + 2a_{n-2}$ , with  $a_0 = 2$ ,  $a_1 = 7$

The *characteristic equation* is

$$r^2 - r - 2 = 0.$$

Two roots are 2 and  $-1$ . So, assume that

$$a_n = \alpha_1 2^n + \alpha_2 (-1)^n.$$



# Solving Linear Recurrence Relations of degree 2

- **Example 1**  $a_n = a_{n-1} + 2a_{n-2}$ , with  $a_0 = 2$ ,  $a_1 = 7$

The *characteristic equation* is

$$r^2 - r - 2 = 0.$$

Two roots are 2 and  $-1$ . So, assume that

$$a_n = \alpha_1 2^n + \alpha_2 (-1)^n.$$

By the two initial conditions, we have

$$a_0 = \alpha_1 + \alpha_2 = 2$$

$$a_1 = 2\alpha_1 - \alpha_2 = 7$$





# Solving Linear Recurrence Relations of degree 2

- **Example 1**  $a_n = a_{n-1} + 2a_{n-2}$ , with  $a_0 = 2$ ,  $a_1 = 7$

The *characteristic equation* is

$$r^2 - r - 2 = 0.$$

Two roots are 2 and  $-1$ . So, assume that

$$a_n = \alpha_1 2^n + \alpha_2 (-1)^n.$$

By the two initial conditions, we have

$$a_0 = \alpha_1 + \alpha_2 = 2$$

$$a_1 = 2\alpha_1 - \alpha_2 = 7$$

We get  $\alpha_1 = 3$  and  $\alpha_2 = -1$ . Thus,

$$a_n = 3 \cdot 2^n - (-1)^n$$



# Solving Linear Recurrence Relations of degree 2

- **Example 2**  $a_n = 7a_{n-1} - 10a_{n-2}$ , with  $a_0 = 2$ ,  $a_1 = 1$



# Solving Linear Recurrence Relations of degree 2

- **Example 2**  $a_n = 7a_{n-1} - 10a_{n-2}$ , with  $a_0 = 2$ ,  $a_1 = 1$

The *characteristic equation* is

$$r^2 - 7r + 10 = 0.$$



# Solving Linear Recurrence Relations of degree 2

- **Example 2**  $a_n = 7a_{n-1} - 10a_{n-2}$ , with  $a_0 = 2$ ,  $a_1 = 1$

The *characteristic equation* is

$$r^2 - 7r + 10 = 0.$$

Two roots are 2 and 5. So, assume that

$$a_n = \alpha_1 2^n + \alpha_2 5^n.$$



# Solving Linear Recurrence Relations of degree 2

- **Example 2**  $a_n = 7a_{n-1} - 10a_{n-2}$ , with  $a_0 = 2$ ,  $a_1 = 1$

The *characteristic equation* is

$$r^2 - 7r + 10 = 0.$$

Two roots are 2 and 5. So, assume that

$$a_n = \alpha_1 2^n + \alpha_2 5^n.$$

By the two initial conditions, we have

$$a_0 = \alpha_1 + \alpha_2 = 2$$

$$a_1 = 2\alpha_1 + 5\alpha_2 = 1$$



# Solving Linear Recurrence Relations of degree 2

- **Example 2**  $a_n = 7a_{n-1} - 10a_{n-2}$ , with  $a_0 = 2$ ,  $a_1 = 1$

The *characteristic equation* is

$$r^2 - 7r + 10 = 0.$$

Two roots are 2 and 5. So, assume that

$$a_n = \alpha_1 2^n + \alpha_2 5^n.$$

By the two initial conditions, we have

$$a_0 = \alpha_1 + \alpha_2 = 2$$

$$a_1 = 2\alpha_1 + 5\alpha_2 = 1$$

We get  $\alpha_1 = 3$  and  $\alpha_2 = -1$ . Thus,

$$a_n = 3 \cdot 2^n - 5^n$$



# Solving Linear Recurrence Relations of degree $k$

- Consider an arbitrary linear homogeneous relation of degree  $k$  with constant coefficients:

$$a_n = \sum_{i=1}^k c_i a_{n-i}.$$



# Solving Linear Recurrence Relations of degree $k$

- Consider an arbitrary linear homogeneous relation of degree  $k$  with constant coefficients:

$$a_n = \sum_{i=1}^k c_i a_{n-i}.$$

The characteristic equation (CE) is:

$$r^k - \sum_{i=1}^k c_i r^{k-i} = 0.$$





# Solving Linear Recurrence Relations of degree $k$

- Consider an arbitrary linear homogeneous relation of degree  $k$  with constant coefficients:

$$a_n = \sum_{i=1}^k c_i a_{n-i}.$$

The characteristic equation (CE) is:

$$r^k - \sum_{i=1}^k c_i r^{k-i} = 0.$$

**Theorem** If this CE has  $k$  distinct roots  $r_i$ , then the solutions to the recurrence are of the form

$$a_n = \sum_{i=1}^k \alpha_i r_i^n$$

for all  $n \geq 0$ , where the  $\alpha_i$ 's are constants.



# Solving Linear Recurrence Relations of degree $k$

- Consider an arbitrary linear homogeneous relation of degree  $k$  with constant coefficients:

$$a_n = \sum_{i=1}^k c_i a_{n-i}.$$

The characteristic equation (CE) is:

$$r^k - \sum_{i=1}^k c_i r^{k-i} = 0.$$

**Theorem** If this CE has  $k$  distinct roots  $r_i$ , then the solutions to the recurrence are of the form

$$a_n = \sum_{i=1}^k \alpha_i r_i^n$$

for all  $n \geq 0$ , where the  $\alpha_i$ 's are constants.

**Example**  $a_n = 2a_{n-1} + 5a_{n-2} - 6a_{n-3}$



# The Case of Degenerate Roots

- **Theorem** If the CE  $r^2 - c_1 r - c_2 = 0$  has **only 1** root  $r_0$ , then

$$a_n = \alpha_1 r_0^n + \alpha_2 n r_0^n,$$

for all  $n \geq 0$  and two constants  $\alpha_1$  and  $\alpha_2$ .



# The Case of Degenerate Roots

- **Theorem** If the CE  $r^2 - c_1 r - c_2 = 0$  has **only 1** root  $r_0$ , then

$$a_n = \alpha_1 r_0^n + \alpha_2 n r_0^n,$$

for all  $n \geq 0$  and two constants  $\alpha_1$  and  $\alpha_2$ .

**Proof?**



# The Case of Degenerate Roots

- **Example**  $a_n = 4a_{n-1} - 4a_{n-2}$ , with  $a_0 = 1$ ,  $a_1 = 0$



# The Case of Degenerate Roots

- **Example**  $a_n = 4a_{n-1} - 4a_{n-2}$ , with  $a_0 = 1$ ,  $a_1 = 0$

The *characteristic equation* is

$$r^2 - 4r + 4 = 0.$$



# The Case of Degenerate Roots

- **Example**  $a_n = 4a_{n-1} - 4a_{n-2}$ , with  $a_0 = 1$ ,  $a_1 = 0$

The *characteristic equation* is

$$r^2 - 4r + 4 = 0.$$

The only root is 2. So, assume that

$$a_n = \alpha_1 2^n + \alpha_2 n 2^n.$$



# The Case of Degenerate Roots

- **Example**  $a_n = 4a_{n-1} - 4a_{n-2}$ , with  $a_0 = 1$ ,  $a_1 = 0$

The *characteristic equation* is

$$r^2 - 4r + 4 = 0.$$

The only root is 2. So, assume that

$$a_n = \alpha_1 2^n + \alpha_2 n 2^n.$$

By the two initial conditions, we have

$$a_0 = \alpha_1 = 1$$

$$a_1 = 2\alpha_1 + 2\alpha_2 = 0$$





# The Case of Degenerate Roots

- **Example**  $a_n = 4a_{n-1} - 4a_{n-2}$ , with  $a_0 = 1$ ,  $a_1 = 0$

The *characteristic equation* is

$$r^2 - 4r + 4 = 0.$$

The only root is 2. So, assume that

$$a_n = \alpha_1 2^n + \alpha_2 n 2^n.$$

By the two initial conditions, we have

$$a_0 = \alpha_1 = 1$$

$$a_1 = 2\alpha_1 + 2\alpha_2 = 0$$

We get  $\alpha_1 = 1$  and  $\alpha_2 = -1$ . Thus,

$$a_n = 2^n - n2^n$$



# The Case of Degenerate Roots in General

- **Theorem** [Theorem 4, p.519] Suppose that there are  $t$  roots  $r_1, \dots, r_t$  with multiplicities  $m_1, \dots, m_t$ . Then

$$a_n = \sum_{i=1}^t \left( \sum_{j=0}^{m_i-1} \alpha_{i,j} n^j \right) r_i^n,$$

for all  $n \geq 0$  and constants  $\alpha_{i,j}$ .



# The Case of Degenerate Roots in General

- **Theorem** [Theorem 4, p.519] Suppose that there are  $t$  roots  $r_1, \dots, r_t$  with multiplicities  $m_1, \dots, m_t$ . Then

$$a_n = \sum_{i=1}^t \left( \sum_{j=0}^{m_i-1} \alpha_{i,j} n^j \right) r_i^n,$$

for all  $n \geq 0$  and constants  $\alpha_{i,j}$ .

## Example

$$a_n = -3a_{n-1} - 3a_{n-2} - a_{n-3} \text{ with } a_0 = 1, a_1 = -2, \\ a_2 = -1$$



# Linear Nonhomogeneous Recurrence Relations

- **Definition** A *linear nonhomogeneous relation* with constant coefficients may contain some terms  $F(n)$  that depend only on  $n$

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k} + F(n).$$

The recurrence relation

$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k}$  is called the *associated homogeneous recurrence relation*.



# Linear Nonhomogeneous Recurrence Relations

- **Theorem** If  $a_n = p(n)$  is any particular solution to the linear nonhomogeneous relation with constant coefficients,

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k} + F(n),$$

Then all its solutions are of the form

$$a_n = p(n) + h(n),$$

where  $a_n = h(n)$  is any solution to the associated homogeneous recurrence relation

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_k a_{n-k}.$$



# Solving Linear Nonhomogeneous Recurrence Relations

- **Example**  $a_n = 3a_{n-1} + 2n$ . Which solution has  $a_1 = 3$ ?



# Solving Linear Nonhomogeneous Recurrence Relations

- **Example**  $a_n = 3a_{n-1} + 2n$ . Which solution has  $a_1 = 3$ ?

The *characteristic equation* of the associated linear homogeneous recurrence relation is  $r^2 - 3r = 0$ . Thus, the solution to the original problem are all of the form

$$a_n = \alpha 3^n + p(n) .$$



# Solving Linear Nonhomogeneous Recurrence Relations

- **Example**  $a_n = 3a_{n-1} + 2n$ . Which solution has  $a_1 = 3$ ?

The *characteristic equation* of the associated linear homogeneous recurrence relation is  $r^2 - 3r = 0$ . Thus, the solution to the original problem are all of the form

$$a_n = \alpha 3^n + p(n) .$$

We try a degree- $t$  polynomial as the particular solution  $p(n)$ .





# Solving Linear Nonhomogeneous Recurrence Relations


- **Example**  $a_n = 3a_{n-1} + 2n$ . Which solution has  $a_1 = 3$ ?

The *characteristic equation* of the associated linear homogeneous recurrence relation is  $r^2 - 3r = 0$ . Thus, the solution to the original problem are all of the form  $a_n = \alpha 3^n + p(n)$ .

We try a degree- $t$  polynomial as the particular solution  $p(n)$ .

Let  $p(n) = cn + d$ , then

$$cn + d = 3(c(n-1) + d) + 2n, \text{ which means } (-2c + 2)n + (3c - 2d) = 0.$$



# Solving Linear Nonhomogeneous Recurrence Relations

- **Example**  $a_n = 3a_{n-1} + 2n$ . Which solution has  $a_1 = 3$ ?

The *characteristic equation* of the associated linear homogeneous recurrence relation is  $r^2 - 3r = 0$ . Thus, the solution to the original problem are all of the form  $a_n = \alpha 3^n + p(n)$ .


We try a degree- $t$  polynomial as the particular solution  $p(n)$ .

Let  $p(n) = cn + d$ , then

$$cn + d = 3(c(n-1) + d) + 2n, \text{ which means } (-2c + 2)n + (3c - 2d) = 0.$$

We get  $c = 1$  and  $d = 3/2$ . Thus,

$$p(n) = n + 3/2$$



# Generating Functions

- We may use *generating functions* to **characterize** sequences.



# Generating Functions

- We may use *generating functions* to **characterize** sequences.
  - ◇ The sequence  $\{a_k\}$  with  $a_k = 3$



# Generating Functions

- We may use *generating functions* to **characterize** sequences.

◇ The sequence  $\{a_k\}$  with  $a_k = 3$

$$\sum_{k=0}^{\infty} 3x^k$$



# Generating Functions

- We may use *generating functions* to **characterize** sequences.

- ◇ The sequence  $\{a_k\}$  with  $a_k = 3$

$$\sum_{k=0}^{\infty} 3x^k$$

- ◇ The sequence  $\{a_k\}$  with  $a_k = 2^k$



# Generating Functions

- We may use *generating functions* to **characterize** sequences.

◇ The sequence  $\{a_k\}$  with  $a_k = 3$

$$\sum_{k=0}^{\infty} 3x^k$$

◇ The sequence  $\{a_k\}$  with  $a_k = 2^k$

$$\sum_{k=0}^{\infty} 2^k x^k$$



# Generating Functions

- We may use *generating functions* to **characterize** sequences.

◇ The sequence  $\{a_k\}$  with  $a_k = 3$

$$\sum_{k=0}^{\infty} 3x^k$$

◇ The sequence  $\{a_k\}$  with  $a_k = 2^k$

$$\sum_{k=0}^{\infty} 2^k x^k$$

**Definition** The *generating function* for the sequence  $a_0, a_1, \dots, a_k, \dots$  of real numbers is the infinite series

$$G(x) = a_0 + a_1x + \dots + a_kx^k + \dots = \sum_{k=0}^{\infty} a_kx^k$$





# Generating Functions for Finite Sequences

- A finite sequence  $a_0, a_1, \dots, a_n$  can be easily extended by setting  $a_{n+1} = a_{n+2} = \dots = 0$



# Generating Functions for Finite Sequences

- A finite sequence  $a_0, a_1, \dots, a_n$  can be easily extended by setting  $a_{n+1} = a_{n+2} = \dots = 0$

The *generating function*  $G(x)$  of this infinite sequence  $\{a_n\}$  is a polynomial of degree  $n$ , i.e.,

$$G(x) = a_0 + a_1x + \dots + a_nx^n$$



# Generating Functions for Finite Sequences

- A finite sequence  $a_0, a_1, \dots, a_n$  can be easily extended by setting  $a_{n+1} = a_{n+2} = \dots = 0$

The *generating function*  $G(x)$  of this infinite sequence  $\{a_n\}$  is a polynomial of degree  $n$ , i.e.,

$$G(x) = a_0 + a_1x + \dots + a_nx^n$$

- ◇ What is the generating function for the sequence  $a_0, a_1, \dots, a_m$ , with  $a_k = C(m, k)$ ?



# Generating Functions for Finite Sequences


- A finite sequence  $a_0, a_1, \dots, a_n$  can be easily extended by setting  $a_{n+1} = a_{n+2} = \dots = 0$

The generating function  $G(x)$  of this infinite sequence  $\{a_n\}$  is a polynomial of degree  $n$ , i.e.,

$$G(x) = a_0 + a_1x + \dots + a_nx^n$$

- ◇ What is the generating function for the sequence  $a_0, a_1, \dots, a_m$ , with  $a_k = C(m, k)$ ?

$$G(x) = C(m, 0) + \dots + C(m, m)x^m = (1 + x)^m$$



# Examples

◇  $G(x) = 1/(1 - x)$  for  $|x| < 1$



# Examples

◇  $G(x) = 1/(1 - x)$  for  $|x| < 1$

$1, 1, 1, 1, 1, \dots$



# Examples

◇  $G(x) = 1/(1 - x)$  for  $|x| < 1$

$1, 1, 1, 1, 1, \dots$

◇  $G(x) = 1/(1 - ax)$  for  $|ax| < 1$



# Examples

◇  $G(x) = 1/(1 - x)$  for  $|x| < 1$

$$1, 1, 1, 1, 1, \dots$$

◇  $G(x) = 1/(1 - ax)$  for  $|ax| < 1$

$$1, a, a^2, a^3, a^4, \dots$$





# Examples


◇  $G(x) = 1/(1 - x)$  for  $|x| < 1$

$$1, 1, 1, 1, 1, \dots$$

◇  $G(x) = 1/(1 - ax)$  for  $|ax| < 1$

$$1, a, a^2, a^3, a^4, \dots$$

◇  $G(x) = 1/(1 - x)^2$  for  $|x| < 1$



# Examples

◇  $G(x) = 1/(1 - x)$  for  $|x| < 1$


$$1, 1, 1, 1, 1, \dots$$

◇  $G(x) = 1/(1 - ax)$  for  $|ax| < 1$

$$1, a, a^2, a^3, a^4, \dots$$

◇  $G(x) = 1/(1 - x)^2$  for  $|x| < 1$

$$1, 2, 3, 4, 5, \dots$$



# Operations of Generating Functions

- **Theorem** Let  $f(x) = \sum_{k=0}^{\infty} a_k x^k$ , and  $g(x) = \sum_{k=0}^{\infty} b_k x^k$ .  
Then

$$f(x) + g(x) = \sum_{k=0}^{\infty} (a_k + b_k) x^k$$

$$f(x)g(x) = \sum_{k=0}^{\infty} \left( \sum_{j=0}^k a_j b_{k-j} \right) x^k$$



# Operations of Generating Functions

- **Theorem** Let  $f(x) = \sum_{k=0}^{\infty} a_k x^k$ , and  $g(x) = \sum_{k=0}^{\infty} b_k x^k$ .  
Then

$$f(x) + g(x) = \sum_{k=0}^{\infty} (a_k + b_k) x^k$$

$$f(x)g(x) = \sum_{k=0}^{\infty} \left( \sum_{j=0}^k a_j b_{k-j} \right) x^k$$

◇  $G(x) = 1/(1-x)^2$  for  $|x| < 1$



# Operations of Generating Functions

- **Theorem** Let  $f(x) = \sum_{k=0}^{\infty} a_k x^k$ , and  $g(x) = \sum_{k=0}^{\infty} b_k x^k$ .  
Then

$$f(x) + g(x) = \sum_{k=0}^{\infty} (a_k + b_k) x^k$$

$$f(x)g(x) = \sum_{k=0}^{\infty} \left( \sum_{j=0}^k a_j b_{k-j} \right) x^k$$

◇  $G(x) = 1/(1-x)^2$  for  $|x| < 1$

$$f(x) = 1/(1-x), \quad g(x) = 1/(1-x)$$

$$G(x) = f(x)g(x) = \sum_{k=0}^{\infty} \left( \sum_{j=0}^k 1 \right) x^k = \sum_{k=0}^{\infty} (k+1) x^k$$



# Operations of Generating Functions

- **Theorem** Let  $f(x) = \sum_{k=0}^{\infty} a_k x^k$ , and  $g(x) = \sum_{k=0}^{\infty} b_k x^k$ .  
Then

$$f(x) + g(x) = \sum_{k=0}^{\infty} (a_k + b_k) x^k$$

$$f(x)g(x) = \sum_{k=0}^{\infty} \left( \sum_{j=0}^k a_j b_{k-j} \right) x^k$$

◇  $G(x) = 1/(1 - ax)^2$  for  $|x| < 1$



# Operations of Generating Functions

- **Theorem** Let  $f(x) = \sum_{k=0}^{\infty} a_k x^k$ , and  $g(x) = \sum_{k=0}^{\infty} b_k x^k$ .  
Then


$$f(x) + g(x) = \sum_{k=0}^{\infty} (a_k + b_k) x^k$$

$$f(x)g(x) = \sum_{k=0}^{\infty} \left( \sum_{j=0}^k a_j b_{k-j} \right) x^k$$

◇  $G(x) = 1/(1 - ax)^2$  for  $|x| < 1$

$$f(x) = 1/(1 - ax), \quad g(x) = 1/(1 - ax)$$

$$G(x) = f(x)g(x) = \sum_{k=0}^{\infty} \left( \sum_{j=0}^k a^j a^{k-j} \right) x^k = \sum_{k=0}^{\infty} (k+1) a^k x^k$$



# Useful Generating Functions

$$(1 + x)^n = \sum_{k=0}^n C(n, k) x^k$$

$$(1 + ax)^n = \sum_{k=0}^n C(n, k) a^k x^k$$

$$(1 + x^r)^n = \sum_{k=0}^n C(n, k) x^{rk}$$





# Useful Generating Functions

$$(1 + x)^n = \sum_{k=0}^n C(n, k) x^k$$

$$(1 + ax)^n = \sum_{k=0}^n C(n, k) a^k x^k$$


$$(1 + x^r)^n = \sum_{k=0}^n C(n, k) x^{rk}$$

$$\frac{1-x^{n+1}}{1-x} = \sum_{k=0}^n x^k = 1 + x + x^2 + \dots + x^n$$

$$\frac{1}{1-x} = \sum_{k=0}^{\infty} x^k = 1 + x + x^2 + \dots$$

$$\frac{1}{1-ax} = \sum_{k=0}^{\infty} a^k x^k = 1 + ax + a^2 x^2 + \dots$$

$$\frac{1}{1-x^r} = \sum_{k=0}^{\infty} x^{rk} = 1 + x^r + x^{2r} + \dots$$



# Useful Generating Functions

$$(1+x)^n = \sum_{k=0}^n C(n, k)x^k$$

$$(1+ax)^n = \sum_{k=0}^n C(n, k)a^k x^k$$

$$(1+x^r)^n = \sum_{k=0}^n C(n, k)x^{rk}$$

$$\frac{1-x^{n+1}}{1-x} = \sum_{k=0}^n x^k = 1 + x + x^2 + \dots + x^n$$

$$\frac{1}{1-x} = \sum_{k=0}^{\infty} x^k = 1 + x + x^2 + \dots$$

$$\frac{1}{1-ax} = \sum_{k=0}^{\infty} a^k x^k = 1 + ax + a^2 x^2 + \dots$$

$$\frac{1}{1-x^r} = \sum_{k=0}^{\infty} x^{rk} = 1 + x^r + x^{2r} + \dots$$

$$\frac{1}{(1-x)^2} = \sum_{k=0}^{\infty} (k+1)x^k = 1 + 2x + 3x^2 + \dots$$




# Useful Generating Functions

$$\frac{1}{(1-x)^n} = \sum_{k=0}^{\infty} C(n+k-1, k)x^k$$

$$\frac{1}{(1+x)^n} = \sum_{k=0}^{\infty} C(n+k-1, k)(-1)^k x^k$$

$$\frac{1}{(1-ax)^n} = \sum_{k=0}^{\infty} C(n+k-1, k)a^k x^k$$



# Useful Generating Functions


$$\frac{1}{(1-x)^n} = \sum_{k=0}^{\infty} C(n+k-1, k)x^k$$

$$\frac{1}{(1+x)^n} = \sum_{k=0}^{\infty} C(n+k-1, k)(-1)^k x^k$$

$$\frac{1}{(1-ax)^n} = \sum_{k=0}^{\infty} C(n+k-1, k)a^k x^k$$

$$e^x = \sum_{k=0}^{\infty} \frac{x^k}{k!} = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$$

$$\ln(1+x) = \sum_{k=0}^{\infty} \frac{(-1)^{k+1} x^k}{k} = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots$$



# Counting and Generating Functions

- **Problem 1** How many solutions are there to the equation

$$x_1 + x_2 + x_3 = 17,$$

where  $x_1, x_2, x_3$  are **nonnegative** integers?



# Counting and Generating Functions

- **Problem 1** How many solutions are there to the equation

$$x_1 + x_2 + x_3 = 17,$$

where  $x_1, x_2, x_3$  are **nonnegative** integers?

This is **equivalent** to the problem of  $r$ -combinations from a set with  $n$  elements when **repetition** is allowed.



# Counting and Generating Functions

- **Problem 1** How many solutions are there to the equation

$$x_1 + x_2 + x_3 = 17,$$

where  $x_1, x_2, x_3$  are **nonnegative** integers?

This is **equivalent** to the problem of  $r$ -combinations from a set with  $n$  elements when **repetition** is allowed.

$$C(n + r - 1, n - 1) = C(19, 2)$$



# Counting and Generating Functions

- **Problem 2** Find the number of solutions of

$$x_1 + x_2 + x_3 = 17,$$

where  $x_1, x_2, x_3$  are **nonnegative** integers with  $2 \leq x_1 \leq 5$ ,  
 $3 \leq x_2 \leq 6$ ,  $4 \leq x_3 \leq 7$ .





# Counting and Generating Functions


- **Problem 2** Find the number of solutions of

$$x_1 + x_2 + x_3 = 17,$$

where  $x_1, x_2, x_3$  are **nonnegative** integers with  $2 \leq x_1 \leq 5$ ,  
 $3 \leq x_2 \leq 6$ ,  $4 \leq x_3 \leq 7$ .

Using *generating functions*, the number is the **coefficient** of  $x^{17}$  in the expansion of

$$(x^2 + x^3 + x^4 + x^5)(x^3 + x^4 + x^5 + x^6)(x^4 + x^5 + x^6 + x^7)$$



# Counting and Generating Functions

- **Problem 3** In how many ways can eight identical cookies be distributed among three distinct children if each child receives at least two cookies and no more than four cookies?



# Counting and Generating Functions

- **Problem 3** In how many ways can eight identical cookies be distributed among three distinct children if each child receives at least two cookies and no more than four cookies?

The coefficient of  $x^8$  in the expansion

$$(x^2 + x^3 + x^4)^3$$



# Counting and Generating Functions

- **Problem 4** Use generating functions to find the number of  $k$ -combinations of a set with  $n$  elements,  $C(n, k)$ .




# Counting and Generating Functions

- **Problem 4** Use **generating functions** to find the number of  **$k$ -combinations of a set with  $n$  elements**,  $C(n, k)$ .

Each of the  $n$  elements in the set contributes the term  $(1 + x)$  to the generating function  $f(x) = \sum_{k=0}^n a^k x^k$ .  
Hence,  $f(x) = (1 + x)^n$ .

Then by the **binomial theorem**, we have  $a_k = \binom{n}{k}$ .



# Cartesian Product

- Let  $A = \{a_1, a_2, \dots, a_m\}$  and  $B = \{b_1, b_2, \dots, b_n\}$ , the *Cartesian product*  $A \times B$  is the set of pairs

$$\{(a_1, b_1), (a_2, b_2), \dots, (a_1, b_n), \dots, (a_m, b_n)\}$$



# Cartesian Product

- Let  $A = \{a_1, a_2, \dots, a_m\}$  and  $B = \{b_1, b_2, \dots, b_n\}$ , the *Cartesian product*  $A \times B$  is the set of pairs

$$\{(a_1, b_1), (a_2, b_2), \dots, (a_1, b_n), \dots, (a_m, b_n)\}$$

**Cartesian product** defines a set of all **ordered** arrangements of elements in the two sets.



# Binary Relation

- **Definition:** Let  $A$  and  $B$  be two sets. A *binary relation from  $A$  to  $B$*  is a **subset** of a **Cartesian product**  $A \times B$ .





# Binary Relation

- **Definition:** Let  $A$  and  $B$  be two sets. A *binary relation from  $A$  to  $B$*  is a **subset** of a **Cartesian product**  $A \times B$ .

Let  $R \subseteq A \times B$  denote  $R$  is a set of ordered pairs of the form  $(a, b)$  where  $a \in A$  and  $b \in B$ .



# Binary Relation

- **Definition:** Let  $A$  and  $B$  be two sets. A *binary relation from  $A$  to  $B$*  is a **subset** of a **Cartesian product**  $A \times B$ .

Let  $R \subseteq A \times B$  denote  $R$  is a set of ordered pairs of the form  $(a, b)$  where  $a \in A$  and  $b \in B$ .

We use the notation  $a R b$  to denote  $(a, b) \in R$ , and  $a \not R b$  to denote  $(a, b) \notin R$ .



# Binary Relation

- **Definition:** Let  $A$  and  $B$  be two sets. A *binary relation from  $A$  to  $B$*  is a **subset** of a **Cartesian product**  $A \times B$ .

Let  $R \subseteq A \times B$  denote  $R$  is a set of ordered pairs of the form  $(a, b)$  where  $a \in A$  and  $b \in B$ .

We use the notation  $a R b$  to denote  $(a, b) \in R$ , and  $a \not R b$  to denote  $(a, b) \notin R$ .

**Example:** Let  $A = \{a, b, c\}$  and  $B = \{1, 2, 3\}$

- ◇ Is  $R = \{(a, 1), (b, 2), (c, 2)\}$  a relation from  $A$  to  $B$ ?
- ◇ Is  $Q = \{(1, a), (2, b)\}$  a relation from  $A$  to  $B$ ?
- ◇ Is  $P = \{(a, a), (b, c), (b, a)\}$  a relation from  $A$  to  $A$ ?



# Representing Binary Relations

- We can **graphically** represent a binary relation  $R$  as:  
if  $a R b$ , then we draw an **arrow** from  $a$  to  $b$ :  $a \rightarrow b$



# Representing Binary Relations

- We can **graphically** represent a binary relation  $R$  as:  
if  $a R b$ , then we draw an **arrow** from  $a$  to  $b$ :  $a \rightarrow b$


**Example:** Let  $A = \{0, 1, 2\}$  and  $B = \{u, v\}$ , and  
 $R = \{(0, u), (0, v), (1, v), (2, u)\}$ . ( $R \subseteq A \times B$ )



# Representing Binary Relations

- We can **graphically** represent a binary relation  $R$  as:  
if  $a R b$ , then we draw an **arrow** from  $a$  to  $b$ :  $a \rightarrow b$

**Example:** Let  $A = \{0, 1, 2\}$  and  $B = \{u, v\}$ , and  $R = \{(0, u), (0, v), (1, v), (2, u)\}$ . ( $R \subseteq A \times B$ )



# Representing Binary Relations

- We can also represent a binary relation  $R$  by a table showing the ordered pairs of  $R$ .



# Representing Binary Relations

- We can also represent a binary relation  $R$  by a table showing the ordered pairs of  $R$ .

**Example:** Let  $A = \{0, 1, 2\}$  and  $B = \{u, v\}$ , and  $R = \{(0, u), (0, v), (1, u), (2, v)\}$ . ( $R \subseteq A \times B$ )






# Representing Binary Relations

- We can also represent a binary relation  $R$  by a table showing the ordered pairs of  $R$ .

**Example:** Let  $A = \{0, 1, 2\}$  and  $B = \{u, v\}$ , and  $R = \{(0, u), (0, v), (1, u), (2, v)\}$ . ( $R \subseteq A \times B$ )

$R$	$u$	$v$
0	×	×
1	×	
2		×




# Relations and Functions

- Relations represent **one to many relationships** between elements in  $A$  and  $B$ .




# Relations and Functions

- Relations represent **one to many relationships** between elements in  $A$  and  $B$ .



# Relations and Functions

- Relations represent **one to many relationships** between elements in  $A$  and  $B$ .



What is the **difference** between a **relation** and a **function** from  $A$  to  $B$ ?



# Relation on the Set

- **Definition:** A relation on the set  $A$  is a relation from  $A$  to itself.



# Relation on the Set

- **Definition:** A relation on the set  $A$  is a relation from  $A$  to itself.

**Example:** Let  $A = \{1, 2, 3, 4\}$  and  $R_{div} = \{(a, b) : a|b\}$ . What does  $R_{div}$  consist of?



# Relation on the Set

- **Definition:** A relation on the set  $A$  is a relation from  $A$  to itself.

**Example:** Let  $A = \{1, 2, 3, 4\}$  and  $R_{div} = \{(a, b) : a|b\}$ . What does  $R_{div}$  consist of?

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$





# Relation on the Set

- **Definition:** A relation on the set  $A$  is a relation from  $A$  to itself.

**Example:** Let  $A = \{1, 2, 3, 4\}$  and  $R_{div} = \{(a, b) : a|b\}$ . What does  $R_{div}$  consist of?

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$






# Relation on the Set

- **Definition:** A relation on the set  $A$  is a relation from  $A$  to itself.

**Example:** Let  $A = \{1, 2, 3, 4\}$  and  $R_{div} = \{(a, b) : a|b\}$ . What does  $R_{div}$  consist of?

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$



$R$	1	2	3	4
1	×	×	×	×
2		×		×
3			×	
4				×



# Number of Binary Relations

- **Theorem** The number of binary relations on a set  $A$ , where  $|A| = n$  is  $2^{n^2}$ .



# Number of Binary Relations

- **Theorem** The number of binary relations on a set  $A$ , where  $|A| = n$  is  $2^{n^2}$ .

## Proof

If  $|A| = n$ , then the cardinality of the Cartesian product  $|A \times A| = n^2$ .



# Number of Binary Relations

- **Theorem** The number of binary relations on a set  $A$ , where  $|A| = n$  is  $2^{n^2}$ .

## Proof

If  $|A| = n$ , then the cardinality of the Cartesian product  $|A \times A| = n^2$ .

$R$  is a binary relation on  $A$  if  $R \subseteq A \times A$  ( $R$  is subset)



# Number of Binary Relations


- **Theorem** The number of binary relations on a set  $A$ , where  $|A| = n$  is  $2^{n^2}$ .

## Proof

If  $|A| = n$ , then the cardinality of the Cartesian product  $|A \times A| = n^2$ .

$R$  is a binary relation on  $A$  if  $R \subseteq A \times A$  ( $R$  is subset)

The number of subsets of a set with  $k$  elements is  $2^k$



# Next Lecture

- relation ...

