



DISCRETE MATHEMATICS FOR COMPUTER SCIENCE

Dr. QI WANG

Department of Computer Science and Engineering

Office: Room903, Nanshan iPark A7 Building

Email: wangqi@sustc.edu.cn

Assignment #1

Please collect your
assignments!



Recall

- **Theorem (Fermat's little theorem)** : Let p be a prime, and let x be an integer such that $x \not\equiv 0 \pmod{p}$. Then

$$x^{p-1} \equiv 1 \pmod{p}.$$


- **Theorem (Euler's theorem)** : Let n be a positive integer, and let x be an integer such that $\gcd(x, n) = 1$. Then

$$x^{\phi(n)} \equiv 1 \pmod{n}.$$


- **Theorem** Let $a \in \mathbb{Z}$ and $n \in \mathbb{N}$ with $\gcd(a, n) = 1$. Then $\text{ord}_n(a)$ exists and divides $\phi(n)$.



Symmetric Cryptography



Asymmetric Cryptography



Public-Key Cryptosystems

R. Rivest, A. Shamir, L. Adleman, "A Method for Obtaining Digital Signatures and Public-Key Cryptosystems",
Communications of the ACM, vol. 21-2, pages 120-126, 1978.



RSA Public-Key Cryptosystem

Pick two **large** primes, p and q . Let $n = pq$, then $\phi(n) = (p - 1)(q - 1)$. Encryption and decryption keys e and d are selected such that

- $\gcd(e, \phi(n)) = 1$
- $ed \equiv 1 \pmod{\phi(n)}$



RSA Public-Key Cryptosystem

Pick two **large** primes, p and q . Let $n = pq$, then $\phi(n) = (p - 1)(q - 1)$. Encryption and decryption keys e and d are selected such that

- $\gcd(e, \phi(n)) = 1$
- $ed \equiv 1 \pmod{\phi(n)}$

- $C = M^e \bmod n$ (RSA **encryption**)
- $M = C^d \bmod n$ (RSA **decryption**)



RSA Public-Key Cryptosystem

$$C = M^e \bmod n \text{ (RSA encryption)}$$

$$M = C^d \bmod n \text{ (RSA decryption)}$$

- **Theorem 12** (Correctness) : Let p and q be two odd primes, and define $n = pq$. Let e be relatively prime to $\phi(n)$ and let d be the multiplicative inverse of e modulo $\phi(n)$. For each integer x such that $0 \leq x < n$,

$$x^{ed} \equiv x \pmod{n}.$$



RSA Public-Key Cryptosystem

$$C = M^e \bmod n \text{ (RSA encryption)}$$

$$M = C^d \bmod n \text{ (RSA decryption)}$$

- **Theorem 12** (Correctness) : Let p and q be two odd primes, and define $n = pq$. Let e be relatively prime to $\phi(n)$ and let d be the multiplicative inverse of e modulo $\phi(n)$. For each integer x such that $0 \leq x < n$,

$$x^{ed} \equiv x \pmod{n}.$$

Q : How to prove this?



RSA Public-Key Cryptosystem: Example

Parameters:	p	q	n	$\phi(n)$	e	d
	5	11	55	40	7	23



RSA Public-Key Cryptosystem: Example

Parameters:	p	q	n	$\phi(n)$	e	d
	5	11	55	40	7	23

Public key: (7, 55)

Private key: 23



RSA Public-Key Cryptosystem: Example

Parameters:

p	q	n	$\phi(n)$	e	d
5	11	55	40	7	23

Public key: $(7, 55)$

Private key: 23

Encryption: $M = 28, C = M^7 \bmod 55 = 52$

Decryption: $M = C^{23} \bmod 55 = 28$



RSA Public-Key Cryptosystem: Parameters

Parameters: p q n $\phi(n)$ e d

Public key: (e, n)

Private key: d

$p, q, \phi(n)$ must be kept **secret**!



RSA Public-Key Cryptosystem: Parameters

Parameters: p q n $\phi(n)$ e d

Public key: (e, n)

Private key: d

$p, q, \phi(n)$ must be kept **secret**!

Q : Why?



The Security of the RSA

Brute-force attack:

Trying all possible private keys.

$$\# = \phi(\phi(n)) = \phi((p-1)(q-1))$$



The Security of the RSA

Brute-force attack:

Trying all possible private keys.

$$\# = \phi(\phi(n)) = \phi((p-1)(q-1))$$

Attack: Factor n into pq .

Attack: Determine $\phi(n)$ directly.

Attack: Determine d directly.



The Security of the RSA

Brute-force attack:

Trying all possible private keys.

$$\# = \phi(\phi(n)) = \phi((p-1)(q-1))$$

Attack: Factor n into pq .

Attack: Determine $\phi(n)$ directly.

Attack: Determine d directly.

Comment: It is believed that determining $\phi(n)$ is **equivalent** to factoring n . Meanwhile, determining d given e and n , appears to be at least as time-consuming as **the integer factoring problem**.



The Security of the RSA

In practice, RSA keys are typically 1024 to 2048 bits long.



The Security of the RSA

In practice, RSA keys are typically 1024 to 2048 bits long.

Remark: There are some suggestions for choosing p and q .

A. Salomaa, *Public-Key Cryptography*, 2nd Edition, Springer, 1996, pp. 134-136.



The Security of the RSA

In practice, RSA keys are typically 1024 to 2048 bits long.

Remark: There are some suggestions for choosing p and q .

A. Salomaa, *Public-Key Cryptography*, 2nd Edition, Springer, 1996, pp. 134-136.

Q : Consider the RSA system, where $n = pq$ is the modulus. Let (e, d) be a key pair for the RSA. Define

$$\lambda(n) = \text{lcm}(p - 1, q - 1)$$

and compute $d' = e^{-1} \bmod \lambda(n)$. Will decryption using d' instead of d still work?



Using RSA for Digital Signature

$$S = M^d \bmod n \text{ (RSA signature)}$$

$$M = S^e \bmod n \text{ (RSA verification)}$$

Why?



The Discrete Logarithm

- **The discrete logarithm** of an integer y to the base b is an integer x , such that

$$b^x \equiv y \pmod{n}.$$



The Discrete Logarithm

- **The discrete logarithm** of an integer y to the base b is an integer x , such that

$$b^x \equiv y \pmod{n}.$$

Discrete Logarithm Problem:

Given n , b and y , find x .



The Discrete Logarithm

- **The discrete logarithm** of an integer y to the base b is an integer x , such that

$$b^x \equiv y \pmod{n}.$$

Discrete Logarithm Problem:

Given n , b and y , find x .

This is very hard!!!



El Gamal Encryption

- **Setup** Let p be a prime, and g be a generator of \mathbb{Z}_p . The **private key** x is an integer with $1 < x < p - 2$. Let $y = g^x \bmod p$. The **public key** for *El Gamal encryption* is (p, g, y) .



El Gamal Encryption

- **Setup** Let p be a prime, and g be a generator of \mathbb{Z}_p . The **private key** x is an integer with $1 < x < p - 2$. Let $y = g^x \bmod p$. The **public key** for *El Gamal encryption* is (p, g, y) .

El Gamal Encryption: Pick a random integer k from \mathbb{Z}_{p-1} ,

$$a = g^k \bmod p$$

$$b = My^k \bmod p$$

The ciphertext C consists of the pair (a, b) .

El Gamal Decryption:

$$M = b(a^x)^{-1} \bmod p$$



Using El Gamal for Digital Signature

$$\begin{aligned}a &= g^k \bmod p \\b &= k^{-1}(M - xa) \bmod (p - 1)\end{aligned}$$

(El Gamal **signature**)

$$y^a a^b \equiv g^M \pmod{p}$$

(El Gamal **verification**)



Using El Gamal for Digital Signature

$$\begin{aligned}a &= g^k \bmod p \\ b &= k^{-1}(M - xa) \bmod (p - 1) \\ &\text{(El Gamal **signature**)}\end{aligned}$$

$$\begin{aligned}y^a a^b &\equiv g^M \pmod{p} \\ &\text{(El Gamal **verification**)}\end{aligned}$$

Q : How to verify it?



An Example

Choose $p = 2579$, $g = 2$, and $x = 765$. Hence
 $y = 2^{765} \bmod 2579 = 949$.



An Example

Choose $p = 2579$, $g = 2$, and $x = 765$. Hence $y = 2^{765} \bmod 2579 = 949$.

- ▶ (Public key) $k_e = (p, g, y) = (2579, 2, 949)$
- ▶ (Private key) $k_d = x = 765$



An Example

Choose $p = 2579$, $g = 2$, and $x = 765$. Hence $y = 2^{765} \bmod 2579 = 949$.

► (Public key) $k_e = (p, g, y) = (2579, 2, 949)$

► (Private key) $k_d = x = 765$

Encryption: Let $M = 1299$ and choose a random $k = 853$,

$$\begin{aligned}(a, b) &= (g^k \bmod p, My^k \bmod p) \\ &= (2^{853} \bmod 2579, 1299 \cdot 949^{853} \bmod 2579) \\ &= (435, 2396).\end{aligned}$$


Decryption:

$$M = b(a^x)^{-1} \bmod p = 2396 \times (435^{765})^{-1} \bmod 2579 = 1299.$$



Security of the El Gamal Cryptosystem

Question 1: Is it feasible to derive x from (p, g, y) ?



Security of the El Gamal Cryptosystem

Question 1: Is it feasible to derive x from (p, g, y) ?

It is equivalent to solving the DLP. It is **believed** that there is **NO** polynomial-time algorithm. p should be large enough, typically 160 bits.



Security of the El Gamal Cryptosystem

Question 1: Is it feasible to derive x from (p, g, y) ?

It is equivalent to solving the DLP. It is **believed** that there is **NO** polynomial-time algorithm. p should be large enough, typically 160 bits.

Question 2: Given a ciphertext (a, b) , is it feasible to derive the plaintext M ?



Security of the El Gamal Cryptosystem

Question 1: Is it feasible to derive x from (p, g, y) ?

It is equivalent to solving the DLP. It is **believed** that there is **NO** polynomial-time algorithm. p should be large enough, typically 160 bits.

Question 2: Given a ciphertext (a, b) , is it feasible to derive the plaintext M ?

Attack 1: Use $M = by^{-k}$. However, k is **randomly** picked.

Attack 2: Use $M = b(a^x)^{-1} \bmod p$, but x is **secret**.



Diffie-Hellman Key Exchange Protocol

User A

User B

Generate random

$$X_A < p$$


calculate

$$Y_A = \alpha^{X_A} \bmod p$$

Calculate

$$k = (Y_B)^{X_A} \bmod p$$

Y_A



Y_B

Generate random


$$X_B < p$$

Calculate

$$Y_B = \alpha^{X_B} \bmod p$$

Calculate

$$k = (Y_A)^{X_B} \bmod p$$




Announcements

■ Homework assignment 3



- ◇ P245 Ex. 37, 38, 39, P255 Ex. 2, 26, P272 Ex. 11*, 12, P273 Ex. 42, P274 Ex. 50, 55, P284 Ex. 7*, P285 Ex. 22, P286 Ex. 39, P305 Ex. 23, 28, 30
- ◇ Due on *Nov. 7th, 2017 at the beginning of class*
- ◇ Please try your best to solve problems marked with *
- ◇ Please write your homework **neatly**, as a courtesy to graders.



Mathematical Induction



Mathematical Induction



Mathematical Induction

- We start by reviewing proof by smallest counterexample to try and understand what it is really doing.



Mathematical Induction

- We start by reviewing proof by smallest counterexample to try and understand what it is really doing.
- This leads us to transform the *indirect proof* of proof by counterexample to *direct proof*. This direct proof technique will be **induction**.



Mathematical Induction

- We start by *reviewing proof by smallest counterexample* to try and understand what it is really doing.
- This leads us to transform the *indirect proof* of *proof by counterexample* to *direct proof*. This direct proof technique will be **induction**.
- We conclude by distinguishing between the *weak principle* of mathematical induction and the *strong principle* of mathematical induction.



Mathematical Induction

- We start by *reviewing proof by smallest counterexample* to try and understand what it is really doing.
- This leads us to transform the *indirect proof* of *proof by counterexample* to *direct proof*. This direct proof technique will be **induction**.
- We conclude by distinguishing between the *weak principle* of mathematical induction and the *strong principle* of mathematical induction.

The *strong principle* can actually be derived from the *weak principle*.



Proof by Smallest Counterexample

- The statement $P(n)$ is true for all $n = 0, 1, 2, \dots$



Proof by Smallest Counterexample

- The statement $P(n)$ is true for all $n = 0, 1, 2, \dots$

We prove this by

- (i) Assume that a counterexample exists, i.e., There is some $n > 0$ for which $P(n)$ is false




Proof by Smallest Counterexample

- The statement $P(n)$ is true for all $n = 0, 1, 2, \dots$

We prove this by

- (i) Assume that a counterexample exists, i.e., There is some $n > 0$ for which $P(n)$ is false
- (ii) Let $m > 0$ be the **smallest** value for which $P(n)$ is false




Proof by Smallest Counterexample

- The statement $P(n)$ is true for all $n = 0, 1, 2, \dots$

We prove this by

- (i) Assume that a counterexample exists, i.e., There is some $n > 0$ for which $P(n)$ is false
- (ii) Let $m > 0$ be the **smallest** value for which $P(n)$ is false
- (iii) Then use the fact that $P(m')$ is true for all $0 \leq m' < m$ to show that $P(m)$ is true, **contradicting** the choice of m .




Proof by Smallest Counterexample

- The statement $P(n)$ is true for all $n = 0, 1, 2, \dots$

We prove this by

- (i) Assume that a counterexample exists, i.e., There is some $n > 0$ for which $P(n)$ is false
- (ii) Let $m > 0$ be the **smallest** value for which $P(n)$ is false
- (iii) Then use the fact that $P(m')$ is true for all $0 \leq m' < m$ to show that $P(m)$ is true, **contradicting** the choice of m .



Example 1

- Use **proof by smallest counterexample** to show that, $\forall n \in \mathbb{N}$,

$$(*) \quad 0 + 1 + 2 + 3 + \cdots + n = \frac{n(n+1)}{2}$$



Example 1

- Use **proof by smallest counterexample** to show that, $\forall n \in \mathbb{N}$,

$$(*) \quad 0 + 1 + 2 + 3 + \cdots + n = \frac{n(n+1)}{2}$$

- ◇ Suppose that **(*)** is not always true



Example 1

- Use **proof by smallest counterexample** to show that, $\forall n \in \mathbb{N}$,

$$(*) \quad 0 + 1 + 2 + 3 + \cdots + n = \frac{n(n+1)}{2}$$

- ◇ Suppose that **(*)** is not always true
- ◇ Then there must be a **smallest $n \in \mathbb{N}$ s.t. (*) does not hold for n**



Example 1

- Use **proof by smallest counterexample** to show that, $\forall n \in \mathbb{N}$,

$$(*) \quad 0 + 1 + 2 + 3 + \cdots + n = \frac{n(n+1)}{2}$$

- ◇ Suppose that **(*)** is not always true
- ◇ Then there must be a **smallest** $n \in \mathbb{N}$ s.t. **(*)** does not hold for n
- ◇ For any nonnegative integer $i < n$,

$$1 + 2 + \cdots + i = \frac{i(i+1)}{2}$$



Example 1

- Use **proof by smallest counterexample** to show that, $\forall n \in \mathbb{N}$,

$$(*) \quad 0 + 1 + 2 + 3 + \cdots + n = \frac{n(n+1)}{2}$$

- ◇ Suppose that **(*)** is not always true
- ◇ Then there must be a **smallest** $n \in \mathbb{N}$ s.t. **(*)** does not hold for n
- ◇ For any nonnegative integer $i < n$,

$$1 + 2 + \cdots + i = \frac{i(i+1)}{2}$$

- ◇ Since $0 = 0 \cdot 1/2$, **(*)** holds for $n = 0$



Example 1

- Use **proof by smallest counterexample** to show that, $\forall n \in \mathbb{N}$,

$$(*) \quad 0 + 1 + 2 + 3 + \cdots + n = \frac{n(n+1)}{2}$$

- ◇ Suppose that **(*)** is not always true
- ◇ Then there must be a **smallest** $n \in \mathbb{N}$ s.t. **(*)** does not hold for n
- ◇ For any nonnegative integer $i < n$,
$$1 + 2 + \cdots + i = \frac{i(i+1)}{2}$$
- ◇ Since $0 = 0 \cdot 1/2$, **(*)** holds for $n = 0$
- ◇ The smallest counterexample n is larger than 0



Example 1

- We now have
 - (i) smallest counterexample n is greater than 0, and
 - (ii) $(*)$ holds for $n - 1$

Example 1

■ We now have

- (i) smallest counterexample n is greater than 0, and
- (ii) $(*)$ holds for $n - 1$

◇ Substituting $n - 1$ for i gives

$$1 + 2 + \cdots + n - 1 = \frac{(n - 1)n}{2}$$



Example 1

■ We now have

- (i) smallest counterexample n is greater than 0, and
- (ii) $(*)$ holds for $n - 1$

◇ Substituting $n - 1$ for i gives

$$1 + 2 + \cdots + n - 1 = \frac{(n - 1)n}{2}$$

◇ Adding n to both sides gives

$$1 + 2 + \cdots + n - 1 + n = \frac{(n - 1)n}{2} + n = \frac{n(n + 1)}{2}$$



Example 1

■ We now have

- (i) smallest counterexample n is greater than 0, and
- (ii) $(*)$ holds for $n - 1$

◇ Substituting $n - 1$ for i gives

$$1 + 2 + \cdots + n - 1 = \frac{(n - 1)n}{2}$$

◇ Adding n to both sides gives

$$1 + 2 + \cdots + n - 1 + n = \frac{(n - 1)n}{2} + n = \frac{n(n + 1)}{2}$$

◇ Thus, n is not a counterexample. Contradiction!



Example 1

■ We now have

- (i) smallest counterexample n is greater than 0, and
- (ii) $(*)$ holds for $n - 1$

◇ Substituting $n - 1$ for i gives

$$1 + 2 + \cdots + n - 1 = \frac{(n - 1)n}{2}$$

◇ Adding n to both sides gives

$$1 + 2 + \cdots + n - 1 + n = \frac{(n - 1)n}{2} + n = \frac{n(n + 1)}{2}$$

◇ Thus, n is not a counterexample. Contradiction!

◇ Therefore, $(*)$ holds for all positive integers n .



Example 1

- What implication did we have to prove?



Example 1

- What implication did we have to prove?

The **key step** was proving that

$$P(n - 1) \rightarrow P(n)$$

where $P(n)$ is the statement

$$1 + 2 + \cdots + n = \frac{n(n + 1)}{2}$$



Example 2

- Use **proof by smallest counterexample** to show that, $\forall n \in \mathbb{N}$,
$$2^{n+1} \geq n^2 + 2.$$



Example 2

- Use **proof by smallest counterexample** to show that, $\forall n \in \mathbb{N}$,
$$2^{n+1} \geq n^2 + 2.$$

Let $P(n) = 2^{n+1} \geq n^2 + 2$. We start by assuming that the statement

$$\forall n \in \mathbb{N} \ P(n)$$

is **false**.



Example 2

- Use **proof by smallest counterexample** to show that, $\forall n \in \mathbb{N}$,
$$2^{n+1} \geq n^2 + 2.$$

Let $P(n) = 2^{n+1} \geq n^2 + 2$. We start by assuming that the statement

$$\forall n \in \mathbb{N} \ P(n)$$

is **false**.

When a **for all** quantifier is false, **there must be some n for which it is false**. Let n be the **smallest nonnegative integer** for which $2^{n+1} \not\geq n^2 + 2$.



Example 2

- Let n be the smallest nonnegative integer for which $2^{n+1} \not\geq n^2 + 2$.

This means that, for all $i \in \mathbb{N}$ with $i < n$,

$$2^{i+1} \geq i^2 + 2$$



Example 2

- Let n be the **smallest nonnegative integer** for which $2^{n+1} \not\geq n^2 + 2$.

This means that, **for all $i \in \mathbb{N}$ with $i < n$,**

$$2^{i+1} \geq i^2 + 2$$

Since $2^{0+1} \geq 0^2 + 2$, we know that $n > 0$. Thus, **$n - 1$ is a nonnegative integer less than n .**



Example 2

- Let n be the smallest nonnegative integer for which $2^{n+1} \not\geq n^2 + 2$.

This means that, for all $i \in \mathbb{N}$ with $i < n$,

$$2^{i+1} \geq i^2 + 2$$

Since $2^{0+1} \geq 0^2 + 2$, we know that $n > 0$. Thus, $n - 1$ is a nonnegative integer less than n .

Then setting $i = n - 1$ gives

$$2^{(n-1)+1} \geq (n-1)^2 + 2.$$

or

$$(*) \quad 2^n \geq n^2 - 2n + 1 + 2 = n^2 - 2n + 3$$



Example 2

- Let n be the **smallest nonnegative integer** for which $2^{n+1} \not\geq n^2 + 2$.

We are now given $2^n \geq n^2 - 2n + 3$. (*)




Example 2

- Let n be the **smallest nonnegative integer** for which $2^{n+1} \not\geq n^2 + 2$.

We are now given $2^n \geq n^2 - 2n + 3$. (*)

Multiply both sides by 2, giving

$$2^{n+1} = 2 \cdot 2^n \geq 2 \cdot (n^2 - 2n + 3) = 2n^2 - 4n + 6.$$



Example 2

- Let n be the **smallest nonnegative integer** for which $2^{n+1} \not\geq n^2 + 2$.

We are now given $2^n \geq n^2 - 2n + 3$. $(*)$

Multiply both sides by 2, giving

$$2^{n+1} = 2 \cdot 2^n \geq 2 \cdot (n^2 - 2n + 3) = 2n^2 - 4n + 6.$$

To get a **contradiction**, we want to convert the right side into $n^2 + 2$ plus an additional nonnegative term.



Example 2

- Let n be the **smallest nonnegative integer** for which $2^{n+1} \not\geq n^2 + 2$.

We are now given $2^n \geq n^2 - 2n + 3$. $(*)$

Multiply both sides by 2, giving

$$2^{n+1} = 2 \cdot 2^n \geq 2 \cdot (n^2 - 2n + 3) = 2n^2 - 4n + 6.$$

To get a **contradiction**, we want to convert the right side into $n^2 + 2$ plus an additional nonnegative term.

Thus, we write

$$\begin{aligned} 2^{n+1} &\geq 2n^2 - 4n + 6 \\ &= (n^2 + 2) + (n^2 - 4n + 4) \\ &= n^2 + 2 + (n - 2)^2 \\ &\geq n^2 + 2. \end{aligned}$$



Example 2

- Let n be the **smallest nonnegative integer** for which $2^{n+1} \not\geq n^2 + 2$.

We are now given $2^n \geq n^2 - 2n + 3$. $(*)$

Multiply both sides by 2, giving

$$2^{n+1} = 2 \cdot 2^n \geq 2 \cdot (n^2 - 2n + 3) = 2n^2 - 4n + 6.$$

To get a **contradiction**, we want to convert the right side into $n^2 + 2$ plus an additional nonnegative term.

Thus, we write

$$\begin{aligned} 2^{n+1} &\geq 2n^2 - 4n + 6 \\ &= (n^2 + 2) + (n^2 - 4n + 4) \\ &= n^2 + 2 + (n - 2)^2 \\ &\geq n^2 + 2. \end{aligned}$$

contradiction!



Example 2

- Let $P(n) = 2^{n+1} \geq n^2 + 2$

We just showed that

(a) $P(0)$ is true

(b) if $n > 0$, then $P(n-1) \rightarrow P(n)$



Example 2

- Let $P(n) = 2^{n+1} \geq n^2 + 2$

We just showed that

- (a) $P(0)$ is true
- (b) if $n > 0$, then $P(n-1) \rightarrow P(n)$
- ◇ Suppose there is some n for which $P(n)$ is false (*)



Example 2

- Let $P(n) - 2^{n+1} \geq n^2 + 2$

We just showed that

- (a) $P(0)$ is true
- (b) if $n > 0$, then $P(n-1) \rightarrow P(n)$
- ◇ Suppose there is some n for which $P(n)$ is false (*)
- ◇ Let n be the smallest counterexample



Example 2

- Let $P(n) - 2^{n+1} \geq n^2 + 2$

We just showed that

- (a) $P(0)$ is true
- (b) if $n > 0$, then $P(n-1) \rightarrow P(n)$
- ◇ Suppose there is some n for which $P(n)$ is false (*)
- ◇ Let n be the smallest counterexample
- ◇ Then, from (a) $n > 0$, so $P(n-1)$ is true



Example 2

- Let $P(n) = 2^{n+1} \geq n^2 + 2$

We just showed that

- (a) $P(0)$ is true
- (b) if $n > 0$, then $P(n-1) \rightarrow P(n)$
- ◇ Suppose there is some n for which $P(n)$ is false (*)
- ◇ Let n be the smallest counterexample
- ◇ Then, from (a) $n > 0$, so $P(n-1)$ is true
- ◇ Therefore, from (b), using direct inference, $P(n)$ is true



Example 2

- Let $P(n) = 2^{n+1} \geq n^2 + 2$

We just showed that

- (a) $P(0)$ is true
- (b) if $n > 0$, then $P(n-1) \rightarrow P(n)$
- ◇ Suppose there is some n for which $P(n)$ is false (*)
- ◇ Let n be the smallest counterexample
- ◇ Then, from (a) $n > 0$, so $P(n-1)$ is true
- ◇ Therefore, from (b), using direct inference, $P(n)$ is true
- ◇ This contradicts (*).



Example 2

- Let $P(n) - 2^{n+1} \geq n^2 + 2$

We just showed that

- (a) $P(0)$ is true
- (b) if $n > 0$, then $P(n-1) \rightarrow P(n)$
- ◇ Suppose there is some n for which $P(n)$ is false (*)
- ◇ Let n be the smallest counterexample
- ◇ Then, from (a) $n > 0$, so $P(n-1)$ is true
- ◇ Therefore, from (b), using direct inference, $P(n)$ is true
- ◇ This contradicts (*).
- ◇ Thus, $P(n)$ is true for all $n \in \mathbb{N}$.



Example 2

- What did we really do?

Let $P(n) - 2^{n+1} \geq n^2 + 2$

We just showed that

(a) $P(0)$ is true

(b) if $n > 0$, then $P(n-1) \rightarrow P(n)$



Example 2

■ What did we really do?

Let $P(n) = 2^{n+1} \geq n^2 + 2$

We just showed that

(a) $P(0)$ is true

(b) if $n > 0$, then $P(n-1) \rightarrow P(n)$

We then used proof by smallest counterexample to derive that $P(n)$ is true for all $n \in \mathbb{N}$.



Example 2

■ What did we really do?

Let $P(n) = 2^{n+1} \geq n^2 + 2$

We just showed that

(a) $P(0)$ is true

(b) if $n > 0$, then $P(n-1) \rightarrow P(n)$

We then used **proof by smallest counterexample** to derive that $P(n)$ is true for all $n \in \mathbb{N}$.

This is an *indirect proof*. Is it possible to prove this fact *directly*?



Example 2

■ What did we really do?

Let $P(n) = 2^{n+1} \geq n^2 + 2$

We just showed that

(a) $P(0)$ is true

(b) if $n > 0$, then $P(n-1) \rightarrow P(n)$

We then used **proof by smallest counterexample** to derive that $P(n)$ is true for all $n \in \mathbb{N}$.

This is an *indirect proof*. Is it possible to prove this fact *directly*?

Since $P(n-1) \rightarrow P(n)$, we see that

$P(0)$ implies $P(1)$, $P(1)$ implies $P(2)$, ...



The Principle of Mathematical Induction

- The *well-ordering* principle permits us to assume that every set of nonnegative integers has a smallest element, allowing us to use the smallest counterexample.



The Principle of Mathematical Induction

- The *well-ordering* principle permits us to assume that every set of nonnegative integers has a smallest element, *allowing us to use the smallest counterexample*.

This is actually **equivalent** to the *principle of mathematical induction*.




The Principle of Mathematical Induction

- The *well-ordering* principle permits us to assume that every set of nonnegative integers has a smallest element, allowing us to use the smallest counterexample.

This is actually **equivalent** to the *principle of mathematical induction*.

Principle. (*the Weak Principle of Mathematical Induction*)

- (a) If the statement $P(b)$ is true
- (b) the statement $P(n-1) \rightarrow P(n)$ is true for all $n > b$,
then $P(n)$ is true for all integers $n \geq b$



The Principle of Mathematical Induction

- The *well-ordering* principle permits us to assume that every set of nonnegative integers has a smallest element, allowing us to use the smallest counterexample.

This is actually **equivalent** to the *principle of mathematical induction*.

Principle. (*the Weak Principle of Mathematical Induction*)

- (a) If the statement $P(b)$ is true
- (b) the statement $P(n-1) \rightarrow P(n)$ is true for all $n > b$,
then $P(n)$ is true for all integers $n \geq b$

(a) – *Basic Step* *Inductive Hypothesis*

(b) – *Inductive Step* *Inductive Conclusion*



Proof by Induction

■ $\forall n \geq 0, 2^{n+1} \geq n^2 + 2$



Proof by Induction

■ $\forall n \geq 0, 2^{n+1} \geq n^2 + 2$

Let $P(n) - 2^{n+1} \geq n^2 + 2$



Proof by Induction

■ $\forall n \geq 0, 2^{n+1} \geq n^2 + 2$

Let $P(n) - 2^{n+1} \geq n^2 + 2$

(i) Note that for $n = 0$, $2^{0+1} = 2 \geq 2 = 0^2 + 2 - P(0)$



Proof by Induction

■ $\forall n \geq 0, 2^{n+1} \geq n^2 + 2$

Let $P(n) - 2^{n+1} \geq n^2 + 2$

(i) Note that for $n = 0$, $2^{0+1} = 2 \geq 2 = 0^2 + 2 - P(0)$

(ii) Suppose that $n > 0$ and that $2^n \geq (n-1)^2 + 2 \quad (*)$



Proof by Induction

■ $\forall n \geq 0, 2^{n+1} \geq n^2 + 2$

Let $P(n) - 2^{n+1} \geq n^2 + 2$

(i) Note that for $n = 0$, $2^{0+1} = 2 \geq 2 = 0^2 + 2 - P(0)$

(ii) Suppose that $n > 0$ and that $2^n \geq (n-1)^2 + 2 \quad (*)$

$$\begin{aligned} 2^{n+1} &\geq 2(n-1)^2 + 4 \\ &= (n^2 + 2) + (n^2 - 4n + 4) \\ &= n^2 + 2 + (n-2)^2 \\ &\geq n^2 + 2 \end{aligned}$$



Proof by Induction

■ $\forall n \geq 0, 2^{n+1} \geq n^2 + 2$

Let $P(n) - 2^{n+1} \geq n^2 + 2$

(i) Note that for $n = 0$, $2^{0+1} = 2 \geq 2 = 0^2 + 2 - P(0)$

(ii) Suppose that $n > 0$ and that $2^n \geq (n-1)^2 + 2 \quad (*)$

$$\begin{aligned} 2^{n+1} &\geq 2(n-1)^2 + 4 \\ &= (n^2 + 2) + (n^2 - 4n + 4) \\ &= n^2 + 2 + (n-2)^2 \\ &\geq n^2 + 2 \end{aligned}$$

Hence, we've just prove that for $n > 0$, $P(n-1) \rightarrow P(n)$.



Proof by Induction

■ $\forall n \geq 0, 2^{n+1} \geq n^2 + 2$

Let $P(n) - 2^{n+1} \geq n^2 + 2$

(i) Note that for $n = 0$, $2^{0+1} = 2 \geq 2 = 0^2 + 2 - P(0)$

(ii) Suppose that $n > 0$ and that $2^n \geq (n-1)^2 + 2 \quad (*)$

$$\begin{aligned} 2^{n+1} &\geq 2(n-1)^2 + 4 \\ &= (n^2 + 2) + (n^2 - 4n + 4) \\ &= n^2 + 2 + (n-2)^2 \\ &\geq n^2 + 2 \end{aligned}$$

Hence, we've just prove that for $n > 0$, $P(n-1) \rightarrow P(n)$.

By mathematical induction, $\forall n > 0, 2^{n+1} \geq n^2 + 2$.



Proof by Induction

■ $\forall n \geq 2, 2^{n+1} \geq n^2 + 3$



Proof by Induction

■ $\forall n \geq 2, 2^{n+1} \geq n^2 + 3$

Let $P(n) - 2^{n+1} \geq n^2 + 3$



Proof by Induction

$$\blacksquare \forall n \geq 2, 2^{n+1} \geq n^2 + 3$$

$$\text{Let } P(n) - 2^{n+1} \geq n^2 + 3$$

$$(i) \text{ Note that for } n = 2, 2^{2+1} = 8 \geq 7 = 2^2 + 3 - P(2)$$



Proof by Induction

■ $\forall n \geq 2, 2^{n+1} \geq n^2 + 3$

Let $P(n) - 2^{n+1} \geq n^2 + 3$

(i) Note that for $n = 2, 2^{2+1} = 8 \geq 7 = 2^2 + 3 - P(2)$

(ii) Suppose that $n > 2$ and that $2^n \geq (n-1)^2 + 3 \quad (*)$



Proof by Induction

■ $\forall n \geq 2, 2^{n+1} \geq n^2 + 3$

Let $P(n) - 2^{n+1} \geq n^2 + 3$

(i) Note that for $n = 2, 2^{2+1} = 8 \geq 7 = 2^2 + 3 - P(2)$

(ii) Suppose that $n > 2$ and that $2^n \geq (n-1)^2 + 3 \quad (*)$

$$\begin{aligned} 2^{n+1} &\geq 2(n-1)^2 + 6 \\ &= n^2 + 3 + n^2 - 4n + 4 + 1 \\ &= n^2 + 3 + (n-2)^2 + 1 \\ &> n^2 + 3 \end{aligned}$$



Proof by Induction

■ $\forall n \geq 2, 2^{n+1} \geq n^2 + 3$

Let $P(n) - 2^{n+1} \geq n^2 + 3$

(i) Note that for $n = 2, 2^{2+1} = 8 \geq 7 = 2^2 + 3 - P(2)$

(ii) Suppose that $n > 2$ and that $2^n \geq (n-1)^2 + 3 \quad (*)$

$$\begin{aligned} 2^{n+1} &\geq 2(n-1)^2 + 6 \\ &= n^2 + 3 + n^2 - 4n + 4 + 1 \\ &= n^2 + 3 + (n-2)^2 + 1 \\ &> n^2 + 3 \end{aligned}$$

Hence, we've just prove that for $n > 2, P(n-1) \rightarrow P(n)$.



Proof by Induction

■ $\forall n \geq 2, 2^{n+1} \geq n^2 + 3$

Let $P(n) - 2^{n+1} \geq n^2 + 3$

(i) Note that for $n = 2, 2^{2+1} = 8 \geq 7 = 2^2 + 3 - P(2)$

(ii) Suppose that $n > 2$ and that $2^n \geq (n-1)^2 + 3 \quad (*)$

$$\begin{aligned} 2^{n+1} &\geq 2(n-1)^2 + 6 \\ &= n^2 + 3 + n^2 - 4n + 4 + 1 \\ &= n^2 + 3 + (n-2)^2 + 1 \\ &> n^2 + 3 \end{aligned}$$

Hence, we've just prove that for $n > 2, P(n-1) \rightarrow P(n)$.

By mathematical induction, $\forall n > 2, 2^{n+1} \geq n^2 + 3$.



Proof by Induction

■ $\forall n \geq 2, 2^{n+1} \geq n^2 + 3$

Let $P(n) - 2^{n+1} \geq n^2 + 3$

Base Step

(i) Note that for $n = 2, 2^{2+1} = 8 \geq 7 = 2^2 + 3 - P(2)$

(ii) Suppose that $n > 2$ and that $2^n \geq (n-1)^2 + 3$ (*)
 $2^{n+1} \geq 2(n-1)^2 + 6$ Inductive Hypothesis

$$= n^2 + 3 + n^2 - 4n + 4 + 1$$

$$= n^2 + 3 + (n-2)^2 + 1$$

$$> n^2 + 3$$

Inductive Step

Hence, we've just prove that for $n > 2, P(n-1) \rightarrow P(n)$.

By mathematical induction, $\forall n > 2, 2^{n+1} \geq n^2 + 3$.

Inductive Conclusion



Next Lecture

- induction II, ...

