



DISCRETE MATHEMATICS FOR COMPUTER SCIENCE

Dr. QI WANG

Department of Computer Science and Engineering

Office: Room903, Nanshan iPark A7 Building

Email: wangqi@sustc.edu.cn

Assignment #3

Please collect your
assignments!



Cartesian Product

- Let $A = \{a_1, a_2, \dots, a_m\}$ and $B = \{b_1, b_2, \dots, b_n\}$, the *Cartesian product* $A \times B$ is the set of pairs

$$\{(a_1, b_1), (a_2, b_2), \dots, (a_1, b_n), \dots, (a_m, b_n)\}$$



Cartesian Product

- Let $A = \{a_1, a_2, \dots, a_m\}$ and $B = \{b_1, b_2, \dots, b_n\}$, the *Cartesian product* $A \times B$ is the set of pairs

$$\{(a_1, b_1), (a_2, b_2), \dots, (a_1, b_n), \dots, (a_m, b_n)\}$$

Cartesian product defines a set of all **ordered** arrangements of elements in the two sets.



Binary Relation

- **Definition:** Let A and B be two sets. A *binary relation from A to B* is a **subset** of a **Cartesian product** $A \times B$.



Binary Relation

- **Definition:** Let A and B be two sets. A *binary relation from A to B* is a **subset** of a **Cartesian product** $A \times B$.

Let $R \subseteq A \times B$ denote R is a set of ordered pairs of the form (a, b) where $a \in A$ and $b \in B$.



Binary Relation

- **Definition:** Let A and B be two sets. A *binary relation from A to B* is a **subset** of a **Cartesian product** $A \times B$.

Let $R \subseteq A \times B$ denote R is a set of ordered pairs of the form (a, b) where $a \in A$ and $b \in B$.

We use the notation $a R b$ to denote $(a, b) \in R$, and $a \not R b$ to denote $(a, b) \notin R$.



Binary Relation

- **Definition:** Let A and B be two sets. A *binary relation from A to B* is a **subset** of a **Cartesian product** $A \times B$.

Let $R \subseteq A \times B$ denote R is a set of ordered pairs of the form (a, b) where $a \in A$ and $b \in B$.

We use the notation $a R b$ to denote $(a, b) \in R$, and $a \not R b$ to denote $(a, b) \notin R$.

Example: Let $A = \{a, b, c\}$ and $B = \{1, 2, 3\}$

- ◇ Is $R = \{(a, 1), (b, 2), (c, 2)\}$ a relation from A to B ?
- ◇ Is $Q = \{(1, a), (2, b)\}$ a relation from A to B ?
- ◇ Is $P = \{(a, a), (b, c), (b, a)\}$ a relation from A to A ?



Representing Binary Relations

- We can **graphically** represent a binary relation R as:
if $a R b$, then we draw an **arrow** from a to b : $a \rightarrow b$



Representing Binary Relations

- We can **graphically** represent a binary relation R as:
if $a R b$, then we draw an **arrow** from a to b : $a \rightarrow b$


Example: Let $A = \{0, 1, 2\}$ and $B = \{u, v\}$, and
 $R = \{(0, u), (0, v), (1, v), (2, u)\}$. ($R \subseteq A \times B$)



Representing Binary Relations

- We can **graphically** represent a binary relation R as:
if $a R b$, then we draw an **arrow** from a to b : $a \rightarrow b$

Example: Let $A = \{0, 1, 2\}$ and $B = \{u, v\}$, and $R = \{(0, u), (0, v), (1, v), (2, u)\}$. ($R \subseteq A \times B$)



Representing Binary Relations

- We can also represent a binary relation R by a table showing the ordered pairs of R .



Representing Binary Relations

- We can also represent a binary relation R by a table showing the ordered pairs of R .

Example: Let $A = \{0, 1, 2\}$ and $B = \{u, v\}$, and $R = \{(0, u), (0, v), (1, u), (2, v)\}$. ($R \subseteq A \times B$)



Representing Binary Relations

- We can also represent a binary relation R by a table showing the ordered pairs of R .

Example: Let $A = \{0, 1, 2\}$ and $B = \{u, v\}$, and $R = \{(0, u), (0, v), (1, u), (2, v)\}$. ($R \subseteq A \times B$)

| R | u | v |
|-----|-----|-----|
| 0 | × | × |
| 1 | × | |
| 2 | | × |




Relations and Functions

- Relations represent **one to many relationships** between elements in A and B .




Relations and Functions

- Relations represent **one to many relationships** between elements in A and B .



Relations and Functions

- Relations represent **one to many relationships** between elements in A and B .



What is the **difference** between a **relation** and a **function** from A to B ?

Relation on the Set

- **Definition:** A relation on the set A is a relation from A to itself.



Relation on the Set

- **Definition:** A relation on the set A is a relation from A to itself.

Example: Let $A = \{1, 2, 3, 4\}$ and $R_{div} = \{(a, b) : a|b\}$. What does R_{div} consist of?



Relation on the Set

- **Definition:** A relation on the set A is a relation from A to itself.

Example: Let $A = \{1, 2, 3, 4\}$ and $R_{div} = \{(a, b) : a|b\}$. What does R_{div} consist of?

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$





Relation on the Set

- **Definition:** A relation on the set A is a relation from A to itself.

Example: Let $A = \{1, 2, 3, 4\}$ and $R_{div} = \{(a, b) : a|b\}$. What does R_{div} consist of?

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$




Relation on the Set

- **Definition:** A relation on the set A is a relation from A to itself.

Example: Let $A = \{1, 2, 3, 4\}$ and $R_{div} = \{(a, b) : a|b\}$. What does R_{div} consist of?

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$



| R | 1 | 2 | 3 | 4 |
|-----|---|---|---|---|
| 1 | × | × | × | × |
| 2 | | × | | × |
| 3 | | | × | |
| 4 | | | | × |



Number of Binary Relations

- **Theorem** The number of binary relations on a set A , where $|A| = n$ is 2^{n^2} .



Number of Binary Relations

- **Theorem** The number of binary relations on a set A , where $|A| = n$ is 2^{n^2} .

Proof

If $|A| = n$, then the cardinality of the Cartesian product $|A \times A| = n^2$.



Number of Binary Relations

- **Theorem** The number of binary relations on a set A , where $|A| = n$ is 2^{n^2} .

Proof

If $|A| = n$, then the cardinality of the Cartesian product $|A \times A| = n^2$.

R is a binary relation on A if $R \subseteq A \times A$ (R is subset)



Number of Binary Relations

- **Theorem** The number of binary relations on a set A , where $|A| = n$ is 2^{n^2} .

Proof

If $|A| = n$, then the cardinality of the Cartesian product $|A \times A| = n^2$.

R is a binary relation on A if $R \subseteq A \times A$ (R is subset)

The number of subsets of a set with k elements is 2^k



Properties of Relations

- **Reflexive Relation:** A relation R on a set A is called *reflexive* if $(a, a) \in R$ for **every** element $a \in A$.



Properties of Relations

- **Reflexive Relation:** A relation R on a set A is called *reflexive* if $(a, a) \in R$ for **every** element $a \in A$.

Example: Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.



Properties of Relations

- **Reflexive Relation:** A relation R on a set A is called *reflexive* if $(a, a) \in R$ for **every** element $a \in A$.

Example: Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.

Is R_{div} **reflexive**?



Properties of Relations

- **Reflexive Relation:** A relation R on a set A is called *reflexive* if $(a, a) \in R$ for **every** element $a \in A$.

Example: Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.

Is R_{div} **reflexive**?

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$



Properties of Relations

- **Reflexive Relation:** A relation R on a set A is called *reflexive* if $(a, a) \in R$ for **every** element $a \in A$.

Example: Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.

Is R_{div} **reflexive**?

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$

Yes. $(1, 1), (2, 2), (3, 3), (4, 4) \in R_{div}$



Reflexive Relation

- **Example:** Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$




Reflexive Relation

- **Example:** Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$

$$MR_{div} = \begin{matrix} & 1 & 2 & 3 & 4 \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} & \begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \end{matrix}$$



Reflexive Relation

- **Example:** Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$

$$MR_{div} = \begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} & \begin{bmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \end{matrix}$$

A relation R is reflexive if and only if MR has 1 in every position on its main diagonal.



Reflexive Relation

- **Example:** Assume that $R = \{(1, 2), (2, 2), (3, 3)\}$ on $A = \{1, 2, 3, 4\}$.



Reflexive Relation

- **Example:** Assume that $R = \{(1, 2), (2, 2), (3, 3)\}$ on $A = \{1, 2, 3, 4\}$.

Is R reflexive?



Reflexive Relation

- **Example:** Assume that $R = \{(1, 2), (2, 2), (3, 3)\}$ on $A = \{1, 2, 3, 4\}$.

Is R reflexive?

No. $(1, 1) \notin R$



Properties of Relations

- **Irreflexive Relation:** A relation R on a set A is called *irreflexive* if $(a, a) \notin R$ for **every** element $a \in A$.



Properties of Relations

- **Irreflexive Relation:** A relation R on a set A is called *irreflexive* if $(a, a) \notin R$ for **every** element $a \in A$.

Example: Assume that $R_{\neq} = \{(a, b) : a \neq b\}$ on $A = \{1, 2, 3, 4\}$.



Properties of Relations

- **Irreflexive Relation:** A relation R on a set A is called *irreflexive* if $(a, a) \notin R$ for **every** element $a \in A$.

Example: Assume that $R_{\neq} = \{(a, b) : a \neq b\}$ on $A = \{1, 2, 3, 4\}$.

Is R_{\neq} **irreflexive**?



Properties of Relations

- **Irreflexive Relation:** A relation R on a set A is called *irreflexive* if $(a, a) \notin R$ for **every** element $a \in A$.

Example: Assume that $R_{\neq} = \{(a, b) : a \neq b\}$ on $A = \{1, 2, 3, 4\}$.

Is R_{\neq} **irreflexive**?

$$R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), \\ (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}.$$



Properties of Relations

- **Irreflexive Relation:** A relation R on a set A is called *irreflexive* if $(a, a) \notin R$ for **every** element $a \in A$.

Example: Assume that $R_{\neq} = \{(a, b) : a \neq b\}$ on $A = \{1, 2, 3, 4\}$.

Is R_{\neq} **irreflexive**?

$$R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), \\ (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}.$$

Yes. $(1, 1), (2, 2), (3, 3), (4, 4) \notin R_{\neq}$



Irreflexive Relation

- **Example:** Assume that $R_{\neq} = \{(a, b) : a \neq b\}$ on $A = \{1, 2, 3, 4\}$.

$$R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), \\ (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}.$$



Irreflexive Relation

- **Example:** Assume that $R_{\neq} = \{(a, b) : a \neq b\}$ on $A = \{1, 2, 3, 4\}$.

$$R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), \\ (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}.$$

$$\text{MR} = \begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} & \begin{bmatrix} 0 & 1 & 1 & 1 \\ 1 & 0 & 1 & 1 \\ 1 & 1 & 0 & 1 \\ 1 & 1 & 1 & 0 \end{bmatrix} \end{matrix}$$




Irreflexive Relation

- **Example:** Assume that $R_{\neq} = \{(a, b) : a \neq b\}$ on $A = \{1, 2, 3, 4\}$.

$$R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), \\ (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}.$$

$$\text{MR} = \begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} & \begin{bmatrix} 0 & 1 & 1 & 1 \\ 1 & 0 & 1 & 1 \\ 1 & 1 & 0 & 1 \\ 1 & 1 & 1 & 0 \end{bmatrix} \end{matrix}$$

A relation R is irreflexive if and only if MR has 0 in every position on its **main diagonal**.



Properties of Relations

- **Symmetric Relation:** A relation R on a set A is called *symmetric* if $(b, a) \in R$ whenever $(a, b) \in R$ for all $a, b \in A$.



Properties of Relations

- **Symmetric Relation:** A relation R on a set A is called *symmetric* if $(b, a) \in R$ whenever $(a, b) \in R$ for all $a, b \in A$.

Example: Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.



Properties of Relations

- **Symmetric Relation:** A relation R on a set A is called *symmetric* if $(b, a) \in R$ whenever $(a, b) \in R$ for all $a, b \in A$.

Example: Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.

Is R_{div} symmetric?



Properties of Relations

- **Symmetric Relation:** A relation R on a set A is called *symmetric* if $(b, a) \in R$ whenever $(a, b) \in R$ for all $a, b \in A$.

Example: Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.

Is R_{div} *symmetric*?

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$



Properties of Relations

- **Symmetric Relation:** A relation R on a set A is called *symmetric* if $(b, a) \in R$ whenever $(a, b) \in R$ for all $a, b \in A$.

Example: Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.

Is R_{div} symmetric?

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$

No. $(1, 2) \in R_{div}$ but $(2, 1) \notin R$



Symmetric Relation

- **Example:** Assume that $R_{\neq} = \{(a, b) : a \neq b\}$ on $A = \{1, 2, 3, 4\}$.

$$R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), \\ (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}.$$



Symmetric Relation

- **Example:** Assume that $R_{\neq} = \{(a, b) : a \neq b\}$ on $A = \{1, 2, 3, 4\}$.

$$R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), \\ (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}.$$

Is R_{\neq} symmetric?



Symmetric Relation

- **Example:** Assume that $R_{\neq} = \{(a, b) : a \neq b\}$ on $A = \{1, 2, 3, 4\}$.

$$R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), \\ (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}.$$

Is R_{\neq} symmetric?

Yes. If $(a, b) \in R_{\neq}$ then $(b, a) \in R_{\neq}$.



Symmetric Relation

- **Example:** Assume that $R_{\neq} = \{(a, b) : a \neq b\}$ on $A = \{1, 2, 3, 4\}$.

$$R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), \\ (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}.$$



Symmetric Relation

- **Example:** Assume that $R_{\neq} = \{(a, b) : a \neq b\}$ on $A = \{1, 2, 3, 4\}$.

$$R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), \\ (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}.$$

$$\text{MR} = \begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} & \begin{bmatrix} 0 & 1 & 1 & 1 \\ 1 & 0 & 1 & 1 \\ 1 & 1 & 0 & 1 \\ 1 & 1 & 1 & 0 \end{bmatrix} \end{matrix}$$



Symmetric Relation

- **Example:** Assume that $R_{\neq} = \{(a, b) : a \neq b\}$ on $A = \{1, 2, 3, 4\}$.

$$R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), \\ (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}.$$

$$\text{MR} = \begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} & \begin{bmatrix} 0 & 1 & 1 & 1 \\ 1 & 0 & 1 & 1 \\ 1 & 1 & 0 & 1 \\ 1 & 1 & 1 & 0 \end{bmatrix} \end{matrix}$$

A relation R is symmetric if and only if **MR is symmetric**.



Properties of Relations

- **Antisymmetric Relation:** A relation R on a set A is called *antisymmetric* if $(b, a) \in R$ and $(a, b) \in R$ implies $a = b$ for all $a, b \in A$.



Properties of Relations

- **Antisymmetric Relation:** A relation R on a set A is called *antisymmetric* if $(b, a) \in R$ and $(a, b) \in R$ implies $a = b$ for all $a, b \in A$.

Example: Assume that $R = \{(1, 2), (2, 2), (3, 3)\}$ on $A = \{1, 2, 3, 4\}$.



Properties of Relations

- **Antisymmetric Relation:** A relation R on a set A is called *antisymmetric* if $(b, a) \in R$ and $(a, b) \in R$ implies $a = b$ for all $a, b \in A$.

Example: Assume that $R = \{(1, 2), (2, 2), (3, 3)\}$ on $A = \{1, 2, 3, 4\}$.

Is R antisymmetric?



Properties of Relations

- **Antisymmetric Relation:** A relation R on a set A is called *antisymmetric* if $(b, a) \in R$ and $(a, b) \in R$ implies $a = b$ for all $a, b \in A$.

Example: Assume that $R = \{(1, 2), (2, 2), (3, 3)\}$ on $A = \{1, 2, 3, 4\}$.

Is R antisymmetric?

Yes.



Antisymmetric Relation

- **Example:** Assume that $R = \{(1, 2), (2, 2), (3, 3)\}$ on $A = \{1, 2, 3, 4\}$.



Antisymmetric Relation

- **Example:** Assume that $R = \{(1, 2), (2, 2), (3, 3)\}$ on $A = \{1, 2, 3, 4\}$.

$$MR = \begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} & \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \end{matrix}$$



Antisymmetric Relation

- **Example:** Assume that $R = \{(1, 2), (2, 2), (3, 3)\}$ on $A = \{1, 2, 3, 4\}$.

$$MR = \begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} & \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \end{matrix}$$

A relation R is antisymmetric if and only if $m_{ij} = 1$ implies $m_{ji} = 0$ for $i \neq j$.



Antisymmetric Relation

- **Example:** Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.



Antisymmetric Relation

- **Example:** Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.

Is R_{div} antisymmetric?



Antisymmetric Relation

- **Example:** Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.

Is R_{div} antisymmetric?

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$



Antisymmetric Relation

- **Example:** Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.

Is R_{div} antisymmetric?

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$

Yes. If $a|b$ and $b|a$, then $a = b$.



Antisymmetric Relation

- **Example:** Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.

Is R_{div} antisymmetric?

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$

Yes. If $a|b$ and $b|a$, then $a = b$.

$$MR_{div} = \begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} & \begin{bmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \end{matrix}$$



Properties of Relations

- **Transitive Relation:** A relation R on a set A is called *transitive* if $(a, b) \in R$ and $(b, c) \in R$ implies $(a, c) \in R$ for **all** $a, b, c \in A$.



Properties of Relations

- **Transitive Relation:** A relation R on a set A is called *transitive* if $(a, b) \in R$ and $(b, c) \in R$ implies $(a, c) \in R$ for **all** $a, b, c \in A$.

Example: Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.



Properties of Relations

- **Transitive Relation:** A relation R on a set A is called *transitive* if $(a, b) \in R$ and $(b, c) \in R$ implies $(a, c) \in R$ for **all** $a, b, c \in A$.

Example: Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.

Is R_{div} *transitive*?



Properties of Relations

- **Transitive Relation:** A relation R on a set A is called *transitive* if $(a, b) \in R$ and $(b, c) \in R$ implies $(a, c) \in R$ for **all** $a, b, c \in A$.

Example: Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.

Is R_{div} *transitive*?

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$



Properties of Relations

- **Transitive Relation:** A relation R on a set A is called *transitive* if $(a, b) \in R$ and $(b, c) \in R$ implies $(a, c) \in R$ for **all** $a, b, c \in A$.

Example: Assume that $R_{div} = \{(a, b) : a|b\}$ on $A = \{1, 2, 3, 4\}$.

Is R_{div} *transitive*?

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$

Yes. If $a|b$ and $b|c$, then $a|c$.



Transitive Relation

- **Example:** Assume that $R_{\neq} = \{(a, b) : a \neq b\}$ on $A = \{1, 2, 3, 4\}$.

$$R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), \\ (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}.$$



Transitive Relation

- **Example:** Assume that $R_{\neq} = \{(a, b) : a \neq b\}$ on $A = \{1, 2, 3, 4\}$.

$$R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), \\ (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}.$$

Is R_{\neq} transitive?



Transitive Relation

- **Example:** Assume that $R_{\neq} = \{(a, b) : a \neq b\}$ on $A = \{1, 2, 3, 4\}$.

$$R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), \\ (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}.$$

Is R_{\neq} transitive?

No. $(1, 2), (2, 1) \in R_{\neq}$ but $(1, 1) \notin R_{\neq}$.



Transitive Relation

- **Example:** Assume that $R = \{(1, 2), (2, 2), (3, 3)\}$ on $A = \{1, 2, 3, 4\}$.



Transitive Relation

- **Example:** Assume that $R = \{(1, 2), (2, 2), (3, 3)\}$ on $A = \{1, 2, 3, 4\}$.

Is R transitive?



Transitive Relation

- **Example:** Assume that $R = \{(1, 2), (2, 2), (3, 3)\}$ on $A = \{1, 2, 3, 4\}$.

Is R transitive?

Yes.



Combining Relations

- **Definition:** Let A and B be two sets. A *binary relation from A to B* is a **subset** of a **Cartesian product** $A \times B$.

Let $R \subseteq A \times B$ denote R is a set of ordered pairs of the form (a, b) where $a \in A$ and $b \in B$.



Combining Relations

- **Definition:** Let A and B be two sets. A *binary relation from A to B* is a **subset** of a **Cartesian product** $A \times B$.

Let $R \subseteq A \times B$ denote R is a set of ordered pairs of the form (a, b) where $a \in A$ and $b \in B$.

Combining Relations: Since **relations are sets**, we can *combine* relations via **set operations**.



Combining Relations

- **Definition:** Let A and B be two sets. A *binary relation from A to B* is a **subset** of a **Cartesian product** $A \times B$.

Let $R \subseteq A \times B$ denote R is a set of ordered pairs of the form (a, b) where $a \in A$ and $b \in B$.

Combining Relations: Since **relations are sets**, we can *combine* relations via **set operations**.

Set operations: **union, intersection, difference, etc.**



Combining Relations

- **Example:** Let $A = \{1, 2, 3\}$, $B = \{u, v\}$, and
 $R_1 = \{(1, u), (2, u), (2, v), (3, u)\}$,
 $R_2 = \{(1, v), (3, u), (3, v)\}$



Combining Relations

- **Example:** Let $A = \{1, 2, 3\}$, $B = \{u, v\}$, and
 $R_1 = \{(1, u), (2, u), (2, v), (3, u)\}$,
 $R_2 = \{(1, v), (3, u), (3, v)\}$

What is $R_1 \cup R_2$, $R_1 \cap R_2$, $R_1 - R_2$, $R_2 - R_1$?



Combining Relations

- **Example:** Let $A = \{1, 2, 3\}$, $B = \{u, v\}$, and
 $R_1 = \{(1, u), (2, u), (2, v), (3, u)\}$,
 $R_2 = \{(1, v), (3, u), (3, v)\}$

What is $R_1 \cup R_2$, $R_1 \cap R_2$, $R_1 - R_2$, $R_2 - R_1$?

We may also combine relations by **matrix operations**.



Composite of Relations

- **Definition:** Let R be a relation from a set A to a set B and S be a relation from B to C . The *composite of R and S* is the relation consisting of the ordered pairs (a, c) where $a \in A$ and $c \in C$ and for which there is a $b \in B$ such that $(a, b) \in R$ and $(b, c) \in S$. We denote the composite of R and S by $S \circ R$.



Composite of Relations

- **Definition:** Let R be a relation from a set A to a set B and S be a relation from B to C . The *composite of R and S* is the relation consisting of the ordered pairs (a, c) where $a \in A$ and $c \in C$ and for which there is a $b \in B$ such that $(a, b) \in R$ and $(b, c) \in S$. We denote the composite of R and S by $S \circ R$.

Example: Let $A = \{1, 2, 3\}$, $B = \{0, 1, 2\}$, and $C = \{a, b\}$




Composite of Relations

- **Definition:** Let R be a relation from a set A to a set B and S be a relation from B to C . The *composite of R and S* is the relation consisting of the ordered pairs (a, c) where $a \in A$ and $c \in C$ and for which there is a $b \in B$ such that $(a, b) \in R$ and $(b, c) \in S$. We denote the composite of R and S by $S \circ R$.

Example: Let $A = \{1, 2, 3\}$, $B = \{0, 1, 2\}$, and $C = \{a, b\}$

$$R = \{(1, 0), (1, 2), (3, 1), (3, 2)\}$$

$$S = \{(0, b), (1, a), (2, b)\}$$



Composite of Relations


- **Definition:** Let R be a relation from a set A to a set B and S be a relation from B to C . The *composite of R and S* is the relation consisting of the ordered pairs (a, c) where $a \in A$ and $c \in C$ and for which there is a $b \in B$ such that $(a, b) \in R$ and $(b, c) \in S$. We denote the composite of R and S by $S \circ R$.

Example: Let $A = \{1, 2, 3\}$, $B = \{0, 1, 2\}$, and $C = \{a, b\}$

$$R = \{(1, 0), (1, 2), (3, 1), (3, 2)\}$$

$$S = \{(0, b), (1, a), (2, b)\}$$

$$S \circ R = \{(1, b), (3, a), (3, b)\}$$



Implementation of Composite

- **Example:** Let $A = \{1, 2\}$, $B = \{1, 2, 3\}$, and $C = \{a, b\}$
 $R = \{(1, 2), (1, 3), (2, 1)\}$ is a relation from A to B
 $S = \{(1, a), (3, b), (3, a)\}$ is a relation from B to C



Implementation of Composite

- **Example:** Let $A = \{1, 2\}$, $B = \{1, 2, 3\}$, and $C = \{a, b\}$
 $R = \{(1, 2), (1, 3), (2, 1)\}$ is a relation from A to B
 $S = \{(1, a), (3, b), (3, a)\}$ is a relation from B to C
 $S \circ R = \{(1, b), (1, a), (2, a)\}$



Implementation of Composite

■ **Example:** Let $A = \{1, 2\}$, $B = \{1, 2, 3\}$, and $C = \{a, b\}$

$R = \{(1, 2), (1, 3), (2, 1)\}$ is a relation from A to B

$S = \{(1, a), (3, b), (3, a)\}$ is a relation from B to C

$S \circ R = \{(1, b), (1, a), (2, a)\}$

$$M_R = \begin{matrix} & \begin{matrix} 1 & 2 & 3 \end{matrix} \\ \begin{matrix} 1 \\ 2 \end{matrix} & \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 0 \end{bmatrix} \end{matrix}$$



Implementation of Composite

■ **Example:** Let $A = \{1, 2\}$, $B = \{1, 2, 3\}$, and $C = \{a, b\}$

$R = \{(1, 2), (1, 3), (2, 1)\}$ is a relation from A to B

$S = \{(1, a), (3, b), (3, a)\}$ is a relation from B to C

$S \circ R = \{(1, b), (1, a), (2, a)\}$

$$M_R = \begin{matrix} & \begin{matrix} 1 & 2 & 3 \end{matrix} \\ \begin{matrix} 1 \\ 2 \end{matrix} & \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 0 \end{bmatrix} \end{matrix} \quad M_S = \begin{matrix} & \begin{matrix} a & b \end{matrix} \\ \begin{matrix} 1 \\ 3 \end{matrix} & \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \end{matrix}$$



Implementation of Composite

■ **Example:** Let $A = \{1, 2\}$, $B = \{1, 2, 3\}$, and $C = \{a, b\}$

$R = \{(1, 2), (1, 3), (2, 1)\}$ is a relation from A to B

$S = \{(1, a), (3, b), (3, a)\}$ is a relation from B to C

$S \circ R = \{(1, b), (1, a), (2, a)\}$

$$M_R = \begin{matrix} & \begin{matrix} 1 & 2 & 3 \end{matrix} \\ \begin{matrix} 1 \\ 2 \end{matrix} & \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 0 \end{bmatrix} \end{matrix} \quad M_S = \begin{matrix} & \begin{matrix} a & b \end{matrix} \\ \begin{matrix} 1 \\ 3 \end{matrix} & \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \end{matrix}$$

$$M_R \odot M_S = \begin{matrix} & \begin{matrix} a & b \end{matrix} \\ \begin{matrix} 1 \\ 2 \end{matrix} & \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} \end{matrix}$$



Composite of Relations

- **Definition** Let R be a relation on A . The *powers* R^n , for $n = 1, 2, 3, \dots$, is defined **inductively** by

$$R^1 = R \text{ and } R^{n+1} = R^n \circ R$$



Composite of Relations

- **Definition** Let R be a relation on A . The *powers* R^n , for $n = 1, 2, 3, \dots$, is defined **inductively** by

$$R^1 = R \text{ and } R^{n+1} = R^n \circ R$$

Example: Let $A = \{1, 2, 3, 4\}$, and
 $R = \{(1, 2), (2, 3), (2, 4), (3, 3)\}$



Composite of Relations

- **Definition** Let R be a relation on A . The *powers* R^n , for $n = 1, 2, 3, \dots$, is defined **inductively** by

$$R^1 = R \text{ and } R^{n+1} = R^n \circ R$$

Example: Let $A = \{1, 2, 3, 4\}$, and
 $R = \{(1, 2), (2, 3), (2, 4), (3, 3)\}$

$$R^1 = R$$



Composite of Relations

- **Definition** Let R be a relation on A . The *powers* R^n , for $n = 1, 2, 3, \dots$, is defined **inductively** by

$$R^1 = R \text{ and } R^{n+1} = R^n \circ R$$

Example: Let $A = \{1, 2, 3, 4\}$, and
 $R = \{(1, 2), (2, 3), (2, 4), (3, 3)\}$

$$R^1 = R$$

$$R^2 = R \circ R = \{(1, 3), (1, 4), (2, 3), (3, 3)\}$$



Composite of Relations

- **Definition** Let R be a relation on A . The *powers* R^n , for $n = 1, 2, 3, \dots$, is defined **inductively** by


$$R^1 = R \text{ and } R^{n+1} = R^n \circ R$$

Example: Let $A = \{1, 2, 3, 4\}$, and
 $R = \{(1, 2), (2, 3), (2, 4), (3, 3)\}$

$$R^1 = R$$

$$R^2 = R \circ R = \{(1, 3), (1, 4), (2, 3), (3, 3)\}$$

$$R^3 = R^2 \circ R = \{(1, 3), (2, 3), (3, 3)\}$$



Composite of Relations

- **Definition** Let R be a relation on A . The *powers* R^n , for $n = 1, 2, 3, \dots$, is defined **inductively** by

$$R^1 = R \text{ and } R^{n+1} = R^n \circ R$$


Example: Let $A = \{1, 2, 3, 4\}$, and
 $R = \{(1, 2), (2, 3), (2, 4), (3, 3)\}$

$$R^1 = R$$

$$R^2 = R \circ R = \{(1, 3), (1, 4), (2, 3), (3, 3)\}$$

$$R^3 = R^2 \circ R = \{(1, 3), (2, 3), (3, 3)\}$$

$$R^4 = R^3 \circ R = \{(1, 3), (2, 3), (3, 3)\}$$



Composite of Relations

- **Definition** Let R be a relation on A . The *powers* R^n , for $n = 1, 2, 3, \dots$, is defined **inductively** by

$$R^1 = R \text{ and } R^{n+1} = R^n \circ R$$

Example: Let $A = \{1, 2, 3, 4\}$, and
 $R = \{(1, 2), (2, 3), (2, 4), (3, 3)\}$

$$R^1 = R$$

$$R^2 = R \circ R = \{(1, 3), (1, 4), (2, 3), (3, 3)\}$$

$$R^3 = R^2 \circ R = \{(1, 3), (2, 3), (3, 3)\}$$

$$R^4 = R^3 \circ R = \{(1, 3), (2, 3), (3, 3)\}$$

$$R^k = ? \text{ for } k > 3$$



Transitive Relation and R^n

- **Theorem** The relation R on a set A is transitive if and only if $R^n \subseteq R$ for $n = 1, 2, 3, \dots$



Transitive Relation and R^n

- **Theorem** The relation R on a set A is transitive if and only if $R^n \subseteq R$ for $n = 1, 2, 3, \dots$

Proof.



Transitive Relation and R^n

- **Theorem** The relation R on a set A is transitive if and only if $R^n \subseteq R$ for $n = 1, 2, 3, \dots$

Proof.

“if” part: In particular, $R^2 \subseteq R$.



Transitive Relation and R^n

- **Theorem** The relation R on a set A is transitive if and only if $R^n \subseteq R$ for $n = 1, 2, 3, \dots$

Proof.

“if” part: In particular, $R^2 \subseteq R$.

If $(a, b) \in R$ and $(b, c) \in R$, then by the definition of composition, we have $(a, c) \in R^2 \subseteq R$.



Transitive Relation and R^n

- **Theorem** The relation R on a set A is transitive if and only if $R^n \subseteq R$ for $n = 1, 2, 3, \dots$

Proof.

“if” part: In particular, $R^2 \subseteq R$.

If $(a, b) \in R$ and $(b, c) \in R$, then by the definition of composition, we have $(a, c) \in R^2 \subseteq R$.

“only if” part: by induction.



Number of Reflexive Relations

- **Theorem** The number of reflexive relations on a set A with $|A| = n$ is $2^{n(n-1)}$.



Number of Reflexive Relations

- **Theorem** The number of reflexive relations on a set A with $|A| = n$ is $2^{n(n-1)}$.

Proof. A reflexive relation R on A must contain all pairs (a, a) for every $a \in A$.



Number of Reflexive Relations

- **Theorem** The number of reflexive relations on a set A with $|A| = n$ is $2^{n(n-1)}$.

Proof. A reflexive relation R on A must contain all pairs (a, a) for every $a \in A$.

All other pairs in R are of the form (a, b) with $a \neq b$ s.t. $a, b \in A$.



Number of Reflexive Relations

- **Theorem** The number of reflexive relations on a set A with $|A| = n$ is $2^{n(n-1)}$.

Proof. A reflexive relation R on A must contain all pairs (a, a) for every $a \in A$.

All other pairs in R are of the form (a, b) with $a \neq b$ s.t. $a, b \in A$.

How many of these pairs are there?



Number of Reflexive Relations

- **Theorem** The number of reflexive relations on a set A with $|A| = n$ is $2^{n(n-1)}$.

Proof. A reflexive relation R on A must contain all pairs (a, a) for every $a \in A$.

All other pairs in R are of the form (a, b) with $a \neq b$ s.t. $a, b \in A$.

How many of these pairs are there?

How many subsets on $n(n-1)$ elements are there?



Summary on Properties of Relations

- **Reflexive Relation:** A relation R on a set A is called *reflexive* if $(a, a) \in R$ for **every** element $a \in A$.



Summary on Properties of Relations

- **Reflexive Relation:** A relation R on a set A is called *reflexive* if $(a, a) \in R$ for **every** element $a \in A$.

Irreflexive Relation: A relation R on a set A is called *irreflexive* if $(a, a) \notin R$ for **every** element $a \in A$.



Summary on Properties of Relations

- **Reflexive Relation:** A relation R on a set A is called *reflexive* if $(a, a) \in R$ for **every** element $a \in A$.

Irreflexive Relation: A relation R on a set A is called *irreflexive* if $(a, a) \notin R$ for **every** element $a \in A$.

Symmetric Relation: A relation R on a set A is called *symmetric* if $(b, a) \in R$ whenever $(a, b) \in R$ for **all** $a, b \in A$.



Summary on Properties of Relations

■ **Reflexive Relation:** A relation R on a set A is called *reflexive* if $(a, a) \in R$ for **every** element $a \in A$.

Irreflexive Relation: A relation R on a set A is called *irreflexive* if $(a, a) \notin R$ for **every** element $a \in A$.

Symmetric Relation: A relation R on a set A is called *symmetric* if $(b, a) \in R$ whenever $(a, b) \in R$ for **all** $a, b \in A$.

Antisymmetric Relation: A relation R on a set A is called *antisymmetric* if $(b, a) \in R$ and $(a, b) \in R$ implies $a = b$ for **all** $a, b \in A$.



Summary on Properties of Relations

■ **Reflexive Relation:** A relation R on a set A is called *reflexive* if $(a, a) \in R$ for **every** element $a \in A$.

Irreflexive Relation: A relation R on a set A is called *irreflexive* if $(a, a) \notin R$ for **every** element $a \in A$.

Symmetric Relation: A relation R on a set A is called *symmetric* if $(b, a) \in R$ whenever $(a, b) \in R$ for **all** $a, b \in A$.

Antisymmetric Relation: A relation R on a set A is called *antisymmetric* if $(b, a) \in R$ and $(a, b) \in R$ implies $a = b$ for **all** $a, b \in A$.

Transitive Relation: A relation R on a set A is called *transitive* if $(a, b) \in R$ and $(b, c) \in R$ implies $(a, c) \in R$ for **all** $a, b, c \in A$.



n -ary Relations

- **Definition** An n -ary relation R on sets A_1, \dots, A_n , written as $R : A_1, \dots, A_n$, is a subset $R \subseteq A_1 \times \dots \times A_n$.



n -ary Relations

- **Definition** An n -ary relation R on sets A_1, \dots, A_n , written as $R : A_1, \dots, A_n$, is a subset $R \subseteq A_1 \times \dots \times A_n$.
 - The sets A_i 's are called the *domains* of R .



n -ary Relations

- **Definition** An n -ary relation R on sets A_1, \dots, A_n , written as $R : A_1, \dots, A_n$, is a subset $R \subseteq A_1 \times \dots \times A_n$.
 - The sets A_i 's are called the *domains* of R .
 - The *degree* of R is n .



n -ary Relations

- **Definition** An n -ary relation R on sets A_1, \dots, A_n , written as $R : A_1, \dots, A_n$, is a subset $R \subseteq A_1 \times \dots \times A_n$.
 - The sets A_i 's are called the *domains* of R .
 - The *degree* of R is n .
 - R is *functional* in domain A_i if it contains **at most one** n -tuple (\dots, a_i, \dots) for any value a_i within domain A_i .



Relational Databases

- A *relational database* is essentially an *n*-ary relation *R*.



Relational Databases

- A *relational database* is essentially an *n*-ary relation R .
- A domain A_i is a *primary key* for the database if the relation R is *functional* in A_i .




Relational Databases

- A *relational database* is essentially an n -ary relation R .
- A domain A_i is a *primary key* for the database if the relation R is *functional* in A_i .
- A *composite key* for the database is a set of domains $\{A_i, A_j, \dots\}$ such that R contains **at most 1 n -tuple** $(\dots, a_i, \dots, a_j, \dots)$ for each composite value $(a_i, a_j, \dots) \in A_i \times A_j \times \dots$.



Relational Databases

E-R Diagram



Selection Operators

- Let A be any *n -ary domain* $A = A_1 \times \cdots \times A_n$, and let $C : A \rightarrow \{T, F\}$ be any *condition* (predicate) on elements (n -tuples) of A .



Selection Operators

- Let A be any *n -ary domain* $A = A_1 \times \cdots \times A_n$, and let $C : A \rightarrow \{T, F\}$ be any *condition* (predicate) on elements (n -tuples) of A .
- The *selection operator* s_C is the operator that maps any (n -ary) relation R on A to the n -ary relation of all n -tuples from R that *satisfy* C .



Selection Operators

- Let A be any *n -ary domain* $A = A_1 \times \cdots \times A_n$, and let $C : A \rightarrow \{T, F\}$ be any *condition* (predicate) on elements (n -tuples) of A .
- The *selection operator* s_C is the operator that maps any (n -ary) relation R on A to the n -ary relation of all n -tuples from R that *satisfy* C .

$$- \forall R \subseteq A,$$

$$\begin{aligned} s_C(R) &= R \cap \{a \in A \mid s_C(a) = T\} \\ &= \{a \in R \mid s_C(a) = T\}. \end{aligned}$$



Selection Operator Example

- Suppose that we have a domain

$$A = \textit{StudentName} \times \textit{Standing} \times \textit{SocSecNos}$$



Selection Operator Example

- Suppose that we have a domain

$$A = \textit{StudentName} \times \textit{Standing} \times \textit{SocSecNos}$$

- Suppose that we have a domain

$$\begin{aligned} &\textit{UpperLevel}(\textit{name}, \textit{standing}, \textit{ssn}) \\ &::= [(\textit{standing} = \textit{junior}) \vee (\textit{standing} = \textit{senior})] \end{aligned}$$



Selection Operator Example

- Suppose that we have a domain

$$A = \textit{StudentName} \times \textit{Standing} \times \textit{SocSecNos}$$

- Suppose that we have a domain

$$\begin{aligned} &\textit{UpperLevel}(\textit{name}, \textit{standing}, \textit{ssn}) \\ &:\equiv [(\textit{standing} = \textit{junior}) \vee (\textit{standing} = \textit{senior})] \end{aligned}$$

- Then, $\textit{SUpperLevel}$ is the selection operator that takes any relation R on A (database of students) and produces a relation consisting of just the upper-level classes (juniors and seniors).



Projection Operators

- Let $A = A_1 \times \cdots \times A_n$ be any n -ary domain, and let $\{i_k\} = (i_1, \dots, i_m)$ be a sequence of indices all falling in the range 1 to n .
i.e., where $1 \leq i_k \leq n$ for all $1 \leq k \leq m$.



Projection Operators

- Let $A = A_1 \times \cdots \times A_n$ be any *n-ary domain*, and let $\{i_k\} = (i_1, \dots, i_m)$ be a sequence of indices all falling in the range 1 to n .

i.e., where $1 \leq i_k \leq n$ for all $1 \leq k \leq m$.

- Then the *projection operator* on n -tuples

$$P_{\{i_k\}} : A \rightarrow A_{i_1} \times \cdots \times A_{i_m}$$

is defined by

$$P_{\{i_k\}}(a_1, \dots, a_n) = (a_{i_1}, \dots, a_{i_m})$$



Projection Example

- Suppose that we have a tenary domain

$$\textit{Cars} = \textit{Model} \times \textit{Year} \times \textit{Color} \ (n = 3)$$



Projection Example

- Suppose that we have a ternary domain

$$Cars = Model \times Year \times Color \quad (n = 3)$$

- Consider the index sequence $\{i_k\} = \{1, 3\} \quad (m = 2)$



Projection Example

- Suppose that we have a ternary domain

$$Cars = Model \times Year \times Color \quad (n = 3)$$

- Consider the index sequence $\{i_k\} = \{1, 3\}$ ($m = 2$)

- Then the projection $P_{\{i_k\}}$ simply maps each tuple $(a_1, a_2, a_3) = (model, year, color)$ to its image:

$$(a_{i_1}, a_{i_2}) = (a_1, a_3) = (model, color)$$



Projection Example

- Suppose that we have a tenary domain

$$Cars = Model \times Year \times Color \quad (n = 3)$$

- Consider the index sequence $\{i_k\} = \{1, 3\}$ ($m = 2$)

- Then the projection $P_{\{i_k\}}$ simply maps each tuple $(a_1, a_2, a_3) = (model, year, color)$ to its image:

$$(a_{i_1}, a_{i_2}) = (a_1, a_3) = (model, color)$$

- This operator can be usefully applied to a whole relation $R \subseteq Cars$ (database of cars) to obtain a list of *model/color* combinations available.



Join Operator

- Puts two relations together to form a sort of *combined relation*.



Join Operator

- Puts two relations together to form a sort of *combined relation*.
- If the tuple (A, B) appears in R_1 , and the tuple (B, C) appears in R_2 , then the tuple (A, B, C) appears in the *join* $J(R_1, R_2)$.



Join Operator

- Puts two relations together to form a sort of *combined relation*.
- If the tuple (A, B) appears in R_1 , and the tuple (B, C) appears in R_2 , then the tuple (A, B, C) appears in the *join* $J(R_1, R_2)$.
- A, B, C can also be sequences of elements rather than single elements.



Join Example

- Suppose that R_1 is a teaching assignment table, relating *Professors* to *Courses*.



Join Example

- Suppose that R_1 is a teaching assignment table, relating *Professors* to *Courses*.
- Suppose that R_2 is a room assignment table relating *Courses* to *Rooms* and *Times*.



Join Example

- Suppose that R_1 is a teaching assignment table, relating *Professors* to *Courses*.
- Suppose that R_2 is a room assignment table relating *Courses* to *Rooms* and *Times*.
- Then $J(R_1, R_2)$ is like your **class schedule**, listing *(professor, course, room, time)*.



Representing Relations

- Some ways to represent *n-ary* relations:
 - with an *explicit list* or *table* of its tuples
 - with a *function* from the domain to $\{T, F\}$



Representing Relations

- Some ways to represent *n-ary* relations:
 - with an *explicit list* or *table* of its tuples
 - with a *function* from the domain to $\{T, F\}$
- Some special ways to represent *binary* relations:
 - with a *zero-one matrix*
 - with a *directed graph*



Next Lecture

- relation II ...

