



DISCRETE MATHEMATICS FOR COMPUTER SCIENCE

Dr. QI WANG

Department of Computer Science and Engineering

Office: Room903, Nanshan iPark A7 Building

Email: wangqi@sustc.edu.cn

Assignment #2

Please collect your
assignments!



Counting

- Assume we have a set of objects with certain properties


Counting

- Assume we have a set of objects with certain properties
Counting is used to determine the number of these objects.

Counting

- Assume we have a set of objects with certain properties
Counting is used to determine the number of these objects.

How many different ways are
there to choose 2 balls from



Counting

- Assume we have a set of objects with certain properties
Counting is used to determine the number of these objects.

How many different ways are there to choose 2 balls from



Counting

- Assume we have a set of objects with certain properties
Counting is used to determine the number of these objects.

How many different ways are there to choose 2 balls from




What about when order counts?


Counting

- Assume we have a set of objects with certain properties
Counting is used to determine the number of these objects.

How many different ways are there to choose 2 balls from



What about when order counts?



Counting

- Assume we have a set of objects with certain properties
Counting is used to determine the number of these objects.



Counting

- Assume we have a set of objects with certain properties
Counting is used to determine the number of these objects.

Examples

- ◇ the number of steps in a computer program
- ◇ the number of passwords between 6 – 10 characters
- ◇ the number of telephone numbers with 8 digits



Counting

- Assume we have a set of objects with certain properties
Counting is used to determine the number of these objects.

Examples

- ◇ the number of steps in a computer program
- ◇ the number of passwords between 6 – 10 characters
- ◇ the number of telephone numbers with 8 digits

Counting may be very hard, not trivial.



Counting

- Assume we have a set of objects with certain properties
Counting is used to determine the number of these objects.

Examples

- ◇ the number of steps in a computer program
- ◇ the number of passwords between 6 – 10 characters
- ◇ the number of telephone numbers with 8 digits

Counting may be very hard, not trivial.

- simplify the solution by decomposing the problem



Basic Counting Rules

- *the Product Rule*

- *the Sum Rule*



Basic Counting Rules

■ *the Product Rule*

- ◇ A count decomposes into a sequence of **dependent** counts
(each element in the first count is associated with all elements of the second count)

■ *the Sum Rule*

- ◇ A count decomposes into a set of **independent** counts
(elements of counts are alternatives)



The Product Rule

- A count decomposes into a sequence of **dependent** counts
(each element in the first count is associated with all elements of the second count)



The Product Rule

- A count decomposes into a sequence of **dependent** counts (each element in the first count is associated with all elements of the second count)

Example

In an auditorium, the seats are labeled by a letter and numbers in between 1 to 50 (e.g., A23). What is the total number of seats?



The Product Rule

- A count decomposes into a sequence of **dependent** counts (each element in the first count is associated with all elements of the second count)

Example

In an auditorium, the seats are labeled by a letter and numbers in between 1 to 50 (e.g., A23). What is the total number of seats?

We may either list all or use the product rule.

$$26 \times 50 = 1300$$



The Product Rule

- **Product Rule:** If a count of elements can be broken down into a **sequence of dependent counts** where the first count yields n_1 elements, the second n_2 elements, and k th count n_k elements, then the total number of elements is

$$n = n_1 \cdot n_2 \cdot \cdots \cdot n_k$$



The Product Rule

- **Product Rule:** If a count of elements can be broken down into a **sequence of dependent counts** where the first count yields n_1 elements, the second n_2 elements, and k th count n_k elements, then the total number of elements is

$$n = n_1 \cdot n_2 \cdot \dots \cdot n_k$$

Example

How many different bit strings of length 7 are there?



The Product Rule


- **Product Rule:** If a count of elements can be broken down into a **sequence of dependent counts** where the first count yields n_1 elements, the second n_2 elements, and k th count n_k elements, then the total number of elements is

$$n = n_1 \cdot n_2 \cdot \dots \cdot n_k$$

Example

How many different bit strings of length 7 are there?

How many different functions are there from a set with m elements to a set with n elements?



The Product Rule

- **Product Rule:** If a count of elements can be broken down into a **sequence of dependent counts** where the first count yields n_1 elements, the second n_2 elements, and k th count n_k elements, then the total number of elements is

$$n = n_1 \cdot n_2 \cdot \dots \cdot n_k$$

Example

How many different bit strings of length 7 are there?

How many different functions are there from a set with m elements to a set with n elements?

How many **one-to-one** functions are there from a set with m elements to a set with n elements?



The Product Rule

- **Product Rule:** If a count of elements can be broken down into a **sequence of dependent counts** where the first count yields n_1 elements, the second n_2 elements, and k th count n_k elements, then the total number of elements is

$$n = n_1 \cdot n_2 \cdot \dots \cdot n_k$$

Example

How many different bit strings of length 7 are there?

How many different functions are there from a set with m elements to a set with n elements?

How many **one-to-one** functions are there from a set with m elements to a set with n elements?

How many **onto** functions?



The Product Rule

- The following loop is a part of program computing the product of two matrices.

```
(1) for i = 1 to r
(2)   for j = 1 to m
(3)     S = 0
(4)     for k = 1 to n
(5)       S = S + A[i,k] * B[k,j]
(6)     C[i,j] = S
```



The Product Rule

- The following loop is a part of program computing the product of two matrices.

```
(1) for i = 1 to r
(2)   for j = 1 to m
(3)     S = 0
(4)     for k = 1 to n
(5)       S = S + A[i,k] * B[k,j]
(6)     C[i,j] = S
```

How many multiplications (in terms of r, m, n) does this program carry out in total among all iterations of line 5?



The Sum Rule

- A count decomposes into a set of **independent** counts
(elements of counts are alternatives)



The Sum Rule

- A count decomposes into a set of **independent** counts
(elements of counts are alternatives)

Example

You need to travel from city A to B. You may either fly, take a train, or a bus. There are 12 different flights, 5 different trains and 10 buses. **How many options do you have to get from A to B?**



The Sum Rule

- A count decomposes into a set of **independent** counts
(elements of counts are alternatives)

Example

You need to travel from city A to B. You may either fly, take a train, or a bus. There are 12 different flights, 5 different trains and 10 buses. **How many options do you have to get from A to B?**

We may **use the sum rule.**

$$12 + 5 + 10$$



The Sum Rule

- **Sum Rule:** If a count of elements can be broken down into a **set of independent counts** where the first count yields n_1 elements, the second n_2 elements, and k th count n_k elements, then the total number of elements is

$$n = n_1 + n_2 + \cdots + n_k$$



The Sum Rule

- The following loop is from [selection sort](#).

```
(1) for i = 1 to n-1
(2)   for j = i+1 to n
(3)     if (A[i] > A[j])
(4)       exchange A[i] and A[j]
```



The Sum Rule

- The following loop is from **selection sort**.

```
(1) for i = 1 to n-1
(2)   for j = i+1 to n
(3)     if (A[i] > A[j])
(4)       exchange A[i] and A[j]
```

How many **comparisons** (in terms of n) does this program carry out in total among all iterations of line 3?



More Complex Counting

- Typically requires a combination of the sum and product rules.



More Complex Counting

- Typically requires a **combination** of the sum and product rules.

Example

Each password is **6 to 8 characters** long, where each character is an lowercase letter or a digit. Each password must contain **at least one digit**. How many possible passwords are there?




More Complex Counting

- Typically requires a combination of the sum and product rules.

Example

Each password is 6 to 8 characters long, where each character is an lowercase letter or a digit. Each password must contain at least one digit. How many possible passwords are there?

$$P = P_6 + P_7 + P_8$$



Inclusion-Exclusion Principle

- Used in counts where the decomposition yields two independent counting tasks with overlapping elements



Inclusion-Exclusion Principle

- Used in counts where the decomposition yields two independent counting tasks with overlapping elements

If we use the sum rule, some elements would be counted twice.



Inclusion-Exclusion Principle

- Used in counts where the decomposition yields two independent counting tasks with overlapping elements

If we use the sum rule, some elements would be counted twice.

Inclusion-Exclusion Principle: uses a sum rule and then corrects for the overlapping elements.



Inclusion-Exclusion Principle

- Used in counts where the decomposition yields two independent counting tasks with overlapping elements

If we use the sum rule, some elements would be counted twice.

Inclusion-Exclusion Principle: uses a sum rule and then corrects for the overlapping elements.

$$|A \cup B| = |A| + |B| - |A \cap B|$$



Inclusion-Exclusion Principle

■ Example

How many bit strings of length 8 either start with a '1' bit or end with the two bits '00'?



Inclusion-Exclusion Principle

■ Example

How many bit strings of length 8 either start with a '1' bit or end with the two bits '00'?

◇ it is easy to count bit strings starting with '1':



Inclusion-Exclusion Principle

■ Example

How many bit strings of length 8 either start with a '1' bit or end with the two bits '00'?

◇ it is easy to count bit strings starting with '1': 2^7



Inclusion-Exclusion Principle

■ Example

How many bit strings of length 8 either start with a '1' bit or end with the two bits '00'?

- ◇ it is easy to count bit strings starting with '1': 2^7
- ◇ it is easy to count bit strings ending with '00':



Inclusion-Exclusion Principle

■ Example

How many bit strings of length 8 either start with a '1' bit or end with the two bits '00'?

- ◇ it is easy to count bit strings starting with '1': 2^7
- ◇ it is easy to count bit strings ending with '00': 2^6



Inclusion-Exclusion Principle

■ Example

How many bit strings of length 8 either start with a '1' bit or end with the two bits '00'?

- ◇ it is easy to count bit strings starting with '1': 2^7
- ◇ it is easy to count bit strings ending with '00': 2^6

Overcounting!!!



Inclusion-Exclusion Principle

■ Example

How many bit strings of length 8 either start with a '1' bit or end with the two bits '00'?

- ◇ it is easy to count bit strings starting with '1': 2^7
- ◇ it is easy to count bit strings ending with '00': 2^6

Overcounting!!!

- ◇ deduct the number of strings starting with '1' and ending with "00":



Inclusion-Exclusion Principle

■ Example

How many bit strings of length 8 either start with a '1' bit or end with the two bits '00'?

◇ it is easy to count bit strings starting with '1': 2^7

◇ it is easy to count bit strings ending with '00': 2^6

Overcounting!!!


◇ deduct the number of strings starting with '1' and ending with "00": 2^5



Inclusion-Exclusion Principle

- Two sets


$$|E \cup F| = |E| + |F| - |E \cap F|$$




Inclusion-Exclusion Principle

■ Two sets

$$|E \cup F| = |E| + |F| - |E \cap F|$$




Three sets



Inclusion-Exclusion Principle


■ Two sets

$$|E \cup F| = |E| + |F| - |E \cap F|$$



Three sets

$$\begin{aligned} &|E \cup F \cup G| \\ &= |E| + |F| + |G| \\ &\quad - |E \cap F| - |E \cap G| - |F \cap G| \\ &\quad + |E \cap F \cap G| \end{aligned}$$



Inclusion-Exclusion Principle



$$|\cup_{i=1}^n E_i| = \sum_{k=1}^n (-1)^{k+1} \sum_{1 \leq i_1 < i_2 < \dots < i_k \leq n} |E_{i_1} \cap E_{i_2} \cap \dots \cap E_{i_k}|$$



Inclusion-Exclusion Principle



$$|\cup_{i=1}^n E_i| = \sum_{k=1}^n (-1)^{k+1} \sum_{1 \leq i_1 < i_2 < \dots < i_k \leq n} |E_{i_1} \cap E_{i_2} \cap \dots \cap E_{i_k}|$$

Proof by induction



Inclusion-Exclusion Principle



$$|\cup_{i=1}^n E_i| = \sum_{k=1}^n (-1)^{k+1} \sum_{1 \leq i_1 < i_2 < \dots < i_k \leq n} |E_{i_1} \cap E_{i_2} \cap \dots \cap E_{i_k}|$$

Proof by induction

Base case ($n = 2$)

$$|E \cup F| = |E| + |F| - |E \cap F|$$



Inclusion-Exclusion Principle



$$|\cup_{i=1}^n E_i| = \sum_{k=1}^n (-1)^{k+1} \sum_{1 \leq i_1 < i_2 < \dots < i_k \leq n} |E_{i_1} \cap E_{i_2} \cap \dots \cap E_{i_k}|$$

Proof by induction

Base case ($n = 2$)

$$|E \cup F| = |E| + |F| - |E \cap F|$$

Inductive Hypothesis

$$|\cup_{i=1}^{n-1} E_i| = \sum_{k=1}^{n-1} (-1)^{k+1} \sum_{1 \leq i_1 < i_2 < \dots < i_k \leq n-1} |E_{i_1} \cap E_{i_2} \cap \dots \cap E_{i_k}|$$



Inclusion-Exclusion Principle

- Inductive step

Set $E = E_1 \cup \cdots \cup E_{n-1}$, and $F = E_n$.



Inclusion-Exclusion Principle

- Inductive step

Set $E = E_1 \cup \cdots \cup E_{n-1}$, and $F = E_n$.

By $|E \cup F| = |E| + |F| - |E \cap F|$



Inclusion-Exclusion Principle

- Inductive step

Set $E = E_1 \cup \cdots \cup E_{n-1}$, and $F = E_n$.

By $|E \cup F| = |E| + |F| - |E \cap F|$

$$|\cup_{i=1}^n E_i| = |\cup_{i=1}^{n-1} E_i| + |E_n| - |(\cup_{i=1}^{n-1} E_i) \cap E_n|$$



Inclusion-Exclusion Principle

- Inductive step

Set $E = E_1 \cup \cdots \cup E_{n-1}$, and $F = E_n$.

By $|E \cup F| = |E| + |F| - |E \cap F|$

$$|\cup_{i=1}^n E_i| = |\cup_{i=1}^{n-1} E_i| + |E_n| - |(\cup_{i=1}^{n-1} E_i) \cap E_n|$$

The first term is given by i.h.



Inclusion-Exclusion Principle

■ Inductive step

Set $E = E_1 \cup \cdots \cup E_{n-1}$, and $F = E_n$.

By $|E \cup F| = |E| + |F| - |E \cap F|$


$$|\cup_{i=1}^n E_i| = |\cup_{i=1}^{n-1} E_i| + |E_n| - |(\cup_{i=1}^{n-1} E_i) \cap E_n|$$

The first term is given by i.h.

For the third term, by distributive law,

$$|(\cup_{i=1}^{n-1} E_i) \cap E_n| = |\cup_{i=1}^{n-1} (E_i \cap E_n)| = |\cup_{i=1}^{n-1} G_i|$$

where $G_i = E_i \cap E_n$.



Inclusion-Exclusion Principle

- So far

$$|\cup_{i=1}^n E_i| = |\cup_{i=1}^{n-1} E_i| + |E_n| - |\cup_{i=1}^{n-1} G_i|$$

where $G_i = E_i \cap E_n$.

Inclusion-Exclusion Principle

- So far

$$|\cup_{i=1}^n E_i| = |\cup_{i=1}^{n-1} E_i| + |E_n| - |\cup_{i=1}^{n-1} G_i|$$

where $G_i = E_i \cap E_n$.

Note that (why?)

$$\begin{aligned} & -(-1)^{k+1} |G_{i_1} \cap G_{i_2} \cap \cdots \cap G_{i_k}| \\ & = (-1)^{k+2} |E_{i_1} \cap E_{i_2} \cap \cdots \cap E_{i_k} \cap E_n| \end{aligned}$$

Inclusion-Exclusion Principle

- So far

$$|\cup_{i=1}^n E_i| = |\cup_{i=1}^{n-1} E_i| + |E_n| - |\cup_{i=1}^{n-1} G_i|$$

where $G_i = E_i \cap E_n$.

Note that (why?)

$$\begin{aligned} & -(-1)^{k+1} |G_{i_1} \cap G_{i_2} \cap \dots \cap G_{i_k}| \\ & = (-1)^{k+2} |E_{i_1} \cap E_{i_2} \cap \dots \cap E_{i_k} \cap E_n| \end{aligned}$$

Some discussion:

first summation sums $(-1)^{k+1} |E_{i_1} \cap E_{i_2} \cap \dots \cap E_{i_k}|$ over **all lists** i_1, i_2, \dots, i_k that **do not contain** n
 $|E_n|$ and **second summation** together sum $(-1)^{k+1} |E_{i_1} \cap E_{i_2} \cap \dots \cap E_{i_k}|$ over **all lists** i_1, i_2, \dots, i_k that **do contain** n

Inclusion-Exclusion Principle



$$|\cup_{i=1}^n E_i| = \sum_{k=1}^n (-1)^{k+1} \sum_{1 \leq i_1 < i_2 < \dots < i_k \leq n} |E_{i_1} \cap E_{i_2} \cap \dots \cap E_{i_k}|$$



Inclusion-Exclusion Principle



$$|\cup_{i=1}^n E_i| = \sum_{k=1}^n (-1)^{k+1} \sum_{1 \leq i_1 < i_2 < \dots < i_k \leq n} |E_{i_1} \cap E_{i_2} \cap \dots \cap E_{i_k}|$$

This can be used to determine the number of onto functions



Inclusion-Exclusion Principle



$$|\cup_{i=1}^n E_i| = \sum_{k=1}^n (-1)^{k+1} \sum_{1 \leq i_1 < i_2 < \dots < i_k \leq n} |E_{i_1} \cap E_{i_2} \cap \dots \cap E_{i_k}|$$

This can be used to determine the number of onto functions

A, B are two sets with $|A| = m$ and $|B| = n$.



Inclusion-Exclusion Principle



$$|\cup_{i=1}^n E_i| = \sum_{k=1}^n (-1)^{k+1} \sum_{1 \leq i_1 < i_2 < \dots < i_k \leq n} |E_{i_1} \cap E_{i_2} \cap \dots \cap E_{i_k}|$$

This can be used to determine the number of onto functions

A, B are two sets with $|A| = m$ and $|B| = n$.

(a) How many onto functions are there from A to B ?

(b) How many functions are there from A to B that map nothing to at least one element of B ?



Inclusion-Exclusion Principle



$$|\cup_{i=1}^n E_i| = \sum_{k=1}^n (-1)^{k+1} \sum_{1 \leq i_1 < i_2 < \dots < i_k \leq n} |E_{i_1} \cap E_{i_2} \cap \dots \cap E_{i_k}|$$


This can be used to determine the number of onto functions

A, B are two sets with $|A| = m$ and $|B| = n$.

(a) How many onto functions are there from A to B ?

(b) How many functions are there from A to B that map nothing to at least one element of B ?

$$\#(a) + \#(b) = n^m$$



Inclusion-Exclusion Principle

- This can be used to determine the number of onto functions

A, B are two sets with $|A| = m$ and $|B| = n$.

(a) How many onto functions are there from A to B ?

(b) How many functions are there from A to B that map nothing to at least one element of B ?

$$\#(a) + \#(b) = n^m$$



Inclusion-Exclusion Principle

- This can be used to determine the number of onto functions

A, B are two sets with $|A| = m$ and $|B| = n$.

(a) How many onto functions are there from A to B ?

(b) How many functions are there from A to B that map nothing to at least one element of B ?

$$\#(a) + \#(b) = n^m$$

Set E_i – set of functions that map nothing to element i of B



Inclusion-Exclusion Principle

- This can be used to determine the number of onto functions

A, B are two sets with $|A| = m$ and $|B| = n$.


(a) How many onto functions are there from A to B ?

(b) How many functions are there from A to B that map nothing to at least one element of B ?

$$\#(a) + \#(b) = n^m$$

Set E_i – set of functions that map nothing to element i of B

$$\#(b) = \left| \bigcup_{i=1}^n E_i \right|$$



Inclusion-Exclusion Principle

- This can be used to determine the number of onto functions

A, B are two sets with $|A| = m$ and $|B| = n$.

(a) How many onto functions are there from A to B ?


(b) How many functions are there from A to B that map nothing to at least one element of B ?

$$\#(a) + \#(b) = n^m$$

Set E_i – set of functions that map nothing to element i of B

$$\#(b) = \left| \bigcup_{i=1}^n E_i \right|$$

$$= \sum_{k=1}^n (-1)^{k+1} \sum_{1 \leq i_1 < i_2 < \dots < i_k \leq n} |E_{i_1} \cap E_{i_2} \cap \dots \cap E_{i_k}|$$



Inclusion-Exclusion Principle

- This can be used to determine the number of onto functions

A, B are two sets with $|A| = m$ and $|B| = n$.

(a) How many onto functions are there from A to B ?

(b) How many functions are there from A to B that map nothing to at least one element of B ?


$$\#(a) + \#(b) = n^m$$

Set E_i – set of functions that map nothing to element i of B

$$\#(b) = \left| \bigcup_{i=1}^n E_i \right|$$

$$= \sum_{k=1}^n (-1)^{k+1} \sum_{1 \leq i_1 < i_2 < \dots < i_k \leq n} |E_{i_1} \cap E_{i_2} \cap \dots \cap E_{i_k}|$$

$$= \sum_{k=1}^n (-1)^{k+1} \binom{n}{k} (n-k)^m$$



Tree Diagrams

- A *tree* is a structure that consists of a *root*, *branches* and *leaves*.



Tree Diagrams

- A *tree* is a structure that consists of a *root*, *branches* and *leaves*.

Can be useful to represent a counting problem and record the choices we made for alternatives. *The count appears on the leaves.*



Tree Diagrams

- A *tree* is a structure that consists of a **root**, **branches** and **leaves**.

Can be useful to represent a counting problem and record the choices we made for alternatives. **The count appears on the leaves.**

Example

What is the number of bit strings of length 4 that **do not have two consecutive 1's**?



Tree Diagrams


- A *tree* is a structure that consists of a *root*, *branches* and *leaves*.

Can be useful to track the choices with the leaves.

Problem and record count appears on

Example

What is the probability of having two consecutive 1s in a 4-bit sequence?



h 4 that do not


Tree Diagram

- How many different ways can a “best 3 of 5” playoff occur?



Tree Diagram

- How many different ways can a “best 3 of 5” playoff occur?



Pigeonhole Principle

- Assume that there are a set of objects and a set of bins to store them.



Pigeonhole Principle

- Assume that there are a set of objects and a set of bins to store them.

The pigeonhole principle states that if there are more objects than bins then there is at least one bin with more than one object.




Pigeonhole Principle

- Assume that there are a set of objects and a set of bins to store them.

The pigeonhole principle states that if there are more objects than bins then there is at least one bin with more than one object.

Example: 7 balls and 5 bins to store them




Pigeonhole Principle

- Assume that there are a set of objects and a set of bins to store them.

The pigeonhole principle states that if there are more objects than bins then there is at least one bin with more than one object.

Example: 7 balls and 5 bins to store them



Pigeonhole Principle

- **Theorem** If there are $k + 1$ objects and k bins, then there is at least one bin with two or more objects.



Pigeonhole Principle

- **Theorem** If there are $k + 1$ objects and k bins, then there is at least one bin with two or more objects.

Proof by contradiction



Pigeonhole Principle

- **Theorem** If there are $k + 1$ objects and k bins, then there is **at least** one bin with two or more objects.

Proof by contradiction

Example

Assume that there are 367 students. Are there any two people who has the same birthday?

There are 5 bins and 12 objects. Then there must be a bin with at least 3 objects. Why?



Generalized Pigeonhole Principle

- If N objects are placed into k bins, then there is at least one bin containing at least $\lceil N/k \rceil$ objects.



Generalized Pigeonhole Principle

- If N objects are placed into k bins, then there is at least one bin containing at least $\lceil N/k \rceil$ objects.

Example

Assume there are 100 students. How many of them were born in the same month?



Bijections and Permutations

- A function that is **both one-to-one and onto** is called a *bijection*, or a *one-to-one correspondence*.



Bijections and Permutations

- A function that is **both one-to-one and onto** is called a *bijection*, or a *one-to-one correspondence*.

How many **bijections** are there?




Bijections and Permutations

- A function that is **both one-to-one and onto** is called a *bijection*, or a *one-to-one correspondence*.

How many **bijections** are there?

$f : \{a, b, c\} \rightarrow \{1, 2, 3\}$ defined by $f(a) = 3, f(b) = 2, f(c) = 1$ is a bijection.




Bijections and Permutations

- A function that is **both one-to-one and onto** is called a *bijection*, or a *one-to-one correspondence*.

How many **bijections** are there?

$f : \{a, b, c\} \rightarrow \{1, 2, 3\}$ defined by $f(a) = 3, f(b) = 2, f(c) = 1$ is a bijection.




A **bijection** from a set **onto itself** is called a *permutation*.

Bijections and Permutations

- A function that is **both one-to-one and onto** is called a *bijection*, or a *one-to-one correspondence*.


How many **bijections** are there?

$f : \{a, b, c\} \rightarrow \{1, 2, 3\}$ defined by $f(a) = 3, f(b) = 2, f(c) = 1$ is a bijection.



A **bijection** from a set **onto itself** is called a *permutation*.

$f : \{1, 2, 3\} \rightarrow \{1, 2, 3\}$ defined by $f(1) = 3, f(2) = 2, f(3) = 1$ is a bijection.





Bijections and Permutations

- A function that is **both one-to-one and onto** is called a *bijection*, or a *one-to-one correspondence*.

A **bijection** from a set **onto itself** is called a *permutation*.

In a *bijection*,

exactly one arrow leaves each item on the left and exactly one arrow arrives at each item on the right.



Bijections and Permutations

- A function that is **both one-to-one and onto** is called a *bijection*, or a *one-to-one correspondence*.



A **bijection** from a set **onto itself** is called a *permutation*.

In a *bijection*,

exactly one arrow leaves each item on the left and exactly one arrow arrives at each item on the right.

Thus,

the left and right sides must have the same size.



The Bijection Principle

- The following loop is a part of program to determine the number of triangles formed by n points in the plane.

```
(1) trianglecount = 0
(2)   for i = 1 to n
(3)     for j = i+1 to n
(4)       for k = j+1 to n
(5)         if points i, j, k are not collinear
(6)           trianglecount = trianglecount + 1
```



The Bijection Principle

- The following loop is a part of program to determine the number of triangles formed by n points in the plane.

```
(1) trianglecount = 0
(2)   for i = 1 to n
(3)     for j = i+1 to n
(4)       for k = j+1 to n
(5)         if points i, j, k are not collinear
(6)           trianglecount = trianglecount + 1
```

Among all iterations of line 5, what is the total number of times this line checks three points to see if they are collinear?




Counting Triangles

- 3 points form a triangle if and only if they are non collinear




Counting Triangles

- 3 points form a triangle if and only if they are non collinear



Counting Triangles


- 3 points form a triangle if and only if they are non collinear



1 – 2 – 7: yes

Counting Triangles

- 3 points form a triangle if and only if they are non collinear




1 – 2 – 7: yes

1 – 2 – 5: yes

Counting Triangles

- 3 points form a triangle if and only if they are non collinear




1 – 2 – 7: yes

1 – 2 – 5: yes

1 – 2 – 3: no

Counting Triangles

- 3 points form a triangle if and only if they are non collinear



1 – 2 – 7: yes


1 – 2 – 5: yes

1 – 2 – 3: no

1 – 5 – 6: yes

Counting Triangles

- 3 points form a triangle if and only if they are non collinear



1 – 2 – 7: yes

1 – 2 – 5: yes


1 – 2 – 3: no

1 – 5 – 6: yes

3 – 4 – 7: yes

Counting Triangles

- 3 points form a **triangle** if and only if **they are non collinear**



1 – 2 – 7: yes

1 – 2 – 5: yes

1 – 2 – 3: **no**


1 – 5 – 6: yes

3 – 4 – 7: yes

4 – 5 – 6: **no**

Counting Triangles

- 3 points form a triangle if and only if they are non collinear



1 – 2 – 7: yes

1 – 2 – 5: yes

1 – 2 – 3: no

1 – 5 – 6: yes

3 – 4 – 7: yes

4 – 5 – 6: no

number of triangles: 33

Counting Triangles

```
(1) trianglecount = 0
(2)   for i = 1 to n
(3)       for j = i+1 to n
(4)           for k = j+1 to n
(5)               if points i, j, k are not collinear
(6)                   trianglecount = trianglecount + 1
```


Counting Triangles

```
(1) trianglecount = 0
(2)   for i = 1 to n
(3)     for j = i+1 to n
(4)       for k = j+1 to n
(5)         if points i, j, k are not collinear
(6)           trianglecount = trianglecount + 1
```

A loop

Counting Triangles

(1) trianglecount = 0

(2) for i = 1 to n

(3) for j = i+1 to n

(4) for k = j+1 to n

(5) if points i, j, k are not collinear

(6) trianglecount = trianglecount + 1

A loop embedded in a loop

Counting Triangles

(1) trianglecount = 0

(2) for *i* = 1 to *n*

(3) for *j* = *i*+1 to *n*

(4) for *k* = *j*+1 to *n*

(5) if points *i*, *j*, *k* are not collinear

(6) trianglecount = trianglecount + 1

A loop embedded in a loop embedded in another loop.

Counting Triangles

```
(1) trianglecount = 0
```

```
(2)   for  $i = 1$  to  $n$ 
```

```
(3)       for  $j = i+1$  to  $n$ 
```

```
(4)           for  $k = j+1$  to  $n$ 
```

```
(5)               if points  $i, j, k$  are not collinear
```

```
(6)                   trianglecount = trianglecount + 1
```

A loop embedded in a loop embedded in another loop.

Second loop begins with $j = i + 1$ and j increases up to n .

Third loop begins with $k = j + 1$ and k increases up to n .

Counting Triangles

```
(1) trianglecount = 0
```

```
(2)   for  $i = 1$  to  $n$ 
```

```
(3)       for  $j = i+1$  to  $n$ 
```

```
(4)           for  $k = j+1$  to  $n$ 
```

```
(5)               if points  $i, j, k$  are not collinear
```

```
(6)                   trianglecount = trianglecount + 1
```

A loop embedded in a loop embedded in another loop.

Second loop begins with $j = i + 1$ and j increases up to n .

Third loop begins with $k = j + 1$ and k increases up to n .

Thus each triple i, j, k with $i < j < k$ is examined exactly once.

Counting Triangles

```
(1) trianglecount = 0
(2)   for i = 1 to n
(3)     for j = i+1 to n
(4)       for k = j+1 to n
(5)         if points i, j, k are not collinear
(6)           trianglecount = trianglecount + 1
```

A loop embedded in a loop embedded in another loop.

Second loop begins with $j = i + 1$ and j increases up to n .

Third loop begins with $k = j + 1$ and k increases up to n .

Thus each triple i, j, k with $i < j < k$ is examined exactly once.

For example, if $n = 4$, then triples (i, j, k) used by algorithm are $(1, 2, 3)$, $(1, 2, 4)$, $(1, 3, 4)$, and $(2, 3, 4)$.

Counting Triangles

- Want to compute the number of *increasing triples* (i, j, k) with $1 \leq i < j < k \leq n$.

Counting Triangles

- Want to compute the number of *increasing triples* (i, j, k) with $1 \leq i < j < k \leq n$.

Claim: Number of increasing triples is **exactly** the same as number of 3-element subsets from $\{1, 2, \dots, n\}$

Counting Triangles

- Want to compute the number of *increasing triples* (i, j, k) with $1 \leq i < j < k \leq n$.

Claim: Number of increasing triples is **exactly** the same as number of 3-element subsets from $\{1, 2, \dots, n\}$

Why? Let $X =$ set of increasing triples and
 $Y =$ set of 3-element subsets from $\{1, 2, \dots, n\}$

Counting Triangles

- Want to compute the number of *increasing triples* (i, j, k) with $1 \leq i < j < k \leq n$.

Claim: Number of increasing triples is **exactly** the same as number of 3-element subsets from $\{1, 2, \dots, n\}$

Why? Let $X =$ set of increasing triples and
 $Y =$ set of 3-element subsets from $\{1, 2, \dots, n\}$

Define: $f : X \rightarrow Y$ by $f((i, j, k)) = \{i, j, k\}$

Claim: f is a **bijection** (why) so $|X| = |Y|$

Counting Triangles

- Want to compute the number of *increasing triples* (i, j, k) with $1 \leq i < j < k \leq n$.

Claim: Number of increasing triples is **exactly** the same as number of 3-element subsets from $\{1, 2, \dots, n\}$

Why? Let X = set of increasing triples and
 Y = set of 3-element subsets from $\{1, 2, \dots, n\}$

Define: $f : X \rightarrow Y$ by $f((i, j, k)) = \{i, j, k\}$

Claim: f is a **bijection** (why) so $|X| = |Y|$

f is a bijection because

f is one-to-one

if $(i, j, k) \neq (i', j', k') \Rightarrow f((i, j, k)) \neq f((i', j', k'))$

f is onto

if γ is a 3-element subset then it can be written as $\gamma = \{i, j, k\}$
where $i < j < k$ so $f((i, j, k)) = \gamma$.

Counting Pairs

- We've already seen something very similar.
The number of
increasing pairs (i, j) with $1 \leq i < j \leq n$
is the same as the number of
2-sets from $\{1, 2, \dots, n\}$



Counting Pairs

- We've already seen something very similar.
The number of
increasing pairs (i, j) with $1 \leq i < j \leq n$
is the same as the number of
2-sets from $\{1, 2, \dots, n\}$

Define $f : X \rightarrow Y$ by $f((i, j)) = \{i, j\}$

Claim: f is a **bijection** so $|X| = |Y|$



Counting Pairs

- We've already seen something very similar.
The number of
increasing pairs (i, j) with $1 \leq i < j \leq n$
is the same as the number of
2-sets from $\{1, 2, \dots, n\}$

Define $f : X \rightarrow Y$ by $f((i, j)) = \{i, j\}$

Claim: f is a **bijection** so $|X| = |Y|$

We actually already saw that $|X| = |Y| = \binom{n}{2}$



The Bijection Principle

- Two sets have the same size if and only if there is a one-to-one function from one set onto the other.



The Bijection Principle

- Two sets **have the same size** if and only if there is a **one-to-one function from one set onto the other**.

A standard first step in counting the size of a set is to use a bijection to show that it has the same size as a 2nd set, and then count the 2nd set instead.



The Bijection Principle

- Two sets **have the same size** if and only if there is a **one-to-one function from one set onto the other**.

A standard first step in counting the size of a set is to use a bijection to show that it has the same size as a 2nd set, and then count the 2nd set instead.

In practice, in real problems we often only *implicitly* use the bijection and don't *explicitly* describe it



The Bijection Principle

- Two sets **have the same size** if and only if there is a **one-to-one function from one set onto the other**.

A standard first step in counting the size of a set is to use a bijection to show that it has the same size as a 2nd set, and then count the 2nd set instead.

In practice, in real problems we often only *implicitly* use the bijection and don't *explicitly* describe it

Currently, we started with the problem of counting the **# of increasing triples** and changed it to the problem of counting the **# of 3-element sets from $\{1, 2, \dots, n\}$**



k -Element Permutations of a Set

- In how many ways can we choose **an ordered triple** of distinct elements from $\{1, 2, \dots, n\}$?



k -Element Permutations of a Set

- In how many ways can we choose **an ordered triple** of distinct elements from $\{1, 2, \dots, n\}$?

More generally, in how many ways can we choose **a list of k distinct elements** from $\{1, 2, \dots, n\}$?



k -Element Permutations of a Set

- In how many ways can we choose **an ordered triple** of distinct elements from $\{1, 2, \dots, n\}$?

More generally, in how many ways can we choose **a list of k distinct elements** from $\{1, 2, \dots, n\}$?

A list of k ***distinct*** elements chosen from a set N is called a **k -element permutation of N**



k -Element Permutations of a Set

- In how many ways can we choose **an ordered triple** of distinct elements from $\{1, 2, \dots, n\}$?

More generally, in how many ways can we choose **a list of k distinct elements** from $\{1, 2, \dots, n\}$?

A list of k ***distinct*** elements chosen from a set N is called a **k -element permutation of N**

Note that the case of $k = n$ is special;

An **n -element permutation** of a **set N** of size $|N| = n$ is what we earlier simply called a **permutation**.



k -Element Permutations of a Set

- How many three-element permutations of $\{1, 2, \dots, n\}$ are there?



k -Element Permutations of a Set

- How many three-element permutations of $\{1, 2, \dots, n\}$ are there?

n choices for first number



k -Element Permutations of a Set

- How many three-element permutations of $\{1, 2, \dots, n\}$ are there?

n choices for first number

For each way of choosing first number there are $n - 1$ choices for the second



k -Element Permutations of a Set

- How many three-element permutations of $\{1, 2, \dots, n\}$ are there?

n choices for first number

For each way of choosing first number there are $n - 1$ choices for the second

For each way of choosing first two numbers, there are $n - 2$ choices for the third number



k -Element Permutations of a Set

- How many three-element permutations of $\{1, 2, \dots, n\}$ are there?

n choices for first number

For each way of choosing first number there are $n - 1$ choices for the second

For each way of choosing first two numbers, there are $n - 2$ choices for the third number

By product principle, there are $n(n - 1)(n - 2)$ ways to choose the permutation



An Example

- By **product principle**, there are $n(n-1)(n-2)$ ways to choose the permutation



An Example

- By **product principle**, there are $n(n-1)(n-2)$ ways to choose the permutation

Ex: When $n = 4$, there are $4 \times 3 \times 2 = 24$
3 -element permutations of $\{1, 2, 3, 4\}$

$$L = \{123, 124, 132, 134, 142, 143, 213, 214, 231, 234, 241, 243, 312, 314, 321, 324, 341, 342, 412, 413, 421, 423, 431, 432\}.$$



An Example

- By **product principle**, there are $n(n-1)(n-2)$ ways to choose the permutation

Ex: When $n = 4$, there are $4 \times 3 \times 2 = 24$
3 -element permutations of $\{1, 2, 3, 4\}$

$L = \{123, 124, 132, 134, 142, 143, 213, 214, 231, 234, 241, 243, 312, 314, 321, 324, 341, 342, 412, 413, 421, 423, 431, 432\}.$

Note: This type of "dictionary" ordering of tuples (assuming that we treat numbers the same as letters) is called a **lexicographic ordering** and is used quite often.



k -Element Permutations of a Set

- **Theorem** If N is a positive integer and k is an integer with $1 \leq k \leq n$, then there are

$$P(n, k) = n(n-1)(n-2) \cdots (n-k+1)$$

k -element permutations with n distinct elements.



k -Element Permutations of a Set

- **Theorem** If N is a positive integer and k is an integer with $1 \leq k \leq n$, then there are
 $P(n, k) = n(n-1)(n-2) \cdots (n-k+1)$
 k -element permutations with n distinct elements.

How does this help us solve our original problem(from triangle program) of counting # of 3-element subsets?



k -Element Permutations of a Set

- **Theorem** If N is a positive integer and k is an integer with $1 \leq k \leq n$, then there are
 $P(n, k) = n(n-1)(n-2) \cdots (n-k+1)$
 k -element permutations with n distinct elements.

How does this help us solve our original problem (from triangle program) of counting # of 3-element subsets?

Note that every 3-element subset $\{i, j, k\}$ can be made into exactly 6 3-element perms



k -Element Permutations of a Set

- **Theorem** If N is a positive integer and k is an integer with $1 \leq k \leq n$, then there are
 $P(n, k) = n(n-1)(n-2) \cdots (n-k+1)$
 k -element permutations with n distinct elements.

How does this help us solve our original problem (from triangle program) of counting # of 3-element subsets?

Note that every 3-element subset $\{i, j, k\}$ can be made into exactly 6 3-element perms

$$(\# \text{ 3-element perms}) = 6 \times (\# \text{ 3-element subsets})$$



k -Element Permutations of a Set

- **Theorem** If N is a positive integer and k is an integer with $1 \leq k \leq n$, then there are
 $P(n, k) = n(n-1)(n-2) \cdots (n-k+1)$
 k -element permutations with n distinct elements.

How does this help us solve our original problem (from triangle program) of counting # of 3-element subsets?

Note that every 3-element subset $\{i, j, k\}$ can be made into exactly 6 3-element perms

$$(\# \text{ 3-element perms}) = 6 \times (\# \text{ 3-element subsets})$$

$$P(n, 3) = 3! \cdot C(n, 3)$$



Binomial Coefficient

- **Theorem** For integers n and k with $0 \leq k \leq n$, the number of k -element subsets of an n -element set is

$$\binom{n}{k} = C(n, k) = \frac{P(n, k)}{k!} = \frac{n!}{k!(n - k)!}.$$

This is the number of k -combinations of a set with n elements.



Some Properties of Binomial Coefficients

- $\binom{n}{k} = \frac{n!}{k!(n-k)!}$ is the number of k -element subsets of an n -element set.

$$\binom{n}{0} = 1 \text{ only one set of size } 0.$$

$$\binom{n}{n} = 1 \text{ only one set of size } n.$$

$\binom{n}{k} = \binom{n}{n-k}$ Obvious from equation. Can you think of a simple bijection that explains this?



Some Properties of Binomial Coefficients (cont.)

$$\sum_{i=0}^n \binom{n}{i} = 2^n$$



Some Properties of Binomial Coefficients (cont.)

$$\sum_{i=0}^n \binom{n}{i} = 2^n$$

Use Sum Rule

Let P = set of all subsets of $\{1, 2, \dots, n\}$

S_i = set of all i subsets of $\{1, 2, \dots, n\}$



Some Properties of Binomial Coefficients (cont.)

$$\sum_{i=0}^n \binom{n}{i} = 2^n$$

Use Sum Rule

Let P = set of all subsets of $\{1, 2, \dots, n\}$

S_i = set of all i subsets of $\{1, 2, \dots, n\}$

$$\Rightarrow |P| = \sum_{i=0}^n |S_i| = \sum_{i=0}^n \binom{n}{i}$$



Some Properties of Binomial Coefficients (cont.)

■ Let $L = L_1 L_2 \dots L_n$ be a list of size n from $\{0, 1\}$

If \mathcal{L} = set of all such lists $\Rightarrow |\mathcal{L}| = 2^n$

There is a *bijection* between \mathcal{L} and P so
 $|P| = 2^n$ and we are done.

Some Properties of Binomial Coefficients (cont.)

■ Let $L = L_1 L_2 \dots L_n$ be a list of size n from $\{0, 1\}$

If \mathcal{L} = set of all such lists $\Rightarrow |\mathcal{L}| = 2^n$

There is a *bijection* between \mathcal{L} and P so
 $|P| = 2^n$ and we are done.

Define the following function $f : \mathcal{L} \rightarrow P$

If $L \in \mathcal{L}$ then $f(L)$ is the set $S \subseteq \{1, 2, \dots, n\}$ defined by

$$i \in S \Leftrightarrow L_i = 1$$

Some Properties of Binomial Coefficients (cont.)

■ Let $L = L_1 L_2 \dots L_n$ be a list of size n from $\{0, 1\}$

If \mathcal{L} = set of all such lists $\Rightarrow |\mathcal{L}| = 2^n$

There is a *bijection* between \mathcal{L} and P so
 $|P| = 2^n$ and we are done.

Define the following function $f : \mathcal{L} \rightarrow P$

If $L \in \mathcal{L}$ then $f(L)$ is the set $S \subseteq \{1, 2, \dots, n\}$ defined by

$$i \in S \Leftrightarrow L_i = 1$$

f is a *bijection* between \mathcal{L} and P (why?) so $|\mathcal{L}| = |P|$

Some Properties of Binomial Coefficients (cont.)

■ Let $L = L_1 L_2 \dots L_n$ be a list of size n from $\{0, 1\}$

If \mathcal{L} = set of all such lists $\Rightarrow |\mathcal{L}| = 2^n$

There is a *bijection* between \mathcal{L} and P so
 $|P| = 2^n$ and we are done.

Define the following function $f : \mathcal{L} \rightarrow P$

If $L \in \mathcal{L}$ then $f(L)$ is the set $S \subseteq \{1, 2, \dots, n\}$ defined by

$$i \in S \Leftrightarrow L_i = 1$$

f is a *bijection* between \mathcal{L} and P (why?) so $|\mathcal{L}| = |P|$

Ex: $n = 5$

$$f(10101) = \{1, 3, 5\}, \quad f(11101) = \{1, 2, 3, 5\}, \quad f(00000) = \emptyset$$

Binomial Coefficients

$n \backslash k$	0	1	2	3	4	5	6
0	1						
1	1	1					
2	1	2	1				
3	1	3	3	1			
4	1	4	6	4	1		
5	1	5	10	10	5	1	
6	1	6	15	20	15	6	1



Binomial Coefficients

$n \backslash k$	0	1	2	3	4	5	6
0	1						
1	1	1					
2	1	2	1				
3	1	3	3	1			
4	1	4	6	4	1		
5	1	5	10	10	5	1	
6	1	6	15	20	15	6	1

Each row begins with a 1
because $\binom{n}{0} = 1$



Binomial Coefficients

$n \backslash k$	0	1	2	3	4	5	6
0	1						
1	1	1					
2	1	2	1				
3	1	3	3	1			
4	1	4	6	4	1		
5	1	5	10	10	5	1	
6	1	6	15	20	15	6	1

Each row begins with a 1
because $\binom{n}{0} = 1$

Each row ends with a 1
because $\binom{n}{n} = 1$.



Binomial Coefficients

$n \backslash k$	0	1	2	3	4	5	6
0	1						
1	1	1					
2	1	2	1				
3	1	3	3	1			
4	1	4	6	4	1		
5	1	5	10	10	5	1	
6	1	6	15	20	15	6	1

Each row begins with a 1
because $\binom{n}{0} = 1$

Each row ends with a 1
because $\binom{n}{n} = 1$.

Each row increases at first
then decreases.



Binomial Coefficients

$n \backslash k$	0	1	2	3	4	5	6
0	1						
1	1	1					
2	1	2	1				
3	1	3	3	1			
4	1	4	6	4	1		
5	1	5	10	10	5	1	
6	1	6	15	20	15	6	1

Each row begins with a 1
because $\binom{n}{0} = 1$

Each row ends with a 1
because $\binom{n}{n} = 1$.

Each row increases at first
then decreases.

Second half of each row is the reverse of the first half.



Binomial Coefficients

$n \backslash k$	0	1	2	3	4	5	6
0	1						
1	1	1					
2	1	2	1				
3	1	3	3	1			
4	1	4	6	4	1		
5	1	5	10	10	5	1	
6	1	6	15	20	15	6	1

Each row begins with a 1
because $\binom{n}{0} = 1$

Each row ends with a 1
because $\binom{n}{n} = 1$.

Each row increases at first
then decreases.

Second half of each row is the reverse of the first half.
Sum of items on n -th row is 2^n



Pascal's Triangle

Take the table

$n \backslash k$	0	1	2	3	4	5	6
0	1						
1	1	1					
2	1	2	1				
3	1	3	3	1			
4	1	4	6	4	1		
5	1	5	10	10	5	1	
6	1	6	15	20	15	6	1



Pascal's Triangle

Take the table

$n \backslash k$	0	1	2	3	4	5	6
0	1						
1	1	1					
2	1	2	1				
3	1	3	3	1			
4	1	4	6	4	1		
5	1	5	10	10	5	1	
6	1	6	15	20	15	6	1

and shift each row slightly
so that middle element is
in middle

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--



Pascal's Triangle

				1				
			1		1			
		1		2		1		
	1		3		3		1	
	1	4		6		4	1	
1		5	10		10	5		1
1	6	15	20	15	6		1	



Pascal's Triangle

				1			
			1		1		
		1		2		1	
	1		3		3		1
	1	4		6		4	1
1		5	10		10	5	1
1	6	15	20	15	6	1	

What is the next row in the table?



Pascal's Triangle

				1				
			1		1			
		1		2		1		
	1		3		3		1	
	1	4		6		4	1	
	1	5	10		10	5	1	
	1	6	15	20		15	6	1
1	7	21	35	35	21	7	1	



Pascal's Triangle

				1				
			1		1			
		1		2		1		
	1		3		3		1	
	1	4		6		4	1	
	1	5	10		10	5	1	
	1	6	15	20		15	6	1
1	7	21	35	35	21	7	1	

Pascal identity

Each (non-1) entry in Pascal's Triangle is the sum of

the two entries directly above it (to left and to right).



Pascal's Triangle

				1				
			1		1			
		1		2		1		
	1		3		3		1	
	1	4		6		4		1
	1	5	10		10	5		1
	1	6	15	20		15	6	1
1	7	21	35	35	21	7		1

$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$$

Pascal identity

Each (non-1) entry in Pascal's Triangle is the sum of the two entries directly above it (to left and to right).



Pascal's Identity



$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$$



Pascal's Identity



$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$$

A purely *algebraic* proof (manipulating formulas) is possible.



Pascal's Identity



$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$$

A purely *algebraic* proof (manipulating formulas) is possible.

We will use a *combinatorial proof*.



A Combinatorial Proof

- $\binom{n}{k}$ is the number of k -element subsets of an n -element set.



A Combinatorial Proof

- $\binom{n}{k}$ is the number of k -element subsets of an n -element set.

$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$$

Therefore, each term (left and right) represents the number of subsets of a particular size chosen from an appropriately sized set.



A Combinatorial Proof

■

$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$$



A Combinatorial Proof

■

$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$$

Number of k -subsets of an n -element set.




A Combinatorial Proof



$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$$

Number of k -subsets of an n -element set.

Number of $(k-1)$ -subsets of an $(n-1)$ -element set.



A Combinatorial Proof



$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$$

Number of k -subsets of an n -element set.

Number of $(k-1)$ -subsets of an $(n-1)$ -element set.

Number of k -subsets of an $(n-1)$ -element set.



A Combinatorial Proof



$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$$

Number of k -subsets of an n -element set.

Number of $(k-1)$ -subsets of an $(n-1)$ -element set.

Number of k -subsets of an $(n-1)$ -element set.

Try to use sum principle to explain relationship among these three terms.

Example: $n = 5, k = 2$

$$\binom{5}{2} = \binom{4}{1} + \binom{4}{2}.$$



A Combinatorial Proof

$$\binom{5}{2} = \binom{4}{1} + \binom{4}{2}.$$



A Combinatorial Proof

$$\binom{5}{2} = \binom{4}{1} + \binom{4}{2}.$$

Consider $S = \{A, B, C, D, E\}$.




A Combinatorial Proof

$$\binom{5}{2} = \binom{4}{1} + \binom{4}{2}.$$

Consider $S = \{A, B, C, D, E\}$.

Set S_1 of 2-subsets of S

$$S_1 = \{\{A, B\}, \{A, C\}, \{A, D\}, \{A, E\}, \{B, C\}, \\ \{B, D\}, \{B, E\}, \{C, D\}, \{C, E\}, \{D, E\}\}.$$



A Combinatorial Proof

$$\binom{5}{2} = \binom{4}{1} + \binom{4}{2}.$$

Consider $S = \{A, B, C, D, E\}$.

Set S_1 of 2-subsets of S can be partitioned into 2 disjoint parts.

S_2 the 2-subsets that contain E and

S_3 , the set of 2-subsets that do not contain E .

$$S_1 = \{\{A, B\}, \{A, C\}, \{A, D\}, \{A, E\}, \{B, C\}, \\ \{B, D\}, \{B, E\}, \{C, D\}, \{C, E\}, \{D, E\}\}.$$



A Combinatorial Proof

$$\binom{5}{2} = \binom{4}{1} + \binom{4}{2}.$$

Consider $S = \{A, B, C, D, E\}$.

Set S_1 of 2-subsets of S can be partitioned into 2 disjoint parts.

S_2 the 2-subsets that contain E and

S_3 , the set of 2-subsets that do not contain E .

$$S_1 = \{\{A, B\}, \{A, C\}, \{A, D\}, \{A, E\}, \{B, C\}, \\ \{B, D\}, \{B, E\}, \{C, D\}, \{C, E\}, \{D, E\}\}.$$

$$S_1 = \{\{A, B\}, \{A, C\}, \{A, D\}, \{A, E\}, \{B, C\}, \\ \{B, D\}, \{B, E\}, \{C, D\}, \{C, E\}, \{D, E\}\}.$$



A Combinatorial Proof

- If n and k are integers satisfying $0 < k < n$, then

$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}.$$




A Combinatorial Proof

- If n and k are integers satisfying $0 < k < n$, then

$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}.$$

Proof: Apply **sum rule**.



A Combinatorial Proof

- If n and k are integers satisfying $0 < k < n$, then

$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}.$$

Proof: Apply **sum rule**.

Let S_1 be set of all k -element subsets.



A Combinatorial Proof

- If n and k are integers satisfying $0 < k < n$, then

$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}.$$

Proof: Apply **sum rule**.

Let S_1 be set of all k -element subsets.

To apply **sum rule**, partition S_1 into S_2 and S_3 .

Let S_2 be set of k -element subsets that **contain** x_n .

Let S_3 be set of k -element subsets that **don't contain** x_n .



Blaise Pascal

Born 1623; Died 1662

French Mathematician

A Founder of Probability Theory

Inventor of one of the first mechanical
calculating machines

Pascal Programming Language named for him



The Binomial Theorem

$$(x + y) = \binom{1}{0}x + \binom{1}{1}y$$



The Binomial Theorem

$$(x + y) = \binom{1}{0}x + \binom{1}{1}y$$

$$(x + y)^2 = x^2 + 2xy + y^2 = \binom{2}{0}x^2 + \binom{2}{1}x^1y^1 + \binom{2}{2}y^2$$



The Binomial Theorem

$$(x + y) = \binom{1}{0}x + \binom{1}{1}y$$

$$(x + y)^2 = x^2 + 2xy + y^2 = \binom{2}{0}x^2 + \binom{2}{1}x^1y^1 + \binom{2}{2}y^2$$

$$\begin{aligned}(x + y)^3 &= x^3 + 3x^2y + 3xy^2 + y^3 \\ &= \binom{3}{0}x^3 + \binom{3}{1}x^2y + \binom{3}{2}xy^2 + \binom{3}{3}y^3\end{aligned}$$



The Binomial Theorem

- Number of k -element subsets of an n -element set is called a **binomial coefficient** because of its role in the algebraic expansion of a binomial $(x + y)^n$.




The Binomial Theorem

- Number of k -element subsets of an n -element set is called a **binomial coefficient** because of its role in the algebraic expansion of a binomial $(x + y)^n$.

The Binomial Theorem For any integer $n \geq 0$,

$$(x+y)^n = \binom{n}{0}x^n + \binom{n}{1}x^{n-1}y + \binom{n}{2}x^{n-2}y^2 + \dots + \binom{n}{n-1}xy^{n-1} + \binom{n}{n}y^n$$



The Binomial Theorem

- Number of k -element subsets of an n -element set is called a **binomial coefficient** because of its role in the algebraic expansion of a binomial $(x + y)^n$.

The Binomial Theorem For any integer $n \geq 0$,

$$(x+y)^n = \binom{n}{0}x^n + \binom{n}{1}x^{n-1}y + \binom{n}{2}x^{n-2}y^2 + \dots + \binom{n}{n-1}xy^{n-1} + \binom{n}{n}y^n$$

$$(x+y)^n = \sum_{i=0}^n \binom{n}{i} x^{n-i} y^i.$$



The Binomial Theorem

- Number of k -element subsets of an n -element set is called a **binomial coefficient** because of its role in the algebraic expansion of a binomial $(x + y)^n$.

The Binomial Theorem For any integer $n \geq 0$,

$$(x+y)^n = \binom{n}{0}x^n + \binom{n}{1}x^{n-1}y + \binom{n}{2}x^{n-2}y^2 + \dots + \binom{n}{n-1}xy^{n-1} + \binom{n}{n}y^n$$

$$(x+y)^n = \sum_{i=0}^n \binom{n}{i} x^{n-i} y^i.$$

Proof?



Application of the Binomial Theorem

- We may use the Binomial Theorem to prove

$$\sum_{i=0}^n \binom{n}{i} = 2^n$$



Labelling and Trinomial Coefficients


- Suppose we have k labels of one kind, e.g., red and $n - k$ labels of another, e.g., blue. In how many different ways can we apply these labels to n objects?



Labelling and Trinomial Coefficients

- Suppose we have k labels of one kind, e.g., red and $n - k$ labels of another, e.g., blue. In how many different ways can we apply these labels to n objects?

Show that if we have k_1 labels of one kind, e.g., red, k_2 labels of a second kind, e.g., blue, and $k_3 = n - k_1 - k_2$ labels of a third kind, then there are $\frac{n!}{k_1!k_2!k_3!}$ ways to apply these labels to n objects




Labelling and Trinomial Coefficients

- Suppose we have k labels of one kind, e.g., red and $n - k$ labels of another, e.g., blue. In how many different ways can we apply these labels to n objects?

Show that if we have k_1 labels of one kind, e.g., red, k_2 labels of a second kind, e.g., blue, and $k_3 = n - k_1 - k_2$ labels of a third kind, then there are $\frac{n!}{k_1!k_2!k_3!}$ ways to apply these labels to n objects

What is the coefficient of $x^{k_1}y^{k_2}z^{k_3}$ in $(x + y + z)^n$?



Labelling and Trinomial Coefficients

- There are $\binom{n}{k_1}$ ways to choose the red items. There are then $\binom{n-k_1}{k_2}$ ways to choose the blue items from the remaining $n - k_1$. The remaining k_3 items get labelled a third color.



Labelling and Trinomial Coefficients

- There are $\binom{n}{k_1}$ ways to choose the red items. There are then $\binom{n-k_1}{k_2}$ ways to choose the blue items from the remaining $n - k_1$. The remaining k_3 items get labelled a third color.

Using the *product rule* the total number of labellings is

$$\begin{aligned}\binom{n}{k_1} \binom{n-k_1}{k_2} &= \frac{n!}{k_1!(n-k_1)!} \frac{(n-k_1)!}{(k_2)!(n-k_1-k_2)!} \\ &= \frac{n!}{k_1!k_2!(n-k_1-k_2)!} = \frac{n!}{k_1!k_2!k_3!}\end{aligned}$$



Labelling and Trinomial Coefficients

- When $k_1 + k_2 + k_3 = n$, we call

$$\frac{n!}{k_1!k_2!k_3!}$$

a *trinomial coefficient* and denote it as

$$\binom{n}{k_1 \ k_2 \ k_3}$$



Labelling and Trinomial Coefficients

- When $k_1 + k_2 + k_3 = n$, we call

$$\frac{n!}{k_1!k_2!k_3!}$$

a *trinomial coefficient* and denote it as

$$\binom{n}{k_1 \ k_2 \ k_3}$$

What is the coefficient of $x^{k_1}y^{k_2}z^{k_3}$ in $(x + y + z)^n$?



The Birthday Paradox

- Suppose that 25 students are in a room. What is the probability that at least two of them share a birthday?



The Birthday Paradox

- Suppose that 25 students are in a room. What is the probability that at least two of them share a birthday?

It's greater than $1/2$! (only need 23)



The Birthday Paradox

- Suppose that 25 students are in a room. What is the probability that at least two of them share a birthday?

It's greater than $1/2$! (only need 23)

A_n – “there are n students in a room and at least two of them share a birthday.”



The Birthday Paradox

- Suppose that 25 students are in a room. What is the probability that at least two of them share a birthday?

It's greater than $1/2$! (only need 23)

A_n – “there are n students in a room and at least two of them share a birthday.”

We may assume that a year has 365 days and there are no twins in the room.



The Birthday Paradox

- Suppose that 25 students are in a room. What is the probability that at least two of them share a birthday?

It's greater than $1/2$! (only need 23)

A_n – “there are n students in a room and at least two of them share a birthday.”

We may assume that a year has 365 days and there are no twins in the room.

This will be very similar to the analysis of hashing n keys into a table of size 365.



The Birthday Paradox

- A_n – “there are n students in a room and at least two of them share a birthday.”

Sample space: $|S| = 365^n$




The Birthday Paradox

- A_n – “there are n students in a room and at least two of them share a birthday.”

Sample space: $|S| = 365^n$

B_n – “there are n students in a room and none of them share a birthday.”



The Birthday Paradox

- A_n – “there are n students in a room and at least two of them share a birthday.”

Sample space: $|S| = 365^n$

B_n – “there are n students in a room and none of them share a birthday.”

$$\#B_n = 365 \times 364 \times \cdots \times (365 - (n - 1))$$



The Birthday Paradox

- A_n – “there are n students in a room and at least two of them share a birthday.”

Sample space: $|S| = 365^n$

B_n – “there are n students in a room and none of them share a birthday.”


$$\#B_n = 365 \times 364 \times \cdots \times (365 - (n - 1))$$

$$\#A_n + \#B_n = 365^n$$



The Birthday Paradox

n	A_n	B_n	n	A_n	B_n
1	0.00000000	1.00000000	16	0.28360400	0.71639599
2	0.00273972	0.99726027	17	0.31500766	0.68499233
3	0.00820416	0.99179583	18	0.34691141	0.65308858
4	0.01635591	0.98364408	19	0.37911852	0.62088147
5	0.02713557	0.97286442	20	0.41143838	0.58856161
6	0.04046248	0.95953751	21	0.44368833	0.55631166
7	0.05623570	0.94376429	22	0.47569530	0.52430469
8	0.07433529	0.92566470	23	0.50729723	0.49270276
9	0.09462383	0.90537616	24	0.53834425	0.46165574
10	0.11694817	0.88305182	25	0.56869970	0.43130029
11	0.14114137	0.85885862	26	0.59824082	0.40175917
12	0.16702478	0.83297521	27	0.62685928	0.37314071
13	0.19441027	0.80558972	28	0.65446147	0.34553852
14	0.22310251	0.77689748	29	0.68096853	0.31903146
15	0.25290131	0.74709868	30	0.70631624	0.29368375



Next Lecture

- counting II, relation, ...

