

---

# MA204: Mathematical Statistics

## Suggested Solutions to Assignment 5

---

**5.1 Solution.** In Example 5.5, let  $\mathbb{C} = \{1, 7, 3, 8, 4\}$ . (a) The Type I error rate is

$$\begin{aligned}\alpha(0) &= \Pr(X \in \mathbb{C} | \theta = 0) \\ &= \Pr(X = 1|0) + \Pr(X = 7|0) + \Pr(X = 3|0) \\ &\quad + \Pr(X = 8|0) + \Pr(X = 4|0) \\ &= 0 + 0.01 + 0.02 + 0.07 + 0.05 \\ &= 0.15.\end{aligned}$$

(b) The acceptance region is  $\mathbb{C}' = \{5, 9, 10, 6, 2\}$  so that the Type II error rate is given by

$$\begin{aligned}\beta(1) &= \Pr(X \in \mathbb{C}' | \theta = 1) \\ &= \Pr(X = 5|1) + \Pr(X = 9|1) + \Pr(X = 10|1) \\ &\quad + \Pr(X = 6|1) + \Pr(X = 2|1) \\ &= 0.03 + 0.02 + 0.04 + 0.01 + 0 \\ &= 0.1.\end{aligned}$$

**5.2 Solution.** Since  $Y \sim \text{Binomial}(n, \theta)$ , we have

$$E(Y) = n\theta \quad \text{and} \quad \text{Var}(Y) = n\theta(1 - \theta).$$

Let  $Z \sim N(0, 1)$ . By the normal approximation, the power function is

$$\begin{aligned}
 p(\theta) &= \Pr(\text{reject } H_0 | \theta) \\
 &= \Pr(Y \geq c | \theta) \\
 &= \Pr \left\{ \frac{Y - n\theta}{\sqrt{n\theta(1-\theta)}} \geq \frac{c - n\theta}{\sqrt{n\theta(1-\theta)}} \middle| \theta \right\} \\
 &\doteq \Pr \left\{ Z \geq \frac{c - n\theta}{\sqrt{n\theta(1-\theta)}} \middle| \theta \right\}.
 \end{aligned}$$

Since  $0.1 = p(0.5)$ , we obtain

$$0.1 \doteq \Pr \left\{ Z \geq \frac{2c - n}{\sqrt{n}} \middle| \theta \right\}$$

so that

$$\frac{2c - n}{\sqrt{n}} = z_{0.10} = 1.2816. \quad (\text{SA5.1})$$

On the other hand,  $0.95 = p(2/3)$ , we obtain

$$0.95 \doteq \Pr \left\{ Z \geq \frac{3c - 2n}{\sqrt{2n}} \middle| \theta \right\}$$

so that

$$\frac{3c - 2n}{\sqrt{2n}} = z_{0.95} = -1.645. \quad (\text{SA5.2})$$

Solving equations (SA5.1) and (SA5.2), we get

$$1.5(n + 1.2816\sqrt{n}) - 2n = -1.645\sqrt{2n},$$

i.e.,

$$\sqrt{n} = 8.49756 \quad \text{or} \quad n = 72.2086.$$

Then

$$c = \frac{n + 1.2816\sqrt{n}}{2} = 41.5495.$$

Approximately,  $n = 72$  and  $c = 42$ .

**5.3 Solution.** (a) The distribution of  $Y = \sum_{i=1}^n X_i$  is Gamma( $2n, \theta$ ).

(b) The likelihood function is

$$L(\theta) = \prod_{i=1}^n \frac{\theta^2}{\Gamma(2)} x_i e^{-\theta x_i} = \theta^{2n} \{\prod_{i=1}^n x_i\} \cdot e^{-\theta \sum_{i=1}^n x_i}.$$

Since  $\theta_1 > 1$ , the ratio

$$\frac{L(1)}{L(\theta_1)} = \theta_1^{-2n} e^{(\theta_1 - 1) \sum_{i=1}^n x_i} \leq k$$

is equivalent to

$$\sum_{i=1}^n x_i \leq c.$$

By Neymann-Pearson Lemma, a test  $\varphi$  of size  $\alpha$  with critical region

$$\mathbb{C} = \{x: L(1)/L(\theta_1) \leq k\} = \left\{x: \sum_{i=1}^n x_i \leq c\right\}$$

is the most powerful test for testing  $H_0$  against  $H_1$ .

**How to determine the  $c$ ?** Under  $H_0$  (i.e.  $\theta = 1$ ),

$$Y = \sum_{i=1}^n X_i \sim \text{Gamma}(2n, 1).$$

Using Property 1 in Appendix A.2.4, under  $H_0$ , we have

$$2Y \sim \chi^2(4n).$$

Hence

$$\begin{aligned} \alpha &= \Pr(X \in \mathbb{C} | H_0) \\ &= \Pr\{\sum_{i=1}^n X_i \leq c | H_0\} \\ &= \Pr\{Y \leq c | H_0\} \\ &= \Pr\{2Y \leq 2c | H_0\} \end{aligned}$$

or

$$1 - \alpha = \Pr\{2Y > 2c\}$$

so that  $2c = \chi^2(1 - \alpha, 4n)$ , i.e.,

$$c = \frac{1}{2}\chi^2(1 - \alpha, 4n).$$

(c) The power function, for  $\theta \geq 1$ ,

$$\begin{aligned} p(\theta) &= \Pr(X \in \mathbb{C}|\theta) \\ &= \Pr\{\sum_{i=1}^n X_i \leq c|\theta\} \\ &= \Pr\{Y \leq c|\theta\} \quad \text{by } Y \sim \text{Gamma}(2n, \theta) \\ &= \int_0^c \frac{\theta^{2n}}{\Gamma(2n)} y^{2n-1} e^{-\theta y} dy. \end{aligned}$$

**5.4 Solution.** (a) Let  $X = (X_1, \dots, X_n)$ ,  $x = (x_1, \dots, x_n)$ ,

$$Q(X) \hat{=} \prod_{i=1}^n (1 - X_i) \quad \text{and} \quad Q(x) \hat{=} \prod_{i=1}^n (1 - x_i),$$

then, the likelihood function is

$$L(\theta) = \prod_{i=1}^n \theta (1 - x_i)^{\theta-1} = \theta^n [Q(x)]^{\theta-1}. \quad (\text{SA5.3})$$

Note that

$$\frac{L(1)}{L(\theta_1)} = \frac{1}{\theta_1^n [Q(x)]^{\theta_1-1}} \leq k$$

is equivalent to

$$\log Q(x) \geq c.$$

By Neymann-Pearson Lemma, a test  $\varphi$  of size  $\alpha$  with critical region

$$\mathbb{C} = \{x: L(1)/L(\theta_1) \leq k\} = \{x: \log Q(x) \geq c\}$$

is the most powerful test for testing  $H_0$  against  $H_1$ . To determine the  $c$ , we note that

$$Y_i = -\log(1 - X_i) \sim \text{Exponential}(\theta) = \text{Gamma}(1, \theta),$$

thus

$$-\log Q(X) = -\sum_{i=1}^n \log(1 - X_i) = \sum_{i=1}^n Y_i \sim \text{Gamma}(n, \theta).$$

Under  $H_0$ ,  $-\log Q(X) \sim \text{Gamma}(n, 1)$ . Using Property 1 in Appendix A.2.4, under  $H_0$ , we have

$$-2\log Q(X) \sim \chi^2(2n). \quad (\text{SA5.4})$$

Hence

$$\begin{aligned} \alpha &= \Pr(X \in \mathbb{C} | H_0) \\ &= \Pr\{\log Q(X) \geq c | H_0\} \\ &= \Pr\{-2\log Q(X) \leq -2c | H_0\} \end{aligned}$$

or

$$1 - \alpha = \Pr\{-2\log Q(X) > -2c\}$$

so that  $-2c = \chi^2(1 - \alpha, 2n)$ , i.e.,  $c = -\chi^2(1 - \alpha, 2n)/2$ .

(b) Now  $\Theta_0 = \{1\}$  and  $\Theta = (0, \infty)$ . To derive the likelihood ratio statistic (LRS), we first need to find the MLE of  $\theta$ . From (SA5.3), we have

$$\begin{aligned} \log L(\theta) &= n \log \theta + (\theta - 1) \log Q(x), \\ \frac{d \log L(\theta)}{d\theta} &= \frac{n}{\theta} + \log Q(x). \end{aligned}$$

Thus, the MLE is

$$\hat{\theta} = -n / \log Q(X)$$

with  $Q(X)$  as a sufficient statistic, and the LRS is

$$\lambda(X) = \frac{L(1)}{L(\hat{\theta})} = \frac{1}{\hat{\theta}^n [Q(X)]^{\hat{\theta}-1}}.$$

Denoting  $Q(x)$  by  $Q$ , we have

$$\lambda(x) = \frac{1}{(-n/\log Q)^n \cdot Q^{-n/\log Q-1}}.$$

Thus, based on (5.29) in Chapter 5 (page 99), the critical region that the  $H_0$  is rejected is

$$\mathbb{C} = \{x: \lambda(x) \leq \lambda_\alpha\} = \{x: (-n/\log Q)^n \cdot Q^{-n/\log Q-1} \geq c\}. \quad (\text{SA5.5})$$

**To determine  $c$** , we let

$$h(Q) = (-n/\log Q)^n \cdot Q^{-n/\log Q-1},$$

then

$$\log h(Q) = n \log n - n \log(-\log Q) - n - \log Q.$$

We have

$$\frac{d \log h(Q)}{dQ} = \frac{n}{Q(-\log Q)} - \frac{1}{Q} = \frac{n + \log Q}{Q(-\log Q)}.$$

Setting  $\frac{d \log h(Q)}{dQ} = 0$ , we obtain  $Q = e^{-n}$ . Note that  $0 < Q < 1$ , then

$$\frac{d \log h(Q)}{dQ} = \frac{n + \log Q}{Q(-\log Q)} < 0, \quad \text{when } Q < e^{-n},$$

and

$$\frac{d \log h(Q)}{dQ} = \frac{n + \log Q}{Q(-\log Q)} > 0, \quad \text{when } Q > e^{-n},$$

Hence,  $Q = e^{-n}$  is the minimum of  $h(Q)$ , and  $h(Q)$  is decreasing when  $Q < e^{-n}$  and increasing when  $Q > e^{-n}$ . Therefore, (SA5.5) is equivalent to

$$\mathbb{C} = \{x: Q \leq c_1 \quad \text{or} \quad Q \geq c_2\}. \quad (\text{SA5.6})$$

Namely, determining  $c$  is equivalent to determining  $c_1$  and  $c_2$ .

**How to determine  $c_1$  and  $c_2$ ?** Based on the Type I error rate

$$\begin{aligned} \alpha &= \Pr\{Q(X) \leq c_1 \quad \text{or} \quad Q(X) \geq c_2 | H_0\} \\ &= \Pr\{Q(X) \leq c_1 | H_0\} + \Pr\{Q(X) \geq c_2 | H_0\}, \end{aligned}$$

we use the equal-tail approach, i.e.,

$$\alpha/2 = \Pr\{Q(X) \leq c_1 | H_0\} \quad (\text{SA5.7})$$

and

$$\alpha/2 = \Pr\{Q(X) \geq c_2 | H_0\}. \quad (\text{SA5.8})$$

Recall (SA5.4), from (SA5.7), we have

$$\begin{aligned} \alpha/2 &= \Pr\{\log Q(X) \leq \log c_1 | H_0\} \\ &= \Pr\{-2 \log Q(X) \geq -2 \log c_1 | H_0\} \end{aligned}$$

so that  $-2 \log c_1 = \chi^2(\alpha/2, 2n)$ , i.e.,

$$c_1 = \exp\{-0.5\chi^2(\alpha/2, 2n)\}. \quad (\text{SA5.9})$$

Similarly, from (SA5.8), we have

$$\begin{aligned} 1 - \alpha/2 &= \Pr\{Q(X) < c_2 | H_0\} \\ &= \Pr\{\log Q(X) < \log c_2 | H_0\} \\ &= \Pr\{-2 \log Q(X) > -2 \log c_2 | H_0\} \end{aligned}$$

so that  $-2 \log c_2 = \chi^2(1 - \alpha/2, 2n)$ , i.e.,

$$c_2 = \exp\{-0.5\chi^2(1 - \alpha/2, 2n)\}. \quad (\text{SA5.10})$$

**5.5 Solution.** We first consider to test

$$H_0: \theta = \theta_0 \quad \text{against} \quad H_1: \theta = \theta_1 (< \theta_0).$$

Let  $\varphi$  be a test with critical region satisfying (5.17). The likelihood function is given by

$$\begin{aligned} L(\theta) &= \prod_{i=1}^n \frac{1}{\sqrt{2\pi}} \exp \left\{ -\frac{1}{2}(x_i - \theta)^2 \right\} \\ &= (2\pi)^{-n/2} \exp \left\{ -\frac{1}{2} \sum_{i=1}^n (x_i - \theta)^2 \right\}. \end{aligned}$$

Then

$$\begin{aligned} \frac{L(\theta_0)}{L(\theta_1)} &= \exp \left\{ -\frac{1}{2} \sum_{i=1}^n [(x_i - \theta_0)^2 - (x_i - \theta_1)^2] \right\} \\ &= \exp \left\{ (\theta_0 - \theta_1) \sum_{i=1}^n x_i - n(\theta_0^2 - \theta_1^2)/2 \right\} \leq k \end{aligned}$$

is equivalent to (by noting that  $\theta_0 - \theta_1 > 0$ )

$$\bar{x} \leq \frac{\log k}{n(\theta_0 - \theta_1)} + \frac{\theta_0 + \theta_1}{2} \triangleq c.$$

To determine  $c$ , we consider the size

$$\begin{aligned} \alpha &= \Pr\{\bar{X} \leq c | \theta = \theta_0\} \\ &= \Pr\{\sqrt{n}(\bar{X} - \theta_0) \leq \sqrt{n}(c - \theta_0)\} \\ &= \Pr\{Z \leq \sqrt{n}(c - \theta_0)\} \\ &= \Pr\{Z \leq -z_\alpha\}. \end{aligned}$$

Then,  $\sqrt{n}(c - \theta_0) = -z_\alpha$  or  $c = \theta_0 - z_\alpha/\sqrt{n}$ . Therefore, the test with critical region

$$\mathbb{C} = \{x: \bar{x} \leq \theta_0 - z_\alpha/\sqrt{n}\}$$



is a most powerful test (MPT) of size  $\alpha$ . Since the  $\mathbb{C}$  depends only on  $n$ ,  $\theta_0$ ,  $\alpha$  and the fact  $\theta_1 < \theta_0$ , but not on the value of  $\theta_1$ , the test  $\varphi$  is also a UMPT of size  $\alpha$  for testing

$$H_0: \theta = \theta_0 \quad \text{against} \quad H_1: \theta < \theta_0.$$

On the other hand, the power function is given by

$$\begin{aligned} p_\varphi(\theta) &= \Pr\{X \in \mathbb{C}|\theta\} \\ &= \Pr\{\bar{X} \leq \theta_0 - z_\alpha/\sqrt{n}|\theta\} \\ &= \Pr\{\sqrt{n}(\bar{X} - \theta) \leq -z_\alpha + \sqrt{n}(\theta_0 - \theta)|\theta\} \\ &= \Pr\{Z \leq -z_\alpha + \sqrt{n}(\theta_0 - \theta)\} \\ &= \Phi(-z_\alpha + \sqrt{n}(\theta_0 - \theta)) \end{aligned}$$

so that

$$\begin{aligned} \sup_{\theta \in \Theta_0} p_\varphi(\theta) &= \sup_{\theta \geq \theta_0} \Phi(-z_\alpha + \sqrt{n}(\theta_0 - \theta)) \\ &= \max_{\theta \geq \theta_0} \Phi(-z_\alpha + \sqrt{n}(\theta_0 - \theta)) \\ &= \Phi(-z_\alpha) \\ &= \alpha = p_\varphi(\theta_0). \end{aligned}$$

Then, the test  $\varphi$  is also a UMPT of size  $\alpha$  for testing

$$H_0: \theta \geq \theta_0 \quad \text{against} \quad H_1: \theta < \theta_0.$$

**5.6 Solution.** (a) We first consider

$$H_0: \sigma^2 = \sigma_0^2 \quad \text{against} \quad H_1: \sigma^2 \neq \sigma_0^2.$$

Note that  $\Theta_0 = \{(\mu, \sigma_0^2): -\infty < \mu < +\infty, \}$  and the whole parameter space  $\Theta = \{(\mu, \sigma^2): -\infty < \mu < +\infty, \sigma^2 > 0\}$ . The likelihood function is given by

$$L(\mu, \sigma^2) = (2\pi\sigma^2)^{-n/2} \exp \left\{ -\frac{1}{2\sigma^2} \sum_{i=1}^n (x_i - \mu)^2 \right\}.$$

so that

$$\log L(\mu, \sigma^2) = -\frac{n}{2} \log(2\pi) - \frac{n}{2} \log(\sigma^2) - \frac{1}{2\sigma^2} \sum_{i=1}^n (x_i - \mu)^2.$$

Hence, the unrestricted maximum likelihood estimates of  $\mu$  and  $\sigma^2$  are given by

$$\hat{\mu} = \bar{x} = \frac{\sum_{i=1}^n x_i}{n}, \quad \text{and} \quad \hat{\sigma}^2 = \frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n} = \frac{(n-1)s^2}{n}.$$

Under  $H_0$ , the restricted maximum likelihood estimate of  $\mu$  is  $\hat{\mu}^R = \bar{x}$ . Thus, the likelihood ratio is

$$\lambda(x) = \frac{L(\hat{\mu}^R, \sigma_0^2)}{L(\hat{\theta}, \hat{\sigma}^2)} = \left[ \frac{(n-1)s^2}{n\sigma_0^2} \right]^{n/2} \exp \left\{ -\frac{(n-1)s^2}{2\sigma_0^2} + \frac{n}{2} \right\}.$$

Define  $f(s^2) = \log \lambda(x)$ , then

$$\begin{aligned} \frac{df(s^2)}{ds^2} = f'(s^2) &= \frac{n}{2s^2} - \frac{n-1}{2\sigma_0^2}, \\ f''(s^2) &= -\frac{n}{2s^4} < 0, \end{aligned}$$

i.e.,  $f(s^2)$  has a maximum at  $s^2 = n\sigma_0^2/(n-1)$ . Since  $\lambda(x) \leq \lambda_\alpha$  is equivalent to

$$\log \lambda(x) = f(s^2) \leq \log \lambda_\alpha,$$

the critical region is given by

$$\mathbb{C}_1 = \{(x_1, \dots, x_n): s^2 \leq c_1 \quad \text{or} \quad s^2 \geq c_2\}.$$

Namely, determining  $\lambda_\alpha$  is equivalent to determining  $c_1$  and  $c_2$ .

**How to determine  $c_1$  and  $c_2$ ?** Based on the Type I error rate

$$\begin{aligned} \alpha &= \Pr\{S^2 \leq c_1 \quad \text{or} \quad S^2 \geq c_2 | H_0\} \\ &= \Pr\{S^2 \leq c_1 | H_0\} + \Pr\{S^2 \geq c_2 | H_0\}, \end{aligned}$$

we use the equal-tail approach, i.e.,

$$\alpha/2 = \Pr\{S^2 \leq c_1 | H_0\} \quad (\text{SA5.11})$$

and

$$\alpha/2 = \Pr\{S^2 \geq c_2 | H_0\}. \quad (\text{SA5.12})$$

Recall that  $(n-1)S^2/\sigma^2 \sim \chi^2(n-1)$ , from (SA5.11), we have

$$\begin{aligned} 1 - \alpha/2 &= \Pr\{S^2 > c_1 | H_0\} \\ &= \Pr\{(n-1)S^2/\sigma^2 > (n-1)c_1/\sigma^2 | \sigma^2 = \sigma_0^2\} \\ &= \Pr\{(n-1)S^2/\sigma_0^2 > (n-1)c_1/\sigma_0^2\} \end{aligned}$$

so that  $(n-1)c_1/\sigma_0^2 = \chi^2(1 - \alpha/2, n-1)$ , i.e.,

$$c_1 = \frac{\sigma_0^2 \chi^2(1 - \alpha/2, n-1)}{n-1}.$$

Similarly, from (SA5.12), we have

$$\begin{aligned} \alpha/2 &= \Pr\{S^2 \geq c_2 | H_0\} \\ &= \Pr\{(n-1)S^2/\sigma^2 \geq (n-1)c_2/\sigma^2 | \sigma^2 = \sigma_0^2\} \\ &= \Pr\{(n-1)S^2/\sigma_0^2 \geq (n-1)c_2/\sigma_0^2\} \end{aligned}$$

so that  $(n-1)c_2/\sigma_0^2 = \chi^2(\alpha/2, n-1)$ , i.e.,

$$c_2 = \frac{\sigma_0^2 \chi^2(\alpha/2, n-1)}{n-1}.$$

Therefore, the critical region is

$$\begin{aligned} \mathbb{C}_1 &= \left\{ (x_1, \dots, x_n): s^2 \leq \frac{\sigma_0^2 \chi^2(1 - \alpha/2, n-1)}{n-1} \right. \\ &\quad \left. \text{or } s^2 \geq \frac{\sigma_0^2 \chi^2(\alpha/2, n-1)}{n-1} \right\}. \end{aligned}$$

(b) We then consider

$$H_0: \sigma^2 = \sigma_0^2 \quad \text{against} \quad H_1: \sigma^2 > \sigma_0^2.$$

The critical region is

$$\mathbb{C}_2 = \left\{ (x_1, \dots, x_n): s^2 \geq \frac{\sigma_0^2 \chi^2(1 - \alpha, n - 1)}{n - 1} \right\}.$$

(c) We finally consider

$$H_0: \sigma^2 = \sigma_0^2 \quad \text{against} \quad H_1: \sigma^2 < \sigma_0^2.$$

The critical region is

$$\mathbb{C}_3 = \left\{ (x_1, \dots, x_n): s^2 \leq \frac{\sigma_0^2 \chi^2(1 - \alpha, n - 1)}{n - 1} \right\}.$$

**5.7 Solution.** Let  $\theta = (\mu_1, \mu_2, \sigma^2)^\top$ ,  $\Theta_0 = \{(0, 0, \sigma^2)^\top: \sigma^2 > 0\}$  and  $\Theta = \{\theta = (\mu_1, \mu_2, \sigma^2)^\top: -\infty < \mu_1, \mu_2 < +\infty, \sigma^2 > 0\}$ , then

$$H_0: \theta \in \Theta_0 \quad \text{against} \quad H_1: \theta \in \Theta_1 = \Theta - \Theta_0.$$

The likelihood function is

$$L(\theta) = (2\pi\sigma^2)^{-n} \exp \left[ -\frac{\sum_{i=1}^n (x_i - \mu_1)^2 + \sum_{i=1}^n (y_i - \mu_2)^2}{2\sigma^2} \right]$$

so that

$$\log L(\theta) = -n \log(2\pi) - n \log(\sigma^2) - \frac{1}{2\sigma^2} \sum_{i=1}^n [(x_i - \mu_1)^2 + (y_i - \mu_2)^2].$$

Hence, the unrestricted maximum likelihood estimates of  $\mu_1$ ,  $\mu_2$  and  $\sigma^2$  are given by

$$\begin{aligned} \hat{\mu}_1 &= \frac{\sum_{i=1}^n x_i}{n} = \bar{x}, \\ \hat{\mu}_2 &= \frac{\sum_{i=1}^n y_i}{n} = \bar{y}, \quad \text{and} \\ \hat{\sigma}^2 &= \frac{\sum_{i=1}^n [(x_i - \bar{x})^2 + (y_i - \bar{y})^2]}{2n} = \frac{(n-1)(s_1^2 + s_2^2)}{2n}. \end{aligned}$$

Under  $H_0$  (i.e.,  $\mu_1 = \mu_2 = 0$ ), the restricted maximum likelihood estimate of  $\sigma^2$  is

$$\begin{aligned}\hat{\sigma}^{2R} &= \frac{\sum_{i=1}^n (x_i^2 + y_i^2)}{2n} \\ &= \frac{\sum_{i=1}^n (x_i - \bar{x})^2 + n\bar{x}^2 + \sum_{i=1}^n (y_i - \bar{y})^2 + n\bar{y}^2}{2n} \\ &= \frac{(n-1)(s_1^2 + s_2^2) + n\bar{x}^2 + n\bar{y}^2}{2n}.\end{aligned}$$

Thus, the likelihood ratio is

$$\begin{aligned}\lambda(x) &= \frac{L(0, 0, \hat{\sigma}^{2R})}{L(\hat{\mu}_1, \hat{\mu}_2, \hat{\sigma}^2)} = \left( \frac{\hat{\sigma}^2}{\hat{\sigma}^{2R}} \right)^n \\ &= \left[ \frac{(n-1)(s_1^2 + s_2^2)}{(n-1)(s_1^2 + s_2^2) + n\bar{x}^2 + n\bar{y}^2} \right]^n \\ &= \left[ \frac{1}{1 + F} \right]^n,\end{aligned}$$

where

$$F = \frac{n\bar{x}^2 + n\bar{y}^2}{(n-1)(s_1^2 + s_2^2)}.$$

Since  $\lambda(x) \leq \lambda_\alpha$  is equivalent to  $F \geq c$ , the critical region is given by

$$\mathbb{C} = \{(x_1, \dots, x_n, y_1, \dots, y_n): F \geq c\}.$$

**How to determine  $c$ ?** Note that

$$\frac{\sqrt{n}(\bar{X} - \mu_1)}{\sigma} \sim N(0, 1), \quad \frac{\sqrt{n}(\bar{Y} - \mu_2)}{\sigma} \sim N(0, 1),$$

$(n-1)S_i^2/\sigma^2 \sim \chi^2(n-1)$ ,  $i = 1, 2$ , and they are independent.

Thus, under  $H_0$ ,

$$\frac{n(\bar{X}^2 + \bar{Y}^2)}{\sigma^2} \sim \chi^2(2) \quad \text{and} \quad \frac{(n-1)(S_1^2 + S_2^2)}{\sigma^2} \sim \chi^2(2n-2)$$

so that

$$(n-1)F = \frac{\frac{n(\bar{X}^2 + \bar{Y}^2)}{\sigma^2}/2}{\frac{(n-1)(S_1^2 + S_2^2)}{\sigma^2}/2(n-1)} \sim F(2, 2n-2).$$

Based on the Type I error rate

$$\begin{aligned}\alpha &= \Pr\{F \geq c | H_0\} \\ &= \Pr\{(n-1)F \geq (n-1)c\},\end{aligned}$$

we have  $(n-1)c = f(\alpha, 2, 2n-2)$ , i.e.,

$$c = \frac{f(\alpha, 2, 2n-2)}{n-1}.$$

**5.8 Solution.** (a) We first consider

$$H_0: \sigma_1^2 = \sigma_2^2 = \sigma^2 \quad \text{against} \quad H_1: \sigma_1^2 \neq \sigma_2^2.$$

Let  $\theta = (\mu_1, \mu_2, \sigma_1^2, \sigma_2^2)^\top$ . The likelihood function is given by

$$\begin{aligned}L(\theta) &= (2\pi\sigma_1^2)^{-n_1/2} \exp\left\{-\frac{1}{2\sigma_1^2} \sum_{i=1}^{n_1} (x_i - \mu_1)^2\right\} \\ &\quad \times (2\pi\sigma_2^2)^{-n_2/2} \exp\left\{-\frac{1}{2\sigma_2^2} \sum_{j=1}^{n_2} (y_j - \mu_2)^2\right\}\end{aligned}$$

so that

$$\begin{aligned}\log L(\theta) &= -\frac{n_1}{2} \log(2\pi) - \frac{n_1}{2} \log(\sigma_1^2) - \frac{1}{2\sigma_1^2} \sum_{i=1}^{n_1} (x_i - \mu_1)^2 \\ &\quad - \frac{n_2}{2} \log(2\pi) - \frac{n_2}{2} \log(\sigma_2^2) - \frac{1}{2\sigma_2^2} \sum_{j=1}^{n_2} (y_j - \mu_2)^2.\end{aligned}$$

Hence, the unrestricted maximum likelihood estimates of the 4 parameters are given by

$$\hat{\mu}_1 = \bar{x}, \quad \hat{\mu}_2 = \bar{y}, \quad \text{and} \quad \hat{\sigma}_k^2 = \frac{(n_k - 1)s_k^2}{n_k}, \quad k = 1, 2.$$

Under  $H_0$ , the restricted maximum likelihood estimates of the 3 parameters are

$$\hat{\mu}_1^R = \bar{x}, \quad \hat{\mu}_2^R = \bar{y}, \quad \text{and} \quad \hat{\sigma}^{2R} = \frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{n_1 + n_2}.$$

Thus, the likelihood ratio is

$$\begin{aligned}
& \lambda(x) \\
&= \frac{L(\hat{\mu}_1^R, \hat{\mu}_2^R, \hat{\sigma}^{2R}, \hat{\sigma}^{2R})}{L(\hat{\mu}_1^R, \hat{\mu}_2^R, \hat{\sigma}_1^{2R}, \hat{\sigma}_2^{2R})} \\
&= \frac{(\hat{\sigma}_1^2)^{n_1/2} (\hat{\sigma}_2^2)^{n_2/2}}{(\hat{\sigma}^{2R})^{(n_1+n_2)/2}} \\
&= \left[ \frac{(n_1+n_2)(n_1-1)/n_1}{n_1-1+(n_2-1)s_2^2/s_1^2} \right]^{\frac{n_1}{2}} \left[ \frac{(n_1+n_2)(n_2-1)[s_2^2/s_1^2]/n_2}{n_1-1+(n_2-1)s_2^2/s_1^2} \right]^{\frac{n_2}{2}}.
\end{aligned}$$

Define  $t = s_2^2/s_1^2$  and

$$f(t) = \frac{t^{n_2/2}}{[n_1-1+(n_2-1)t]^{\frac{n_1+n_2}{2}}},$$

then

$$\frac{d \log f(t)}{dt} = \frac{n_2}{2t} - \frac{n_1+n_2}{2} \cdot \frac{n_2-1}{n_1-1+(n_2-1)t}$$

Setting  $\frac{d \log f(t)}{dt} = 0$ , we obtain  $t = n_2(n_1-1)/[n_1(n_2-1)] \triangleq t_0$ .

Note that

$$\frac{d \log f(t)}{dt} = \frac{n_2(n_1-1) - n_1(n_2-1)t}{2t[n_1-1+(n_2-1)t]} < 0, \quad \text{when } t > t_0,$$

and

$$\frac{d \log f(t)}{dt} = \frac{n_2(n_1-1) - n_1(n_2-1)t}{2t[n_1-1+(n_2-1)t]} > 0, \quad \text{when } t < t_0.$$

Hence,  $t = t_0$  is the maximum of  $f(t)$ , and  $f(t)$  is increasing when  $t < t_0$  and decreasing when  $t > t_0$ . Therefore,  $\lambda(x) \leq \lambda_\alpha$  is equivalent to  $f(t) \leq c$ , resulting in the following critical region

$$\mathbb{C}_1 = \{x: s_2^2/s_1^2 \leq c_1 \quad \text{or} \quad s_2^2/s_1^2 \geq c_2\}.$$

**How to determine  $c_1$  and  $c_2$ ?** Based on the Type I error rate

$$\begin{aligned}
\alpha &= \Pr\{S_2^2/S_1^2 \leq c_1 \quad \text{or} \quad S_2^2/S_1^2 \geq c_2 | H_0\} \\
&= \Pr\{S_2^2/S_1^2 \leq c_1 | H_0\} + \Pr\{S_2^2/S_1^2 \geq c_2 | H_0\},
\end{aligned}$$

we use the equal-tail approach, i.e.,

$$\alpha/2 = \Pr\{S_2^2/S_1^2 \leq c_1 | H_0\}$$

and

$$\alpha/2 = \Pr\{S_2^2/S_1^2 \geq c_2 | H_0\}.$$

Note that under  $H_0$ ,  $S_2^2/S_1^2 \sim F(n_2 - 1, n_1 - 1)$ , we have

$$c_1 = f(1 - \alpha/2, n_2 - 1, n_1 - 1) \quad \text{and} \quad c_2 = f(\alpha/2, n_2 - 1, n_1 - 1).$$

(b) We then consider

$$H_0: \sigma_1^2 = \sigma_2^2 \quad \text{against} \quad H_1: \sigma_1^2 > \sigma_2^2.$$

The critical region is

$$\mathbb{C}_2 = \{(x_1, \dots, x_n): s_2^2/s_1^2 \leq f(1 - \alpha, n_2 - 1, n_1 - 1)\}.$$

(c) We finally consider

$$H_0: \sigma_1^2 = \sigma_2^2 \quad \text{against} \quad H_1: \sigma_1^2 < \sigma_2^2.$$

The critical region is

$$\mathbb{C}_3 = \{(x_1, \dots, x_n): s_2^2/s_1^2 \geq f(\alpha, n_2 - 1, n_1 - 1)\}.$$

**5.9 Solution.** We consider

$$H_0: \theta = 0.5 \quad \text{against} \quad H_1: \theta \neq 0.5.$$

Let  $X \sim \text{Binomial}(n, \theta)$ , the likelihood function is given by

$$L(\theta) = \binom{n}{x} \theta^x (1 - \theta)^{n-x}$$



so that

$$\log L(\theta) = \log \binom{n}{x} + x \log \theta + (n - x) \log(1 - \theta).$$

Hence, the maximum likelihood estimate of  $\theta$  is  $\hat{\theta} = x/n$ .

(a) The likelihood ratio is

$$\lambda(x) = \frac{L(0.5)}{L(\hat{\theta})} = \frac{(0.5n)^n}{x^x (n - x)^{n-x}}$$

so that the LR statistic is

$$\lambda(X) = \frac{(0.5n)^n}{X^X (n - X)^{n-X}}.$$

(b) Note that  $\lambda(x) \leq \lambda_\alpha$  is equivalent to  $x^x (n - x)^{n-x} \geq k$ , or

$$x \log(x) + (n - x) \log(n - x) \geq c.$$

(c) Define

$$f(x) = x \log(x) + (n - x) \log(n - x),$$

then

$$\frac{df(x)}{dx} = \log(x) - \log(n - x) = \log \frac{x}{n - x}.$$

Setting  $\frac{df(x)}{dx} = 0$ , we obtain  $x = n/2$ . Note that

$$\frac{df(x)}{dx} = \log \frac{x}{n - x} < 0, \quad \text{when } x < n/2,$$

and

$$\frac{df(x)}{dx} = \log \frac{x}{n - x} > 0, \quad \text{when } x > n/2.$$

Hence,  $x = n/2$  is the minimum of  $f(x)$ , and  $f(x)$  is decreasing when  $x < n/2$  and increasing when  $x > n/2$ . Therefore,  $f(x) \geq c^*$ , resulting in the following critical region

$$\mathbb{C} = \{x: 0 \leq x \leq c_1 \quad \text{or} \quad c_2 \leq x \leq n\},$$

where  $c_1 < n/2 < c_2$ .

Note that

$$\begin{aligned} f(0) &= f(n) = n \log(n), \\ f(1) &= f(n-1) = (n-1) \log(n-1), \\ f(2) &= f(n-2) = 2 \log(2) + (n-2) \log(n-2), \\ &\vdots \\ f(n/2) &= n \log(n/2), \end{aligned}$$

i.e., the function  $f(x)$  is symmetrical about  $x = n/2$ . Therefore, the critical region  $\mathbb{C}$  can be written as

$$\mathbb{C} = \{x: |x - n/2| \geq c\}.$$

**5.10 Solution.** Now  $n = 556$ ,  $np_{10} = 556 \times 9/16 = 312.75$ ,  $np_{20} = 556 \times 3/16 = 104.25$ ,  $np_{30} = 556 \times 3/16 = 104.25$ , and  $np_{40} = 556 \times 1/16 = 34.75$ . According to (5.40) and (5.42), we have

$$\begin{aligned} Q_n &= \sum_{i=1}^4 \frac{(N_i - np_{i0})^2}{np_{i0}} \\ &= \frac{(315 - 312.75)^2}{312.75} + \frac{(108 - 104.25)^2}{104.25} \\ &\quad + \frac{(101 - 104.25)^2}{104.25} + \frac{(32 - 34.75)^2}{34.75} \\ &= 0.470 < \chi^2(0.05, 3) = 7.8147, \end{aligned}$$

we cannot reject  $H_0$ , so there is a good agreement with null hypothesis or there is a good fit of the data to the model.

**5.11 Solution.** The null hypothesis is  $H_0: p_1 = \cdots = p_6 = 1/6$ .  
Now  $n = 300$ ,  $np_{i0} = 300 \times 1/6 = 50$ ,  $i = 1, \dots, 6$ . According to (5.40) and (5.42), we have

$$\begin{aligned} Q_n &= \sum_{i=1}^6 \frac{(N_i - np_{i0})^2}{np_{i0}} \\ &= \frac{(43 - 50)^2}{50} + \frac{(49 - 50)^2}{50} + \frac{(56 - 50)^2}{50} \\ &\quad + \frac{(45 - 50)^2}{50} + \frac{(66 - 50)^2}{50} + \frac{(41 - 50)^2}{50} \\ &= 8.96 < \chi^2(0.05, 5) = 11.07, \end{aligned}$$

we cannot reject  $H_0$ .