$$\Psi(k) = \mathcal{N}_\Psi \exp(-ik\mu) \exp\left(-\frac{(k-k_0)^2}{4\sigma^2}\right)$$

Adopted

$$b^\dagger_\Psi = \sum_k \Psi(k) b^\dagger_k$$

$$\Psi(k=-\tfrac{\pi}{3})\, b^\dagger_{k=-\frac{\pi}{3}} \qquad \Psi(k=0)\, b^\dagger_{k=0} \qquad \Psi(k=\tfrac{\pi}{3})\, b^\dagger_{k=\frac{\pi}{3}}$$

$$\eta(p,q) = \mathcal{N}_\eta \exp\left(\frac{i(p-q)\mu_k^A}{2}\right) \exp\left(-\frac{(p-q)^2}{4{\sigma_k^A}^2}\right)$$

Optimized

$$\eta(-\tfrac{\pi}{3},\tfrac{\pi}{3})\mathcal{B}(-\tfrac{\pi}{3},\tfrac{\pi}{3}) \qquad \eta(0,0)\mathcal{B}(0,0) \qquad \eta(\tfrac{\pi}{3},-\tfrac{\pi}{3})\mathcal{B}(\tfrac{\pi}{3},-\tfrac{\pi}{3})$$



$$p=-\tfrac{\pi}{3}, q=\tfrac{\pi}{3} \qquad p=0, q=0 \qquad p=\tfrac{\pi}{3}, q=-\tfrac{\pi}{3}$$