

1	2	3	4	5	6	Sum
/20	/20	/20	/20	/20	/20	/100

Name _____

Dec. 29, 2005 Thursday

Department _____

SID _____

University Physics
Practice Final Examination
School of Intensive Instruction in Sciences and Arts, Nanjing University

Physical Constants

Electron charge magnitude	$e = 1.60 \times 10^{-19} \text{ C}$
Dielectric constant of vacuum	$\epsilon_0 = 8.85 \times 10^{-12} \text{ F/m}$
Permeability of vacuum	$\mu_0 = 4\pi \times 10^{-7} \text{ N/A}^2$
Vacuum speed of light	$c = 3.00 \times 10^8 \text{ m/s}$
Electron rest mass	$m_e = 9.11 \times 10^{-31} \text{ kg} = 0.511 \text{ MeV}/c^2$
Proton rest mass	$m_p = 1.672 \times 10^{-27} \text{ kg} = 938.3 \text{ MeV}/c^2$
Neutron rest mass	$m_n = 1.675 \times 10^{-27} \text{ kg} = 939.6 \text{ MeV}/c^2$
Atomic mass unit	$1 \text{ amu} = 1.660 \times 10^{-27} \text{ kg} = 931.5 \text{ MeV}/c^2$
Electron volt (eV)	$1 \text{ eV} = 1.60 \times 10^{-19} \text{ J}$
Planck constant	$h = 6.63 \times 10^{-34} \text{ Js}$ $\hbar = 2.11 \times 10^{-34} \text{ Js} = 0.66 \times 10^{-15} \text{ eVs}$
Boltzmann constant	$k_B = 1.38 \times 10^{-23} \text{ J/K} = 8.62 \times 10^{-5} \text{ eV/K}$
Bohr radius	$a_0 = 0.529 \text{ Angstroms}$

Select five out of following six problems.

1. (20pts). (a) Show that the fractional loss of energy of a photon during a Compton collision is given by

$$\frac{\Delta E}{E} = \frac{h\nu'}{m_e c^2} (1 - \cos \phi)$$

where ν' is the frequency of scattered photon, ϕ is the angle between the momentum of incident photon and the momentum of scattered photon..

(b) Show that when a photon of frequency ν scatters from a free (rest) electron, the maximum kinetic energy of the electron is given by

$$K_{\max} = \frac{2(h\nu)^2}{2h\nu + m_e c^2}.$$

2.(20 pts) An electron is moving on a circular mesoscopic ring of perimeter l .

- (a) Write the Schrödinger's equation of the electron and the boundary condition for the wave function of the electron.
- (b) Obtain the energy levels and the corresponding stationary wave functions of the electron.
- (c) Find the degeneracy of the energy levels.

3.(20pts) The asymmetry term of the empirical nuclear binding energy can be obtained through the Fermi gas model of the nucleus. The model is analogous to electron gas model of the metal. It assumes that the nucleus consists of independent proton and neutron gases. Consider a nucleus consisting of N neutrons and Z protons respectively, where $N + Z = A$, the mass number of the nucleus. The radius of the nucleus is given by $R = r_0 A^{1/3}$ and $r_0 = 1.2$ fm.

- (a) Derive the expressions of Fermi energies for both neutron and proton gases.
- (b) Find the total kinetic energy of the nucleons when the system is in its ground state.
- (c) Show that the kinetic energy is minimized when $Z = N = A/2$.
- (d) Find the difference of the kinetic energy from its minimum value when the proton number is deviated from $A/2$.

4.(20pts) The half-life of ^{235}U is $7.04 \times 10^8 \text{ y}$.

(a) What is the decay constant of ^{235}U ?

(b) A sample of rock, which solidified with the earth 4.55×10^9 years ago, contains N atoms of ^{235}U . How many ^{235}U atoms did the same rock have at the time it solidified?

5. The quark combinations of following particles are

$$K^0 - d\bar{s} \quad \pi^- - d\bar{u} \quad K^- - s\bar{u} \quad p - uud \quad \Lambda^0 - uds \quad \Sigma^- - dds.$$

(a) Find the charge number Q , the baryon number B and the strangeness S of these particles.

(b) Check the conservations of baryon number, charge and strangeness in the following processes.

(c) Determine if these processes are mediated by the strong or weak interaction, or are forbidden. Explain why.

(1) $\Lambda^0 \rightarrow \pi^- + p$

(2) $\pi^- + p \rightarrow \Lambda^0 + K^0$

(3) $K^- + p \rightarrow \Sigma^- + \pi^+$

(4) $K^- + p \rightarrow \Lambda^0 + K^0$

Quantum numbers of three quarks

Quark symbol	Charge number	Spin	Baryon number	Strangeness
u	$2/3$	$1/2$	$1/3$	0
d	$-1/3$	$1/2$	$1/3$	0
s	$-1/3$	$1/2$	$1/3$	-1


6.(20pts) In the ^{60}Co beta-decay experiment, the electrons emitted in the direction parallel to the applied magnetic field are found to be more than those emitted in the direction anti-parallel to the field. The original process is shown in the figure.

(a) Draw a figure to show the mirror image process of the beta-decay.

(b) Can the mirror image process occur in nature? Why?

(c) What conclusion can you draw if the mirror process can/cannot occur in nature?

Solution: (a)



Mirror image process