

Halo Abundance Matching for eBOSS Galaxies

Jiaxi Yu


May 12, 2020

Background:


Study the Universe:

- **Theories : the framework - LCDM**
- **Observations :** in the redshift space
- **Simulations :** in the real space


Why do we need HAM?




Why do we need HAM?




Why do we need HAM?



Why do we need HAM?



Why do we need HAM?



What is Halo Abundance Matching?

- A method to test the galaxy bias models
- By investigating galaxy distribution in DM halos

density contrast: $\delta(\mathbf{x}) = \frac{\rho(\mathbf{x}) - \bar{\rho}(\mathbf{x})}{\bar{\rho}(\mathbf{x})}$

correlation function: $\xi(\mathbf{s}) = \langle \delta(\mathbf{x})\delta(\mathbf{x} - \mathbf{s}) \rangle$

linear galaxy bias: $b = \frac{\delta_{gal}(\mathbf{x})}{\delta_{DM}(\mathbf{x})} \Rightarrow b^2 = \frac{\xi_{gal}(\mathbf{s})}{\xi_{DM}(\mathbf{s})}$

What is Halo Abundance Matching?

- Select halos that can host a galaxy so that:
 - Selected halos (i.e., galaxies) match empirical distribution
 - Galaxy correlation functions match the observation






Fig 1. Galaxy P.D.F. for halos



How to use HAM?




How to use HAM?



Scattering
with



How to use HAM?



Progress:

Thesis topic:

- Test the galaxy bias models with HAM

The specific question:

- How do galaxies distribute in DM haloes?

Solutions:

- ✓ Select halos to assign galaxies (HAM)
- ✓ Fit the galaxy $\xi_l(r)$ to observations to get bias models (test)

Progress:

- ✓ the Redshift Space Distortion effect (fitting in the redshift space)




Fig 2. The Redshift Space Distortion effects (diagrams from Jean-Paul's AstroIV slides)

Progress:

✓ the Redshift Space Distortion effect (fitting in the redshift space)

$$\xi_l(r, \mu) = a_0 P_0(\mu) + a_2 P_2(\mu) + a_4 P_4(\mu) + \dots \quad (P_n(\mu) \text{ is the Legendre polynomials})$$




Fig 3. The correlation function multipoles of DM halo simulations in the real space (black) and in the redshift space (red)

Progress:

- ✓ Redshift Space Distortion effect
- ✓ Parameters' effects on $\xi_l(r)$ and galaxy distributions




Fig 4. The correlation function multipoles for different scattering parameters

Progress:

- ✓ Redshift Space Distortion effect
- ✓ Parameters' effects on $\xi_l(r)$ and galaxy distributions




Fig 5. The galaxy P.D.F for different model parameters


Progress:

- ✓ Redshift Space Distortion effect
- ✓ Parameters' effects on $\xi_l(r)$ and galaxy distributions
- ✓ χ^2 minimisation for mono+quadru -poles
 - ✓ scripts
 - ✓ statistical instability of HAM in $\xi_l(r)$

Progre

- ✓ Red
- ✓ Para
- ✓ χ^2 Γ_{ξ}^2

itions



Progress:

- ✓ Redshift Space Distortion effect
- ✓ Parameters' effects on $\xi_l(r)$ and galaxy distributions
- ✓ χ^2 minimisation for mono+quadru -poles
 - ✓ scripts
 - ✓ statistical fluctuations of HAM in $\xi_l(r)$




Fig 7. χ^2 - σ relations with different number of random seeds being averaged

Progress:

- ✓ Redshift Space Distortion effect
- ✓ Parameters' effects on $\xi_l(r)$ and galaxy distributions
- ✓ χ^2 minimisation for mono+quadru -poles
 - ✓ scripts
 - ✓ statistical fluctuations of HAM in $\xi_l(r)$

To do:

- Remove the fluctuation completely with more seeds
- Use MCMC sampler to get reliable parameter errors

Outlook:

- ✓ Reliable LRG & ELG bias models
- ❑ Multi-tracer HAM
 - ❑ 'generate' multiple tracers simultaneously
 - ❑ difficulty: overlapped distribution function
- ❑ Cross-Correlation between tracers

Thanks!