



Journal club

Papers to be selected by 18.00 on 18th October; please discuss it with Hubert and I before submitting PR.

Start: Oct 25

Journal Club schedule to be finalized by 25th October

Given the number of students, groups of 2 are recommended.

Use the #journal-clubs channel (e.g., to find a group member). I will put some suggestions there.

Sign up by pull request to the 'material' repo. "First come first served"

18.10.2021	Mark	limma + friends	linear model simulation + design matrices		
25.10.2021	Hubert	RNA-seq quantification	RSEM	X	X
01.11.2021	Mark	edgeR+friends 1	basic edgeR/voom	OUTRIDER: A Statistical Method for Detecting Aberrantly Expressed Genes in RNA Sequencing Data (BT, KN)	Powerful and robust non-parametric association testing for microbiome data via a zero-inflated quantile approach (ZINQ) (RM, DS)
08.11.2021	Mark	edgeR+friends 2	advanced edgeR/voom	ZeitZeiger: supervised learning for high-dimensional data from an oscillatory system (TB, OF)	SnapHiC: a computational pipeline to identify chromatin loops from single-cell Hi-C data (JS NH)
15.11.2021	TBA	hands-on session #1: RNA-seq	FASTQC/Salmon/etc.	Differential abundance testing on single-cell data using k-nearest neighbor graphs (VW,	TedSim: temporal dynamics simulation of single cell RNA-sequencing data and cell division



Journal Club procedure

- During/after journal clubs: give the presenters some constructive feedback
- Giving feedback (via Google form) is part of your JC grade! Feedback forms must be submitted within 1 week of presentation; comments will be sent to presenters (anonymously)
- Note that they will be part of the recordings (tube.switch.ch videos only shared with registered students)

Feedback form: 14.10. Redefining CpG islands using hidden Markov models

Presenters:

* Required

How would you rate the presenters' coverage of the topic? *

- ☐ Poor
- ☐ Fair
- ☐ Good
- ☐ Very Good
- ☐ Excellent

How would you rate the presenters' knowledge of the topic? *

- ☐ Poor
- ☐ Fair
- ☐ Good
- ☐ Very Good
- ☐ Excellent

From the feed: “Over-optimism” + Terry’s IMS Bulletin

We will see a lot of methods in this course – **how do we evaluate what works well in practice ?**

BIOINFORMATICS ORIGINAL PAPER

Vol. 26 no. 16 2010, pages 1990–1998
doi:10.1093/bioinformatics/btq323

Gene expression

Advance Access publication June 26, 2010

Over-optimism in bioinformatics: an illustration

Monika Jelizarow¹, Vincent Guillemot^{1,2}, Arthur Tenenhaus², Korbinian Strimmer³ and Anne-Laure Boulesteix^{1,*}

¹Department of Medical Informatics, Biometry and Epidemiology, University of Munich, Marchioninstr. 15, 81377 Munich, Germany, ²SUPELEC Sciences des Systèmes (E3S)-Department of Signal Processing and Electronics Systems - 3, rue Joliot Curie, Plateau de Moulon, 91192 Gif-sur-Yvette Cedex, France and ³Department of Medical Informatics, Statistics and Epidemiology, University of Leipzig, Härtelstr. 16-18, 04107 Leipzig, Germany

Associate Editor: John Quackenbush

“if the improvement of a quantitative criterion such as the error rate is the main contribution of a paper, the superiority of new algorithms should always be demonstrated on independent validation data.”



EDITORIAL

Ten Simple Rules for Reducing Overoptimistic Reporting in Methodological Computational Research

Anne-Laure Boulesteix*


Institute for Medical Informatics, Biometry and Epidemiology, Ludwig Maximilians University, Munich, Germany

* boulesteix@ibe.med.uni-muenchen.de

REVIEW

Open Access

Essential guidelines for computational method benchmarking

Lukas M. Weber^{1,2}, Wouter Saelens^{3,4}, Robrecht Cannoodt^{3,4}, Charlotte Soneson^{1,2,8}, Alexander Hapfelmeier⁵, Paul P. Gardner⁶, Anne-Laure Boulesteix⁷, Yvan Saeys^{3,4*} and Mark D. Robinson^{1,2*} 





In class discussion

- **5 minutes:** read (the excerpt from “Terence’s Stuff”).
- **10 minutes:** Discuss with your neighbour/row and answer the following 4 questions:
 1. How do we tell what works in practice?
 2. What problems arise using simulated (synthetic) data?
 3. What problems arise using real data?
 4. What are positive/negative controls?
- **n.b. include this (method comparison) context in your Journal Club talks**



limma fundamentals

The simplistic view: Differential expression, small sample inference

- Table of data (e.g., microarray gene expression data with replicates of each of condition A, condition B)
 - rows = features (e.g., genes), columns = experimental units (samples)
- Most common problem in statistical bioinformatics: want to infer whether there is a change in the response
—> a statistical test for each row of the table.

What test might you use? Why is this hard? What issues arise? How much statistical power is there [1] ?


```
> head(y)
      group0 group0 group0 group1 group1 group1
gene1 -0.1874854 0.2584037 -0.05550717 -0.4617966 -0.3563024 -0.03271432
gene2 -3.5418798 -2.4540999 0.11750996 -4.3270442 -5.3462622 -5.54049106
gene3 -0.1226303 0.9354707 -1.10537767 -0.1037990 0.5221678 -1.72360854
gene4 -2.3394536 -0.3495697 -3.47742610 -3.2287093 6.1376670 -2.23871974
gene5 -3.7978820 1.4545702 -7.14796503 -4.0500796 4.7235714 10.00033769
gene6 1.4627078 -0.3096070 -0.26230124 -0.7903434 0.8398769 -0.96822312
```

[1] <http://www.stat.ubc.ca/~rollin/stats/ssize/n2.html>



Microarray expression measures

Two-colour



$$y_{ga} = \log_2(R/G)$$

array

probe or gene

Affymetrix



$$y_{ga} = \text{log-intensity (summarized over probes)}$$

Illumina




$$y_{ga} = \text{log-intensity (summarized over beads)}$$


Normalization: one-colour



Similarly for single channel data, adjustments need to be made for all samples to be comparable.



Quantile normalization



```
x <- rnorm(10000, mean=0, sd=1)
y <- rnorm(10000, mean=3)
z <- cbind(x,y)
```

```
# create "reference" distribution
s <- apply(z,2,sort)
sm <- rowMeans(s)
```

```
# impose ref. distribution by ranks
r <- apply(z,2,rank)
n <- apply(r,2,function(u) sm[u])
```

```
boxplot( data.frame(x=x,y=y,n) )
```

```
#> library(limma)
#> zn <- normalizeQuantiles(z)
#> all(zn==n)
#[1] TRUE
```



Preprocessing: additive + multiplicative error model

Observe intensity for one probe on one array

Intensity = background + signal

$$I = B + S$$

additive
errors

multiplicative errors

This idea underlies variance stabilizing transformations vsn (two colour data) and vst (for Illumina data)




normexp convolution model

$$\text{Intensity} = \text{Background} + \text{Signal}$$

$N(\mu, \sigma^2)$

$\text{Exponential}(\alpha)$



Microarray background correction: maximum likelihood estimation for the normal-exponential convolution

JEREMY D. SILVER

*Bioinformatics Division, Walter and Eliza Hall Institute, Parkville 3050, Victoria, Australia and
Department of Biostatistics, University of Copenhagen, Øster Farimagsgade 5, Entrance B,
PO Box 2099, DK-1014 Copenhagen K, Denmark
j.silver@biostat.ku.dk*

MATTHEW E. RITCHIE

Department of Oncology, University of Cambridge, Cambridge CB2 0RE, UK


GORDON K. SMYTH*

*Bioinformatics Division, Walter and Eliza Hall Institute, Parkville 3050, Victoria, Australia
smyth@wehi.edu.au*



Quality assessments / spot checks

Multidimensional scaling plot



```
sd <- 0.3*sqrt(4/rchisq(1000,df=4))  
x <- matrix(rnorm(1000*6,sd=sd),1000,6)  
x[1:50,4:6] <- x[1:50,4:6] + 2
```

```
mds <- plotMDS(x)
```

```
> round(mds$distance.matrix,3)  
      [,1] [,2] [,3] [,4] [,5] [,6]  
[1,] 0.000 0.000 0.000 0.000 0.00 0  
[2,] 0.835 0.000 0.000 0.000 0.00 0  
[3,] 0.850 0.793 0.000 0.000 0.00 0  
[4,] 1.089 1.068 1.058 0.000 0.00 0  
[5,] 1.050 1.058 1.072 0.863 0.00 0  
[6,] 0.991 1.047 1.046 0.865 0.85 0
```




“To consult the statistician after an experiment is finished is often merely to ask him[her] to conduct a post mortem examination. He[She] can perhaps say what the experiment died of.” R. A. Fisher

Motivation for exploratory data analysis: Case Study

(from Stefano, a former M.Sc. student in my Institute)

He is studying gene expression in fruitfly and is interested in transcriptional responses following “heat shock”.

Basic schematic of experiment:





~4 replicates for each condition



```
library(limma)  
plotMDS(d) # 'd' is a matrix
```

Take a close look at where the replicates are to each other relative to the X- and Y-axes



22 samples x
~20,000 genes

reduced to 22
samples x 2
dimensions




Limma concept: borrowing information across genes


- **Small data sets**: few samples, generally under-powered for 1 gene
- **Curse of dimensionality**: many tests, need to adjust for multiple testing (= loss of power)
- **Benefit of parallelism**: same model is fit for every gene. Can borrow information from one gene to another
 - **Hard**: assume parameters are constant across genes
 - **Soft**: smooth genewise parameters towards a common value in a graduated way, e.g., Bayes, empirical Bayes, Stein shrinkage ...

A very common experiment (1-colour)

Mutant x 2



WT x 2



Gene X



Which genes are differentially expressed?

$n_1 = n_2 = 2$ Affymetrix arrays

~30,000 probe-sets



Ordinary t-tests (1-colour)

$$t_g = \frac{\overline{y}_{\text{mu}} - \overline{y}_{\text{wt}}}{s_g c}$$

give very high false discovery rates

$$c = \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}$$

Residual df = 2



t-tests with common variance

$$t_{g,\text{pooled}} = \frac{\bar{y}_{\text{mu}} - \bar{y}_{\text{wt}}}{s_0 c}$$

with residual standard deviation s_0 pooled
across genes

More stable, but ignores gene-specific variability

$$c = \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}$$

A better compromise


Shrink standard deviations towards common value

$$\tilde{s}_g^2 = \frac{d_0 s_0^2 + d_g s_g^2}{d_0 + d_g}$$


d = degrees of
freedom

Moderated t-statistics

$$\tilde{t}_g = \frac{\bar{y}_{\text{mu}} - \bar{y}_{\text{wt}}}{\tilde{s}_g u}$$



Shrinkage of standard deviations



The **data decides** whether \tilde{t}_g should be closer to $t_{g,pooled}$ or t_g

Why does it work?

- We learn what is the **typical** variability level by looking at all genes, but allow some **flexibility** from this for individual genes
- Adaptive – data (through hyperparameter estimates, d_0 and s_0) suggests how much to “squeeze”/„moderate“ toward common value



Hierarchical model for variances

Data

$$s_g^2 \sim \sigma_g^2 \frac{\chi_{d_g}^2}{d_g}$$

Prior

$$\frac{1}{\sigma_g^2} \sim s_0^2 \frac{\chi_{d_0}^2}{d_0}$$

Posterior

$$E\left(\frac{1}{\sigma_g^2} \mid s_g^2\right) = \frac{d_0 + d_g}{s_0^2 d_0 + s_g^2 d_g}$$

Posterior Statistics

Posterior variance estimators

$$\tilde{s}_g^2 = \frac{s_0^2 d_0 + s_g^2 d_g}{d_0 + d_g}$$

Moderated t-statistics

$$\tilde{t}_{gj} = \frac{\hat{\beta}_{gj}}{\tilde{s}_g \sqrt{c_{gj}}}$$

Baldi & Long 2001, Wright & Simon 2003, Smyth 2004



Exact distribution for moderated t

An unexpected piece of mathematics shows that, under the null hypothesis,

$$\tilde{t}_g \sim t_{d_0 + d_g}$$

The degrees of freedom add!

The Bayes prior in effect adds d_0 extra arrays for estimating the variance.

Wright and Simon 2003, Smyth 2004



Aside: Marginal Distributions to calculate

Under usual likelihood model, s_g is independent of the estimated coefficients.

Under the hierarchical model, s_g is independent of the moderated t-statistics instead

$$s_g^2 \sim s_0^2 F_{d, d_0}$$




Multiple testing and adjusted p-values

- Each statistical test has an associated false positive rate
- Traditional method in statistics is to control family wise error rate, e.g., by Bonferroni.
- Controlling the false discovery rate (FDR) is more **appropriate** in microarray studies
- Benjamini and Hochberg method controls expected FDR for independent or weakly dependent test statistics. Simulation studies support use for genomic data.
- All methods can be implemented in terms of adjusted p-values.

Linear Models

- In general, need to specify:
 - Dependent variable
 - Explanatory variables (experimental design, covariates, etc.)
- More generally:

$$y = X\beta + \epsilon$$



vector of
observed
data

design
matrix


Vector of
parameters to
estimate

Design → Linear models

WT x 2



Mutant x 2



$$\begin{bmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 1 & 0 \\ 1 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix}$$

β_1 = wt log-expression

β_2 = mutant – wt

$$E[y_1] = E[y_2] = \beta_1$$

$$E[y_3] = E[y_4] = \beta_1 + \beta_2$$



Layers to add ..

- Where does the moderated variance come from?
- Why the degrees of freedom add: $d_0 + d$
- empirical Bayes: how to estimate the hyperparameters (d_0 and s_0)
- Design matrices + contrast matrices in practice

Unexpected mathematics: Why do degrees of freedom add?

The construction of the classical t-statistic:

$$Z = (\bar{X}_n - \mu) \frac{\sqrt{n}}{\sigma}$$

$$V = (n - 1) \frac{S_n^2}{\sigma^2}$$

$$T \equiv \frac{Z}{\sqrt{V/\nu}} = (\bar{X}_n - \mu) \frac{\sqrt{n}}{S_n},$$

Bonus Exercise Part a (optional): what are a, b above?

If T is distributed as $(a/b)^{1/2} Z/U$ where $Z \sim N(0, 1)$ and $U \sim \chi_\nu$, then T has density function

$$p(t) = \frac{a^{\nu/2} b^{1/2}}{B(1/2, \nu/2) (a + bt^2)^{1/2 + \nu/2}}$$

Optional exercise: Derive the posterior

Data

$$s_g^2 \sim \sigma_g^2 \frac{\chi_{d_g}^2}{d_g}$$

Prior

$$\frac{1}{\sigma_g^2} \sim s_0^2 \frac{\chi_{d_0}^2}{d_0}$$

Posterior

$$E\left(\frac{1}{\sigma_g^2} \mid s_g^2\right) = \frac{d_0 + d_g}{s_0^2 d_0 + s_g^2 d_g}$$

$$p(\theta|x) = \frac{f(x|\theta)p(\theta)}{\int f(x|\theta)p(\theta)d\theta}$$

Bonus exercise Part b

Sketch: i) Let $x=s^2$, $\theta=\sigma^{-2}$; ii) Using the functional form of chi-squared distribution, calculate only the numerator (since denominator does not contain θ); iii) collect terms and see if you can identify the distribution and the parameters of it; iv) What is the mean of this distribution?

Unexpected mathematics: Why do degrees of freedom add?


$$p(\hat{\beta}, s^2 | \beta = 0) = \int p(\hat{\beta} | \sigma^{-2}, \beta = 0) p(s^2 | \sigma^{-2}) p(\sigma^{-2}) d(\sigma^{-2})$$

The integrand is

$$\begin{aligned} & \frac{1}{(2\pi v \sigma^2)^{1/2}} \exp\left(-\frac{\hat{\beta}^2}{2v\sigma^2}\right) \\ & \times \left(\frac{d}{2\sigma^2}\right)^{d/2} \frac{s^{2(d/2-1)}}{\Gamma(d/2)} \exp\left(-\frac{ds^2}{2\sigma^2}\right) \\ & \times \left(\frac{d_0 s_0^2}{2}\right)^{d_0/2} \frac{\sigma^{-2(d_0/2-1)}}{\Gamma(d_0/2)} \exp\left(-\sigma^{-2} \frac{d_0 s_0^2}{2}\right) \\ & = \frac{(d_0 s_0^2/2)^{d_0/2} (d/2)^{d/2} s^{2(d/2-1)}}{(2\pi v)^{1/2} \Gamma(d_0/2) \Gamma(d/2)} \\ & \quad \sigma^{-2(1/2+d_0/2+d/2-1)} \exp\left\{-\sigma^{-2} \left(\frac{\hat{\beta}^2}{2v} + \frac{ds^2}{2} + \frac{d_0 s_0^2}{2}\right)\right\} \end{aligned}$$

Unexpected mathematics: Why do degrees of freedom add?

$$\begin{aligned}
 p(\hat{\beta}, s^2 \mid \beta = 0) &= \int p(\hat{\beta} \mid \sigma^{-2}, \beta = 0) p(s^2 \mid \sigma^{-2}) p(\sigma^{-2}) d(\sigma^{-2}) \\
 &= \frac{(d_0 s_0^2 / 2)^{d_0/2} (d/2)^{d/2} s^{2(d/2-1)}}{(2\pi v)^{1/2} \Gamma(d_0/2) \Gamma(d/2)} \\
 &\quad \sigma^{-2(1/2+d_0/2+d/2-1)} \exp \left\{ -\sigma^{-2} \left(\frac{\hat{\beta}^2}{2v} + \frac{ds^2}{2} + \frac{d_0 s_0^2}{2} \right) \right\}
 \end{aligned}$$



 σ^{-2} is chi-squared (or gamma)

$$f(x; k) = \begin{cases} \frac{x^{(k/2)-1} e^{-x/2}}{2^{k/2} \Gamma(\frac{k}{2})}, & x \geq 0; \\ 0, & \text{otherwise.} \end{cases}$$

http://en.wikipedia.org/wiki/Chi-squared_distribution



Unexpected mathematics: Why do degrees of freedom add?

$$p(\hat{\beta}, s^2 | \beta = 0) = \int p(\hat{\beta} | \sigma^{-2}, \beta = 0) p(s^2 | \sigma^{-2}) p(\sigma^{-2}) d(\sigma^{-2})$$

$$\begin{aligned} p(\hat{\beta}, s^2 | \beta = 0) \\ = \frac{(1/2v)^{1/2} (d_0 s_0^2/2)^{d_0/2} (d/2)^{d/2} s^{2(d/2-1)}}{D(1/2, d_0/2, d/2)} \left(\frac{\hat{\beta}^2/v + d_0 s_0^2 + d s^2}{2} \right)^{-(1+d_0+d)/2} \end{aligned}$$

Unexpected mathematics: Why do degrees of freedom add?

$$p(\hat{\beta}, s^2 \mid \beta = 0) = \frac{(1/2v)^{1/2} (d_0 s_0^2/2)^{d_0/2} (d/2)^{d/2} s^{2(d/2-1)}}{D(1/2, d_0/2, d/2)} \left(\frac{\hat{\beta}^2/v + d_0 s_0^2 + d s^2}{2} \right)^{-(1+d_0+d)/2}$$

The null joint distribution of \tilde{t} and s^2 is

$$p(\tilde{t}, s^2 \mid \beta = 0) = \tilde{s} v^{1/2} p(\hat{\beta}, s^2 \mid \beta = 0)$$

http://en.wikipedia.org/wiki/Random_variable#Distribution_functions_of_random_variables

$$f_Y(y) = f_X(g^{-1}(y)) \left| \frac{dg^{-1}(y)}{dy} \right|$$

Unexpected mathematics: Why do degrees of freedom add?

If T is distributed as $(a/b)^{1/2}Z/U$ where $Z \sim N(0, 1)$ and $U \sim \chi_\nu$, then T has density function

$$p(t) = \frac{a^{\nu/2} b^{1/2}}{B(1/2, \nu/2) (a + bt^2)^{1/2 + \nu/2}}$$

$$p(\tilde{t}, s^2 | \beta = 0) = \frac{(d_0 s_0^2)^{d_0/2} d^{d/2} s^{2(d/2-1)}}{B(d/2, d_0/2) (d_0 s_0^2 + ds^2)^{d_0/2 + d/2}} \times \frac{(d_0 + d)^{-1/2}}{B(1/2, d_0/2 + d/2)} \left(1 + \frac{\tilde{t}^2}{d_0 + d}\right)^{-(1+d_0+d)/2}$$

This shows that \tilde{t} and s^2 are independent with

$$s^2 \sim s_0^2 F_{d, d_0}$$

and

$$\tilde{t} | \beta = 0 \sim t_{d_0 + d}.$$

Linear Models

- In general, need to specify:
 - Dependent variable
 - Explanatory variables (experimental design, covariates, etc.)
- More generally:

$$y = X\alpha + \epsilon$$

Diagram illustrating the components of the linear model equation $y = X\alpha + \epsilon$:

- y : vector of observed data (indicated by a blue arrow)
- X : design matrix (indicated by a red arrow)
- α : Vector of parameters to estimate (indicated by an orange arrow)

Obtain a linear model for each gene g

$$E(\underline{y}_g) = X\alpha_g$$
$$\text{var}(\underline{y}_g) = W_g^{-1}\sigma_g^2$$

Contrasts -- `contrasts.fit()`

A *contrast* is any linear combination of the coefficients α_j which we want to test equal to zero.

Define contrasts

$$\underset{\sim}{\beta}_g = C^T \underset{\sim}{\alpha}_g$$

where C is the contrast matrix.

Want to test


$$H_0 : \beta_{gj} = 0$$

vs


$$H_a : \beta_{gj} \neq 0$$

Analysis of Variance → Linear model


WT x 2



Cond A x 2



Cond B x 2



$$\begin{bmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \\ y_5 \\ y_6 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \\ 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} \alpha_1 \\ \alpha_2 \\ \alpha_3 \end{bmatrix} + \begin{bmatrix} \epsilon_1 \\ \epsilon_2 \\ \epsilon_3 \\ \epsilon_4 \\ \epsilon_5 \\ \epsilon_6 \end{bmatrix}$$

α_1 = wt log-expression

α_2 = Cond A - wt

α_3 = Cond B - wt


$$E[y_1] = E[y_2] = \alpha_1$$

$$E[y_3] = E[y_4] = \alpha_1 + \alpha_2$$


$$E[y_5] = E[y_6] = \alpha_1 + \alpha_3$$

Analysis of Variance → Linear model, alternative parameterization


WT x 2



Cond A x 2



Cond B x 2



$$\begin{bmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \\ y_5 \\ y_6 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \alpha_1 \\ \alpha_2 \\ \alpha_3 \end{bmatrix} + \begin{bmatrix} \epsilon_1 \\ \epsilon_2 \\ \epsilon_3 \\ \epsilon_4 \\ \epsilon_5 \\ \epsilon_6 \end{bmatrix}$$

$\alpha_1 = \text{wt log-expression}$
 $\alpha_2 = \text{Cond A log-expression}$
 $\alpha_3 = \text{Cond B log-expression}$

$$E[y_1] = E[y_2] = \alpha_1$$

$$E[y_3] = E[y_4] = \alpha_2$$

$$E[y_5] = E[y_6] = \alpha_3$$

An example use of design and contrast matrices

design matrix

$$\begin{bmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \\ y_5 \\ y_6 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \alpha_1 \\ \alpha_2 \\ \alpha_3 \end{bmatrix} + \begin{bmatrix} \epsilon_1 \\ \epsilon_2 \\ \epsilon_3 \\ \epsilon_4 \\ \epsilon_5 \\ \epsilon_6 \end{bmatrix}$$

$$\begin{aligned} E[y_1] &= E[y_2] = \alpha_1 \\ E[y_3] &= E[y_4] = \alpha_2 \\ E[y_5] &= E[y_6] = \alpha_3 \end{aligned}$$

contrast matrix

$$\beta = C\alpha = \begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} \alpha_1 \\ \alpha_2 \\ \alpha_3 \end{bmatrix} = \begin{bmatrix} \alpha_2 - \alpha_1 \\ \alpha_3 - \alpha_2 \end{bmatrix}$$

Contrasts -- `contrasts.fit()`

A *contrast* is any linear combination of the coefficients α_j that we want to test equal to zero.

Define contrasts

$$\beta_g = C^T \alpha_g$$

where C is the contrast matrix.

Want to test

$$H_0 : \beta_{gj} = 0$$

vs

$$H_a : \beta_{gj} \neq 0$$

Limma / Analysis of Variance

$$F = \frac{\text{variance between treatments}}{\text{variance within treatments}}$$

$$F = \frac{MS_{\text{Treatments}}}{MS_{\text{Error}}} = \frac{SS_{\text{Treatments}}/(I-1)}{SS_{\text{Error}}/(n_T - I)}$$

The moderated t -statistics also lead naturally to moderated F -statistics which can be used to test hypotheses about any set of contrasts simultaneously. Appropriate quadratic forms of moderated t -statistics follow F -distributions just as do quadratic forms of ordinary t -statistics. Suppose that we wish to test all contrasts for a given gene equal to zero, i.e., $H_0 : \beta_g = 0$. The correlation matrix of $\hat{\beta}_g$ is $R_g = U_g^{-1}C^TV_gCU_g^{-1}$ where U_g is the diagonal matrix with unscaled standard deviations $(v_{gj})^{1/2}$ on the diagonal. Let r be the column rank of C . Let Q_g be such that $Q_g^TR_gQ_g = I_r$ and let $\mathbf{q}_g = Q_g^T\mathbf{t}_g$. Then

$$F_g = \mathbf{q}_g^T\mathbf{q}_g/r = \mathbf{t}_g^TQ_gQ_g^T\mathbf{t}_g/r \sim F_{r,d_0+d_g}$$

Aside: Marginal Distributions to calculate

Fun fact: Under usual likelihood model, s_g is independent of the estimated coefficients.

Under the hierarchical model, s_g is independent of the moderated t-statistics instead

$$s_g^2 \sim s_0^2 F_{d, d_0}$$

Thus, the set of s_g can be used to estimate d_0 and s_0

Section 6.2 limma paper: other tricks, such as Fisher's z distribution to estimate d_0 and s_0



Relate to limma objects

$$\begin{bmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \\ y_5 \\ y_6 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \alpha_1 \\ \alpha_2 \\ \alpha_3 \end{bmatrix} + \begin{bmatrix} \epsilon_1 \\ \epsilon_2 \\ \epsilon_3 \\ \epsilon_4 \\ \epsilon_5 \\ \epsilon_6 \end{bmatrix}$$

$$E[y_1] = E[y_2] = \alpha_1$$

$$E[y_3] = E[y_4] = \alpha_2$$

$$E[y_5] = E[y_6] = \alpha_3$$

$$\beta = C\alpha = \begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} \alpha_1 \\ \alpha_2 \\ \alpha_3 \end{bmatrix} = \begin{bmatrix} \alpha_2 - \alpha_1 \\ \alpha_3 - \alpha_2 \end{bmatrix}$$

```
> design
  alpha1 alpha2 alpha3
1      1      0      0
2      1      0      0
3      0      1      0
4      0      1      0
5      0      0      1
6      0      0      1
> cont.matrix <- makeContrasts(beta1="alpha2-alpha1",
                               beta2="alpha3-alpha2", levels=design)
> cont.matrix
      Contrasts
Levels  beta1 beta2
alpha1    -1     0
alpha2     1    -1
alpha3     0     1

fit <- lmFit(y, design)

fit.c <- contrasts.fit(fit, cont.matrix)
fit.c <- eBayes(fit.c)

> head(round(y, 2), 3)
      [,1] [,2] [,3] [,4] [,5] [,6]
[1,] -1.62  1.49  2.50  1.57 -0.71  0.38
[2,] -4.50 -4.95 -3.66 -7.83 -1.59  6.94
[3,] -10.17 -21.90 14.03  3.66 -12.21 -15.26

> head(round(fit$coef, 2), 3)
      alpha1 alpha2 alpha3
[1,]  -0.07   2.03  -0.16
[2,]  -4.73  -5.75   2.67
[3,] -16.04   8.85 -13.74

> head(round(fit.c$coef, 2), 3)
      Contrasts
      beta1  beta2
[1,]   2.10  -2.20
[2,]  -1.02   8.42
[3,]  24.89 -22.59
```



Bonus exercise

- Alongside Exercise 5 (if you choose to do it, submit together with Exercise 5)
- Optional!
- Counts up to 3 points towards your Exercise mark