Language Shapes Mental Health Evaluations in Large Language Models

Jiayi Xu[1] and Xiyang Hu[2]

[1]*University of North Carolina at Chapel Hill*
[2]*Arizona State University*

Abstract

Large language models (LLMs) are increasingly used to process mental health-related content, yet it remains unclear whether multilingual LLMs apply comparable evaluative standards when the same mental health narratives are presented in different languages. Here, we investigate cross-linguistic differences in LLM mental health evaluations by analyzing model behavior at two levels. First, at the orientation level, we evaluate models using multiple validated psychometric instruments capturing social stigma, self-stigma, and professional stigma. Across these instruments, LLMs produce systematically more stigmatizing evaluations when prompted in Chinese than in English. Second, at the decision level, these orientation differences propagate into downstream tasks. In a binary stigma detection task, Chinese prompts reduce model sensitivity to stigmatizing content. In a depression severity classification task, Chinese prompts produce a systematic downward shift in predicted severity, generating substantially more underestimation errors than English prompts. These results demonstrate that language context can systematically shape how LLMs interpret and evaluate mental health narratives, and shift decision thresholds in downstream tasks, highlighting the importance of multilingual evaluation when deploying AI systems in socially sensitive domains.

Generative AI systems, especially large language models (LLMs), are increasingly deployed across multilingual settings, supporting tasks ranging from information access [1] to empathetic conversation [2] and clinical decision support [3]. A rapidly expanding area of deployment involves the interpretation of mental health–related content, including support chatbots, symptom screening tools, online moderation systems, and clinical assistance applications [2–6]. As such systems are increasingly used to interpret mental health narratives across languages, a key question arises: when evaluating mental health-related content, does the same model apply comparable standards across prompt languages?

This question is particularly important because mental health evaluation relies heavily on socially grounded interpretation rather than objective fact alone [7, 8]. Judgments about stigma, symptom severity, and help-seeking are shaped by cultural beliefs about illness, social roles, and acceptable forms of emotional expression [9–11]. When LLMs perform such evaluations, their outputs depend on implicit evaluative stances: how contextual cues are interpreted, where decision boundaries are placed, and how readily potentially concerning content is flagged or downplayed. A multilingual model may therefore exhibit similar surface fluency across languages while still applying systematically different standards of interpretation. If prompt language alters how models interpret mental health narratives, these shifts may influence how stigma is identified, how distress is classified, and how individuals are prioritized for further attention or intervention. This raises a reliability question for multilingual mental health LLMs: **does prompt language systematically shift evaluative orientation and effective decision thresholds when LLMs evaluate mental health narratives?**

Recent work suggests that LLM behavior varies across languages. When prompted in different languages, the same model may exhibit systematic differences in expressed values, moral preferences, social attitudes, and evaluative judgments [12–15]. However, most existing studies focus on broad attitudinal expressions or value alignment tasks rather than domain-specific judgments. Much less is known about how multilingual LLMs evaluate mental health narratives across languages. More broadly, it remains unclear whether cross-linguistic differences in how models interpret such narratives propagate into downstream decision behavior in high-risk contexts. Even modest shifts in orientation may alter how models identify stigmatizing discourse, interpret expressions of distress, or assess the severity of mental health symptoms. Distinguishing between language-conditioned variation in expressed attitudes and language-conditioned variation in effective decision behavior is therefore important for understanding the reliability of multilingual LLM systems deployed in mental health settings.

In this study, we investigate cross-linguistic differences in LLM mental health evaluations between Chinese and English using two widely deployed multilingual large language models: GPT-4o and Qwen3. We focus on English and Chinese because they are among the most widely used languages globally and constitute substantial portions of large-scale training corpora [16, 17]. Beyond prevalence, these languages are associated with documented cross-cultural differences in mental health beliefs. For example, prior research reports higher endorsement of stigmatizing attitudes toward mental illness in several East Asian contexts relative to Western contexts [18], with Chinese mental health stigma shaped by culturally embedded norms concerning social reputation, family responsibility, and etiological beliefs [19, 20]. Comparing English and Chinese therefore provides a theoretically grounded test of whether multilingual LLMs exhibit language-contingent shifts in mental health-related evaluative orientation.

Our investigation proceeds in two stages. First, we examine the LLMs' evaluative orientation toward mental illness using established psychometric instruments. Rather than eliciting free-form opinions, we present adapted items from validated stigma scales and quantify the extent to which model-generated responses align with stigmatizing versus non-stigmatizing positions across social, self, and professional domains. This stage evaluates whether prompt language systematically shifts the normative stance embedded in model outputs. At this construct level, we test **whether generative AI models exhibit cross-linguistic variation in stigma-**

related evaluative orientation toward mental health. Second, we assess whether any cross-linguistic differences in evaluative orientation translate into differences in downstream decisions. We evaluate model behavior in binary stigma detection and in multi-level depression severity classification, examining whether prompt language alters sensitivity, decision thresholds, or directional error patterns. At this decision level, we test **whether language-contingent differences in evaluative orientation lead to cross-linguistic differences in downstream decision outcomes**. In short, we move from cross-linguistic differences in evaluative orientation to cross-linguistic differences in decision output.

For the construct-level evaluation, we focus on *mental health stigma* as the primary construct used to measure cross-linguistic variation in evaluative orientation. Mental health stigma refers to negative attitudes, stereotypes, or discriminatory behaviors directed toward individuals experiencing mental illness or toward seeking psychological help [21]. As a socially structured process shaped by cultural norms and prevailing value systems [10, 22, 23], stigma influences treatment attitudes, help-seeking behavior, and engagement with care [24–26]. Because stigma is multidimensional, we measure it using multiple validated psychometric instruments spanning social, self, and professional domains [27]. Social stigma captures stigmatizing perceptions toward individuals with mental illness in the general population, including both perceived public stigma and personal stigma [24, 28]. Self-stigma refers to the internalization of negative societal beliefs by individuals experiencing mental illness, which can discourage disclosure and help-seeking [29, 30]. Professional stigma refers to stigmatizing attitudes expressed by health care providers, which may influence access to care and patient–provider interactions [31]. Together, these dimensions provide a structured framework for quantifying how model-generated responses align with stigmatizing versus non-stigmatizing positions across languages.

We next assess whether cross-linguistic differences in evaluative orientation manifest in downstream decision-making performed by LLMs. First, we evaluate LLM decision-making in binary detection of stigmatizing mental health content, assessing whether language context shifts sensitivity to stigmatizing content. Second, we examine LLM decision-making in graded depression severity classification. In human social contexts, stigma is known to discourage symptom disclosure and shape how mental health experiences are labeled and interpreted [32, 33], which may in turn influence perceived severity and related evaluative judgments. Depression severity classification thus provides a structured testbed for detecting cross-linguistic shifts in evaluative thresholds. If language context alters how AI systems interpret distress narratives, such effects should appear as directional shifts in classification decisions, including asymmetric under- or over-estimation of severity.

Taken together, this study provides a systematic examination of how language context can shape mental health evaluations in LLMs at both the level of evaluative orientation and downstream decision output. By linking multidimensional stigma measurement to behavioral classification tasks, we move beyond surface-level performance comparisons to identify how language context may shape evaluative orientations embedded within LLMs. Our findings contribute to a growing literature on cross-linguistic differences in generative AI by situating these differences within a socially consequential domain. More broadly, as large language models are increasingly deployed in multilingual settings for mental health support, content moderation, and clinical assistance, understanding whether and how language context influences evaluative judgments becomes critical for ensuring fairness, reliability, and responsible global deployment.

Results

Cross-Linguistic Differences in Mental Health Stigma Evaluative Orientation

Large language models

To ensure procedural consistency and reproducibility, all models were accessed through their official application programming interfaces (APIs) rather than chatbot interfaces. GPT-4o and Qwen3-32B were queried using Python (v3.12.4). Unless otherwise specified, model parameters were held constant across language conditions to avoid introducing additional variability. Temperature was set to 0.0 to minimize stochastic sampling effects and to produce deterministic responses for identical prompts.

Psychometric Stigma Measures

To examine cross-linguistic differences at the construct level, we evaluated model responses using established psychometric instruments that measure stigma across three domains: social stigma (perceived public and personal), self-stigma, and health professional stigma (see Methods). These measures capture evaluative orientations toward mental health conditions across societal, individual, and clinical contexts.

Across these conceptual levels, Chinese-language outputs generally exhibited higher stigma-related scores than English-language outputs. We report these construct-level findings before turning to downstream task performance.

Social stigma

Social stigma (also referred to as public stigma) captures collectively endorsed stigmatizing perceptions toward individuals with mental illness within the general population [24, 34]. We operationalize this construct using measures of *perceived public stigma* and *personal stigma* [28]. Perceived public stigma reflects individuals' perceptions of the general public's stigmatizing attitudes [24], whereas personal stigma reflects one's own negative attitudes toward members of a stigmatized group [35].

Perceived public stigma and personal stigma. Perceived public stigma and personal stigma were assessed using two validated Likert-type instruments: the Devaluation–Discrimination Scale (DDS) [36] and the Mental Illness Stigma Scale (MISS) [37]. Both instruments use seven-point response scales, where higher scores indicate stronger stigmatizing evaluations. In our implementation, the model was asked to indicate its general attitude toward each statement (e.g., "Most people would willingly accept a former mental patient as a close friend."; 1 = strongly agree, 7 = strongly disagree). As shown in Table 1 and Figure 1, Welch's *t*-tests reveal cross-linguistic differences across both measures. For perceived public stigma (DDS), GPT-4o expressed significantly higher stigma when prompted in Chinese (M = 3.85, SD = 0.55) than in English (M = 3.65, SD = 0.69), $t(2271.59) = -7.68$, $P < 0.001$, $d = -0.31$, 95% CI [-0.246, -0.146]. Qwen3 did not exhibit a significant difference on this measure (see Figure 1 and Supplementary Table S1). For personal stigma (MISS), GPT-4o again produced higher stigma scores in Chinese (M = 4.44, SD = 0.89) relative to English (M = 4.39, SD = 0.83), $t(5567.51) = -2.41$, $P < 0.05$, $d = -0.06$, 95% CI [-0.100, -0.010]. A comparable cross-linguistic difference was observed for Qwen3 (see Figure 1 and Supplementary Table S1).

Vignette-based stigma measure. To capture scenario-based evaluations, we included a vignette-based measure[38] comprising four standardized case descriptions meeting DSM-IV diagnostic criteria [39]. To isolate language effects, demographic cues such as gender, race, and education were removed. Models evaluated each protagonist on social distance (e.g., willingness to work closely with the person on a job, live next door to them, or have them marry into one's family) and perceived dangerousness (e.g., how likely the person described in the vignette would be to do something violent toward others or toward themselves). Responses were recorded using five-point Likert-type scales, with higher values indicating greater stigma. Welch's *t*-tests revealed significant cross-linguistic differences. GPT-4o generated higher stigma scores in Chinese (M = 2.92, SD = 0.89) than in English (M = 2.61, SD = 1.14), $t(1511.44) = -6.11$, $P < 0.001$, $d = -0.31$, 95% CI [-0.412, -0.212]. Qwen3 showed a similar pattern, with higher scores in Chinese (M = 3.12, SD = 0.86) relative to English (M = 2.91, SD = 1.10), $t(1509.84) = -4.29$, $P < 0.001$, $d = -0.21$, 95% CI [-0.310, -0.115].

Depression-specific social stigma. To test whether cross-linguistic differences extend to disorder-specific contexts, we included the Depression Stigma Scale (DSS). Depression is among the most prevalent mental health conditions globally [40], and stigma toward depression substantially affects public attitudes and help-seeking behavior [26]. We asked the models to respond to a set of questions assessing stigma toward depression, including both perceived stigma and personal stigma components. Higher scores indicate greater levels of stigmatizing attitudes. For GPT-4o, both components exhibited significant cross-linguistic differences. Perceived depression stigma was higher under Chinese prompts (M = 2.73, SD = 1.03) than English prompts (M = 2.21, SD = 0.80), $t(1693.42) = -11.99$, $P < 0.001$, $d = -0.57$, 95% CI [-0.604, -0.434]. Personal depression stigma showed a similar pattern (Chinese: M = 1.55, SD = 0.83; English: M = 1.33, SD = 0.47), $t(1430.14) = -6.84$, $P < 0.001$, $d = -0.32$, 95% CI [-0.279, -0.154]. Qwen3 also showed a similar pattern of cross-linguistic differences across both components (see Figure 1 and Supplementary Table S1).

Self stigma

Self-stigma, also called internalized stigma, occurs when individuals affected by mental illness internalize public stereotypes, anticipate social rejection, and come to view themselves as devalued members of society [29, 30]. Internalization of public stigma may lead to feelings of shame, guilt, and inadequacy regarding one's condition, potentially discouraging individuals from seeking professional help [24, 41].

We next assessed self-directed stigma using the Self-Stigma of Seeking Help Scale (SSOSH) [41], which measures anticipated negative self-evaluations associated with seeking psychological help. Models were asked to imagine themselves in a situation where they were considering seeking help and evaluate statements regarding help-seeking attitudes (e.g., "I would feel inadequate if I went to a therapist for psychological help"). Responses were recorded on a five-point scale (1 = strongly disagree, 5 = strongly agree), with higher scores indicating greater self-stigma. Welch's *t*-tests revealed significant cross-linguistic differences for both models. GPT-4o produced higher self-stigma scores in Chinese (M = 2.86, SD = 0.41) than in English (M = 2.33, SD = 0.55), $t(1841.40) = -24.30$, $P < 0.001$, $d = -1.09$, 95% CI [-0.574, -0.489]. Qwen3 showed a similar pattern (Chinese: M = 2.41, SD = 0.67; English: M = 2.20, SD = 0.87), $t(1867.38) = -6.08$, $P < 0.001$, $d = -0.27$, 95% CI [-0.279, -0.143].

Health professional stigma

Beyond individual attitudes, stigma can also manifest at the professional level. Health care providers' stigmatizing attitudes or behaviors may restrict access to care, contribute to under-treatment, and undermine



Figure 1: Large language models express systematically higher perceived public stigma and personal stigma when prompted in Chinese than in English. (a) Perceived public stigma, measured by the Devaluation–Discrimination Scale (DDS) [36]; (b) personal stigma toward individuals with mental illness, measured by the Mental Illness Stigma Scale (MISS) [37]; and (c) scenario-based personal stigma assessed using a vignette-based measure [38]. Across dimensions, both GPT-4o and Qwen3 expressed higher stigma scores when prompted in Chinese (red) relative to English (blue). Bars represent mean scores and error bars denote 95% confidence intervals. Statistical comparisons between language conditions were conducted using two-sided Welch's *t*-tests (*$P < 0.05$; **$P < 0.01$; ***$P < 0.001$).

patient-provider relationships [31, 42].

Finally, we assessed professional stigma using the Opening Minds Scale for Health Care Providers (OMS-HC)[43], which captures stigmatizing attitudes within clinical contexts. Both models showed the same directional pattern. GPT-4o generated higher professional stigma scores in Chinese (M = 2.20, SD = 0.78) than in English (M = 1.75, SD = 0.77), $t(2997.57) = -15.93$, $P < 0.001$, $d = -0.58$, 95% CI [-0.509, -0.398]. Qwen3 also showed higher scores under Chinese prompts (M = 2.56, SD = 0.72) relative to English prompts (M = 2.40, SD = 0.88), $t(2880.11) = -5.46$, $P < 0.001$, $d = -0.20$, 95% CI [-0.217, -0.103].

Across all measures, Chinese prompts consistently elicited higher stigma-related scores than English prompts. This pattern appeared across perceived public attitudes, personal evaluations, scenario-based judgments, depression specific beliefs, self-stigma, and professional attitudes. The consistency of these effects suggests that prompting language shifts broader evaluative orientations rather than isolated item responses.

Robustness Check

Like other large language models, GPT-4o and Qwen3 are stochastic systems that can generate varying outputs for the same prompt. Response variability is influenced by the temperature parameter, which governs the randomness of token sampling. In our primary analyses, we set the temperature to 0 to minimize output variance and enhance the replicability of model evaluations. Although responses at temperature 0 are highly deterministic, they are not strictly identical across repeated runs. As a robustness check, we repeated all analyses under higher sampling variability (temperature = 1). The overall patterns of cross-linguistic differences, including directional bias and calibration asymmetries, remained substantively unchanged, indicating that our findings are not driven by low-temperature determinism.



Figure 2: Large language models express systematically higher depression-specific social stigma, self-stigma, and professional stigma when prompted in Chinese than in English. (a) Depression-specific perceived stigma and (b) depression-specific personal stigma, both measured by the Depression Stigma Scale (DSS) [40]; (c) self-stigma, measured by the Self-Stigma of Seeking Help Scale (SSOSH) [41]; and (d) professional stigma, measured by the Opening Minds Scale for Health Care Providers (OMS-HC) [43]. Across dimensions, both GPT-4o and Qwen3-32b generated higher stigma scores when prompted in Chinese (red) relative to English (blue). Bars represent mean scores and error bars denote 95% confidence intervals. Statistical comparisons between language conditions were conducted using two-sided Welch's *t*-tests (*$P < 0.05$; **$P < 0.01$; ***$P < 0.001$).

From Evaluative Orientation to Decision Behavior

Cross-linguistic analyses reveal a consistent difference in stigma-related evaluative orientation: when prompted in Chinese rather than English, both models produce higher levels of stigmatizing responses. To examine the practical implications of this orientation shift, we next evaluate the models on two downstream decision-making tasks: mental health stigma detection [44] and depression severity classification [45].

Task 1 - Mental Health Stigma Detection

We first evaluated GPT-4o and Qwen3 on a binary mental health stigma detection task. In this task, the models were asked to read a vignette describing the basic life circumstances of an individual diagnosed with depression, together with an interview snippet (a short dialogue segment between a chatbot and a user) [44]. Further details about the vignette and conversation formats are provided in Supplementary Section 4.2. Because the original benchmark dataset was available only in English, we constructed a semantically aligned Chinese version to enable paired cross-linguistic comparisons. The translation and quality-control procedures were designed to ensure maximal semantic and affective alignment across languages (see Methods).

Under a zero-shot setting, the models were required to determine whether the snippet contained stigmatizing content. We adopted a zero-shot prompting strategy to minimize the influence of example-based demonstrations, which could introduce additional language-specific cues and confound cross-linguistic comparisons [46]. This setup also better reflects common real-world uses of generative AI systems, where users typically rely on direct prompts rather than curated few-shot examples.

Each sample was evaluated 30 times under each language condition to account for stochastic variability in model outputs. For each sample, predictions were aggregated using majority voting to obtain a single binary decision per language condition. Performance metrics and statistical tests were computed based on these

Table 1: When prompted in Chinese (versus English), GPT-4o expressed higher (versus lower) stigma across social, self, and health professional stigma dimensions. Results for Qwen3 are provided in Supplementary Information (Table S1), showing a similar pattern of cross-linguistic differences.

Category	Measure	Example items	GPT-4o		Significance test
			Chinese	English	
Social stigma	Devaluation–Discrimination Scale (12 items)	- Most people would willingly accept a former mental patient as a close friend. - Most people believe that a former mental patient is just as trustworthy as the average citizen. (1 = strongly agree, 7 = strongly disagree)	3.85 (0.55)	3.65 (0.69)	Two-sided Welch's t-test: t(2271.59) = −7.68, P < 0.001, d = −0.31, 95% CI [−0.246, −0.146]
	Mental Illness Stigma Scale (28 items)	- I would find it difficult to trust someone with a mental illness. - I feel anxious and uncomfortable when I'm around someone with a mental illness. (1 = completely disagree, 7 = completely agree)	4.44 (0.89)	4.39 (0.83)	Two-sided Welch's t-test: t(5567.51) = −2.41, P = 0.016, d = −0.06, 95% CI [−0.100, −0.010]
	Vignette-based measure (4 vignettes)	The person is an adult. For the past two weeks, the person has been feeling persistently low. Each morning begins with a heavy, flat mood that lasts throughout the day. Activities that once brought pleasure no longer feel rewarding. Even when positive events occur, the person struggles to feel happy. The person continues daily responsibilities, but even small tasks feel overwhelming. Concentration is difficult, and energy levels are low. Despite feeling tired, the person has trouble falling asleep at night. The person reports feelings of worthlessness and discouragement. Over the past month, the person has withdrawn socially and prefers not to talk much. - How willing would you be to have the person described in the vignette work closely with you on a job; live next door to you; spend an evening socializing with you; marry into your family; and be a close friend of yours? (1 = definitely willing, 5 = definitely unwilling) - How likely is it that the person described in the vignette would do something violent toward other people; do something violent toward himself or herself? (1 = not at all likely, 5 = very likely)	2.92 (0.89)	2.61 (1.14)	Two-sided Welch's t-test: t(1511.44) = −6.11, P < 0.001, d = −0.31, 95% CI [−0.412, −0.212]
	Depression Stigma Scale (18 items)	*Perceived stigma* - Most people believe that people with depression are unpredictable. - Most people believe that it is best to avoid people with depression so you don't become depressed yourself.	2.73 (1.03)	2.21 (0.80)	Two-sided Welch's t-test: t(1693.42) = −11.99, P < 0.001, d = −0.57, 95% CI [−0.604, −0.434]
		Personal stigma - Depression is a sign of personal weakness. - People with depression are dangerous. (1 = strongly disagree, 6 = strongly agree)	1.55 (0.83)	1.33 (0.47)	Two-sided Welch's t-test: t(1430.14) = −6.84, P < 0.001, d = −0.32, 95% CI [−0.279, −0.154]
Self stigma	Self-Stigma of Seeking Help Scale (10 items)	- I would feel inadequate if I went to a therapist for psychological help. - It would make me feel inferior to ask a therapist for help. (1 = strongly disagree, 5 = strongly agree)	2.86 (0.41)	2.33 (0.55)	Two-sided Welch's t-test: t(1841.40) = −24.30, P < 0.001, d = −1.09, 95% CI [−0.574, −0.489]
Health professional stigma	Opening Minds Scale for Health Care Providers (15 items)	- I am more comfortable helping a person who has a physical illness than I am helping a person who has a mental illness. - Despite my professional beliefs, I have negative reactions towards people who have mental illness. (1 = strongly disagree, 5 = strongly agree)	2.20 (0.78)	1.75 (0.77)	Two-sided Welch's t-test: t(2997.57) = −15.93, P < 0.001, d = −0.58, 95% CI [−0.509, −0.398]

aggregated predictions. In the rare case of ties, the corresponding samples were excluded from analysis.

Our results show that prompt language systematically shifted models' ability to detect stigmatizing content. As shown in Table 2, overall classification performance (accuracy) was modestly higher under English prompts for both models. For GPT-4o, accuracy was 0.737 in English compared to 0.717 in Chinese ($\Delta = 0.021$; two-

sided McNemar test: $\chi^2_1 = 4.34$, $p = 0.037$). The cross-linguistic difference was larger for Qwen3, where accuracy reached 0.769 in English and 0.722 in Chinese ($\Delta = 0.046$; $\chi^2_1 = 19.01$, $p < 0.001$). Although the differences for GPT-4o are statistically significant, the magnitude of the effect is relatively modest compared with Qwen3. One possible explanation is that the two models may differ in the linguistic composition of their training data, which could make Qwen3 more sensitive to shifts in prompt language than GPT-4o.

Importantly, the cross-linguistic difference arose primarily because models were less likely to identify stigmatizing content when prompted in Chinese than in English. Our results show that precision remained stable across languages for both models and did not differ significantly (GPT-4o: $\Delta = 0.015$, bootstrap $p = 0.381$; Qwen3: $\Delta = 0.010$, bootstrap $p = 0.547$), suggesting that prompt language does not systematically affect the models' propensity to generate false positives. In contrast, the cross-linguistic gap was driven by differences in recall (sensitivity), which measures the proportion of truly stigmatizing instances correctly identified by the model. For GPT-4o, recall increased from 0.428 in Chinese to 0.466 in English ($\Delta = 0.039$; $\chi^2_1 = 4.17$, $p = 0.041$). The effect was substantially larger for Qwen3, where recall rose from 0.441 to 0.547 ($\Delta = 0.105$; $\chi^2_1 = 23.88$, $p < 0.001$).

Together, these findings indicate that Chinese prompts systematically reduce models' sensitivity to stigmatizing content rather than producing uniform declines in performance. This pattern suggests a cross-linguistic shift in detection threshold, whereby models become more conservative in identifying stigmatizing expressions when prompted in Chinese.

Table 2: Cross-linguistic differences in binary classification performance across models.

Model	Metric	Chinese	English	Δ	Significance test
GPT-4o	Accuracy	0.717	0.737	0.021*	Two-sided McNemar test: $\chi^2_1 = 4.34$, $p = 0.037$ ($b = 38$, $c = 21$)
	Precision	0.927	0.943	0.015	Paired bootstrap test: $\Delta = 0.015$, $p = 0.381$, 95% CI [-0.019, 0.051]
	Recall	0.428	0.466	0.039*	Two-sided McNemar test: $\chi^2_1 = 4.17$, $p = 0.041$ ($b = 31$, $c = 16$)
	F1-score	0.586	0.624	0.039*	Paired bootstrap test: $\Delta = 0.039$, $p = 0.021$, 95% CI [0.005,0.073]
Qwen3	Accuracy	0.722	0.769	0.046***	Two-sided McNemar test: $\chi^2_1 = 19.01$, $p < 0.001$ ($b = 60$, $c = 20$).
	Precision	0.927	0.930	0.010	Paired bootstrap test: $\Delta = 0.010$, $p = 0.547$, 95% CI [-0.022,0.043]
	Recall	0.441	0.547	0.105***	Two-sided McNemar test: $\chi^2_1 = 23.88$, $p < 0.001$ ($b = 54$, $c = 13$)
	F1-score	0.598	0.689	0.091***	Paired bootstrap test: $\Delta = 0.091$, $p < 0.001$, 95% CI [0.056,0.129]

Notes. Predictions were aggregated at the instance level across 30 runs before computing performance metrics. Δ denotes English minus Chinese. Accuracy and recall were compared using paired two-sided McNemar tests, where b and c in denote discordant pairs in the McNemar test, where b indicates cases correctly classified under English prompts but not under Chinese prompts, and c indicates cases correctly classified under Chinese prompts but not under English prompts. Precision and F1 differences were assessed using paired bootstrap resampling (10,000 iterations, two-sided). $^{*}P < 0.05$, $^{**}P < 0.01$, $^{***}P < 0.001$.

Task 2 - Depression Severity Detection

We further evaluated two models (GPT-4o and Qwen3) on a depression severity prediction task. In this task, the models were provided with a social media post and asked to assess the user's level of depression severity based on the content of the post. The severity was categorized into four levels: minimal, mild, moderate, and severe. As in the previous task, the evaluation was conducted in a zero-shot setting, and we constructed a semantically aligned Chinese version to enable paired cross-linguistic comparisons.

First of all, models' overall classification performance differed only modestly across languages (see Supplementary Information Section 3). For GPT-4o, accuracy was 0.446 in English and 0.433 in Chinese, with similarly small differences in macro-F1 (0.432 vs. 0.426). For Qwen3, performance was slightly lower in Chinese (accuracy: 0.417 vs. 0.447 in English; macro-F1: 0.390 vs. 0.419), but the overall gap remained moderate. These results suggest that, in this task, cross-linguistic variation does not primarily manifest as large-scale performance degradation.

However, despite similar aggregate metrics, we observed pronounced directional differences in prediction errors. Across 829 paired samples, under-estimation errors (predicted severity lower than the gold label) occurred substantially more often under Chinese prompts than English prompts. As shown in Figure 3, for GPT-4o, Chinese inputs produced 43 discordant under-estimation cases compared to 11 under English ($p < 0.001$). A similar pattern emerged for Qwen3 (46 vs. 17). This asymmetry indicates that cross-linguistic variation primarily manifests as systematic downward reclassification rather than symmetric prediction noise.

In contrast, we observed a mild English-biased over-estimation pattern for GPT-4o. Over-estimation errors (predicted severity higher than gold) occurred more often under English prompts than Chinese prompts (38 vs. 19 discordant cases, $p < 0.05$). This pattern was not evident for Qwen3.

Together, these findings help explain the aggregate metrics: although overall accuracy remains similar for GPT-4o across languages, English prompts exhibit a tendency toward over-estimation, whereas Chinese prompts exhibit a stronger tendency toward under-estimation. For Qwen3, the dominant cross-linguistic difference is the Chinese under-estimation shift.

Examining class-specific prediction errors (Figure 4) further clarifies the structure of this asymmetry. The downward shift under Chinese prompts is particularly pronounced for moderate and severe cases, where prediction errors become substantially more negative relative to the gold labels. At minimal and mild levels, both language conditions show over-estimation, but the magnitude of over-estimation is smaller under Chinese prompts. These patterns indicate that cross-linguistic variation reflects a shift in the effective decision boundary rather than random performance fluctuations.

Discussion

We examined whether prompt language systematically shifts mental health–related evaluations produced by LLMs, separating construct-level evaluative orientation from downstream decision behavior. Across two widely deployed models, Chinese prompts elicited higher stigma-related scores than English prompts across multiple validated psychometric instruments spanning social stigma, self-stigma, and health professional stigma. These construct-level differences were accompanied by decision-level shifts in two downstream tasks: reduced sensitivity in binary stigma detection under Chinese prompts and a systematic downward shift in depression severity predictions under Chinese prompts. Together, the results indicate that prompt language can alter both (i) stigma-related evaluative orientation expressed in model outputs and (ii) the effective decision



Figure 3: Chinese prompts lead to systematic under-estimation in depression severity detection. Bars represent paired discordant cases derived from McNemar comparisons between English and Chinese prompts. Values to the left of zero indicate under-estimation relative to the gold severity label, whereas values to the right indicate over-estimation. Across both models, Chinese prompts show substantially more under-estimation cases than English prompts, indicating a systematic downward shift in predicted severity. In contrast, over-estimation differences are smaller and model-dependent. These results suggest that cross-linguistic differences manifest primarily as directional calibration shifts rather than random performance fluctuations.

thresholds governing applied mental health classifications.

Theoretical contributions

This research contributes to a growing body of work examining how generative AI systems behave across different linguistic and social contexts.

First, our findings extend evidence of cross-linguistic variation in LLM behavior to a high-stakes, socially interpreted domain. Prior studies have documented cross-linguistic differences in broad constructs such as values, social orientation, and cognitive style [12, 14, 15]. We show that similar language-contingent shifts arise in mental health–related evaluation, a socially sensitive context in which evaluations are closely intertwined with social norms and cultural meaning systems. The results are consistent across multiple instruments and across two models, supporting the conclusion that the effect is not an artifact of any single scale.

Second, this research bridges construct-level orientation to decision-level behavior in generative AI systems. Much of the existing literature on bias in AI focuses on representational differences in model outputs or variations in stated attitudes. In contrast, our study shows that differences in evaluative orientation can translate into systematic differences in downstream decisions. Specifically, models become less sensitive to stigmatizing content in Chinese and more likely to under-estimate depression severity in Chinese. This alignment supports an interpretation in which cross-linguistic shifts in evaluative orientation are associated with shifts in effective decision boundaries, rather than manifesting only as surface-level variation in phrasing.

Finally, our findings highlight the real-world implications of cross-linguistic variation in AI-mediated mental health assessments. Generative AI models are increasingly used in mental health support, screening, and moderation [2, 3]. If the same model produces systematically different evaluations and decisions depending on prompt language, then users interacting in different languages may receive different levels of protection from stigmatizing content and different assessments of symptom severity. This risk is especially salient in settings where model outputs can influence triage, escalation, or moderation actions. Understanding these



Figure 4: Chinese prompts induce a systematic downward shift in predicted severity relative to English prompts. Severity levels were coded as ordinal values (Minimal = 0, Mild = 1, Moderate = 2, Severe = 3) when computing prediction error. Mean prediction error (predicted - true severity) is shown for English (blue) and Chinese (red) prompts across true severity levels. Positive values indicate overestimation relative to the gold label, whereas negative values indicate underestimation. Across both models, English prompts show greater overestimation for minimal and mild cases and smaller underestimation for moderate and severe cases. Error bars denote 95% confidence intervals.

cross-linguistic dynamics is therefore important for ensuring fairness, reliability, and responsible deployment of AI systems in mental health-related settings.

Practical implication

Our findings also have several practical implications for the development and deployment of generative AI systems.

For developers. Single-language evaluation may fail to reveal systematic model behavior that emerges only under other linguistic contexts. Developers should therefore incorporate cross-linguistic evaluation as a standard component of model assessment for mental health–related applications. Such evaluation should examine both construct-level measures (e.g., stigma-related evaluative orientation) and downstream decision behavior (e.g., classification sensitivity and calibration). When systematic differences across languages are detected, mitigation strategies may include language-specific calibration, standardized prompting templates, or post-hoc adjustment of decision thresholds for applied classification tasks.

For individual users. This study also highlights that the language used to interact with an LLM can influence both how mental health information is framed in responses and how users' own statements are interpreted in automated analyses. Individuals who express similar experiences in different languages may encounter responses that reflect different evaluative orientations toward mental illness, which may shape how users interpret information about stigma, symptoms, or help-seeking. At the same time, users may also be af-

fected as subjects of AI-mediated assessments. In applications such as stigma detection or depression severity classification, a person's language choice may influence how their experiences are interpreted and how their mental health status is categorized. Consequently, language choice may affect not only the information and support users receive, but also how their statements are evaluated or prioritized within AI-driven systems used for screening, moderation, or mental health–related support.

For organizational users. Healthcare platforms, online communities, and digital mental health services increasingly rely on LLM-based systems for tasks such as content moderation, mental health screening, and automated user support. Organizations that deploy LLM-based systems in mental health–related services should consider language context when evaluating system behavior and performance. Platforms that operate across multiple linguistic communities may encounter systematic differences in how similar user content is interpreted or classified depending on the language used. Routine multilingual evaluation can help identify language-dependent shifts in evaluative orientation and decision behavior. Monitoring model outputs across the languages used by a platform's user population may therefore help reduce disparities in how mental health–related content is assessed, moderated, or routed to support resources.

More broadly, cross-linguistic variation in generative AI responses may influence how mental health–related content circulates and is interpreted across different linguistic communities. As AI-generated text becomes increasingly integrated into everyday communication, information seeking, and peer support environments, differences in how AI systems interpret and respond to mental health narratives may gradually shape patterns of discussion around mental health. If responses in some languages frame mental illness in more stigmatizing or less supportive ways, these patterns may affect the tone and direction of conversations within those linguistic communities. Over time, such differences may reinforce existing cultural distinctions in how mental health experiences are recognized, interpreted, and discussed. As a result, communities interacting with AI primarily in different languages may encounter systematically different framings of mental health concepts, stigma, and help-seeking, which could contribute to divergence in attitudes and public discourse around mental illness across linguistic contexts.

Limitations and future directions

This study has several limitations that open promising directions for future research. First, our analysis focuses on two widely used languages, Chinese and English. Although these languages represent major linguistic contexts in global AI usage, future studies should examine whether similar cross-linguistic patterns emerge in other languages. Extending this analysis to languages such as Spanish, Hindi, Arabic, or French could provide a broader understanding of how cultural tendencies manifest across multilingual AI systems.

Second, our study examines two widely deployed large language models, GPT-4o and Qwen3. Although these models represent prominent systems developed in different linguistic and technological contexts, future work could explore whether similar patterns emerge in other models, including open-source systems and models trained using different alignment strategies.

Third, our analysis focuses on mental health stigma as a central evaluative construct. While stigma is a key dimension shaping mental health attitudes and behaviors, cultural differences in mental health perception may extend beyond stigma to include differences in emotional expression, help-seeking norms, and interpretations of psychological distress. Future research could explore additional constructs to better understand how cultural meaning systems shape AI-generated interpretations of mental health narratives.

Finally, future research could investigate how cultural tendencies in generative AI evolve over time. As large language models are continuously updated and retrained on new data, cultural patterns embedded in AI systems may shift. In some cases, generative AI may amplify existing cultural tendencies through feedback loops between AI-generated content and human communication. In other cases, alignment and safety interventions may attenuate such tendencies. Understanding these dynamics will be important for ensuring that generative AI systems remain fair, reliable, and culturally aware as they are increasingly deployed across diverse linguistic environments.

Methods

Evaluation of Mental Health Stigma Orientation in Large Language Models

To systematically evaluate mental health stigma in large language models (GPT-4o and Qwen3), we employed multi-dimensional measures spanning social stigma, self-stigma, and health professional stigma, and additionally incorporated a vignette-based measure as well as a depression-specific stigma measure.

To enable cross-linguistic comparisons, we first translated the original scales and conducted back-translation procedures to ensure semantic equivalence across languages [47]. Furthermore, in the prompts, we avoided including any cues related to nationality, race, or other identity-specific information, ensuring that language was the only manipulated factor. For example, in the vignette-based measure, we replaced the original gender, race, and education information with the neutral third-person reference "this person," so that the only difference in the model input was the language.

In addition, to minimize potential learning effects, such as models inferring that they were completing a psychometric assessment, and to better capture their baseline attitudes, we required the models to respond to each item independently, without contextual continuity or awareness of previous responses [48]. We conducted 100 iterations per language, consistent with prior studies [12]. Thus, for each item in each measure, we ran 100 independent iterations for the English version and 100 for the Chinese version. Each iteration was executed independently via the API, ensuring that the model's response to a given item could not influence its response to subsequent items.

Mental Health Stigma Detection

To evaluate whether cross-linguistic variation influences downstream decision-making behavior in large language models, we conducted a binary mental health stigma detection task. In a zero-shot setting (i.e., without providing in-context labeled demonstrations) [49], models were asked to determine whether a given social media snippet contained stigmatizing content.

Since the original benchmark dataset was available only in English, we constructed a parallel Chinese version to enable paired cross-linguistic comparisons. English texts were translated using the DeepL API. To ensure semantic fidelity, we performed automated consistency verification using GPT-4.1 to confirm that translated texts preserved the meaning and contextual intent of the original content. Finally, we randomly sampled 100 instances for manual inspection to further validate translation quality. All instances were aligned at the item level to enable paired statistical testing.

Using the bilingual dataset, we evaluated two large language models (GPT-4o and Qwen3-32B). We adopted a

zero-shot prompting strategy to avoid task-specific fine-tuning and to ensure comparability across language conditions. Temperature was fixed at 0.0 to minimize stochastic variability. For each instance and language condition, we generated 30 independent predictions. Final performance metrics were computed based on the aggregated predictions across the 30 runs. In the rare case of ties, the corresponding samples were excluded from the analysis.

Cross-linguistic differences in overall classification correctness (accuracy) and sensitivity (recall, computed on gold-positive instances) were assessed using paired two-sided McNemar tests. The McNemar test is appropriate in this context because predictions under English and Chinese prompts were paired at the instance level, and the test specifically evaluates whether discordant classification outcomes differ systematically across conditions.

Differences in precision and F1-score were evaluated using paired non-parametric bootstrap resampling over instances (10,000 iterations). In each iteration, instances were sampled with replacement, metrics were recomputed for both language conditions, and the distribution of metric differences (English minus Chinese) was obtained. Two-sided *p*-values were derived from the empirical bootstrap distribution, and 95% confidence intervals were calculated using the percentile method.

Depression Severity Detection

To enable paired cross-linguistic comparisons, we constructed a semantically aligned Chinese version of the original English benchmark dataset following the same translation and quality-control procedures described in Task 1 (Section 5.2). The original dataset contained over 41,000 annotated social media posts labeled for four levels of depression severity (minimal, mild, moderate, and severe). Due to computational and budget constraints, we randomly sampled 1,000 instances for evaluation using a fixed random seed to ensure reproducibility. The four severity categories were approximately balanced in the sampled subset, thereby avoiding substantial class imbalance.

Consistent with Task 1, all generative AI models were queried via their official APIs in a zero-shot setting with temperature fixed at 0.0 to minimize stochastic variability. For each post, we generated 30 independent responses per language condition. Final predictions were obtained by averaging across the 30 runs, thereby reducing random variation and stabilizing performance estimates.

Overall classification performance was evaluated using accuracy, macro-precision, macro-recall, and macro-F1 (see Supplementary Information Section 3). These aggregate metrics allow comparison of general predictive performance across language conditions.

To examine whether cross-linguistic differences manifested as systematic directional shifts, we computed a signed prediction error for each instance:

$$e_i = \hat{y}_i - y_i,$$

where $\hat{y}_i \in \{0, 1, 2, 3\}$ denotes the predicted severity level and $y_i \in \{0, 1, 2, 3\}$ denotes the gold label. Negative values indicate under-estimation, whereas positive values indicate over-estimation.

To assess whether directional errors differed between English and Chinese prompts, we converted signed errors into paired binary indicators. Specifically, under-estimation was defined as

$$U_i = \mathbb{I}e_i < 0,$$

and over-estimation as

$$O_i = \mathbb{I}e_i > 0,$$

where $\mathbb{1}\cdot$ denotes the indicator function. McNemar-style paired comparisons were then conducted on $\{U_i\}$ and $\{O_i\}$ separately to test whether the frequency of directional errors differed across language conditions.

Finally, to further characterize potential decision-boundary shifts, we computed mean signed prediction error separately for each gold severity category under both language conditions. This class-specific analysis enables examination of whether cross-linguistic variation varies systematically across minimal, mild, moderate, and severe cases.

References

[1] Zhu, Y. *et al.* Large language models for information retrieval: A survey. *ACM Transactions on Information Systems* **44**, 1–54 (2025).

[2] Sorin, V. *et al.* Large language models and empathy: systematic review. *Journal of medical Internet research* **26**, e52597 (2024).

[3] Gaber, F. *et al.* Evaluating large language model workflows in clinical decision support for triage and referral and diagnosis. *npj Digital Medicine* **8**, 263 (2025).

[4] Sharma, A., Lin, I. W., Miner, A. S., Atkins, D. C. & Althoff, T. Human–ai collaboration enables more empathic conversations in text-based peer-to-peer mental health support. *Nature Machine Intelligence* **5**, 46–57 (2023).

[5] Kumar, A. *et al.* When large language models are reliable for judging empathic communication. *Nature Machine Intelligence* 1–13 (2026).

[6] Emotional risks of ai companions demand attention. *Nature Machine Intelligence* **7**, 981–982 (2025).

[7] D ALARCÓN, R. Culture, cultural factors and psychiatric diagnosis: review and projections. *World psychiatry* **8**, 131 (2009).

[8] Corrigan, P. W. & Watson, A. C. Understanding the impact of stigma on people with mental illness. *World psychiatry* **1**, 16 (2002).

[9] Bredström, A. Culture and context in mental health diagnosing: Scrutinizing the dsm-5 revision. *Journal of Medical Humanities* **40**, 347–363 (2019).

[10] Kleinman, A. Anthropology and psychiatry: The role of culture in cross-cultural research on illness. *The British Journal of Psychiatry* **151**, 447–454 (1987).

[11] Brijnath, B., Croy, S. & Antoniades, J. Mental illness. In *Oxford Research Encyclopedia of Anthropology* (2023).

[12] Lu, J. G., Song, L. L. & Zhang, L. D. Cultural tendencies in generative ai. *Nature Human Behaviour* 1–10 (2025).

[13] Schramowski, P., Turan, C., Andersen, N., Rothkopf, C. A. & Kersting, K. Large pre-trained language models contain human-like biases of what is right and wrong to do. *Nature Machine Intelligence* **4**, 258–268 (2022).

[14] Xu, S., Dong, W., Guo, Z., Wu, X. & Xiong, D. Exploring multilingual concepts of human values in large language models: Is value alignment consistent, transferable and controllable across languages? In *Findings of the Association for Computational Linguistics: EMNLP 2024*, 1771–1793 (2024).

[15] Zhou, X. *et al.* Fairness or fluency? an investigation into language bias of pairwise llm-as-a-judge (2026). URL `https://arxiv.org/abs/2601.13649`. `2601.13649`.

[16] Ethnologue. What are the top 200 most spoken languages? `https://www.ethnologue.com/insights/ethnologue200/` (2025). Insight report; accessed 12 May 2025.

[17] Dave, P. Chatgpt is cutting non-english languages out of the ai revolution. *Wired* (2023). URL `https://www.wired.com/story/chatgpt-nonenglish-languages-ai-revolution/`. Accessed 2026-02-25.

[18] Krendl, A. C. & Pescosolido, B. A. Countries and cultural differences in the stigma of mental illness: the east–west divide. *Journal of Cross-Cultural Psychology* **51**, 149–167 (2020).

[19] Yang, L. H. Application of mental illness stigma theory to chinese societies: synthesis and new direction. *Singapore medical journal* **48**, 977 (2007).

[20] Yang, L. H. & Kleinman, A. 'face'and the embodiment of stigma in china: The cases of schizophrenia and aids. *Social science & medicine* **67**, 398–408 (2008).

[21] Goffman, E. *Stigma: Notes on the management of spoiled identity* (Simon and schuster, 2009).

[22] Rao, D., Feinglass, J. & Corrigan, P. Racial and ethnic disparities in mental illness stigma. *The Journal of nervous and mental disease* **195**, 1020–1023 (2007).

[23] Yang, L. H. *et al.* Culture and stigma: Adding moral experience to stigma theory. *Social science & medicine* **64**, 1524–1535 (2007).

[24] Corrigan, P. How stigma interferes with mental health care. *American psychologist* **59**, 614 (2004).

[25] Corrigan, P. W., Druss, B. G. & Perlick, D. A. The impact of mental illness stigma on seeking and participating in mental health care. *Psychological science in the public interest* **15**, 37–70 (2014).

[26] Clement, S. *et al.* What is the impact of mental health-related stigma on help-seeking? a systematic review of quantitative and qualitative studies. *Psychological medicine* **45**, 11–27 (2015).

[27] Ahmedani, B. K. Mental health stigma: society, individuals, and the profession. *Journal of social work values and ethics* **8**, 4–1 (2011).

[28] Schnyder, N., Panczak, R., Groth, N. & Schultze-Lutter, F. Association between mental health-related stigma and active help-seeking: systematic review and meta-analysis. *The British Journal of Psychiatry* **210**, 261–268 (2017).

[29] Corrigan, P. W. & Watson, A. C. The paradox of self-stigma and mental illness. *Clinical psychology: Science and practice* **9**, 35 (2002).

[30] Livingston, J. D. & Boyd, J. E. Correlates and consequences of internalized stigma for people living with mental illness: A systematic review and meta-analysis. *Social science & medicine* **71**, 2150–2161 (2010).

[31] Thornicroft, G. Stigma and discrimination limit access to mental health care. *Epidemiology and Psychiatric Sciences* **17**, 14–19 (2008).

[32] Corrigan, P. W. & Rao, D. On the self-stigma of mental illness: Stages, disclosure, and strategies for change. *The Canadian Journal of Psychiatry* **57**, 464–469 (2012).

[33] Kleinman, A. & Hall-Clifford, R. Stigma: a social, cultural and moral process (2009).

[34] Penn, D. & Wykes, T. Stigma, discrimination and mental illness. *Journal of Mental Health* **12**, 203–208 (2003).

[35] Corrigan, P. W., Kerr, A. & Knudsen, L. The stigma of mental illness: Explanatory models and methods for change. *Applied and preventive Psychology* **11**, 179–190 (2005).

[36] Link, B. G. Understanding labeling effects in the area of mental disorders: An assessment of the effects of expectations of rejection. *American sociological review* 96–112 (1987).

[37] Day, E. N., Edgren, K. & Eshleman, A. Measuring stigma toward mental illness: Development and application of the mental illness stigma scale 1. *Journal of applied social psychology* **37**, 2191–2219 (2007).

[38] Pescosolido, B. A., Halpern-Manners, A., Luo, L. & Perry, B. Trends in public stigma of mental illness in the us, 1996-2018. *JAMA network open* **4**, e2140202–e2140202 (2021).

[39] Guze, S. B. Diagnostic and statistical manual of mental disorders, (dsm-iv). *American Journal of Psychiatry* **152**, 1228–1228 (1995).

[40] Santomauro, D. F. *et al.* Global prevalence and burden of depressive and anxiety disorders in 204 countries and territories in 2020 due to the covid-19 pandemic. *The Lancet* **398**, 1700–1712 (2021).

[41] Vogel, D. L., Wade, N. G. & Haake, S. Measuring the self-stigma associated with seeking psychological help. *Journal of counseling psychology* **53**, 325 (2006).

[42] Oliver, M. I., Pearson, N., Coe, N. & Gunnell, D. Help-seeking behaviour in men and women with common mental health problems: cross-sectional study. *The British Journal of Psychiatry* **186**, 297–301 (2005).

[43] Modgill, G., Patten, S. B., Knaak, S., Kassam, A. & Szeto, A. C. Opening minds stigma scale for health care providers (oms-hc): examination of psychometric properties and responsiveness. *BMC psychiatry* **14**, 120 (2014).

[44] Meng, H. *et al.* What is stigma attributed to? a theory-grounded, expert-annotated interview corpus for demystifying mental-health stigma. *arXiv preprint arXiv:2505.12727* (2025).

[45] Priyadarshana, Y., Liang, Z. & Piumarta, I. Heladepdet: A novel multi-class classification model for detecting the severity of human depression. In *International Conference on Collaboration Technologies and Social Computing*, 3–18 (Springer, 2023).

[46] Hida, R., Kaneko, M. & Okazaki, N. Social bias evaluation for large language models requires prompt variations. *arXiv preprint arXiv:2407.03129* (2024).

[47] Brislin, R. W. Back-translation for cross-cultural research. *Journal of cross-cultural psychology* **1**, 185–216 (1970).

[48] Kratzke, N., Beuter, N., Drews, A. & Janneck, M. Psych—psychometric assessment of large language model characters: An exploration of the german language. *Analytics* **5**, 5 (2026).

[49] Brown, T. *et al.* Language models are few-shot learners. *Advances in neural information processing systems* **33**, 1877–1901 (2020).