# Quark Matter Phase Boundary Evolution in Neutron Star Mergers

M. G. Alford [1,*] L. Brodie [1,†] A. Haber [2,1,‡] and J. Shao [1,3,§]

[1] *Department of Physics, Washington University in St. Louis, St. Louis, MO 63130, USA*
[2] *Mathematical Sciences and STAG Research Centre, University of Southampton, Southampton SO17 1BJ, United Kingdom*
[3] *McDonnell Center for the Space Sciences, Washington University in St. Louis, St. Louis, MO 63130, USA*

(Dated: September 9, 2026)

Place holder

## I. INTRODUCTION

[MGA I plan to do significant editing of the Intro once the rest of the paper is in good shape. Until then let's leave it as it is.] Neutron star mergers provide a promising opportunity to study the dynamics of neutron star matter on the millisecond timescale. It is hoped that this information will help us to map the phases of ultra-dense matter. In particular, if there were a first-order phase transition from nuclear matter to quark matter at some critical pressure, it could produce signatures in the gravitational or electromagnetic waves emitted, either during the inspiral or after the merger. AH cite both PRLs already here? In both scenarios, the speed at which the boundary moves is an essential factor which in current treatments is treated by extreme approximations, such as setting it to zero or infinity, as we describe below. In this work we estimate how fast the boundary is likely to propagate under conditions typically found during the inspiral and the merger phase, and draw conclusions about the validity of those approximations.

During inspiral, a phase boundary between nuclear and quark matter may support interface modes whose excitation produce signatures that are potentially detectable with the current LVK network, and certainly within reach of next-generation detectors like Cosmic Explorer or the Einstein Telescope [1, 2]. The existence and frequency of the interface mode critically depend on the conversion timescale, i.e., the speed with which the phase boundary between nuclear and quark matter can move via conversion of metastable nuclear matter to quark matter, or vice versa. If the phase boundary can move too fast, the restoring force for the interface mode and thus the mode itself vanish. All recent analyses [1–3] set this velocity to zero and assume that on the millisecond timescale of the oscillation, the phase content is frozen.

[MGA Broaden this part to include nuclear/quark transitions in supernovas? e.g. https://arxiv.org/abs/2108.00196 https://arxiv.org/abs/2603.18153] The same phase transition may also power core-collapse supernovae: if the collapsing core of a massive star reaches the density where nuclear matter converts to quark matter, the resulting quark core can trigger a delayed second shock and explode stars that neutrino heating alone fails to explode [4–6]. This scenario has a gravitational-wave signature of its own: the formation of a quark matter core inside a newly born proto-neutron star has been proposed as a source of gravitational waves in the MHz band, a supernova analog of the interface mode discussed above for the inspiral [7]. The boundary speed calculated here bears on both the explosion mechanism and this gravitational-wave signature, though core collapse reaches densities and temperatures beyond the range we consider below. In the post-merger phase there are also possible signatures of a nuclear-quark phase boundary. Several recent works have addressed this question using model equations of state that include a nuclear/quark transition [8–17], looking for features of the gravitational wave emission arising from the redistribution of mass as regions of the denser phase grow or shrink in response to pressure gradients in the violent collision between the stars. Current simulations assume that if a region of nuclear matter is compressed to a pressure above the critical pressure $p_{\rm crit}$ where quark matter becomes favored, it will immediately convert to quark matter, so there will be no significant regions of metastable nuclear matter or out-of-equilibrium quark matter AH we need to verify this as Veronica's EOS that the Most PRL uses does have out-of-equilibrium QM. This assumes that the phase boundary can travel distances of order kilometers in milliseconds, i.e., at speeds of order $10^6\,\mathrm{ms}^{-1}$. Our results will show that this assumption is questionable. AH small sentence about DM induced conversion works, and glitch mechanism via mtastable phase as further science cases of relevance AH microscopic = strong interaction time scale? In both situations described above, the key question is how fast a region of metastable nuclear matter could be converted to quark matter via movement of an existing phase boundary arising from a gradual (compared to microscopic timescales) increase in pressure. The propagation of a nuclear/quark phase boundary has been analyzed before, mostly for the more extreme case of the conversion of hadronic stars into strange stars in the context of the strange matter hypothesis [MGA For now I'm not separating ones that allow for a pressure-induced transition] [18–28] This paper complements earlier work by studying a slowly varying pressure driven transition to quark matter (rather than the global instability assumed

* alford@physics.wustl.edu
† b.liam@wustl.edu
‡ a.haber@soton.ac.uk
§ jan.shao@wustl.edu

by the strange matter hypothesis), studying the temperature and density ranges typical of neutron star mergers and supernovas, giving a survey of how the boundary speed varies as a function of the density and temperature of the nuclear phase, and providing an approximate analytic expression for the boundary speed. To achieve this we adopt a simplified model, focusing on the dominant flavor equilibration channel in quark matter. We do not consider the reverse process, conversion of quark matter to nuclear matter, although this may also occur in situations where there are pressure oscillations. We aim to set an approximate upper bound on the propagation speed of the boundary, so our approximations and simplifications are chosen to favor fast propagation.

[AH]mention neutron matter here as approximation? Our treatment assumes a deflagration-type conversion process, in which the pressure has risen high enough so that nuclear matter becomes metastable relative to 3-flavor quark matter, so weak interactions are required for the phase boundary to move. There is a more extreme possibility, only realized if the pressure becomes so high (e.g. in an externally induced shock front) that quark matter with the same flavor content as nuclear matter is energetically preferred. This opens up the possibility of a detonation-type conversion [25, 29] that does not require weak interactions and hence can proceed much faster. Relativistic jump-condition analyses classify detonation by a supersonic upstream velocity and typically find front speeds of order 0.1 times the speed of light or greater [30]. Such speeds are many orders of magnitude above our phase boundary velocity results and are therefore inaccessible when conversion is limited by weak-interaction-driven flavor equilibration. Detonation would require a separate shock-driven mechanism outside the regime considered in this work. [AH]this paragraph seems somehwat duplicated, big jump in technicality, and takes away results

A more violent version of this process, the quark-nova, is the hypothesized runaway conversion of an entire neutron star into a quark star, invoked as an energy source for superluminous supernovae and other high-energy transients [22, 28]. The microphysical boundary speed derived here is the starting point for such global conversion scenarios; extending our steady, planar treatment to the turbulent, whole-star regime of a quark-nova is left for future work. [AH]also duplicate content partially In Sec. II, we present our assumptions and the steady hydrodynamic equations. Sections III and IV describe the isothermal and adiabatic calculations of the boundary speed, respectively. In Sec. V we present our results, and in Sec. VI we summarize our conclusions and describe directions for future work. Our calculations use natural units, $\hbar = c = k_B = 1$.





FIG. 1: Pressure of chemically equilibrated nuclear matter and *uds* quark matter, and also *ud* quark matter. We study the regime of pressure $P$ between $P_{uds}$ and $P_{ud}$, where *uds* quark matter is favored over nuclear matter, but *ud* quark matter is not.





FIG. 2: [MGA Now that we are doing isothermal and adiabatic cases, we either need different versions of this figure, or drop it.] Schematic illustration of the expected boundary profile. Note the discontinuities in some thermodynamic properties at the phase boundary located at $x = 0$, followed by relaxation to beta equilibrium as $x \to \infty$.

## II. FORMALISM AND ASSUMPTIONS

### A. Main assumptions

We assume the phase boundary is well approximated as a flat plane which can therefore be modeled as a one-dimensional system. This is a reasonable approximation for neutron star mergers, where the curvature of the boundary is on the kilometer scale, while its thickness is of order micrometers (see Eq. (41)). We defer any discussion of instabilities that would lead to a more complicated front geometry. [AH]As far as I know, people assume

that instabilities work on the cm scale, which would still give us many orders of magnitude compared to micrometer. So the planar assumption should be fairly solid We assume that the surface tension of the phase boundary is large enough that mixed phases are not favored. We model the nuclear phase as pure neutron matter described by the QMC-RMF3 mean field theory [31–33]. We model quark matter as a Fermi gas of massless noninteracting up, down, and strange quarks [34, 35].

We will follow the standard approach to nuclear to quark conversion [18, 26, 27, 30, 36, 37] which exploits the clear separation between strong-interaction-driven processes such as local thermal equilibration and deconfinement (which can be treated as instantaneous on the time scales relevant to neutron star mergers) and flavor equilibration, which occurs via weak interactions and is therefore much slower and must be treated explicitly. We focus on the fastest flavor equilibration channel, the nonleptonic process

$$d\ u \leftrightarrow u\ s\ . \tag{1}$$

This equilibrates the quantum number that we will call "K", which measures the imbalance between down and strange quarks. We ignore electromagnetism, electrons, and neutrinos. We therefore neglect Urca processes, which are generally slower than the non-leptonic process [38–40], so their inclusion will make the boundary speed even lower.

As noted above, we assume local thermal equilibration is fast enough so that each fluid element has a well-defined temperature $T$. We will first write down the hydrodynamic equations that describe the movement of the fluid, including radiative loss and diffusion of flavor (the $K$ quantum number) and heat. In the body of the paper, however, we will not explicitly treat the radiation or diffusion of heat via thermal conduction. Instead, we will calculate the boundary propagation speed for two extreme cases: adiabatic (no thermal conduction) and isothermal (the whole system is thermally equilibrated to the same temperature).

### B. Formalism

To calculate the speed at which the nuclear/quark phase boundary propagates through neutron matter, we will look for a steady state solution to the equations that describe the propagation of a deflagration front.

We work in the rest frame of the phase boundary, which is located at $x = 0$, as illustrated schematically in Fig. 2. At $x < 0$ there is pure neutron matter which we take to be homogeneous, neglecting any disturbance propagating ahead of the phase boundary. The neutron matter flows towards the boundary with a velocity $u(x = 0^-)$, which we aim to calculate. Matter at $x = 0^-$ immediately (via strong interactions that are instantaneous on the timescales we consider) deconfines into denser quark matter at $x = 0^+$ with enough strangeness to be created from neutron matter while conserving energy. (In the example shown in Fig. 2 this is an endothermic reaction, so the temperature of freshly created quark matter at $x = 0^+$ is less than in the neutron matter arriving at $x = 0^-$.) [AH] I don't understand the enough strangeness part? [MGA] If the quark matter doesn't have enough strangeness then it won't be energetically favored (extreme case: udd matter).]

At $x > 0$ there is a conversion region where strange quarks are created via the weak interaction, leading to a gradient in the K fraction that drives diffusion of strange quarks upstream towards the boundary. This is how the freshly created quark matter at $x = 0^+$ can immediately have a larger strangeness fraction than was present in the incoming neutron matter. As $x \to \infty$ the quark matter approaches a fully beta-equilibrated state.

The parameters that describe the state of the matter at any given position $x$ are:

$$\begin{array}{ll} n_B & \text{baryon number density} \\ n_K \equiv \dfrac{n_d - n_s}{2} & \text{"K" density} \\ u & \text{velocity in the } x \text{ direction} \\ T & \text{temperature} \end{array} \tag{2}$$

From these one can, via the nuclear and quark equations of state (see Appendix A), calculate other properties such as pressure $P$, baryon chemical potential $\mu_B$, energy density, and the K chemical potential

$$\mu_K = \mu_d - \mu_s\ , \tag{3}$$

which relaxes towards zero via the weak interaction (1).

We also define the local K fraction

$$a(x) \equiv \frac{n_K(x)}{n_B(x)}\ . \tag{4}$$

### C. Hydrodynamic equations

We work in the phase boundary rest frame and include heat transport while neglecting viscous stresses, so the stress-energy tensor is [AH] what would a dissipative correction look like as this would also be a result of the same weak interactions I think? du us BV is resonant at very low temperatures

$$\begin{aligned} T^{\mu\nu} = (P+\epsilon)U^\mu U^\nu + P g^{\mu\nu} \\ + U^\mu q^\nu_{\rm th} + q^\mu_{\rm th} U^\nu\ , \end{aligned} \tag{5}$$

and baryon current

$$J^\mu_B = n_B U^\mu\ , \tag{6}$$

where $U^\mu = (\gamma, u, 0, 0)$ is the four-velocity with $u = \gamma v$ and Lorentz factor $\gamma = 1/\sqrt{1-v^2}$. The heat four current $q^\mu_{\rm th}$ satisfies $U_\mu q^\mu_{\rm th} = 0$ and is purely spatial in the local fluid rest frame [41]. Baryon number is conserved, while

escaping neutrinos can carry energy and momentum out of the fluid:

$$\begin{aligned} \partial_\mu T^{\mu\nu} &= -\mathcal{S}^\nu \,, \\ \partial_\mu J^\mu_B &= 0 \,. \end{aligned} \tag{7}$$

[MGA Is this where we introduce terms corresponding to thermal conduction and radiative energy loss?]

Here $\mathcal{S}^\nu$ is the neutrino energy-momentum loss per unit spacetime volume, with $Q_\nu \equiv \mathcal{S}^0 \geq 0$ the energy loss per unit volume and time in the phase boundary rest frame. We impose steady planar flow, so all fields depend only on $x$. For the slow flows considered here, we model the conductive heat flux by Fourier's law,

$$q^x_{\rm th} = -\kappa_{\rm th} \frac{dT}{dx} \,, \tag{8}$$

where $\kappa_{\rm th} \geq 0$ is the thermal conductivity. The minus sign directs heat toward lower temperature. This approximation assumes that the heat current relaxes on a time scale short compared with changes experienced by the fluid [42]. We retain the leading conductive energy flux and neglect the acceleration correction to the relativistic heat law and the heat-current contribution to the momentum flux. We also neglect neutrino momentum loss, which is suppressed by the fluid speed for isotropic emission in the local rest frame. In quark matter, $\kappa_{\rm th}$ is determined by quark scattering [43].

Weak interactions violate the conservation of the K current,

$$\partial_\mu J^\mu_K = -\Gamma_K, \tag{9}$$

[LB the diffusion term should indeed stay in $J^\mu_K$ because it does not lead to the non conservation of K current unlike the weak interactions.]

where $\Gamma_K$ is the rate of down to strange quark conversion (see Appendix B) and

$$J^\mu_K = n_K U^\mu + D^\mu_K, \tag{10}$$

where $D^\mu_K$ is the strong-interaction-driven K diffusion current

$$D^\mu_K = -D_K (g^{\mu\nu} + U^\mu U^\nu) \partial_\nu n_K. \tag{11}$$

See Appendix C for our approach to estimating the diffusion coefficient $D_K$. This definition ensures that $U_\mu D^\mu_K = 0$, i.e. that there is no time component of diffusion in the local fluid rest frame.

The spatial component of the K current $J_K \equiv J^1_K$ is

$$J_K = u n_K - \gamma^2 D_K \frac{d}{dx} n_K \,, \tag{12}$$

[LB in the relativistic formalism, there is a $\gamma^2$ factor in front of the $D_K$ coefficient so this may need to be propagated forward in the paper.] where $D_K$ is the strong-interaction-driven K diffusion coefficient, described in Appendix C. In the transport equations below we use $\gamma^2 \simeq 1$, appropriate to $u \ll 1$, while retaining the relativistic form of the ideal-fluid fluxes.

The time component of the energy-momentum balance gives $dT^{x0}/dx = -Q_\nu$. To leading order in the heat-transport corrections, $T^{x0}$ is the sum of the advective flux $(P+\epsilon)u\gamma$ and the conductive flux $-\kappa_{\rm th} T'$. Weak equilibration converts chemical energy into heat within the fluid; this energy is included in the equation of state, so $\Gamma_K$ does not enter as an additional source of total energy. The steady hydrodynamic equations, with heat transport treated in this slow-flow approximation, are [MGA add in thermal diffusion and radiation terms]

$$\frac{d}{dx}(u n_B) = 0 \,, \tag{13a}$$

$$\frac{d}{dx}\left((P+\epsilon)u\gamma - \kappa_{\rm th}\frac{dT}{dx}\right) = -Q_\nu \,, \tag{13b}$$

$$\frac{d}{dx}\left((P+\epsilon)u^2 + P\right) = 0 \,, \tag{13c}$$

$$\frac{d}{dx} J_K = -\Gamma_K \,. \tag{13d}$$

We use two limits of Eq. (13) to study the effect of heat transport. For $\kappa_{\rm th} \to \infty$ and $Q_\nu = 0$ , $dT/dx \to 0$ in each phase. Assuming thermal contact with no temperature jump at the interface gives a single temperature across both phases, the isothermal case of Sec. III. The conductive flux can remain finite in this limit and balance changes in the advective flux. For $\kappa_{\rm th} = 0$ and $Q_\nu = 0$, Eq. (13b) instead gives $d[(P+\epsilon)u\gamma]/dx = 0$, the adiabatic case of Sec. IV. Here *adiabatic* excludes conductive and radiative heat loss; weak reactions can still produce entropy. We set $Q_\nu = 0$ in both calculations, consistent with our omission of neutrinos and Urca processes.

## III. BOUNDARY PROPAGATION SPEED: ISOTHERMAL CASE

[MGA Need to write this]

We now describe how to calculate $u(0^-)$, the speed at which the nuclear/quark phase boundary propagates through neutron matter, in the limit where there is a single uniform temperature across both phases.

This is the $\kappa_{\rm th} \to \infty$ limit described in Sec. II C, with finite heat transport maintaining a uniform temperature.[JS citations] [LB Are we sure we cannot impose energy conservation $hu = const$ as well as have constant temperature? I am not sure what the additional unknown quantity would be though.] The energy balance in Eq. (13b) still includes conductive transport, so it does not require the advective flux $(P+\epsilon)u\gamma$ to be constant. This idealization allows heat exchange with the surroundings to maintain the common temperature. We solve for the matter profile using $T(x) = T(0^-)$ in place of the energy equation:





FIG. 3: Caption

$$\frac{d}{dx}(un_B) = 0\,, \tag{14a}$$

$$\frac{d}{dx}T(x) = 0\,, \tag{14b}$$

$$\frac{d}{dx}((P+\epsilon)u^2 + P) = 0\,, \tag{14c}$$

$$\frac{d}{dx}J_K = -\Gamma_K. \tag{14d}$$

Most of the piecewise-constant derivation of Sec. IV C survives this replacement, with one subtlety being that $\mu_B$ and $n_B$ for $x > 0$ are no longer constant, but their change is still quite small. So figuratively, see Fig. 3, when we go from $x = 0^+$ to $x = \infty$ in the isothermal limit, instead of moving along the isobar of quark matter on $(T, \mu_K)$ plane for constant $\mu_B$ , we move along the isobar on $(\mu_B, \mu_K)$ plane for constant $T$ .

Along this isobar, $\mu_B$ and $\mu_K$ are not independent. At fixed $T$, the Gibbs–Duhem relation $dP = n_B d\mu_B + n_K d\mu_K$ gives

$$\left.\frac{d\mu_B}{d\mu_K}\right|_{P,T} = -\frac{n_K}{n_B} = -a\,, \tag{15}$$

Therefore, $\mu_B$ increases as weak reactions reduce $\mu_K$, but the change becomes smaller as the matter approaches flavor equilibrium. For the representative profile with $T = 10\,\mathrm{MeV}$, $n_B(0^-) = 3n_0$, $B^{1/4} = 189.2\,\mathrm{MeV}$, and $a(0^+) = 0.146$, $\mu_B$ and $n_B$ increase by only 0.24% between $0^+$ and $x \to \infty$. Since $a = n_K/n_B$, its exact derivative is $a' = n_K'/n_B - an_B'/n_B$ . Along the same trajectory, the approximation $a' \simeq n_K'/n_B$ differs from the exact derivative by at most 0.47%. We therefore neglect the small variation of $n_B$ in the analytical approximation below.

Baryon and momentum flux conservation are unchanged, so the interface gradient in Eq. (29), the integrated K-ness balance in Eq. (30), the estimate of $I_1$ in Eq. (33), and the general speed relation in Eq. (34) carry over to the isothermal case. The only remaining change is the evaluation of $I_2$.

As in the piecewise-constant treatment, we fix $\mu_q$ at a representative value. The coefficients $D_K$, $\gamma_K$, and $\eta$ are then constant, and Eq. (32) reduces to

$$\begin{aligned} I_2 &= D_K\gamma_K \int_0^{a(0^+)} \left(a^3 + \eta a\right) da \\ &= \frac{D_K\gamma_K\, a^2(0^+)\left[a^2(0^+) + 2\eta\right]}{4}\,. \end{aligned} \tag{16}$$

This integral is simpler than its counterpart in Sec. IV C, where $T(x)$ follows the isobar in Eq. (36). In that case, $D_K[T(a)]$ remains inside the integral and produces the $A^{7/3}$ prefactor and the fractional powers in Eq. (38).

Substituting Eq. (16) into Eq. (34) gives

$$u^2(0^-) = \frac{D_K\gamma_K\, a^2(0^+)\left[a^2(0^+) + 2\eta\right]}{2\lambda_n^2\left[1 - a(0^+)\right]\left[1 + \xi a(0^+)\right]}\,, \tag{17}$$

where $\eta = 9\pi^2T^2/\mu_q^2$, and $\lambda_n$ and $\xi$ are defined in Eqs. (27) and (33), respectively. The two terms in brackets arise from the cubic and linear contributions to the weak source in Eq. (B3). The cubic contribution dominates when $a(0^+) \gg \sqrt{\eta}$, while the thermal contribution dominates at smaller $a(0^+)$ .

The parameter $A$ does not appear in the isothermal result because its defining condition, $T(0^+) = 0$, belongs to the temperature-varying isobar used in Sec. IV C.

Unlike in the energy-conserving case Eq. (20) and consequently Eq. (38), where we can maximize the velocity by setting $a(0^+) = A$ where $T(0^+) = 0$ , in the isothermal limit there's only one obvious upper limit on $a(0^+)$, which is $a(0^+) \le 1$, corresponding to no strangeness in quark matter at $x = 0^+$ , where the velocity blows up. Therefore, we cannot maximize the velocity simply by setting $a(0^+) = 1$ .

We do not know the value of $a(0^+)$ , but $u(0^-)$ is a monotonically increasing function of $a(0^+)$ in the domain $0 < a(0^+) < a(0^-) = 1$ for $\xi = 0$ , so to obtain an upper bound on the isothermal boundary speed $u_{\max}(0^-)$ we need to find the $a_{\max}(0^+)$ in the isothermal limit, which we denote by $A'$, and it's the value of $a(0^+)$ at which the quarks at the boundary have the same average chemical potential $\mu_B(0^+) + n_K(0^+)\mu_K(0^+)/n_B(0^+)$ as the quarks in the baryons. Quark matter with $a(0^+) > A'$ would have insufficient strangeness to make it favored over neutron matter [24].

Taylor-expanding the quark matter EoS around its beta-equilibrated state, we can obtain an approximate formula that in the parameter range that we explored agrees with the numerical result to within 1%,

$$A' \approx \sqrt{\frac{2\big(\mu_B(0^-) - \mu_B(\infty)\big)}{n_B(\infty)}\left.\frac{\partial^2 P_{\mathrm{QM}}}{\partial\mu_K^2}\right|_\infty}\,. \tag{18}$$

Since Eq. (17) has monotonic dependence on $a(0^+)$, and we know that $0 < a(0^+) < A' < 1$, similar to

Eq. (39), we can do the same thing for Eq. (17)

$$u_{\max}(0^-) \simeq (98\,\mathrm{m/s})\frac{n_B(\infty)}{n_B(0^-)} \times \left(\frac{\mu_B(0^+)}{1200\ \mathrm{MeV}}\right)^{\frac{17}{6}} \left(\frac{10\ \mathrm{MeV}}{T}\right)^{\frac{5}{6}} \times \frac{A'\sqrt{(A')^2 + 162\pi^2 T^2/\mu_B^2(0^+)}}{\alpha_s^{5/6}\sqrt{1-A'}\sqrt{1+\xi A'}}\,. \tag{19}$$

## IV. BOUNDARY PROPAGATION SPEED: ADIABATIC CASE

### A. Adiabatic hydrodynamic equations

Setting $\kappa_{\rm th} = 0$ and $Q_\nu = 0$ in Eq. (13) gives the adiabatic steady-front equations

$$\frac{d}{dx}(un_B) = 0\,, \tag{20a}$$

$$\frac{d}{dx}((P+\epsilon)u\gamma) = 0\,, \tag{20b}$$

$$\frac{d}{dx}((P+\epsilon)u^2 + P) = 0\,, \tag{20c}$$

$$\frac{d}{dx}J_K = -\Gamma_K. \tag{20d}$$

Eqs. (20a), (20b), (20c) are flux conservation laws, stating that the baryon, momentum, and energy fluxes

$$\left.\begin{aligned} J_B &= n_B u\,, \\ \Pi &= hu^2 + P\,, \\ E &= hu\sqrt{1+u^2}\,. \end{aligned}\right\} \text{Independent of } x \tag{21}$$

have fixed values at all $x$, while other thermodynamic and compositional properties will vary through the boundary and relaxation region.

To obtain the profile of the moving phase boundary we only need to solve the remaining Eq. (20d), a second-order differential equation that can be recast as two first-order differential equations for the K density and current

$$\frac{d}{dx}n_K = \frac{un_K - J_K}{D_K}\,, \quad x > 0\,, \tag{22a}$$

$$\frac{d}{dx}J_K = -\Gamma_K\,. \tag{22b}$$

However, it's worth noting that diffusion of quarks and hence K-ness is only possible in deconfined quark matter at $x > 0$ , so Eq. (22a) is only defined for $x > 0$ . For $x < 0$ , we have $dn_K/dx = 0$ and $J_K = un_K$ . [MGA Our assumption that the nuclear matter is all neutrons makes K diffusion nonexistent there.] There is now a well defined procedure for constructing a solution to these differential equations. Assume we have solved them up to some position $x$, giving $n_K(x)$ and $J_K(x)$. We can combine this with the already-specified values of the conserved fluxes (21) and the quark matter equation of state (Appendix A) to solve for the remaining properties of the matter at $x$, such as $n_B(x)$, $u(x)$, $\mu_B(x)$, $\mu_K(x)$, $T(x)$, etc. This allows us to calculate $D_K(x)$ and $\Gamma_K(x)$ (see Appendices B and C) and thereby obtain $n_K(x+\delta x)$ and $J_K(x+\delta x)$.

### B. Boundary conditions

To construct the boundary profile we need 5 initial conditions specifying the state of the incoming nuclear matter: $n_B(0^-)$, $n_K(0^-)$, $dn_K/dx(0^-)$, $T(0^-)$, $u(0^-)$. However, since we assumed the incoming matter is neutron matter, for which $n_K = n_d/2$, we have

$$n_K(0^-) = n_B(0^-)\,, \tag{23}$$

and since we have assumed the neutron matter is homogeneous,

$$\frac{dn_K}{dx}(0^-) = 0\,, \tag{24}$$

so we only need to specify $n_B(0^-)$, $T(0^-)$, $u(0^-)$, which fixes the conserved fluxes $J_B$, $\Pi$, and $E$ for the whole solution.

Since the solution must yield equilibrated quark matter as $x \to \infty$, we also impose the boundary condition $\mu_K(\infty) = 0$, which for our model of quark matter is equivalent to $n_K(\infty) = 0$. This turns (22) into an eigenvalue problem, where for given $n_B(0^-)$ and $T(0^-)$ there is only one value of the incoming neutron matter velocity $u(0^-)$ that yields a valid solution. This will yield our desired result, the speed at which the boundary propagates through neutron matter, as a function of the density and temperature of the neutron matter.

The first step in solving the equations is to relate the K density and current at $x = 0^+$ to the initial conditions at $x = 0^-$, after which we can evolve the solution to $x = +\infty$ using (20).

[MGA I don't think this is an essential difference. It's still an equation of the form $dF/dx = RHS$. In order for there to be a discontinuity in $F$ there must be a delta-function in $RHS$. I think the essential point is that we know the RHS of (22b) doesn't have a delta fn, but we can't rule it out for the RHS of (22a)] [JS I think the problem is $D_K$ in Eq. (22a) is not even defined for $x \le 0$, so I don't think we can claim $n_K(0^-) \neq n_K(0^+)$ implies $J_K$ have delta fn part. The only meaningful equation that can be integrated across $x = 0$ is Eq. (22b), and we need to make it clear that Eq. (22a) is only defined for $x > 0$ , and that $J_K = n_K u$ for $x < 0$ and $J_K = n_K u - D_K n_K'$ for $x > 0$ .] [MGA You are focusing on the right side of (13a), saying it only takes that form in quark matter. That's true, but it avoids the key question, which is whether the true right side of (13a) could have a delta function at $x = 0$. If there is a

discontinuity in $n_K$ then the true RHS of (13a) must have such a delta function.]

Integrating Eqs. (22) across the sharp interface yields the hydrodynamic jump condition

$$u(0^+)n_K(0^+) - \left[D_K \frac{dn_K}{dx}\right]_{0^+} = u(0^-)n_K(0^-)\,. \quad (25)$$

We do not need to know the K diffusion coefficient on the nuclear side because we have assumed there is no gradient there. Eq. (25) fixes $dn_K/dx(0^+)$ in terms of $n_K(0^+)$ but we have no constraint on $n_K(0^+)$ so we have to treat it as an additional parameter of the system. In our calculations we will choose the value of $n_K(0^+)$ that maximizes the speed of the boundary.

We can now obtain the boundary speed in terms of the density and temperature of the neutron matter and the K fraction in the freshly created quark matter. We do this by solving (22) as described in Sec. IV A using the boundary conditions described above.

One approach is to solve the differential equations numerically, as described in Appendix E, and this is how most of our results in Sec. V were obtained. However, as we now describe, one can also formulate an approximate analytic solution, which is helpful for interpreting the results.

### C. Piecewise-constant analytical approximation

[MGA Could we think of a more informative name for this approximation? E.g. "constant pressure" or similar.]

If the boundary propagation speed is much less than the speed of light, then, as we see in our full numerical calculations (see Fig. 5) the boundary profile exhibits noticeable simplifying features. One can treat $n_B$ , $u$ , $\mu_B$ , $P$ , and $h$ as independent of $x$ on the quark matter side, while retaining the $x$ dependence of temperature $T(x)$ , and K density $n_K(x)$ .As we will now show, this allows us to find an analytic expression for the speed of the boundary.

We write the $x$-independent baryon density in the quark matter as

$$n_B^{(+)} \equiv n_B(x) \quad \forall x > 0\,, \quad (26)$$

[MGA I am open to alternative notations for the constant quantities] and define $\lambda_n$ (which will always be less than 1) as the ratio of neutron matter to quark matter baryon density at the transition,

$$\lambda_n \equiv \frac{n_B(0^-)}{n_B^{(+)}}\,, \quad (27)$$

so the $x$-independent velocity of the quark matter is

$$u^{(+)} = \lambda_n u(0^-)\,. \quad (28)$$

Since $n_B$ is independent of $x$, it is natural to work in terms of the local K fraction $a(x)$ defined in Eq. (4).

To find the speed $u(0^-)$ we use Eq. (25) along with (26), (27), (28) to obtain

$$\begin{aligned} u(0^-) &= -\frac{D_K(0^+)a'(0^+)}{\lambda_n\big(1-a(0^+)\big)}\,, \\ a'(x) &\equiv \frac{da}{dx} = \frac{1}{n_B^{(+)}}\frac{dn_K}{dx}\,. \end{aligned} \quad (29)$$

The boundary conditions determine $\lambda_n$ and $a(0^+)$, but $D_K(0^+)a'(0^+)$ is determined by the requirement that $a(\infty) = 0$. To impose this exactly we would have to evolve $a(x)$ from $x = 0^+$ to infinity using the K current equation Eq. (22).

We cannot solve this equation exactly in full generality, but we can make a good estimate of the result. Multiplying Eq. (22b) by $D_K(x)a'(x)$, using Eq. (22a) to eliminate $J_K$, and integrating from $x = 0^+$ to infinity, we find

$$\frac{1}{2}\Big(D_K(0^+)\,a'(0^+)\Big)^2 = -u^{(+)}I_1 + I_2\,, \quad (30)$$

where $I_1$ and $I_2$ are positive dimensionless integrals over $x$ which can be rewritten as integrals over $a$, since $a(x)$ is expected to be a monotonically decreasing function over $0^+ \leqslant x < \infty$ as the quark matter equilibrates in the conversion region behind the boundary.

$$\begin{aligned} I_1 &= \int_{0^+}^{\infty} D_K(x)\big(a'(x)\big)^2\,dx\,, \\ &= -\int_0^{a(0^+)} D_K a'\,da\,. \end{aligned} \quad (31)$$

$$\begin{aligned} I_2 &= -\frac{1}{n_B^{(+)}}\int_{0^+}^{\infty} D_K\Gamma_K a'\,dx\,, \\ &= \frac{1}{n_B^{(+)}}\int_0^{a(0^+)} D_K\Gamma_K\,da\,. \end{aligned} \quad (32)$$

In an exact calculation both integrands should be obtained from the solution to Eq. (22), but we will now show how one can obtain reasonable approximations to both integrals without knowing the solution.

We first estimate $I_1$ because the estimate will turn out to depend on our desired result $u(0^-)$, requiring regrouping of Eq. 29. We assume that, as we see in the numerical solutions, the actual profile has $D_K(x)a'(x)$ increasing monotonically from its value $-\lambda_n u(0^-)[1-a(0^+)]$ at $x = 0^+$ (from Eq. (29)) to zero at $x = \infty$. This is expected because $a(x)$ decreases monotonically to zero, and any variation in the diffusion constant $D_K$, which depends mainly on the temperature, is too weak to overcome that. We can then parameterize the integral in terms of a "sharpness" factor $\xi$:

$$\begin{aligned} I_1 &= -\frac{1+\xi}{2}\,D_K(0^+)a'(0^+)\,a(0^+) \\ &= \frac{1+\xi}{2}\,\lambda_n u(0^-)\,[1-a(0^+)]\,a(0^+)\,. \end{aligned} \quad (33)$$





FIG. 4: Constant-pressure trajectory through the quark conversion layer. The red curve is the energy-conserving numerical profile for $T(0^-) = 10$ MeV , $B^{1/4} = 180$ MeV , $n_B(0^-) = 3n_0$ , and $a(0^+) = 0.55$; the dashed curve is the quarter ellipse in Eq. (36). The profile runs from the just-deconfined state at $x = 0^+$ to the equilibrated state at $x \to \infty$ . The point $(A, 0)$ is the zero-temperature endpoint of the extrapolated isobar and need not coincide with the state at $x = 0^+$ .

Here $-1 < \xi < 1$ characterizes the shape of the diffusion profile. For details of this sharpness factor $\xi$ and derivation of Eq. (33), see Appendix D. Substituting this relation into (30) and (29) yields

$$u^2(0^-) = \frac{2I_2}{\lambda_n^2[1-a(0^+)][1+\xi a(0^+)]}\,, \tag{34}$$

The final step is to evaluate $I_2$, for which we need to know how the K diffusion coefficient $D_K$ and K creation rate $\Gamma_K$ vary though the conversion region. In the piecewise constant approximation the pressure and baryon chemical potential have constant values $P^{(+)} \approx \Pi$ and $\mu_B^{(+)}$ in that region, so the system follows an isobar in the $\mu_K$-$T$ plane. Moreover, since $\mu_K/\mu_B^{(+)}$ is typically of order 0.1 and $T/\mu_B^{(+)}$ is of order 0.03, we can expand the quark matter EoS to quadratic order (A3), so the isobar takes the form of a quadrant of an ellipse in the $\mu_K$-$T$ plane. To this order there is a linear relationship between the strangeness fraction $a$ and $\mu_K$ (Eq. (A6)) so the quark matter follows an elliptical quadrant in the $a$-$T$ plane, as shown in Fig. 4.

Using Fig. 4 we can describe the conversion region as follows. At $x = 0$ the quark matter has some initial K fraction $a(0^+)$ which determines the initial temperature $T(0^+)$ (the red star in the figure ). As $x$ increases, the system moves upwards along the elliptical isobar, ending at $a = 0$ and some final temperature when $x \to \infty$ (the red square in the figure).

This gives us a natural way to set a limit on the previously unconstrained parameter $n_K(0^+)$, or equivalently $a(0^+)$. For given initial conditions $(n_B(0^-), T(0^-), u(0^-))$, the largest possible value for $a(0^+)$ is the one that would give the lowest allowed initial quark matter temperature, $T(0^+) = 0$. Thus, $a(0^+)$ cannot be larger than $A$ [MGA I would like to have a more obvious symbol for this, but not sure what would be better], the value where the isobar meets the $T = 0$ axis in Fig. 4, defined by $P_{\rm QM}(A, T = 0) = P^{(+)}$, so to leading order,

$$\begin{aligned} a(0^+) \;&\leqslant A \\ A \;&\approx \frac{3}{2}\frac{\sqrt{108\pi^2\big(P^{(+)}+U_{\rm bag}\big)-\big(\mu_B^{(+)}\big)^4}}{\big(\mu_B^{(+)}\big)^2} \end{aligned} \tag{35}$$

To understand the physics behind this limit, note that larger values of $a(0^+)$ correspond to smaller strangeness fraction in the quark matter that is first created immediately behind the phase boundary,

The isobar in Fig. 4 defines the quark matter temperature $T(x)$ as a function of $a(x)$; solving $P_{\rm QM}(a, T) = P_{\rm QM}(A, 0)$ we find

$$T^2 = \frac{2(\mu_B^{(+)})^2}{81\pi^2}(A^2 - a^2) \tag{36}$$

We can now evaluate the integral $I_2$ (32) by using the results from Appendices B and C) for $\Gamma_K(a, T)$ and $D_K(a, T)$ (keeping only the first term), and using (36) to express $T$ as a function of $a$. We find

$$\begin{aligned} I_2 =& \frac{27\pi^{7/3}}{56\sqrt{2}}\frac{A^{7/3}\gamma_K}{\alpha_s^{5/3}h_D\mu_B^{(+)}} \times \\ & \left[8+\left(1-\frac{a(0^+)^2}{A^2}\right)^{1/6}\left(\frac{a(0^+)^2}{A^2}-8\right)\right]. \end{aligned} \tag{37}$$

So from (34),

$$\begin{aligned} u^2(0^-) =& \frac{9\pi^{7/3}}{4\sqrt{2}\,h_D\alpha_s^{5/3}}\frac{\gamma_K A^{7/3}}{\mu_B^{(+)}\lambda_n^2\big(1-a(0^+)\big)\big(1+\xi a(0^+)\big)} \\ & \times\left[\frac{24}{7}-3\left[1-\frac{a^2(0^+)}{A^2}\right]^{1/6}-\frac{3}{7}\left[1-\frac{a^2(0^+)}{A^2}\right]^{7/6}\right]. \end{aligned} \tag{38}$$

We want to find out the fastest speed at which the phase boundary could move, $u_{\rm max}$, so we will set $a(0^+)$, the K fraction in freshly created quark matter, to the value that maximizes the speed. We see from (38) that the speed of the boundary grows as $a(0^+)$ gets larger. This follows because the factor in square brackets has a weak dependence on $a(0^+)$, so the speed is determined by the denominator $\big(1-a(0^+)\big)\big(1+\xi a(0^+)\big)$. For any reasonably smooth profile we expect $|\xi| \ll 1$ After seeing the numerical profile for various $T(0^-)$ and $n_B(0^-)$ we find out $\xi = -0.5$ is a good approximation (Appendix D), but for any value of $\xi$ in its allowed range $(-1, 1)$ the denominator shrinks (so the speed grows) as $a(0^+)$ grows from 0 towards 1.

This means that to achieve the maximum phase boundary speed for given initial conditions, we need larger $a(0^+)$, which makes sense because that corresponds to a lower strangeness fraction in the freshly created quark matter, closer to the flavor content of the neutron matter. As noted above, the largest possible value of $a(0^+)$ is $A$, so setting $a(0^+) = A$ we find the maximum speed

$$u_{\max}(0^-) = (82\,\mathrm{ms}^{-1})\,\frac{n_B^{(+)}}{n_B(0^-)}\left(\frac{\mu_B^{(+)}}{1200\,\mathrm{MeV}}\right)^2 \times \frac{A^{7/6}}{\alpha_s^{5/6}\sqrt{1-A}\sqrt{1+\xi A}}\,. \tag{39}$$

Thus the typical speed of the boundary is expected to be of order $100\mathrm{m\,s}^{-1}$. In this expression the only factor that could drive the speed to a significantly higher value is $1/\sqrt{1-A}$. To see when $A$ could approach 1 we recall that $A$ is the largest possible K fraction in the freshly created quark matter at $x = 0^+$. The K fraction in neutron matter is 1, so if $A = 1$ then we have such a high pressure that nuclear matter can convert to quark matter with the same flavor content while conserving energy. In this case no weak interactions are needed, so it is not surprising that our approach breaks down.

We can also use the piecewise constant approximation to estimate the thickness $L_{\mathrm{conv}}$ of the conversion region, which determines the range of curvature that is compatible with our assumption of a planar geometry. If the boundary moves slowly then when $a \ll 9\pi T/\mu_B^{(+)}$, which is guaranteed to occur at large $x$ where $a \to 0$, the K current equation reduces to

$$\frac{d^2a}{dx^2} = \frac{\gamma_K}{D_K}\left(\frac{9\pi T}{\mu_B^{(+)}}\right)^2 a\,. \tag{40}$$

At large $x$ the temperature reaches an asymptotic value $T(\infty)$, so we can estimate the thickness of the conversion region as

$$L_{\mathrm{conv}} \approx \frac{\mu_B^{(+)}}{9\pi T(\infty)}\sqrt{\frac{D_K}{\gamma_K}} = (0.2\,\mu\mathrm{m})\,\frac{1}{\alpha_s^{5/6}}\left(\frac{1200\,\mathrm{MeV}}{\mu_B^{(+)}}\right)^{7/6}\left(\frac{10\,\mathrm{MeV}}{T(\infty)}\right)^{11/6}, \tag{41}$$

where we evaluated $D_K$ using (C1) with $T = T(\infty)$. and $\gamma_K$ from Eq. (B3). This supports our use of the planar approximation, since the typical curvature of a quark/nuclear boundary in neutron stars will be of order kilometers, much larger than the thickness of the boundary.

[MGA Edited up to here]

### D. Thermal conductivity

Currently we are setting $T(0^+) = 0$ exactly to maximize velocity. Although this can be done in the analytical or semi-analytical approach, numerically I kept running into difficulties. I think now I have a better understanding: $T(0^+) = 0$ leads to $D_K(0^+) \to \infty$ and $D_K^{-1}(0^+) = 0$, which according to Eq. (22), leads to $n_K'(0^+) = 0$. Although there's no infinities and everything seems fine, zero initial condition in derivatives is what people call non-Lipschitz initial condition.

Why this is problematic? To break it down, this is what the numerical solver sees for $T(0^+) = 0$ :

$$\frac{dn_K}{dx}(0) \equiv \frac{n_K(\delta) - n_K(0)}{\delta} = \frac{un_K - j_K}{D_K}(0) = 0$$
$$\frac{dj_K}{dx}(0) \equiv \frac{j_K(\delta) - j_K(0)}{\delta} = -\Gamma_K(0) \neq 0$$

after some rearrangements, we get

$$n_K(\delta) = n_K(0)\,, \quad j_K(\delta) = j_K(0) - \Gamma_K \cdot \delta$$

So $n_K(\delta) = n_K(0)$ , which is sensitive to the step size $\delta$ . This means the solver can return two solutions that numerically solves the equations, but they look different: their $n_K$ profile have different plateau length in the beginning. These two solutions can have $u(0^-)$ and $j_B$ differ by 20% .

So to resolve this, I have several plans:

- cheapest: when numerically solving the set of equations, don't put in exactly zero $T(0^+)$ ; instead choose a sufficiently small quark matter initial temperature, say $T(0^+) = 0.01$ MeV and spend a sentence or two maybe at the $z(t)$ part explaining thermal conductivity is an interesting future research direction. [MGA Yes I prefer this approach. You need to make $T(0^+)$ small enough so that the speed no longer depends on it noticeably. Maybe that already happens for values above 0.01 MeV.]
- minimal thermal conductivity, where we don't add another full ODE to solve, but just model the thermal conductivity ODE, and just integrate it across $x = 0$ to place a lower bound on $T(0^+)$
- full thermal conductivity modeling. To do this we need to replace the current energy conservation equation $d(h\gamma u)/dx = 0$ by a second order ODE similar to what we have for $n_K$ right now:

$$\frac{d}{dx}\left(h\gamma u - \kappa_{\mathrm{th}}\frac{dT}{dx}\right) = -Q_\nu$$

where Eq. (62) in [43] gives a very similar expression for $\kappa_{\mathrm{th}}$ compared with $D_K$ , and $Q_\nu$ is the energy taken away by neutrinos per unit time. So judging by the similarity, I assume every method: analytical, numerical, whatever we have developed for solving the $n_K$ ODE would also apply here. But the draw back of the full thermal conductivity modeling is that we are not including Urca, so this $Q_\nu$ is essentially zero in our model. We need to explain that away.

## V. RESULTS

- first, the set up

We discuss the energy conserving front system Eq. (20) first. We choose the bag constant by requiring cold neutron matter and cold equilibrated strange quark matter to coexist at neutron matter critical baryon number density $n_B^{\rm crit}(T{=}0) = 3n_0$. This gives $U_{\rm bag}^{1/4} = 189.2\,\mathrm{MeV}$. Since we are not including Urca processes, we set strange quark mass to zero. We also adopt the conventional phenomenological choice $\alpha_s = 0.3$ for bag-model quark matter [35, 44]. For each pair of upstream variables $n_B(0^-)$ and $T(0^-)$, we evolve the K density and current, reconstruct the local fraction $a(x) = n_K(x)/n_B(x)$, and enforce the conserved baryon, momentum, and energy fluxes across the quark phase.

But to get a unique solution to Eq. (20), hence a unique value for $u(0^-)$, we still need a proscribed $a(0^+)$, or equivalently, $T(0^+)$. As mentioned in Sec. IV C, we can utilize this freedom to maximize the velocity by choosing $T(0^+) = 0$, but practically that would make $D_K^{-1} = 0$ hence giving us a non-Lipschitz initial condition for the ODE system Eq. (22), which gives us a family of possible solutions instead of one unique solution. So in practice, when using analytical approximation Eq. (34) or Eq. (38) we can set $a(0^+) = A$ or $T(0^+) = 0$, but when numerically solving the full system Eq. (20) we need to set a small but non-zero $T(0^+)$ to avoid the non-Lipschitz initial condition. From our numerical results Fig. 8, we find that the velocity doesn't change over two times for $T(0^+) < 1$ MeV, so we choose $T(0^+) = 0.1$ MeV as a representative value for the numerical results.

### A. Phase boundary speed

- about contour plot:
  - existence of stable neutron matter region, and why it takes that shape
  - $A > 1$ region and why that shape
  - velocity in between, also mention it diverge as the input goes close to $A = 1$

[AH]I think if we have Fig1 as schematic, we should have an equivalent figure here with an actual result?

### B. Approach to equilibrium

- about the $z(4)$ plot:
  - basics, set up, what it means, why we choose that slope
  - talk about the plot, the fact that time is over tens of seconds





FIG. 5: [LBIn the isothermal case does the upper grey region curve back down at high T(0-)? I suspect not.][JSNo it doesn't] Full numerical phase-boundary speed $u(0^-)$ in $\mathrm{m\,s^{-1}}$ as a function of the upstream baryon density and temperature, with $T(0^+) = 0.1\,\mathrm{MeV}$ and $n_B^{\rm crit}(T{=}0) = 3n_0$. The white band is the domain in which the moving-front solution exists. Below the lower boundary neutron matter is stable against the assumed conversion, while the region marked $A \geq 1$ lies outside the weak-interaction-limited branch described by our model. The contours are drawn at 50, 100, 200, 500, 1000, 2000, and 5000 $\mathrm{m\,s^{-1}}$.

- important: the $z \to 0$ behavior

[MGA I would like this paper to have a section like the one we had in our previous version, where we calculate the time taken for the boundary to reach its equilibrium position. Maybe we could add to that some estimate of the impact on gravitational wave signals, but let's discuss this at our next meeting before writing anything more elaborate.][JSDone, see Fig. 6]

## VI. CONCLUSION

- Impact of instabilities, Urca processes, ... on phase boundary speed

Our results allow us to speculate on the fate of interface modes in the inspiral. Typical frequencies of interface modes depend on the EoS of nuclear and quark matter as well as the star's mass, where heavier stars show lower frequency resonances. The frequencies themselves range from a couple hundred to around $f = 2000$ Hz (see





FIG. 6: Remaining distance $z(t)$ to the zero-speed surface, obtained from the full numerical speed map at fixed $T(0^+) = 5\,\mathrm{MeV}$. The initial distance is $z_0 = 1\,\mathrm{km}$, and the assumed density gradient is $0.3 n_0\,\mathrm{km}^{-1}$. The points mark the fitted catch-up times for $T(0^-) = 0$, 20, 40, and 60 MeV.

Fig. 2 of Ref. [1]). Depending on the energy absorbed by the mode, the amplitude ranges from a few millimeters to centimeters in the inspiral, and might reach several meters in the postmerger. Assuming a frequency of $f = 700$ Hz, we can calculate the resonant velocity where the timescale of the oscillation matches the time given by the amplitude divided by the phase boundary velocity. In this resonant scenario, we expect viscous effects as described in [24] become important. At even larger velocities, we expect the mode to lose its support (see discussion in the nuclear case in Ref. [45]) and vanish. In the inspiral, the resonant velocity ranges from $v_{\mathrm{res}} = 1$ m/s to $v_{\mathrm{res}} = 15$ m/s, which is within the range found in our calculation shown in Fig. 5. In the postmerger scenario, the resonant velocity rises to $\mathcal{O}(1000\,\mathrm{m/s})$, which is reached in the strongly metastable scenario (see Fig. 5). This shows that any accurate calculation of the interface mode must include the finite phase boundary velocity and viscous effects, potentially via the dynamical sound-speed formalism developed in Ref. [45] and applied to quark matter in [cite Zhao Han Haber].

This also motivates a more detailed calculation of the phase boundary velocity:

- In this paper, we model the baryonic side as pure neutron matter. However, in compact stars, baryonic matter is composed of chemically equilibrated nuclear matter, which includes not only neutrons, but also protons, electrons, and muons in a composition determined by charge neutrality and chemical equilibration. Even hyperonic matter could be considered on the baryonic side of the phase boundary.

- The presence of leptonic particles opens up the possibility of proton or strangeness-changing Urca rates. The phase boundary velocity would then potentially be slowed down even further, as we now need to equilibrate multiple particle fractions on the quark matter side via the generally slower Urca processes. Our results thus likely present an upper bound on the velocity.

- At high temperatures, the physics of neutrinos might become important as neutrinos could cool down the conversion region, or if they get trapped, will alter the hydrodynamic description of the phase boundary. However, given our recent work in Refs. [**?** **?** ], neutrino trapping on the microscopic scale of the phase boundary seems unfeasible, but could potentially play a role in the equilibrated baryonic or quark matter

- At the temperatures occurring in the inspiral, baryonic and quark matter might be found in a (color-)superconducting/superfluid state. Cooper pairing will not only alter the hydrodynamics of the problem, but suppresses the flavor-changing rates used in this work, and will also alter diffusive transport, thus furder slowing down the phase boundary.

- In this work, we have assumed that thermal equilibration across the phase boundary is happening on a timescale slower than the evolution of the phase boundary. Towards the end of the equilibration process, however, the phase boundary slows down ([AH]do we have a section explaining this?. To not only reach flavor, but also thermal equilibration, thermal equilibration across the boundary has to be taken into account. Otherwise, the final state of the evolution will be wrongly described as a state where the baryonic and the quark matter phase stay at different temperatures.

- In this work, we assume that all profiles on the nuclear matter side are constant, and neglect any disturbance propagating ahead of the phase boundary.

- Furthermore, the assumption of a steady state, which we deem reasonable, should be further investigated.

- Our hydrodynamic description is based on an ideal fluid description. Dissipative corrections might be taken into account in the stress-energy tensor, or be included via weak interactions as described in Ref. [**?** ].

- If the phase boundary is driven by large-scale oscillations (as we expect it in the inspiral and the merger), the opposite scenario, where quark matter recombines to nuclear matter, has to be calculated. This will also allow us to estimate the strength of the so-called "phase conversion dissipation", a dissipative effect stemming from the out-of-phase movement of the actual phase boundary with the critical pressure [24].

- Cooper pairing: influence on diffusion, weak interaction rates, anything else?

- surface tension?

## ACKNOWLEDGEMENTS

We thank XXX (Nils Andersson? Suprovo Ghosh?) for valuable feedback and discussions. MGA, LB, and AH

are partly supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics, under Award No. #DE-FG02-05ER41375. A.H. furthermore acknowledges financial support by the UKRI under the Horizon Europe Guarantee project EP/Z000939/1.

## Appendix A: Bag model of quark matter

[MGA If we decide to impose electrical neutrality then this will need some rewriting.] In the approximation that we use for quark matter, ignoring electromagnetism and leptons, the pressure of quark matter is the sum of the pressures of the individual flavors,

$$\begin{aligned} P_{\rm QM}(\mu_q,\mu_K,T) &= P_q(\mu_q,T) + P_q(\mu_q{+}\mu_K/2,T) \\ &\quad + P_q(\mu_q{-}\mu_K/2,T) - U_{\rm bag}\,, \end{aligned} \tag{A1}$$

where $\mu_q = \mu_B/3$ is the quark chemical potential and $P_q$ is the pressure of a free Fermi gas of three colors of a single quark flavor. In this approximation the individual flavors have chemical potentials $\mu_u = \mu_B/3$, $\mu_d = \mu_B/3 + \mu_K/2$, and $\mu_s = \mu_B/3 - \mu_K/2$. We expect $\mu_K, T \ll \mu_q$ so we expand in powers of these to obtain

$$\begin{aligned} P_{\rm QM} &= -U_{\rm bag} + \frac{\mu_B^4}{108\pi^2} + \frac{\mu_B^2}{12\pi^2}\mu_K^2 + \frac{\mu_K^4}{32\pi^2} \\ &\quad + \frac{\mu_B^2}{6}T^2 + \frac{1}{4}\mu_K^2 T^2 + \frac{19\pi^2}{36}T^4 + \cdots\,, \end{aligned} \tag{A2}$$

[MGA Do we use this 4th order form of the EoS anywhere?][JS No, we never use it again: only the quadratic truncation (A3) is cited later (line 480 and the analytic derivation in Sec. IV C). Worth keeping, though: these are exactly the terms we'd need if we later want to estimate how far the analytic formula should deviate from the full numerical result, since they're what the quadratic truncation drops. We don't have that comparison anywhere in Sec. V yet.] or to quadratic order,

$$P_{\rm QM} = -U_{\rm bag} + \frac{\mu_B^4}{108\pi^2} + \frac{\mu_B^2}{12\pi^2}\mu_K^2 + \frac{\mu_B^2}{6}T^2\,, \tag{A3}$$

which implies that to leading order,

$$n_K = \left.\frac{\partial P_{\rm QM}}{\partial \mu_K}\right|_{\mu_B,T} = \frac{\mu_B^2\mu_K}{6\pi^2}\,, \tag{A4}$$

$$n_B = \left.\frac{\partial P_{\rm QM}}{\partial \mu_B}\right|_{\mu_K,T} = \frac{\mu_B^3}{27\pi^2}\,. \tag{A5}$$

We can also express the quark matter properties in terms of the K fraction $a$. To this order,

$$a \equiv \frac{n_K}{n_B} = \frac{9\mu_K}{2\mu_B} = \frac{3\mu_K}{2\mu_q}\,. \tag{A6}$$

## Appendix B: Strangeness creation

For positive $\mu_K$, the nonleptonic process relaxes the K density at the rate [46]

$$\Gamma_K(\mu_B,\mu_K,T) = \mathcal{A}_K \mu_u^5 \mu_K \left(\mu_K^2 + 4\pi^2 T^2\right)\,, \tag{B1}$$

[JS $u$ is a real participant, not an average: the reaction is $d+u \leftrightarrow u+s$ (Eq. (1)), with $u$ a spectator on both sides, like the spectator nucleon in modified Urca. Madsen's rate has the same $\mu_u^5$ prefactor for the same reason.][MGA In that case one would expect factors of $\mu_d$ and $\mu_s$ as well. I think Madsen is using $\mu_u$ as a symbol for something else, probably some average of chemical potentials. In our calculation everything should end up as a function of $\mu_B$, $\mu_K$, and $T$.] where

$$\mathcal{A}_K = \frac{16}{5\pi^5} G_F^2 \sin^2\theta_C \cos^2\theta_C\,, \tag{B2}$$

where $G_F = 1.16637\times10^{-11}\,{\rm MeV}^{-2}$ is the Fermi constant and $\theta_C = 13.02^\circ$ is the Cabibbo angle. We use $\mu_u = \mu_B/3 + 2\mu_Q/3$, where electric charge chemical potential $\mu_Q$ is fixed by charge neutrality, $2n_u - n_d - n_s = 0$. [MGA But in the previous section we said $\mu_u = \mu_B/3$. Which are we actually using? Anyway, as noted above $\mu_u$ shouldn't be needed.] [JS Both, in different places: the exact rate used in the numerical code uses the full $\mu_u = \mu_B/3 + 2\mu_Q/3$ with $\mu_Q$ solved exactly from charge neutrality, while only the analytic approximation in Sec. IV C drops $\mu_Q$ and uses $\mu_u \simeq \mu_q$. Not an inconsistency, just two different calculations.] [MGA (1) In general we should show the same formulas that we use in the numerical calculation. If we are making some approximation beyond the ones we use in our numerics then we should say so explicitly. (2) In Appendix A we didn't impose electric neutrality. Our overall approach is to ignore electrons and electromagnetism, so it seems inconsistent to impose electric neutrality.]

For the analytic approximation in Sec. IV C, we use the quadratic approximation of Sec. A to express the rate in terms of the K fraction. Using (A6),

$$\begin{aligned} \Gamma_K(a,T) &= n_B \gamma_K \left(a^3 + \eta a\right)\,, \\ \gamma_K &= \frac{128}{27\times5\pi^3} G_F^2 \sin^2\theta_C \cos^2\theta_C\, \mu_q^5\,, \\ \eta &= \frac{9\pi^2 T^2}{\mu_q^2}\,. \end{aligned} \tag{B3}$$

## Appendix C: Strangeness diffusion

The diffusion of the K flavor in quark matter is described by a diffusion coefficient $D_K$, which, since the quarks are relativistic, is closely related to the mean free path. This is determined by strong interactions, so we model it using the single gluon exchange perturbative expression [43, 47]

$$D_K^{-1} = \frac{8N_f\alpha_s^2}{\pi}\left(h_D \frac{T^{5/3}}{q_D^{2/3}} + \frac{\pi^3 T^2}{12 q_D}\right)\,, \tag{C1}$$

Here $N_f = 3$ is the number of active quark flavors, $\alpha_s = 0.3$ is the strong coupling, $h_D = \Gamma(\frac{8}{3})\zeta(\frac{5}{3})(2\pi)^{2/3} \simeq 1.81$ is a numerical prefactor, $q_D$ is the Debye wavenumber defined by $q_D^2 = g^2 N_f \mu_q^2/(2\pi^2)$, and $g^2 = 4\pi\alpha_s$. In our numerical calculations we set $\alpha_s = 0.3$ [MGA Can we find some standard citation about the running of the QCD coupling that would show that this is an appropriate value at energy scales of order 400 MeV?][35, 44] .

In (C1) the first term comes from the interaction mediated by Landau-damped transverse gluons, and the second term comes from Debye-screened longitudinal [MGA and transverse?] gluons. The first term is larger at temperatures up to $0.5\sqrt{\alpha_s}\mu_q$ which is of order 70 MeV, but because both terms have such similar $T$ dependence they are already of comparable size (to within at factor of 2) when the temperature rises above $0.04\sqrt{\alpha_s}\mu_q$ which is of order 7 MeV. In neutron stars mergers the temperature can range from below 1 MeV (early inspiral) to about 70 MeV, so for our numerical calculations we use the full expression, but for the analytic approximation of Sec. IV C we only use the Landau damping term.

[AH shall I add the derivation of K diffusion from individual quark currents that I have typed up in the notes?]





FIG. 7: Comparing three methods for $T(0^+) = 5$ MeV.

### Appendix D: Shape of the diffusive flux profile

The diffusive flux $g(x) \equiv -D_K(x)a'(x)$ enters $I_1$ (Eq. (31)) only through its shape as a function of $a$ rather than $x$: since $a(x)$ is monotonic, we can trade $x$ for $a$ and write $I_1 = \int_0^{a(0^+)} g(a)\, da$, the area under $g$ between the equilibrated downstream value $a = 0$, where $g = 0$, and the interface value $a = a(0^+)$, where the jump condition Eq. (29) fixes $g(a(0^+)) = \lambda_n u(0^-)[1 - a(0^+)]$.

Write $G \equiv g(a(0^+)) = \lambda_n u(0^-)[1 - a(0^+)]$ for this interface value. Three limiting shapes bracket the possibilities for the area under $g$. If $g$ rises linearly from 0 to $G$ over $0 \leq a \leq a(0^+)$, the area is a triangle, $I_1 = \frac{1}{2}Ga(0^+)$. If $g$ instead jumps to $G$ immediately and stays there, the area is the full rectangle, $I_1 = Ga(0^+)$. If $g$ instead stays near 0 over almost the whole range and only rises to $G$ at the very end, the area shrinks toward 0. These are the $\xi = 0$, $\xi = 1$, and $\xi = -1$ limits of

$$I_1 = \frac{1+\xi}{2} Ga(0^+)\,, \tag{D1}$$

which is just Eq. (33) written in terms of $G$: $\xi$ measures whether the diffusive flux builds up early ($\xi > 0$) or late ($\xi < 0$) across the conversion region, relative to a linear ramp ($\xi = 0$). It's worth noting that mathematically, for each input $T(0^-)$ and $n_B(0^-)$, there is one unique $\xi$ that make Eq. (33) exact.

The full numerical solutions to Eq. (22) are smooth between the two endpoints, and after comparing the actual plots of $g(a)$ using profiles that numerically solve Eq. (20), we find out that $\xi = -0.5$ is a good approximation, meaning that $g(a)$ is mostly convex. We use $\xi = -0.5$ in Eq. (34) and Eq. (38) throughout.

### Appendix E: Numerical method

[MGA If we want to keep this appendix then we should flesh it out with some info about how to solve the equations, e.g. about shooting with trial values of $u(0^-)$ to find the right value.]


We compactify the semi-infinite quark region using

$$x = -L_0 \ln(1 - s)\,, \qquad 0 \leq s \leq 1 - \epsilon_{\rm tail}\,, \tag{E1}$$


where $L_0$ is fixed from the initial interface diffusion–reaction scale. The result is checked for stability under changes of both $L_0$ and $\epsilon_{\rm tail}$. We first attempt the requested $a(0^+)$ directly; if necessary, we continue from a smaller interface fraction and use each converged profile as the initial guess for the next step. States with negative temperature, nonpositive density, a failed equation-of-state closure, or nonphysical transport coefficients are rejected. These profiles provide the numerical reference against which the piecewise-constant result in Eq. (38) is tested.

### Appendix F: Local thermal equilibrium

However, $T(0^+) = 0$ lies outside the validity of the present diffusion treatment. As $T(0^+) \to 0$, the quark mean free path $\lambda_Q^{\rm mfp}$ diverges, so quark matter cannot remain in local thermal equilibrium (LTE) across a conversion layer with a finite variation scale.[JS Think about where to put this LTE. Doesn't seem it belong to piecewise constant subsection, not in numerical] Requiring the mean free path to remain shorter than the scale on which the




(a)




(b)

FIG. 8: Dependence of the numerical phase-boundary speed on the prescribed interface temperature $T(0^+)$. Panel (a) shows $u(0^-)$ as a function of $n_B(0^-)$ at fixed $T(0^-) = 20\,\mathrm{MeV}$, while panel (b) shows $u(0^-)$ as a function of $T(0^-)$ at fixed $n_B(0^-) = 3.5n_0$. The velocity axis is logarithmic, the markers denote the numerical slice samples, and the line color runs from colder (purple and blue) to hotter (red) values of $T(0^+)$. The gray regions are excluded in the corresponding slice by stable neutron matter or by $A \geq 1$.

diffusion coefficient changes,

$$\lambda_Q^{\mathrm{mfp}} = 3D_K < \frac{D_K}{|dD_K/dx|}\,, \tag{F1}$$

and using the constant-pressure trajectory and interface gradient gives

$$5u(0^-)\lambda_n a(0^+)[1-a(0^+)] < A^2 - a^2(0^+)\,. \tag{F2}$$

The LTE-limited composition is obtained by saturating this inequality,

$$A^2 - \left[a^{\mathrm{LTE}}(0^+)\right]^2 = 5\lambda_n u(0^-) a^{\mathrm{LTE}}(0^+)\left[1 - a^{\mathrm{LTE}}(0^+)\right]\,, \tag{F3}$$

with $u(0^-)$ evaluated at $a(0^+)$ from Eq. (38).

Finally, the full numerical solutions confirm the scale estimate: across the conversion layer, the relative variations of $n_B$ and $P$ are typically no larger than $10^{-10}$ and $10^{-12}$, respectively. The piecewise-constant approximation removes variation that is numerically invisible while retaining the composition and temperature profiles that set the conversion rate.


[1] A. R. Counsell, F. Gittins, N. Andersson and I. Tews, *Interface Modes in Inspiralling Neutron Stars: A Gravitational-Wave Probe of First-Order Phase Transitions*, *Phys. Rev. Lett.* **135** (2025) 081402, [2504.06181].

[2] J. P. Pereira, L. Tonetto, M. Bejger, J. L. Zdunik and P. Haensel, *Dynamical Tides in Neutron Stars with First-Order Phase Transitions: The Role of the Discontinuity Mode*, *Phys. Rev. Lett.* **135** (2025) 231401, [2504.16911].

[3] L. Tonetto and G. Lugones, *Discontinuity gravity modes in hybrid stars: assessing the role of rapid and slow phase conversions*, *Phys. Rev. D* **101** (2020) 123029, [2003.01259].

[4] T. Fischer, I. Sagert, G. Pagliara, M. Hempel, J. Schaffner-Bielich, T. Rauscher et al., *Core-Collapse Supernova Explosions Triggered by a Quark-Hadron Phase Transition During the Early Post-Bounce Phase*,

*Astrophys. J. Suppl.* **194** (2011) 39, [1011.3409].

[5] T. Fischer, N.-U. F. Bastian, M.-R. Wu, P. Baklanov, E. Sorokina, S. Blinnikov et al., *Quark deconfinement as a supernova explosion engine for massive blue-supergiant stars*, *Nature Astron.* **2** (2018) 980–986, [1712.08788].

[6] T. Fischer, *QCD phase transition drives supernova explosion of a very massive star*, *Eur. Phys. J. A* **57** (2021) 270, [2108.00196].

[7] K. Bleau, J. Kopp, J. Lee and J. van de Vis, *High-Frequency Gravitational Waves from Phase Transitions in Nascent Neutron Stars*, 2603.18153.

[8] E. R. Most, L. J. Papenfort, V. Dexheimer, M. Hanauske, S. Schramm, H. Stöcker et al., *Signatures of quark-hadron phase transitions in general-relativistic neutron-star mergers*, *Phys. Rev. Lett.* **122** (2019) 061101, [1807.03684].

[9] A. Bauswein, N.-U. F. Bastian, D. B. Blaschke, K. Chatziioannou, J. A. Clark, T. Fischer et al., *Identifying a first-order phase transition in neutron star mergers through gravitational waves*, *Phys. Rev. Lett.* **122** (2019) 061102, [1809.01116].

[10] P. L. Espino, A. Prakash, D. Radice and D. Logoteta, *Revealing phase transition in dense matter with gravitational wave spectroscopy of binary neutron star mergers*, *Phys. Rev. D* **109** (2024) 123009, [2301.03619].

[11] L. R. Weih, M. Hanauske and L. Rezzolla, *Postmerger Gravitational-Wave Signatures of Phase Transitions in Binary Mergers*, *Phys. Rev. Lett.* **124** (2020) 171103, [1912.09340].

[12] R. Harada, K. Cannon, K. Hotokezaka and K. Kyutoku, *Testability of the quark-hadron transition using gravitational waves from merging binary neutron stars*, *Phys. Rev. D* **110** (2024) 123005, [2310.13603].

[13] A. Bauswein and S. Blacker, *Impact of quark deconfinement in neutron star mergers and hybrid star mergers*, *Eur. Phys. J. ST* **229** (2020) 3595–3604, [2006.16183].

[14] S. Haque, R. Mallick and S. K. Thakur, *Effects of onset of phase transition on binary neutron star mergers*, *Mon. Not. Roy. Astron. Soc.* **527** (2023) 11575–11586, [2207.14485].

[15] Y. Fujimoto, K. Fukushima, K. Hotokezaka and K. Kyutoku, *Gravitational Wave Signal for Quark Matter with Realistic Phase Transition*, *Phys. Rev. Lett.* **130** (2023) 091404, [2205.03882].

[16] J.-E. Christian, J. Schaffner-Bielich and S. Rosswog, *Which first order phase transitions to quark matter are possible in neutron stars?*, *Phys. Rev. D* **109** (2024) 063035, [2312.10148].

[17] Y. Fujimoto, K. Fukushima, K. Hotokezaka and K. Kyutoku, *Signature of hadron-quark crossover in binary-neutron-star mergers*, *Phys. Rev. D* **111** (2025) 063054, [2408.10298].

[18] A. V. Olinto, *On the Conversion of Neutron Stars Into Strange Stars*, *Phys. Lett. B* **192** (1987) 71.

[19] J. E. Horvath and O. G. Benvenuto, *On the Stability of Slow Neutron Combustion in Astrophysical Objects*, *Phys. Lett. B* **213** (1988) 516–520.

[20] H. Heiselberg, G. Baym and C. J. Pethick, *Burning of strange quark matter and transport properties of QCD plasmas*, *Nucl. Phys. B Proc. Suppl.* **24** (1991) 144–147.

[21] G. Lugones, O. G. Benvenuto and H. Vucetich, *Combustion of nuclear matter into strange matter*, *Phys. Rev. D* **50** (1994) 6100–6109.

[22] D. Leahy and R. Ouyed, *Supernova SN2006gy as a first ever Quark Nova?*, *Mon. Not. Roy. Astron. Soc.* **387** (2008) 1193, [0708.1787].

[23] G. Pagliara, M. Herzog and F. K. Röpke, *Combustion of a neutron star into a strange quark star: The neutrino signal*, *Phys. Rev. D* **87** (2013) 103007, [1304.6884].

[24] M. G. Alford, S. Han and K. Schwenzer, *Phase conversion dissipation in multicomponent compact stars*, *Phys. Rev. C* **91** (2015) 055804, [1404.5279].

[25] A. Drago and G. Pagliara, *Combustion of a hadronic star into a quark star: the turbulent and the diffusive regimes*, *Phys. Rev. C* **92** (2015) 045801, [1506.08337].

[26] S. Furusawa, T. Sanada and S. Yamada, *Hydrodynamical study on the conversion of hadronic matter to quark matter. II. Diffusion-induced conversion*, *Phys. Rev. D* **93** (2016) 043019, [1511.08153].

[27] B. Niebergal, R. Ouyed and P. Jaikumar, *Numerical Simulation of the Hydrodynamical Combustion to Strange Quark Matter*, *Phys. Rev. C* **82** (2010) 062801, [1008.4806].

[28] R. Ouyed, *The Macro-Physics of the Quark-Nova: Astrophysical Implications*, *Universe* **8** (2022) 322.

[29] S. Furusawa, T. Sanada and S. Yamada, *Hydrodynamical study on the conversion of hadronic matter to quark matter: I. Shock-induced conversion*, *Phys. Rev. D* **93** (2016) 043018, [1511.08148].

[30] A. Bhattacharyya, S. K. Ghosh, P. S. Joardar, R. Mallick and S. Raha, *The conversion of Neutron star to Strange star: A two step process*, *Phys. Rev. C* **74** (2006) 065804, [astro-ph/0606523].

[31] M. G. Alford, L. Brodie, A. Haber and I. Tews, *Relativistic mean-field theories for neutron-star physics based on chiral effective field theory*, *Phys. Rev. C* **106** (2022) 055804, [2205.10283].

[32] M. G. Alford, L. Brodie, A. Haber and I. Tews, *Tabulated equations of state from models informed by chiral effective field theory*, *Phys. Scripta* **98** (2023) 125302, [2304.07836].

[33] M. G. Alford, L. Brodie, A. Haber and I. Tews, "ABHT(QMC-RMF3) Equation of State." https://compose.obspm.fr/eos/299.

[34] A. Chodos, R. L. Jaffe, K. Johnson, C. B. Thorn and V. F. Weisskopf, *A new extended model of hadrons*, *Phys. Rev. D* **9** (1974) 3471–3495.

[35] E. Farhi and R. L. Jaffe, *Strange matter*, *Phys. Rev. D* **30** (1984) 2379–2390.

[36] I. Bombaci and B. Datta, *Conversion of neutron stars to strange stars as the central engine of gamma-ray bursts*, *Astrophys. J. Lett.* **530** (2000) L69, [astro-ph/0001478].

[37] J. A. Rosero-Gil and G. Lugones, *Electroweak decay of quark matter within dense astrophysical combustion flames*, *Phys. Rev. C* **103** (2021) 055816, [2010.04861].

[38] M. G. Alford and A. Schmitt, *Bulk viscosity in 2SC quark matter*, *J. Phys. G* **34** (2007) 67–102, [nucl-th/0608019].

[39] M. Alford, A. Harutyunyan, A. Sedrakian and S. Tsiopelas, *Bulk viscosity of two-flavor color superconducting quark matter in neutron star mergers*, *Front. Astron. Space Sci.* **12** (2025) 1648066, [2506.08144].

[40] B. A. Sa'd, I. A. Shovkovy and D. H. Rischke, *Bulk viscosity of strange quark matter: Urca versus non-leptonic processes*, *Phys. Rev. D* **75** (2007) 125004, [astro-ph/0703016].

[41] C. Eckart, *The Thermodynamics of irreversible processes. 3.. Relativistic theory of the simple fluid*, *Phys. Rev.* **58** (1940) 919–924.

[42] S. K. Lander and N. Andersson, *Heat conduction in rotating relativistic stars*, *Mon. Not. Roy. Astron. Soc.* **479** (2018) 4207–4215, [1804.01520].

[43] H. Heiselberg and C. J. Pethick, *Transport and relaxation in degenerate quark plasmas*, *Phys. Rev. D* **48** (1993) 2916–2928.

[44] J. Madsen, *Physics and astrophysics of strange quark matter*, *Lect. Notes Phys.* **516** (1999) 162–203, [astro-ph/9809032].

[45] T. Zhao, P. B. Rau, A. Haber, S. P. Harris, C. Constantinou and S. Han, *Suppression of Composition g-modes in Chemically Equilibrating Warm Neutron Stars*, *Astrophys. J.* **993** (2025) 161, [2504.12230].

[46] J. Madsen, *Rate of the weak reaction $s + u$ to $u + d$ in quark matter*, *Phys. Rev. D* **47** (1993) 325–330.

[47] P. Haensel and R. Schaeffer, *Bulk viscosity of hot-neutron-star matter from direct URCA processes*, *Phys. Rev. D* **45** (1992) 4708–4712.