

# Kalman Filter

## Setup

Observable equation is described by

$$y_t = x_t + \eta_t \quad (1)$$

where  $y_t$  can be observed at time  $t$  and  $\eta_t \sim N(0, \sigma_\eta^2)$  is a noise shock which prevent to correctly observe state  $x_t$ .

State transition equation is described by

$$x_t = x_{t-1} + \varepsilon_t \quad (2)$$

where  $\varepsilon_t \sim N(0, \sigma_\varepsilon^2)$  is a structural shock which affects the transition from  $x_{t-1}$  to  $x_t$ .

## Procedure

Goal is to optimize the forecast of  $x_{t+1}$  using available information at time  $t$ , i.e.  $y_t$ . Assume an initial value for  $x_{1,0}$  and  $Var(x_{1,0} - x_{0,0}) = P_{1,0}$ . Procedure can be summarized as follows,

1. Forecast  $y_t$  using information at time  $t - 1$  and evaluate the error variance of this prediction.
2. Infer  $x_t$  using information at time  $t$  and evaluate the error variance of this inference.
3. Forecast  $x_{t+1}$  using information at time  $t$  and evaluate the error variance of this prediction.
4. Find the steady state variance  $P_{t,t}$ .

Given  $x_{1,0}$  the three initial steps are

1.  $y_{1,0} = x_{1,0}$  and the forecast error variance is  $\Omega_{1,0}^y = \sigma_\eta^2$ .
2.  $x_{1,1} = y_1$  and the forecast error variance is  $P_{1,1} = \sigma_\eta^2$ .
3.  $x_{2,1} = x_{1,1}$  and the forecast error variance is  $P_{2,1} = \sigma_\eta^2 + \sigma_\varepsilon^2$ .

As a generalization, the three steps are

1.  $y_{t,t-1} = x_{t,t-1}$  and the forecast error variance is  $\Omega_{t,t-1}^y = Var(x_{t,t-1}) + \sigma_\eta^2 = P_{t,t-1} + \sigma_\eta^2$ .
2. We want to forecast  $x_{t,t}$  using all the available information up to time  $t$ . As a simplification we try to forecast  $x_t - x_{t,t-1}$  using  $y_t - x_{t,t-1}$ . Coefficient  $\beta^{PROJ}$  is derived as follows

$$\begin{aligned} \beta^{KG} &= \frac{Cov(x_t - x_{t,t-1}, y_t - x_{t,t-1})}{Var(y_t - x_{t,t-1})} \\ &= \frac{Cov(x_t - x_{t,t-1}, x_t + \eta_t - x_{t,t-1})}{Var(x_t + \eta_t - x_{t,t-1})} \\ &= \frac{P_{t,t-1}}{(P_{t,t-1} + \sigma_\eta^2)} \\ &= \frac{P_{t,t-1}}{P_{t,t-1} + \sigma_\eta^2} \end{aligned} \quad (3)$$

This implies that

$$x_{t,t} - x_{t,t-1} = \beta^{KG}(y_t - x_{t,t-1}) \quad (4)$$

which is

$$\begin{aligned} x_{t,t} &= x_{t,t-1} + \beta^{KG}(y_t - x_{t,t-1}) \\ &= (1 - \beta^{KG})x_{t,t-1} + \beta^{KG}y_t \end{aligned} \quad (5)$$

and this is the reason why  $\beta^{KG}$  is called Kalman gain, i.e. it defines how much to weight new information at time  $t$  to infer  $x_{t,t}$ .

Now we need to figure out the forecast error variance of  $x_t - x_{t,t}$ , i.e.  $P_{t,t} = \text{Var}(x_t - x_{t,t})$

$$\begin{aligned} P_{t,t} &= \text{Var} \left[ x_{t,t} - x_{t,t-1} - \beta^{KG}(y_t - x_{t,t-1}) \right] \\ &= \text{Var} \left[ x_{t,t} - x_{t,t-1} - \beta^{KG}(x_t + \eta_t - x_{t,t-1}) \right] \\ &= \text{Var} \left[ x_{t,t} - x_{t,t-1} - \frac{P_{t,t-1}}{P_{t,t-1} + \sigma_\eta^2}(x_t + \eta_t - x_{t,t-1}) \right] \\ &= E \left\{ \left[ x_{t,t} - x_{t,t-1} - \frac{P_{t,t-1}}{P_{t,t-1} + \sigma_\eta^2}(x_t + \eta_t - x_{t,t-1}) \right]^2 \right\} \\ &= E \left\{ \left[ x_{t,t} - x_{t,t-1} \right]^2 \right\} - 2E \left\{ \left( x_{t,t} - x_{t,t-1} \right) \left( \frac{P_{t,t-1}}{P_{t,t-1} + \sigma_\eta^2}(x_t + \eta_t - x_{t,t-1}) \right) \right\} \\ &\quad + E \left\{ \left[ \frac{P_{t,t-1}}{P_{t,t-1} + \sigma_\eta^2}(x_t + \eta_t - x_{t,t-1}) \right]^2 \right\} \\ &= P_{t,t-1} - 2P_{t,t-1} \frac{P_{t,t-1}}{P_{t,t-1} + \sigma_\eta^2} + P_{t,t-1} \frac{P_{t,t-1}}{P_{t,t-1} + \sigma_\eta^2} \\ &= P_{t,t-1} - P_{t,t-1} \frac{P_{t,t-1}}{P_{t,t-1} + \sigma_\eta^2} \\ &= P_{t,t-1} \left[ 1 - \frac{P_{t,t-1}}{P_{t,t-1} + \sigma_\eta^2} \right] \\ &= P_{t,t-1}(1 - \beta^{KG}) \end{aligned} \quad (6)$$

3.  $x_{t+1,t} = x_{t,t}$  and the forecast error variance is  $P_{t+1,t} = P_{t,t} + \sigma_\varepsilon^2$  which is

$$\begin{aligned} P_{t+1,t} &= P_{t,t} + \sigma_\varepsilon^2 \\ &= P_{t,t-1} \left[ 1 - \frac{P_{t,t-1}}{P_{t,t-1} + \sigma_\eta^2} \right] + \sigma_\varepsilon^2 \end{aligned} \quad (7)$$

4. Find the steady state value of  $P$  as follows

$$\begin{aligned} P &= P \left[ 1 - \frac{P}{P + \sigma_\eta^2} \right] + \sigma_\varepsilon^2 \\ &= P - \frac{P^2}{P + \sigma_\eta^2} + \sigma_\varepsilon^2 \end{aligned} \quad (8)$$

which is

$$P^2 + \sigma_\eta^2 P = P^2 + \sigma_\eta^2 P - P^2 + \sigma_\varepsilon^2 P + \sigma_\varepsilon^2 \sigma_\eta^2 \quad (9)$$

which is

$$P^2 - \sigma_\eta^2 P - \sigma_\varepsilon^2 \sigma_\eta^2 = 0 \quad (10)$$

which is

$$P = \frac{1}{2} \left( \sigma_\varepsilon^2 + \sqrt{\sigma_\varepsilon^4 + 4\sigma_\varepsilon^2 \sigma_\eta^2} \right) \quad (11)$$