

A Framework Toward the Millennium Prize Problem on the Navier–Stokes Equations

Thomas Birnie + ChatGPT

February 26, 2025

Abstract

We provide a comprehensive and rigorously detailed L^AT_EX document outlining a potential framework for solving the global regularity problem of the three-dimensional incompressible Navier–Stokes equations. While this document describes a complete *strategy* with all intermediate steps, we emphasize that the final resolution of this famous open problem remains a subject of ongoing investigation.

Contents

1 Introduction

The Navier–Stokes equations lie at the heart of fluid mechanics, describing the motion of viscous, incompressible fluids. Despite their widespread use in physical applications, the fundamental mathematical question of whether smooth solutions exist for all time in three spatial dimensions remains open. This challenge is recognized as one of the *Millennium Prize Problems* posed by the Clay Mathematics Institute.

In this document, we present a framework containing the standard techniques employed in partial regularity proofs, global energy estimates, compactness arguments, and convergence analyses. Although many of these methods have yielded deep insights, a definitive solution remains elusive. The structure here reflects a best-practice approach to writing a rigorous manuscript that, if completed successfully, would meet the high standards required by the Clay Institute.

2 Statement of the Problem

We consider the three-dimensional incompressible Navier–Stokes equations:

$$\begin{cases} \partial_t \mathbf{u} + (\mathbf{u} \cdot \nabla) \mathbf{u} = -\nabla p + \nu \Delta \mathbf{u}, \\ \nabla \cdot \mathbf{u} = 0, \end{cases} \quad (2.1)$$

where $\mathbf{u}(t, \mathbf{x}) \in \mathbb{R}^3$ is the velocity field, $p(t, \mathbf{x}) \in \mathbb{R}$ is the pressure, and $\nu > 0$ is the kinematic viscosity. The domain is typically $\mathbb{R}^3$ or a subset with appropriate boundary conditions (e.g., periodic or Dirichlet). Initial data is given by

$$\mathbf{u}(0, \mathbf{x}) = \mathbf{u}_0(\mathbf{x}), \quad \nabla \cdot \mathbf{u}_0 = 0.$$

Definition 2.1 (Global Existence and Smoothness Problem). *Show that for any sufficiently smooth, divergence-free initial data $\mathbf{u}_0$ with finite energy, there exists a unique, smooth solution $\mathbf{u}(t, \mathbf{x})$ to (??) for all $t \geq 0$, and that this solution remains smooth for all time.*

3 Preliminaries

Here, we gather fundamental results, tools, and definitions needed throughout the paper. This includes Sobolev spaces, embeddings, the classical Aubin–Lions compactness lemma, and other widely used PDE methods.

3.1 Sobolev spaces and embeddings

Let $\Omega \subset \mathbb{R}^3$. Denote by $H^s(\Omega)$ the Sobolev space of functions with square-integrable weak derivatives up to order s . Key inequalities:

- *Gagliardo–Nirenberg Inequality*
- *Ladyzhenskaya Inequality*
- *Poincaré Inequality*

We recall that for $1 \leq p < \infty$, the norm in $W^{s,p}(\Omega)$ is

$$\|f\|_{W^{s,p}} = \left(\sum_{|\alpha| \leq s} \int_{\Omega} |D^{\alpha} f|^p d\mathbf{x} \right)^{1/p}.$$

All relevant inequalities will be stated precisely where needed.

3.2 Aubin–Lions lemma

We state a version of the Aubin–Lions lemma crucial for compactness arguments:

Lemma 3.1 (Aubin–Lions). *Let $B_0 \subset B_1 \subset B_2$ be Banach spaces with compact embedding $B_1 \hookrightarrow B_0$ and continuous embedding $B_2 \hookrightarrow B_1$. Then for $1 \leq p \leq \infty$, if $\{u_n\}$ is a sequence such that*

$$u_n \in L^p(0, T; B_2), \quad \partial_t u_n \in L^1(0, T; B_0),$$

then there exists a strongly convergent subsequence in $L^p(0, T; B_1)$.

Proof. See [?, ?] for detailed proofs and variants. □

4 Discrete Formulations

In many modern approaches, we rely on discretization schemes (e.g., Galerkin methods) to approximate solutions and then pass to the limit to show existence of smooth solutions. Here we outline the discrete operators, the resulting system, and key properties such as stability.

4.1 Galerkin approximation

Let $\{\mathbf{w}_j(\mathbf{x})\}_{j=1}^{\infty}$ be a suitable orthonormal basis for divergence-free vector fields in $L^2(\Omega)$. For each $n \in \mathbb{N}$, define the finite-dimensional subspace

$$V_n = \text{span}\{\mathbf{w}_1, \dots, \mathbf{w}_n\}.$$

We seek approximations $\mathbf{u}_n(t) \in V_n$ such that

$$\begin{aligned} \frac{d}{dt}(\mathbf{u}_n, \mathbf{w}_j) + ((\mathbf{u}_n \cdot \nabla) \mathbf{u}_n, \mathbf{w}_j) + \nu(\nabla \mathbf{u}_n, \nabla \mathbf{w}_j) &= 0, & \text{for all } j = 1, \dots, n, \\ (\mathbf{u}_n(0), \mathbf{w}_j) &= (\mathbf{u}_0, \mathbf{w}_j). \end{aligned} \tag{4.1}$$

Here $(\cdot, \cdot)$ denotes the $L^2(\Omega)$ inner product.

4.2 Discrete energy estimates

The usual procedure is to show that these approximations satisfy a discrete energy inequality. For instance:

$$\frac{1}{2} \frac{d}{dt} \|\mathbf{u}_n\|_{L^2}^2 + \nu \|\nabla \mathbf{u}_n\|_{L^2}^2 \leq 0.$$

Lemma 4.1 (Discrete Energy Inequality). *Let $\mathbf{u}_n$ be a Galerkin approximation solving (??). Then for $0 \leq t \leq T$,*

$$\|\mathbf{u}_n(t)\|_{L^2}^2 + 2\nu \int_0^t \|\nabla \mathbf{u}_n(\tau)\|_{L^2}^2 d\tau \leq \|\mathbf{u}_0\|_{L^2}^2.$$

Proof. Test (??) by $\mathbf{u}_n$ and use incompressibility to handle the nonlinear term. The Grönwall lemma is then applied. $\square$

5 Mapping Between Discrete and Continuous Formulations

A key step is to rigorously connect the finite-dimensional approximations to the infinite-dimensional PDE. We show that in the limit $n \rightarrow \infty$, $\mathbf{u}_n$ converges to a function $\mathbf{u}$ that satisfies the Navier–Stokes equations in the weak or strong sense.

5.1 Weak formulation

In the usual weak formulation, one tests with smooth divergence-free fields. Convergence of $\mathbf{u}_n$ to $\mathbf{u}$ must preserve all relevant properties:

- (i) $\mathbf{u}_n \rightharpoonup \mathbf{u}$ in appropriate Sobolev spaces,
- (ii) $\nabla \cdot \mathbf{u} = 0$ almost everywhere,
- (iii) The nonlinear term $(\mathbf{u}_n \cdot \nabla) \mathbf{u}_n$ converges to $(\mathbf{u} \cdot \nabla) \mathbf{u}$ in the sense of distributions.

Remark 5.1. *Compactness results such as the Aubin–Lions lemma (see Lemma ??) are instrumental.*

6 Proof of Compactness

We now state and prove the main compactness theorem for the sequence $\{\mathbf{u}_n\}$. The general template involves showing that $\{\mathbf{u}_n\}$ is bounded in a sufficiently strong norm and that its time derivative is bounded in a weaker norm, thus allowing an application of Aubin–Lions.

Theorem 6.1 (Compactness of Discrete Solutions). *Let $\{\mathbf{u}_n\}$ be the Galerkin approximations. Assume that the initial data $\mathbf{u}_0 \in H^1(\Omega)$ with $\nabla \cdot \mathbf{u}_0 = 0$. Then there exists a subsequence (still denoted $\mathbf{u}_n$) and a function $\mathbf{u}$ such that*

$$\mathbf{u}_n \rightarrow \mathbf{u} \quad \text{strongly in } L^2(0, T; H^1(\Omega)),$$

and $\mathbf{u}$ satisfies the incompressible Navier–Stokes equations in the weak sense.

Proof Sketch. **Step 1: Uniform bounds.** From the discrete energy inequality, we have uniform boundedness of $\mathbf{u}_n$ in $L^\infty(0, T; L^2(\Omega))$ and $\nabla \mathbf{u}_n$ in $L^2(0, T; L^2(\Omega))$.

Step 2: Time derivative bounds. Differentiating (??) and using orthogonality properties, one obtains $\partial_t \mathbf{u}_n$ bounded in a negative-order Sobolev space $H^{-1}(\Omega)$, ensuring that

$$\partial_t \mathbf{u}_n \in L^2(0, T; H^{-1}(\Omega)).$$

Step 3: Application of Aubin–Lions. We invoke Lemma ?? with

$$B_2 = H^1(\Omega), \quad B_1 = L^2(\Omega), \quad B_0 = H^{-1}(\Omega).$$

Hence, $\{\mathbf{u}_n\}$ is relatively compact in $L^2(0, T; H^1(\Omega))$. A diagonal argument yields a strongly convergent subsequence.

Step 4: Identification of the limit. Passing to the limit in the Galerkin scheme, we verify that the limit $\mathbf{u}$ satisfies the weak form of the Navier–Stokes equations. $\square$

7 Proof of Energy Bounds

Beyond the basic L^2 energy estimate, one typically seeks higher-order energy bounds (in H^s norms) to argue for smoothness. In 2D, such bounds can be made global in time, but in 3D the bounding technique meets well-known difficulties.

7.1 Higher Sobolev norm estimates

We outline the derivation of an H^1 estimate for $\mathbf{u}$. Testing the Navier–Stokes equations with $-\Delta \mathbf{u}$ can yield:

$$\frac{1}{2} \frac{d}{dt} \|\nabla \mathbf{u}\|_{L^2}^2 + \nu \|\Delta \mathbf{u}\|_{L^2}^2 \leq C \|\nabla \mathbf{u}\|_{L^2}^2 \|\mathbf{u}\|_{L^4}^2,$$

where one applies Sobolev embedding to handle the nonlinear term. The constant C may depend on domain geometry and embedding constants. Similar estimates can be attempted for higher derivatives.

Remark 7.1. *The key unresolved issue is whether these estimates can be closed globally in 3D (leading to a blowup criterion if not). This is precisely the crux of the Millennium problem.*

8 Proof of Convergence

Once compactness and uniform (in n) energy bounds are established, one shows weak and then strong convergence of the discrete solutions to a smooth solution $\mathbf{u}$. Typical steps:

Theorem 8.1 (Weak Convergence). *With the hypotheses of Theorem ??, the limit function $\mathbf{u}$ is a weak solution, meaning:*

$$\int_0^T \int_\Omega \left(\mathbf{u} \cdot \partial_t \phi + (\mathbf{u} \otimes \mathbf{u}) : \nabla \phi + \nu \nabla \mathbf{u} : \nabla \phi \right) d\mathbf{x} dt = 0,$$

for all smooth, divergence-free test functions ϕ that vanish at $t = 0$ and $t = T$.

Proof. Standard limiting procedure: pass to the limit in the Galerkin formulation (??). Use the compactness to handle the nonlinear terms, and identify the weak limit in each term. $\square$

Theorem 8.2 (Strong Convergence). *Under additional assumptions (e.g., higher regularity of initial data), $\mathbf{u}_n \rightarrow \mathbf{u}$ strongly in $L^2(0, T; H^1(\Omega))$, and $\mathbf{u}$ is a strong solution if it meets the PDE in the classical sense and has all required derivatives in L^2 .*

Proof Sketch. A more refined compactness argument using improved estimates on higher Sobolev norms of $\mathbf{u}_n$. Under these assumptions, standard energy methods plus the Aubin–Lions lemma yield strong convergence. $\square$

9 Toward a Proof of Global Smoothness

9.1 Heuristic outline

A classical approach to global smoothness would argue as follows:

- (1) Show that the sequence of approximate solutions remains bounded in arbitrarily high Sobolev norms.
- (2) Use bootstrapping and Sobolev embeddings to promote the solution to classical regularity (smoothness).
- (3) Conclude no finite-time singularity can develop because the blow-up scenario is precluded by the energy estimates and regularity arguments.

9.2 Open problems and partial results

Despite substantial progress:

- Known partial regularity results (Caffarelli–Kohn–Nirenberg, Scheffer, etc.) guarantee that potential singularities, if they exist, occupy a set of measure zero in space-time.
- Scaling arguments and blow-up criteria (e.g., Beale–Kato–Majda for 2D Euler) highlight the difficulty in controlling the nonlinear term in 3D.

Remark 9.1. *Currently, no known method closes the final gap for 3D global regularity. However, the argument structure provided here—energy inequalities, compactness, and Sobolev regularity—is the cornerstone of potential future proofs.*

10 Conclusion

We have presented a rigorous *framework* for addressing the Millennium Prize Problem on the Navier–Stokes equations, mirroring the structure that any final proof would likely require:

- **Preliminaries:** Sobolev spaces, key inequalities, Aubin–Lions lemma.
- **Discrete Formulation:** Galerkin approximations, discrete energy estimates.
- **Compactness and Convergence:** Passing to the limit to obtain weak and strong solutions.
- **Higher Regularity and Smoothness:** Energy bounds in higher Sobolev spaces, plus the conceptual blueprint for proving global smoothness.

While the final, decisive step to show that *no singularities form in finite time* for the 3D case remains open, the methods outlined here represent the rigorous backbone that any successful solution would build upon. It is our hope that clarifying this structure in a cohesive manner fosters deeper understanding and catalyzes future breakthroughs in the Navier–Stokes regularity question.

A Additional Technical Details

A.1 Inequalities and their proofs

A compendium of classical inequalities used (Poincaré, Gagliardo–Nirenberg, etc.) can be found in standard references [?, ?].

A.2 Compatibility conditions and boundary terms

For domain-boundary constraints (e.g., no-slip conditions), further details on elliptic regularity and boundary-layer phenomena must be added for a full treatment.

References

- [1] P. Constantin, C. Foias. *Navier–Stokes Equations*. Chicago Lectures in Mathematics, 1988.
- [2] L. C. Evans. *Partial Differential Equations*. American Mathematical Society, 2010.
- [3] R. Temam. *Navier–Stokes Equations, Theory and Numerical Analysis*. North-Holland, 1977.