

## How the Proof Works

For the global three-dimensional Navier–Stokes regularity problem, take one smooth finite-energy flow with fixed viscosity  $\nu > 0$ :

$$\partial_t u + (u \cdot \nabla)u + \nabla p = \nu \Delta u, \quad \nabla \cdot u = 0, \quad u(0) = u_0.$$

The danger is that this flow could sharpen one of its own structures beyond smooth continuation in finite time. A vortex can concentrate velocity gradients while it stretches. As surrounding strain pulls the tube along its axis, incompressibility forces it to contract across its width: a longer piece of fluid cannot keep its old cross-section without gaining volume. Aligned stretching can also strengthen the rotation. The core narrows, the velocity turns more sharply across it, and the gradients rise. This nonlinear sharpening is opposed by viscosity, which spreads the concentrated vorticity into its surroundings. Both effects can grow stronger as the core gets smaller.

To shrink a width by a factor  $\lambda > 1$  at the Navier–Stokes scale, multiply velocity by  $\lambda$  and shorten time by  $\lambda^2$ :

$$u_\lambda(x, t) = \lambda u(\lambda x, \lambda^2 t), \quad p_\lambda(x, t) = \lambda^2 p(\lambda x, \lambda^2 t).$$

Nonlinear transport and viscosity both gain a factor  $\lambda^3$ : viscosity has not gained a scaling advantage over the sharpening. Equal scaling weight does not tell us which effect wins dynamically, or give a bound on either one.

The faster motion occupies less volume. Velocity squared gains  $\lambda^2$ , while volume loses  $\lambda^3$ , so kinetic energy scales down by  $\lambda^{-1}$ . It can remain finite as velocity and gradients grow. The Sobolev norms register this concentration at different spatial heights:

$$\|u_\lambda(\cdot, t)\|_{\dot{H}^s} = \lambda^{s-1/2} \|u(\cdot, \lambda^2 t)\|_{\dot{H}^s}.$$

Below  $s = \frac{1}{2}$ , the scaled norm falls; above it, the norm rises. At  $s = \frac{1}{2}$ , its size is unchanged by this concentration. That is the critical height:  $\dot{H}^{1/2}$  is scaling-critical, not conserved in time.

If the vortex kept repeating this narrowing and acceleration, its width, velocity scale, and duration at stage  $j$  would follow

$$\begin{aligned} \ell_j &= \lambda^{-j} \ell_0, & U_j &= \lambda^j U_0, & \tau_j &= \lambda^{-2j} \tau_0, \\ \sum_{j=0}^{\infty} \tau_j &= \frac{\tau_0}{1 - \lambda^{-2}} < \infty. \end{aligned}$$

The widths collapse and the velocity scale diverges, yet infinitely many stages fit into a finite interval. This is the hypothetical Zeno cascade: finite energy can survive while the core becomes arbitrarily narrow and fast.

In order for a singularity to happen, velocity has to blow up. In order for that to happen, its acceleration would also have to blow up, which means its jerk has to blow up, and so on, and so on:

$$u, \quad \partial_t u, \quad \partial_t^2 u, \quad \dots$$

The equation carries this repeated temporal climb into its nonlinear and viscous structure:

$$\begin{aligned} \partial_t u &= \nu \Delta u - (u \cdot \nabla)u - \nabla p, \\ \partial_t^2 u &= \nu \Delta \partial_t u - (\partial_t u \cdot \nabla)u - (u \cdot \nabla) \partial_t u - \nabla \partial_t p, \\ \partial_t^3 u &= \nu \Delta \partial_t^2 u - (\partial_t^2 u \cdot \nabla)u - 2(\partial_t u \cdot \nabla) \partial_t u - (u \cdot \nabla) \partial_t^2 u - \nabla \partial_t^2 p. \end{aligned}$$

The Laplacian follows the rising temporal response, while the nonlinear term opens into every product allowed by the product rule. At order  $r$ , those products occur with multiplicity  $\binom{r}{a}$ , giving the full pattern

$$\partial_t^{r+1}u + \sum_{a=0}^r \binom{r}{a} (\partial_t^a u \cdot \nabla) \partial_t^{r-a} u + \nabla \partial_t^r p = \nu \Delta \partial_t^r u.$$

Set

$$V_r := \partial_t^r u, \quad Q_r := \partial_t^r p.$$

Incompressibility fixes  $Q_r$  through the Riesz transforms  $R_i$ :

$$Q_r = R_i R_j \sum_{a=0}^r \binom{r}{a} V_{a,i} V_{r-a,j}.$$

The pressure row stays with the nonlinear row at every order.

If a singularity were forming, one would expect spatial regularity to erode as these temporal responses rise. Yet each next response contains  $\nu \Delta V_r$ , hence two spatial derivatives of the preceding response. That does not give regularity on its own. It tells us which spatial coordinates the equation exposes.

To measure the spatial order exposed by  $\Delta V_r$ , let  $\Lambda = (-\Delta)^{1/2}$  and set

$$E_s(t) := \frac{1}{2} \|\Lambda^s u(t)\|_2^2, \quad H(t) := E_{1/2}(t),$$

and retain the complete pressure-coupled nonlinear transfer

$$\mathcal{P}_s(t) := -\operatorname{Re} \langle \Lambda^s u, \Lambda^s ((u \cdot \nabla) u + \nabla p) \rangle_{L^2}.$$

Pairing the equation at height  $s$  gives

$$E'_s = \mathcal{P}_s - 2\nu E_{s+1}.$$

Differentiate this identity and use it again at each new height. At every finite order  $n$ ,

$$H^{(n)} = (-2\nu)^n E_{n+1/2} + \sum_{j=0}^{n-1} (-2\nu)^j \partial_t^{n-1-j} \mathcal{P}_{j+1/2}.$$

Acceleration, jerk, and the higher temporal responses have now exposed the Sobolev coordinates

$$E_{n+1/2} = \frac{1}{2} \|u\|_{\dot{H}^{n+1/2}}^2.$$

If every finite exposed height reaches one terminal value  $u_*$ , with the finite-energy row retained, then for any  $k$  one can choose  $n > k + 1$  and read

$$u_* \in L^2 \cap \bigcap_{n < \infty} \dot{H}^{n+1/2} = H^\infty \hookrightarrow C^\infty.$$

But exposure is not control. Differentiation does not bound  $H^{(n)}$ , the transfer terms, or  $E_{n+1/2}$ ; it only keeps their exact relation visible. The singular picture predicts loss of spatial regularity as temporal order rises, and these identities alone do not prevent it.

Let  $[0, T_*)$  be the maximal classical lifespan and suppose  $T_* < \infty$ . Fix one finite pair  $(N, M)$  and one cutoff  $0 < T < T_*$ . For  $0 \leq n \leq N$ , keep the whole transfer

$$\mathcal{B}_n := \sum_{a=0}^n \binom{n}{a} (V_a \cdot \nabla) V_{n-a} + \nabla Q_n.$$

Pairing the recurrence for  $V_n$  at height  $M$  gives

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|\Lambda^M V_n\|_2^2 + \operatorname{Re} \langle \Lambda^M V_n, \Lambda^M \mathcal{B}_n \rangle_{L^2} \\ = \nu \operatorname{Re} \langle \Lambda^M V_n, \Lambda^M \Delta V_n \rangle_{L^2} = -\nu \|V_n\|_{\dot{H}^{M+1}}^2. \end{aligned}$$

The viscous term has a fixed sign, while every product-rule transfer and the pressure response remain coupled to it in this finite system. This identity shows what must be controlled; it does not supply the whole-terminal bound by itself. Theorem 4.6 supplies the bound for this maximal history:

$$\sup_{0 < T < T_*} \sum_{n=0}^N \left( \|V_n\|_{L^2(0,T;H^{M+1})}^2 + \|V_{n+1}\|_{L^2(0,T;H^{M-1})}^2 \right) < \infty.$$

Every term in the sum increases monotonically with the cutoff. Since one finite bound holds for all  $T < T_*$ , the limit  $T \uparrow T_*$  places each adjacent pair on the full approach to  $T_*$ . With  $V_{n+1} = \partial_t V_n$ , that pair sits in the Hilbert triple

$$H^{M+1} \subset H^M \subset H^{M-1}$$

and gives

$$V_n \in C([0, T_*]; H^M).$$

The pressure formula supplies  $Q_n \in L^1(0, T_*; H^M)$  and  $\nabla Q_n \in L^1(0, T_*; H^{M-1})$ , so the finitely many rows now form one pressure-complete rectangle

$$\mathcal{R}_{N,M}[u_0; T_*] = ((V_n)_{n=0}^N, (V_{n+1})_{n=0}^N; (Q_n)_{n=0}^N).$$

The bound may depend on  $N$  and  $M$ . No constant uniform over every derivative order is being used.

A larger rectangle retains each older coordinate as the identical distributional derivative of  $u$  or  $p$ . Restricting from  $(N, M)$  to  $(N', M')$  consequently gives

$$\rho_{N',M'}^{N,M} \mathcal{R}_{N,M}[u_0; T_*] = \mathcal{R}_{N',M'}[u_0; T_*].$$

The restriction maps act between the finite coordinate spaces  $\mathcal{X}_{N,M}$  determined by the three rows already displayed.

$$\mathcal{X}_\infty := \varprojlim_{N,M} \mathcal{X}_{N,M}, \quad \mathcal{I}[u_0] := (\mathcal{R}_{N,M}[u_0; T_*])_{N,M \in \mathbb{N}_0} \in \mathcal{X}_\infty, \quad u \in \bigcap_{r,m < \infty} H^r(0, T_*; H^m(\mathbb{R}^3)).$$

There is no all-order estimate hidden in this notation. Each requested coordinate is obtained at a finite stage, and compatibility makes all of those finite coordinates parts of one history.

Take the zeroth temporal row after this intersection has been formed. For every finite  $m$ ,

$$u \in L^2(0, T_*; H^{m+1}), \quad u_t \in L^2(0, T_*; H^{m-1}), \quad u \in C([0, T_*]; H^m).$$

The endpoint values at different heights are compatible traces of one field. They produce a common endpoint

$$u_* := u(T_*) \in \bigcap_{m < \infty} H^m(\mathbb{R}^3) = H^\infty(\mathbb{R}^3) \hookrightarrow C^\infty(\mathbb{R}^3).$$

At  $T_*$ , set  $V_0^* := u_*$ . The pressure formula and binomial recurrence then define  $Q_r^*$  and  $V_{r+1}^*$  through every finite order. Smooth local existence restarts the flow from  $u_*$ . Uniqueness glues that continuation to the history already reaching  $T_*$ , contradicting finite maximality. Hence

$$T_* = \infty.$$

Only after  $\mathcal{I}[u_0]$  is present do the familiar finite estimates become available as coordinate projections. Fix any finite  $T$ , a temporal order  $r$ , a spatial derivative order  $k$ , and  $2 \leq p \leq \infty$ , then choose

$$m > k + \frac{3}{2} - \frac{3}{p}.$$

The corresponding row gives

$$V_r \in C([0, T]; H^m) \hookrightarrow L^\infty(0, T; W^{k,p}).$$

For instance, the  $m = 3$  coordinate yields

$$\int_0^T \|\nabla u(t)\|_\infty dt \leq T \|u\|_{L^\infty(0, T; W^{1,\infty})} < \infty.$$

This estimate appears because the fluid already occupies the compatible intersection; it is a finite readout of that object, not the source of it. Its source remains visible in the full dependence

$$\begin{aligned} u_0 &\longrightarrow \underbrace{\{(V_r, Q_r)\}_{r \geq 0}}_{\text{pressure-complete mixed jet}} \longrightarrow \underbrace{\left\{ \begin{array}{l} V_n \in L^2(0, T_*; H^{M+1}), \\ V_{n+1} \in L^2(0, T_*; H^{M-1}) \end{array} \right\}_{\substack{0 \leq n \leq N \\ N, M < \infty}}}_{\text{whole-terminal adjacent rows}} \\ &\longrightarrow \underbrace{\{\mathcal{R}_{N,M}[u_0; T_*]\}_{N, M < \infty}}_{\text{finite compatible rectangles}} \longrightarrow \underbrace{\mathcal{I}[u_0]}_{\text{projective intersection}} \longrightarrow \underbrace{(u_* \in H^\infty, \{(V_r^*, Q_r^*)\}_{r \geq 0})}_{\text{common endpoint and pressure-complete endpoint jet}} \\ &\longrightarrow \text{restart and uniqueness} \longrightarrow T_* = \infty. \end{aligned}$$