

# Global Smoothness for the Three-Dimensional Incompressible Navier–Stokes Equations

Thomas Birnie

August 2026

## Abstract

For every  $\nu > 0$  and every divergence-free  $u_0 \in H^\infty(\mathbb{R}^3; \mathbb{R}^3)$ , we prove that the unforced three-dimensional incompressible Navier–Stokes equations have a global pressure-normalized smooth solution. It satisfies the kinetic-energy identity and agrees with every Leray–Hopf solution having the same datum and viscosity. Let  $[0, T_*)$  be the lifespan of the maximal classical solution, and suppose that  $T_* < \infty$ . The central estimate is uniform on the full approach to  $T_*$ : for each fixed finite  $N, M$ ,

$$\sup_{0 < T < T_*} \sum_{n=0}^N \left( \|\partial_t^n u\|_{L^2(0, T; H^{M+1})}^2 + \|\partial_t^{n+1} u\|_{L^2(0, T; H^{M-1})}^2 \right) < \infty.$$

The constant may depend on  $N$  and  $M$ , but not on the cutoff  $T$ . All entries in these finite estimates are derivatives of the same solution, so estimates with different depths agree wherever they overlap. Given any finite time order  $r$  and spatial order  $m$ , one sufficiently large pair  $(N, M)$  yields  $u \in H^r(0, T_*; H^m)$ . In particular,  $u \in L^2(0, T_*; H^{m+2})$  for every  $m$ ; the equation and Sobolev multiplication then give  $u_t \in L^1(0, T_*; H^m)$ . Hence  $u(t)$  converges in every  $H^m$ , as  $t \uparrow T_*$ , to a single  $u_* \in H_\sigma^\infty$ . The differentiated equation and pressure formula recover the endpoint velocity–pressure jet. A local solution starting from  $u_*$  has that same jet; late-slice uniqueness joins it to the preterminal solution and continues the latter past  $T_*$ . This contradicts maximality, so  $T_* = \infty$ .

# AI-Assistance Disclosure and Referee Orientation

**AI-assistance disclosure.** OpenAI generative language models, including ChatGPT and Codex, assisted throughout the preparation of this manuscript with mathematical organization, drafting and revision, L<sup>A</sup>T<sub>E</sub>X production, source and citation checks, calculation checks, and adversarial review. Thomas Birnie developed the proof and directed that collaboration; he chose the claims and arguments presented here and is the paper's sole author.

**One history, one jet.** Throughout the paper,  $(T_*, u, p)$  is the pressure-normalized maximal classical history generated by one pair  $(u_0, \nu)$ . Every coordinate below belongs to that history; changing  $(N, M)$  only changes the finite view of it.

**Compact derivation map for Theorem 4.6.** Definition 3.3 sets  $V_n = \partial_t^n u$  and  $Q_n = \partial_t^n p$ . Pressure normalization, the differentiated equation, and the initial-jet induction give

$$\begin{aligned} V_0(0) &= u_0, & Q_n &= R_i R_j \sum_{a=0}^n \binom{n}{a} V_{a,i} V_{n-a,j}, \\ V_{n+1} &= \nu \Delta V_n - \sum_{a=0}^n \binom{n}{a} (V_a \cdot \nabla) V_{n-a} - \nabla Q_n. \end{aligned}$$

These are (44)–(45); at  $t = 0$ , Lemma 3.7 recursively determines the complete initial jet from  $u_0$ . The same equations then govern the one mixed jet on its maximal history. Theorem 4.6 supplies its datum-side whole-terminal adjacent rows, coordinate by coordinate:

$$V_n \in L^2(0, T_*; H^{m+1}), \quad \partial_t V_n = V_{n+1} \in L^2(0, T_*; H^{m-1}) \quad (n, m < \infty).$$

This is exactly (72). Each rectangle is the finite assembly of these rows, not a new estimate or a new solution. Proposition 4.7 records that final finite summation and its monotone realization (73):

$$\begin{aligned} &\sum_{n=0}^N \left( \|V_n\|_{L^2(0, T_*; H^{M+1})}^2 + \|V_{n+1}\|_{L^2(0, T_*; H^{M-1})}^2 \right) \\ &= \lim_{T \uparrow T_*} \mathfrak{R}_{N,M}(u; T) = \sup_{0 < T < T_*} \mathfrak{R}_{N,M}(u; T) = \mathfrak{R}_{N,M}^*[u_0, \nu, T_*] < \infty. \end{aligned}$$

Thus the proof order is

$$\begin{aligned} u_0 &\longrightarrow \{V_n(0), Q_n(0)\}_{n \geq 0} \longrightarrow \{V_n, Q_n\}_{n \geq 0} \\ &\longrightarrow \text{whole-terminal adjacent rows} \xrightarrow{\text{finite sum}} \{\mathcal{R}_{N,M}\}_{N,M < \infty}. \end{aligned}$$

**Closure and audit order.** The compatible rectangles pass to their projective intersection, one common  $H^\infty$  endpoint, the pressure-complete endpoint jet, and the same-history restart with overlap uniqueness, contradicting  $T_* < \infty$ . Any arrow may be challenged at its cited payment, but the review must keep fixed this one history and this dependency order. Proposition 4.7 packages the already obtained adjacent rows; reading it as their upstream source reverses the proof.

# Contents

|          |                                                               |           |
|----------|---------------------------------------------------------------|-----------|
| <b>1</b> | <b>Introduction</b>                                           | <b>4</b>  |
| <b>2</b> | <b>Analytic framework</b>                                     | <b>6</b>  |
| <b>3</b> | <b>The pressure-complete setting and mixed jet</b>            | <b>17</b> |
| <b>4</b> | <b>Datum-generated compatible rectangles</b>                  | <b>21</b> |
| <b>5</b> | <b>Compatible intersection, endpoint jet, and restart</b>     | <b>25</b> |
| <b>6</b> | <b>Matrix forms of the completed intersection</b>             | <b>32</b> |
| <b>7</b> | <b>Quantitative consequences of the compatible rectangles</b> | <b>35</b> |
| 7.1      | Completion in the global H1 mild class . . . . .              | 35        |
| 7.2      | Canonical coordinates and finite majorants . . . . .          | 38        |
| 7.3      | Mixed velocity and pressure rows . . . . .                    | 45        |
| 7.4      | Frequency and critical-scale consequences . . . . .           | 47        |
| 7.5      | Continuation and quantitative restart . . . . .               | 49        |
| <b>8</b> | <b>Three-dimensional applications</b>                         | <b>50</b> |
| 8.1      | Explicit closure for small critical data . . . . .            | 50        |
| 8.2      | Low-rectangle and scale-local bounds . . . . .                | 55        |
| 8.3      | Vorticity and material flow . . . . .                         | 56        |
| 8.4      | Relative energy and the Leray–Hopf class . . . . .            | 58        |
| <b>9</b> | <b>Conclusion</b>                                             | <b>63</b> |
| <b>A</b> | <b>Canonical coordinate catalogue</b>                         | <b>64</b> |
|          | <b>References</b>                                             | <b>68</b> |

**Theorem.** For every  $\nu > 0$  and every  $u_0 \in H^\infty_\sigma(\mathbb{R}^3)$ , there exists a pressure-normalized pair

$$(u, p) \in C^\infty([0, \infty) \times \mathbb{R}^3; \mathbb{R}^3 \times \mathbb{R})$$

satisfying

$$\partial_t u + (u \cdot \nabla)u + \nabla p = \nu \Delta u, \quad \nabla \cdot u = 0, \quad u(0) = u_0, \quad \lim_{|x| \rightarrow \infty} p(t, x) = 0,$$

and

$$\frac{1}{2} \|u(t)\|_{L^2}^2 + \nu \int_0^t \|\nabla u(s)\|_{L^2}^2 ds = \frac{1}{2} \|u_0\|_{L^2}^2 \quad (t \geq 0).$$

Every global Leray–Hopf pair  $(v, q)$  at viscosity  $\nu$  with datum  $u_0$ , in the sense of Definition 8.9, has  $v = u$ , and  $q = p + c(t)$  distributionally for a scalar distribution  $c$  in time. If  $q$  is represented by a locally integrable function with  $q(t, \cdot) \in C_0(\mathbb{R}^3)$  for almost every  $t$ , then  $q = p$  almost everywhere.

Every solenoidal Schwartz datum belongs to  $H^\infty_\sigma(\mathbb{R}^3)$ . Restricted to that subclass, the theorem gives the global smooth pair and bounded kinetic energy required by the whole-space positive formulation in [4]; the theorem’s datum class is broader.

## 1 Introduction

Fix  $\nu > 0$  and a divergence-free datum  $u_0 \in H^\infty(\mathbb{R}^3; \mathbb{R}^3)$ . Smooth-data local theory gives a unique classical solution on a maximal interval  $[0, T_*)$  [5, 9]. Throughout the paper,  $(u, p)$  denotes that solution. The problem is to decide whether  $T_*$  can be finite. On each strict subinterval  $[0, T]$ , smoothness already makes every finite Sobolev norm finite. That fact alone says nothing about what happens as  $T \uparrow T_*$ : the bound may depend on  $T$  and diverge at the endpoint. The proof must instead control the approach to  $T_*$  with constants that do not depend on the cutoff.

Leray constructed global finite-energy weak solutions [7], and Caffarelli, Kohn, and Nirenberg bounded the size of the possible singular set for suitable weak solutions [2]. Serrin proved regularity under scale-balanced spacetime integrability hypotheses [12]; Escauriaza, Seregin, and Šverák treated the endpoint  $L_t^\infty L_x^3$  case [3]. Fujita–Kato and Kato supply the strong local theory used below at the terminal restart [5, 9], and Koch–Tataru place small-data global well-posedness in the critical space  $BMO^{-1}$  [10]. Tao gives a localization and compactness formulation of the global problem [14]. The regularity criteria show that control of an appropriate critical quantity excludes singularity. The argument below does not assume such control. It derives it from one family of estimates for all finite time and space derivatives of the maximal solution.

Pressure is part of that derivative structure. With the decay normalization, incompressibility determines it from the current velocity:

$$-\Delta p = \partial_i \partial_j (u_i u_j), \quad p = R_i R_j (u_i u_j).$$

Here  $R_i$  are the Riesz transforms, and repeated indices are summed. Write

$$V_n := \partial_t^n u, \quad Q_n := \partial_t^n p.$$

Differentiating the equation does not introduce a new solution. It gives

$$Q_n = R_i R_j \sum_{a=0}^n \binom{n}{a} V_{a,i} V_{n-a,j}, \tag{1}$$

$$V_{n+1} = \nu \Delta V_n - \sum_{a=0}^n \binom{n}{a} (V_a \cdot \nabla) V_{n-a} - \nabla Q_n. \tag{2}$$

Thus the time derivatives, spatial derivatives, and pressure derivatives are coordinates of one solution. A time derivative raises the order of the viscous term by two spatial derivatives and distributes the transport and pressure terms among earlier time derivatives. The resulting estimates must track temporal depth and spatial depth independently.

For finite  $N, M \in \mathbb{N}_0$ , set

$$\mathfrak{R}_{N,M}(u; T) := \sum_{n=0}^N \|V_n\|_{L^2(0,T;H^{M+1})}^2 + \sum_{n=0}^N \|V_{n+1}\|_{L^2(0,T;H^{M-1})}^2. \quad (3)$$

The central estimate is that, under the contradiction hypothesis  $T_* < \infty$ , every fixed finite pair satisfies

$$\sup_{0 < T < T_*} \mathfrak{R}_{N,M}(u; T) < \infty. \quad (4)$$

This is Theorem 4.6 below. The bound is coordinatewise: its constant may depend on  $N$  and  $M$ , but it does not depend on  $T$ . No uniform estimate over all derivative orders is asserted or needed.

Every rectangle contains derivatives of the same solution. If  $N' \leq N$  and  $M' \leq M$ , restriction deletes the higher rows and applies Sobolev embeddings to the rows that remain; a shared derivative does not change. Write  $\mathcal{R}_{N,M}[u_0; T_*]$  for the resulting whole-terminal finite rectangle, and denote the compatible family by

$$\mathcal{I}[u_0] := (\mathcal{R}_{N,M}[u_0; T_*])_{N,M \in \mathbb{N}_0}. \quad (5)$$

Given finite  $r, m$ , choose one rectangle deep enough to contain the requested time and space derivatives. Compatibility guarantees that this coordinate is the same one in every larger rectangle, and hence

$$u \in \bigcap_{r,m \in \mathbb{N}_0} H^r(0, T_*; H^m). \quad (6)$$

The zeroth time row of (6) gives, for each finite  $m$ ,  $u \in L^2(0, T_*; H^{m+2})$ . Sobolev multiplication gives

$$\|\mathbb{P}\nabla \cdot (u \otimes u)\|_{H^m} \leq C_m \|u\|_{H^{m+2}}^2.$$

Since  $T_* < \infty$ , the nonlinear term and  $\Delta u$  belong to  $L^1(0, T_*; H^m)$ ; for the linear term, explicitly,

$$\|\Delta u\|_{L^1(0, T_*; H^m)} \leq T_*^{1/2} \|u\|_{L^2(0, T_*; H^{m+2})}.$$

The equation

$$u_t = \nu \Delta u - \mathbb{P}\nabla \cdot (u \otimes u)$$

now gives

$$u \in W^{1,1}(0, T_*; H^m) \hookrightarrow C([0, T_*]; H^m) \quad (m < \infty).$$

The endpoint values agree under the Sobolev inclusions because they are limits of the same function  $u(t)$ . They define one terminal state

$$u_* := u(T_*) \in \bigcap_{m < \infty} H^m.$$

Since  $u_* \in H^\infty$ , local well-posedness restarts the equation at  $T_*$ . Equations (1)–(2) recover the endpoint pressure and time derivatives. Uniqueness joins the restarted solution to  $u$ , producing an extension past  $T_*$ . This contradicts maximality and proves global smoothness.

## 2 Analytic framework

The argument has two analytic joints. Products and Fourier multipliers must keep every differentiated velocity–pressure row in a common Sobolev scale; at the terminal time, an  $H^3$  local theory must restart that same pressure-normalized history. We record the estimates needed at those two joints, with the constants later used in the quantitative bounds.

**Definition 2.1.** For  $f \in \mathcal{S}(\mathbb{R}^3)$ , set

$$\widehat{f}(\xi) := (2\pi)^{-3/2} \int_{\mathbb{R}^3} e^{-ix \cdot \xi} f(x) dx, \quad \langle \xi \rangle := (1 + |\xi|^2)^{1/2}.$$

The same integral defines  $\widehat{f}$  for  $f \in L^1(\mathbb{R}^3)$ . The Fourier transform on  $\mathcal{S}'(\mathbb{R}^3)$  is its extension by duality. For  $s \in \mathbb{R}$ , define

$$H^s(\mathbb{R}^3) := \left\{ f \in \mathcal{S}'(\mathbb{R}^3) : \langle \xi \rangle^s \widehat{f} \in L^2(\mathbb{R}^3) \right\},$$

and define the finite homogeneous-norm domain by

$$\dot{H}^s(\mathbb{R}^3) := \left\{ f \in \mathcal{S}'(\mathbb{R}^3) : \begin{array}{l} \widehat{f} \text{ is represented by a measurable function on } \mathbb{R}^3, \\ |\xi|^s \widehat{f} \in L^2(\mathbb{R}^3) \end{array} \right\}.$$

On these domains set

$$\|f\|_{H^s}^2 := \int_{\mathbb{R}^3} \langle \xi \rangle^{2s} |\widehat{f}(\xi)|^2 d\xi, \quad \|f\|_{\dot{H}^s}^2 := \int_{\mathbb{R}^3} |\xi|^{2s} |\widehat{f}(\xi)|^2 d\xi.$$

Set  $A := (1 - \Delta)^{1/2}$ ,  $\Lambda := (-\Delta)^{1/2}$ ,

$$H_\sigma^s := \{v \in H^s(\mathbb{R}^3; \mathbb{R}^3) : \nabla \cdot v = 0\}, \quad H^\infty := \bigcap_{m \in \mathbb{N}_0} H^m, \quad H_\sigma^\infty := \bigcap_{m \in \mathbb{N}_0} H_\sigma^m.$$

$$\mathcal{S}_\sigma(\mathbb{R}^3) := \{v \in \mathcal{S}(\mathbb{R}^3; \mathbb{R}^3) : \nabla \cdot v = 0\}.$$

For a vector or tensor  $F = (F_I)_I$ , all Sobolev norms are componentwise:

$$\|F\|_{H^s}^2 := \sum_I \|F_I\|_{H^s}^2, \quad \|F\|_{\dot{H}^s}^2 := \sum_I \|F_I\|_{\dot{H}^s}^2.$$

Repeated Latin indices are summed from 1 to 3.

**Lemma 2.2** (Sobolev nesting and embedding). *If  $s_2 \geq s_1$ , then*

$$\|f\|_{H^{s_1}} \leq \|f\|_{H^{s_2}}.$$

*If  $k \in \mathbb{N}_0$  and  $s > k + 3/2$ , then*

$$\max_{|\alpha| \leq k} \|\partial^\alpha f\|_{L^\infty} \leq C_{s,k}^{\text{emb}} \|f\|_{H^s},$$

where

$$C_{s,k}^{\text{emb}} := (2\pi)^{-3/2} \max_{0 \leq j \leq k} \left( \int_{\mathbb{R}^3} |\xi|^{2j} \langle \xi \rangle^{-2s} d\xi \right)^{1/2} < \infty.$$

In particular,

$$H^\infty(\mathbb{R}^3) \hookrightarrow C_b^\infty(\mathbb{R}^3).$$

*Proof.* The first assertion follows from  $\langle \xi \rangle^{2s_1} \leq \langle \xi \rangle^{2s_2}$ . For  $|\alpha| \leq k$ , Fourier inversion and Cauchy–Schwarz give

$$|\partial^\alpha f(x)| \leq (2\pi)^{-3/2} \left( \int_{\mathbb{R}^3} |\xi|^{2|\alpha|} \langle \xi \rangle^{-2s} d\xi \right)^{1/2} \|f\|_{H^s}.$$

The integral is finite exactly when  $s > |\alpha| + 3/2$ . Approximation by Schwartz functions and dominated convergence give continuous representatives. For each  $k$ , choose  $s = k + 2$ .  $\square$

**Lemma 2.3** (Explicit three-dimensional homogeneous Sobolev constants). *For  $3 \leq p \leq 6$ , set*

$$s(p) := 3 \left( \frac{1}{2} - \frac{1}{p} \right)$$

and

$$S(p) := 2^{-s(p)} \pi^{-s(p)/2} \left( \frac{\Gamma((3-2s(p))/2)}{\Gamma((3+2s(p))/2)} \right)^{1/2} \left( \frac{\Gamma(3)}{\Gamma(3/2)} \right)^{s(p)/3}. \quad (7)$$

Then every scalar, vector, or finite tensor field  $F$  in  $\dot{H}^{s(p)}(\mathbb{R}^3)$  satisfies

$$\|F\|_{L^p(\mathbb{R}^3)} \leq S(p) \|F\|_{\dot{H}^{s(p)}(\mathbb{R}^3)}. \quad (8)$$

In particular,

$$S(3) = 2^{-1/6} \pi^{-1/3}, \quad S(6) = \frac{2^{2/3}}{\sqrt{3} \pi^{2/3}}, \quad S(6)S(3) = \frac{1}{\pi} \sqrt{\frac{2}{3}}.$$

*Proof.* For scalar fields, (8) is the sharp three-dimensional homogeneous Sobolev inequality obtained from the sharp Hardy–Littlewood–Sobolev inequality [8]. After duality, insertion of the Riesz-potential kernel constant, and conversion to the unitary Fourier transform fixed above, the squared sharp Sobolev constant for the multiplier  $\widehat{\Lambda^s f}(\xi) = |\xi|^s \widehat{f}(\xi)$  is

$$2^{-2s} \pi^{-s} \frac{\Gamma((3-2s)/2)}{\Gamma((3+2s)/2)} \left( \frac{\Gamma(3)}{\Gamma(3/2)} \right)^{2s/3}. \quad (9)$$

Taking the square root in (9) and putting  $p = 6/(3-2s)$  gives (7)–(8). For a finite component array  $F = (F_I)_I$ , Minkowski’s inequality in  $L^{p/2}$  gives

$$\|F\|_p^2 = \left\| \sum_I |F_I|^2 \right\|_{p/2} \leq \sum_I \|F_I\|_p^2 \leq S(p)^2 \sum_I \|F_I\|_{\dot{H}^{s(p)}}^2.$$

The two endpoint formulas follow by evaluating the gamma factors at  $s = 1/2$  and  $s = 1$ .  $\square$

**Lemma 2.4** (Sobolev multiplication). *Set*

$$H_4 := (2\pi)^{-3/4} \left( \int_{\mathbb{R}^3} \langle \xi \rangle^{-4} d\xi \right)^{1/4}, \quad C_0^{\text{alg}} := H_4^2.$$

For  $m \geq 1$ , define

$$\begin{aligned} E_m &:= \max_{0 \leq j \leq \lfloor m/2 \rfloor} C_{j+2,j}^{\text{emb}}, & C_m^{\text{mix}} &:= 13^{m/2} E_m, \\ C_m^{\text{F}} &:= 2^m (2\pi)^{-3/2} \left( \int_{\mathbb{R}^3} \langle \xi \rangle^{-2m} d\xi \right)^{1/2} & (m \geq 2), \end{aligned}$$

and set

$$C_1^{\text{alg}} := C_1^{\text{mix}}, \quad C_m^{\text{alg}} := \max\{C_m^{\text{mix}}, C_m^{\text{F}}\} \quad (m \geq 2).$$

All these constants are finite and explicit. For every  $m \in \mathbb{N}_0$ ,

$$\|fg\|_{H^m} \leq C_m^{\text{alg}} \|f\|_{H^{m+1}} \|g\|_{H^{m+1}}.$$

For  $m \geq 2$ ,

$$\|fg\|_{H^m} \leq C_m^{\text{alg}} \|f\|_{H^m} \|g\|_{H^m}.$$

*Proof.* Hausdorff–Young and Hölder give

$$\|f\|_4 \leq (2\pi)^{-3/4} \left\| \widehat{f} \right\|_{4/3} \leq H_4 \|f\|_{H^1}.$$

Thus  $\|fg\|_2 \leq C_0^{\text{alg}} \|f\|_{H^1} \|g\|_{H^1}$ .

Let  $m \geq 1$ . The exact integer-order identity

$$\|h\|_{H^m}^2 = \sum_{|\alpha| \leq m} \frac{m!}{(m - |\alpha|)! \alpha!} \|\partial^\alpha h\|_2^2$$

follows by expanding  $(1 + |\xi|^2)^m$ . In each Leibniz term for  $\partial^\alpha(fg)$ , place the factor carrying at most  $\lfloor |\alpha|/2 \rfloor$  derivatives in  $L^\infty$ . The definition of  $E_m$ , Sobolev nesting, and  $\sum_{\beta \leq \alpha} \binom{\alpha}{\beta} = 2^{|\alpha|}$  give

$$\|\partial^\alpha(fg)\|_2 \leq E_m 2^{|\alpha|} \|f\|_{H^{m+1}} \|g\|_{H^{m+1}}.$$

The multinomial identity

$$\sum_{|\alpha| \leq m} \frac{m!}{(m - |\alpha|)! \alpha!} 4^{|\alpha|} = 13^m$$

proves the first estimate with  $C_m^{\text{mix}}$ .

For  $m \geq 2$ , the inequalities

$$\langle \eta + \zeta \rangle^m \leq 2^{m-1} (\langle \eta \rangle^m + \langle \zeta \rangle^m), \quad \left\| \widehat{f} \right\|_1 \leq \left( \int_{\mathbb{R}^3} \langle \xi \rangle^{-2m} d\xi \right)^{1/2} \|f\|_{H^m},$$

combined with  $\widehat{fg} = (2\pi)^{-3/2} \widehat{f} * \widehat{g}$  and Young's convolution inequality give

$$\|fg\|_{H^m} \leq C_m^{\text{F}} \|f\|_{H^m} \|g\|_{H^m}.$$

The chosen maximum proves both asserted estimates with the same constant.  $\square$

**Lemma 2.5** (Riesz and Leray multipliers). *Let*

$$r_j(\xi) := \begin{cases} i\xi_j/|\xi|, & \xi \neq 0, \\ 0, & \xi = 0, \end{cases} \quad P(\xi) := \begin{cases} I - \frac{\xi \otimes \xi}{|\xi|^2}, & \xi \neq 0, \\ I, & \xi = 0, \end{cases}$$

and

$$\widehat{R_j f}(\xi) = r_j(\xi) \widehat{f}(\xi), \quad \widehat{\mathbb{P}f}(\xi) = P(\xi) \widehat{f}(\xi).$$

For all  $s \in \mathbb{R}$ ,

$$\|R_j f\|_{H^s} \leq \|f\|_{H^s}, \quad \|\mathbb{P}f\|_{H^s} \leq \|f\|_{H^s}.$$

If  $m \in \mathbb{N}_0$  and  $v \in H_\sigma^{m+1}$ , then

$$q := \sum_{i,j=1}^3 R_i R_j (v_i v_j) \in H^m, \quad -\Delta q = \sum_{i,j=1}^3 \partial_i \partial_j (v_i v_j),$$

and

$$\mathbb{P} \nabla \cdot (v \otimes v) = (v \cdot \nabla) v + \nabla q.$$

Here

$$(v \otimes v)_{ij} := v_i v_j, \quad (\nabla \cdot (v \otimes v))_i := \sum_{j=1}^3 \partial_j (v_i v_j), \quad ((v \cdot \nabla) v)_i := \sum_{j=1}^3 v_j \partial_j v_i.$$

*Proof.* The multiplier symbols have operator norm at most one. Plancherel proves the first two estimates. Lemma 2.4 gives  $v_i v_j \in H^m$ . The remaining identities follow after multiplication by  $|\xi|^2$  and by the symbol of  $\mathbb{P}$ .  $\square$

**Lemma 2.6** (Mixed Leibniz identities). *For  $N \in \mathbb{N}_0$  and  $\alpha \in \mathbb{N}_0^3$ ,*

$$\partial_t^N \partial_x^\alpha (fg) = \sum_{a=0}^N \sum_{\beta \leq \alpha} \binom{N}{a} \binom{\alpha}{\beta} (\partial_t^a \partial_x^\beta f) (\partial_t^{N-a} \partial_x^{\alpha-\beta} g).$$

*Proof.* Apply the one-variable Leibniz identity successively in  $t, x_1, x_2, x_3$ .  $\square$

**Lemma 2.7** (Adjacent Hilbert-scale trace). *Let  $0 < T < \infty$ ,  $m \in \mathbb{R}$ , and*

$$v \in L^2(0, T; H^{m+1}), \quad \partial_t v \in L^2(0, T; H^{m-1}).$$

*Then  $v$  has a unique representative in  $C([0, T]; H^m)$ , and, for  $0 \leq r \leq t \leq T$ ,*

$$\|v(t)\|_{H^m}^2 - \|v(r)\|_{H^m}^2 = 2 \operatorname{Re} \int_r^t \langle A^{m-1} \partial_t v(s), A^{m+1} v(s) \rangle_{L^2} ds, \quad (10)$$

$$\left| \|v(t)\|_{H^m}^2 - \|v(r)\|_{H^m}^2 \right| \leq 2 \|\partial_t v\|_{L^2(r, t; H^{m-1})} \|v\|_{L^2(r, t; H^{m+1})}, \quad (11)$$

$$\sup_{0 \leq t \leq T} \|v(t)\|_{H^m}^2 \leq T^{-1} \|v\|_{L^2(0, T; H^m)}^2 + 2 \|\partial_t v\|_{L^2(0, T; H^{m-1})} \|v\|_{L^2(0, T; H^{m+1})}. \quad (12)$$

*Proof.* Let  $S_R$  denote Fourier restriction to  $\{|\xi| \leq R\}$ . For  $v_R = S_R v$ ,

$$\frac{d}{dt} \|v_R(t)\|_{H^m}^2 = 2 \operatorname{Re} \langle A^{m-1} \partial_t v_R(t), A^{m+1} v_R(t) \rangle_{L^2}.$$

Integration and Cauchy-Schwarz prove (10)–(11) for  $v_R$ . Applied to  $S_R v - S_S v$ , the same estimate shows that  $(v_R)_R$  is Cauchy in  $C([0, T]; H^m)$ ; choose one Lebesgue time for the initial Cauchy term. Its limit equals  $v$  in  $L^2(0, T; H^m)$ . Passage to the limit proves the first two formulas. Choose  $r \in (0, T)$  with  $\|v(r)\|_{H^m}^2 \leq T^{-1} \|v\|_{L^2 H^m}^2$  and apply (11).  $\square$

**Lemma 2.8** (Explicit integer transport commutator). *Set*

$$\begin{aligned} \Gamma_{1,1} &:= 1, \\ \Gamma_{\ell,k} &:= (2\ell - 4 + \sqrt{3})^{(k-1)(\ell-k)} \quad (\ell \geq 2, 1 \leq k \leq \ell), \end{aligned} \quad (13)$$

$$K_m := \sum_{\ell=1}^m 3^\ell \binom{m}{\ell} \sum_{k=1}^{\ell} \binom{\ell}{k} \Gamma_{\ell,k} \quad (m \geq 3). \quad (14)$$

*Every  $v \in H_\sigma^{m+1}(\mathbb{R}^3)$  satisfies*

$$|\operatorname{Re} \langle A^m \mathbb{P} \nabla \cdot (v \otimes v), A^m v \rangle_{L^2}| \leq K_m \|\nabla v\|_{L^\infty} \|v\|_{H^m}^2. \quad (15)$$

*In particular,*

$$K_m \leq (7^m - 1)(2m - 4 + \sqrt{3})^{(m-1)^2/4}. \quad (16)$$

*Proof.* We first prove the interpolation inequality used in the commutator. For the full ordered derivative tensor  $\nabla^j F$ , fix  $M \geq 1$  and set

$$p_{M,0} := \infty, \quad p_{M,j} := \frac{2M}{j} \quad (1 \leq j \leq M), \quad A_M := 2M - 2 + \sqrt{3}.$$

Then

$$\|\nabla^j F\|_{L^{p_{M,j}}} \leq A_M^{j(M-j)/2} \|F\|_\infty^{1-j/M} \|\nabla^M F\|_2^{j/M} \quad (0 \leq j \leq M). \quad (17)$$

The endpoint cases are identities. For  $1 \leq j \leq M-1$ , put

$$G := \nabla^{j-1} F, \quad W := \nabla G, \quad p := p_{M,j}.$$

All contractions below are the Euclidean contractions of the corresponding ordered tensors. Integration by parts gives

$$\begin{aligned} \int_{\mathbb{R}^3} |W|^p &= - \int_{\mathbb{R}^3} |W|^{p-2} G : \Delta G \\ &\quad - (p-2) \int_{\mathbb{R}^3} |W|^{p-4} \sum_{q=1}^3 (G : W_{\cdot q})(W : \partial_q W). \end{aligned}$$

The pointwise bounds

$$|\Delta G| \leq \sqrt{3} |\nabla^2 G|, \quad \left| \sum_{q=1}^3 (G : W_{\cdot q})(W : \partial_q W) \right| \leq |G| |W|^2 |\nabla^2 G|$$

give

$$\int_{\mathbb{R}^3} |W|^p \leq (p-2 + \sqrt{3}) \int_{\mathbb{R}^3} |G| |W|^{p-2} |\nabla^2 G|.$$

Since

$$\frac{1}{p_{M,j-1}} + \frac{p-2}{p} + \frac{1}{p_{M,j+1}} = 1,$$

Hölder's inequality, with  $X_j := \|\nabla^j F\|_{L^{p_{M,j}}}$ , gives

$$X_j^2 \leq \left( \frac{2M}{j} - 2 + \sqrt{3} \right) X_{j-1} X_{j+1} \leq A_M X_{j-1} X_{j+1}. \quad (18)$$

The numbers  $d_j := j(M-j)/2$  obey

$$2d_j - d_{j-1} - d_{j+1} = 1, \quad d_0 = d_M = 0.$$

The discrete maximum principle applied to (18) gives

$$X_j \leq A_M^{d_j} X_0^{1-j/M} X_M^{j/M},$$

which is (17). The integration by parts is justified first with  $(|W|^2 + \varepsilon)^{(p-2)/2}$ , followed by  $\varepsilon \downarrow 0$ ; smooth approximation handles zero tensor norms.

For  $|\alpha| \leq m$ , set

$$\alpha! := \alpha_1! \alpha_2! \alpha_3!, \quad c_{m,\alpha} := \frac{m!}{(m-|\alpha|)! \alpha!}.$$

The multiplier  $A^{2m} = (1 - \Delta)^m$  and the multinomial identity give the exact pairing formula

$$\langle A^m f, A^m g \rangle_{L^2} = \sum_{|\alpha| \leq m} c_{m,\alpha} \langle \partial^\alpha f, \partial^\alpha g \rangle_{L^2}. \quad (19)$$

The Leray projection is self-adjoint, commutes with  $A^m$ , and fixes  $A^m v$ . Since  $\nabla \cdot (v \otimes v) = (v \cdot \nabla) v$ , it follows that

$$\langle A^m \mathbb{P} \nabla \cdot (v \otimes v), A^m v \rangle_{L^2} = \sum_{|\alpha| \leq m} c_{m,\alpha} \langle \partial^\alpha ((v \cdot \nabla) v), \partial^\alpha v \rangle_{L^2}. \quad (20)$$

For each  $\alpha$ ,

$$\partial^\alpha ((v \cdot \nabla) v) = (v \cdot \nabla) \partial^\alpha v + \sum_{0 < \beta \leq \alpha} \binom{\alpha}{\beta} (\partial^\beta v \cdot \nabla) \partial^{\alpha-\beta} v.$$

The divergence-free transport term has zero real pairing with  $\partial^\alpha v$ , because

$$\operatorname{Re} \langle (v \cdot \nabla) \partial^\alpha v, \partial^\alpha v \rangle_{L^2} = \frac{1}{2} \int_{\mathbb{R}^3} v \cdot \nabla |\partial^\alpha v|^2 \, dx = 0.$$

For a remaining term, put

$$\ell := |\alpha|, \quad k := |\beta|.$$

If  $\ell \geq 2$ , apply (17) to  $F = \nabla v$ , with

$$M = \ell - 1, \quad j = k - 1.$$

The two Hölder exponents are  $p_{M,j}$  and  $p_{M,M-j}$ , whose reciprocal sum is  $1/2$ . The two factors are subarrays of the corresponding ordered derivative tensors, so

$$\begin{aligned} \left\| (\partial^\beta v \cdot \nabla) \partial^{\alpha-\beta} v \right\|_2 &\leq \left\| \partial^\beta v \right\|_{L^{p_{M,j}}} \left\| \nabla \partial^{\alpha-\beta} v \right\|_{L^{p_{M,M-j}}} \\ &\leq \Gamma_{\ell,k} \left\| \nabla v \right\|_\infty \left\| \nabla^\ell v \right\|_2. \end{aligned} \tag{21}$$

For  $\ell = 1$ , the same formula follows from the direct  $L^\infty$ - $L^2$  estimate and  $\Gamma_{1,1} = 1$ . Since

$$\left\| \nabla^\ell v \right\|_2 = \|v\|_{\dot{H}^\ell} \leq \|v\|_{H^m}, \quad \|\partial^\alpha v\|_2 \leq \|v\|_{H^m},$$

each remaining pairing is bounded by

$$\Gamma_{\ell,k} \left\| \nabla v \right\|_\infty \|v\|_{H^m}^2.$$

The multinomial identities

$$\sum_{|\alpha|=\ell} c_{m,\alpha} = 3^\ell \binom{m}{\ell}, \quad \sum_{\substack{\beta \leq \alpha \\ |\beta|=k}} \binom{\alpha}{\beta} = \binom{\ell}{k}$$

now give (15) with the constant (14). Finally,

$$\sum_{k=1}^{\ell} \binom{\ell}{k} \leq 2^\ell, \quad (k-1)(\ell-k) \leq \frac{(\ell-1)^2}{4},$$

and summation of  $\binom{m}{\ell} 3^\ell 2^\ell$  over  $1 \leq \ell \leq m$  proves (16). Approximation by divergence-free Schwartz fields justifies all integrations by parts for  $v \in H_\sigma^{m+1}$ .  $\square$

The preceding estimates control the differentiated rows as they approach an endpoint. To continue their common limit as the same solution, we now fix a time-translation-invariant local class that admits both late preterminal slices and the endpoint restart, and in which uniqueness holds on every overlap.

**Definition 2.9** (Shifted pressure-normalized  $H^3$  mild class). Let  $0 \leq s < b < \infty$  and  $a \in H_\sigma^3(\mathbb{R}^3)$ . Define  $\mathcal{M}_\nu^3(s, b; a)$  to be the set of pairs  $(w, q)$  such that

$$w \in C([s, b]; H_\sigma^3(\mathbb{R}^3)), \quad q \in C([s, b]; H^3(\mathbb{R}^3)), \tag{22}$$

$$w(s) = a, \quad q(t) = R_i R_j (w_i(t) w_j(t)), \tag{23}$$

The remaining defining condition is the Duhamel identity: for every  $t \in [s, b]$ ,

$$w(t) = e^{\nu(t-s)\Delta}a - \int_s^t e^{\nu(t-r)\Delta}\mathbb{P}\nabla \cdot (w \otimes w)(r) \, dr \quad \text{in } H_\sigma^3(\mathbb{R}^3). \quad (24)$$

The integral is an  $H_\sigma^3$ -valued Bochner integral: multiplication in  $H^3(\mathbb{R}^3)$  places  $\mathbb{P}\nabla \cdot (w \otimes w)$  in  $C([s, b]; H_\sigma^2)$ , and the heat multiplier from  $H^2$  to  $H^3$  has an integrable  $(t-r)^{-1/2}$  singularity. The maximal lifespan in this class is

$$\tau_{\nu,3}^{\max}(s, a) := \sup \{b - s : b > s, \mathcal{M}_\nu^3(s, b; a) \neq \emptyset\} \in [0, \infty]. \quad (25)$$

**Lemma 2.10** (Restriction and classical admission). *If  $(w, q) \in \mathcal{M}_\nu^3(s, b; a)$  and  $s \leq r < c \leq b$ , then*

$$(w, q)|_{[r, c]} \in \mathcal{M}_\nu^3(r, c; w(r)). \quad (26)$$

*If  $(u, p)$  is a pressure-complete classical history and  $[s, b]$  is contained in its interval of existence, then*

$$(u, p)|_{[s, b]} \in \mathcal{M}_\nu^3(s, b; u(s)). \quad (27)$$

*Proof.* Split (24) at  $r$  and use the heat semigroup identity. This gives, for  $r \leq t \leq c$ ,

$$w(t) = e^{\nu(t-r)\Delta}w(r) - \int_r^t e^{\nu(t-\rho)\Delta}\mathbb{P}\nabla \cdot (w \otimes w)(\rho) \, d\rho.$$

The pressure formula is unchanged under restriction. For a classical history, apply  $\mathbb{P}$  to the momentum equation and use variation of constants; the decaying pressure formula gives (23).  $\square$

**Theorem 2.11** (Quantitative local restart). *Set  $C_q := \sqrt{3}C_3^{\text{alg}}$ . For  $\nu > 0$ , set*

$$L_\nu := 4(1 + \nu + \nu^{-1}) \exp\left(\frac{\nu + \nu^{-1}}{2}\right), \quad c_\nu := (8C_q L_\nu^2)^{-2}.$$

*For every  $s \geq 0$ , every  $a \in H_\sigma^3(\mathbb{R}^3)$ , and every*

$$0 < \delta \leq \delta_\nu(a) := \min\{1, c_\nu(1 + \|a\|_{H^3})^{-2}\},$$

*the class  $\mathcal{M}_\nu^3(s, s + \delta; a)$  contains exactly one pair  $(w, q)$ . Its velocity satisfies*

$$w \in C([s, s + \delta]; H_\sigma^3) \cap L^2(s, s + \delta; H^4), \quad w_t \in L^2(s, s + \delta; H^2).$$

*If  $a \in H_\sigma^\infty$ , then*

$$\partial_t^n w, \partial_t^n q \in C([s, s + \delta]; H^m) \quad (n, m \in \mathbb{N}_0).$$

*For every finite  $b > s$ , the class  $\mathcal{M}_\nu^3(s, b; a)$  contains at most one pair. Its compatible members on strict subintervals of  $[s, s + \tau_{\nu,3}^{\max}(s, a))$  define the maximal shifted pressure-normalized  $H^3$  mild history from  $a$ .*

*Proof.* By time translation it is enough to prove the assertions with  $s = 0$ . Let  $0 < \delta \leq 1$ , let

$$z_t - \nu\Delta z = F, \quad z(0) = a, \quad a \in H_\sigma^3, \quad F \in L^2(0, \delta; H_\sigma^2),$$

and set

$$A_0 := \|a\|_{H^3}, \quad G := \|F\|_{L^2(0,\delta;H^2)}.$$

The  $H^3$  energy identity is

$$\frac{1}{2} \frac{d}{dt} \|z\|_{H^3}^2 + \nu \|\nabla z\|_{H^3}^2 = \operatorname{Re} \langle A^2 F, A^4 z \rangle_{L^2}.$$

Since  $\|z\|_{H^4}^2 = \|z\|_{H^3}^2 + \|\nabla z\|_{H^3}^2$ , Young's inequality gives

$$\frac{d}{dt} \|z\|_{H^3}^2 + \nu \|\nabla z\|_{H^3}^2 \leq \nu \|z\|_{H^3}^2 + \nu^{-1} \|F\|_{H^2}^2.$$

Consequently,

$$\|z\|_{C([0,\delta];H^3)} \leq e^{\nu/2} (A_0 + \nu^{-1/2} G). \quad (28)$$

Pairing  $A^2 z_t - \nu \Delta A^2 z = A^2 F$  with  $-\Delta A^2 z$  gives

$$\frac{1}{2} \frac{d}{dt} \|\nabla A^2 z\|_2^2 + \nu \|\Delta A^2 z\|_2^2 = \operatorname{Re} \langle A^2 F, -\Delta A^2 z \rangle_{L^2}.$$

Integration and Young's inequality yield

$$\|\Delta z\|_{L^2(0,\delta;H^2)} \leq \nu^{-1/2} A_0 + \nu^{-1} G. \quad (29)$$

The pointwise multiplier inequality

$$\langle \xi \rangle^8 \leq 2(\langle \xi \rangle^6 + |\xi|^4 \langle \xi \rangle^4)$$

and  $\delta \leq 1$  imply

$$\|z\|_{L^2(0,\delta;H^4)} \leq \sqrt{2} \left( \|z\|_{C([0,\delta];H^3)} + \|\Delta z\|_{L^2(0,\delta;H^2)} \right). \quad (30)$$

The equation and (29) give

$$\|z_t\|_{L^2(0,\delta;H^2)} \leq \nu^{1/2} A_0 + 2G. \quad (31)$$

Equations (28)–(31) give

$$\begin{aligned} & \|z\|_{CH^3} + \|z\|_{L^2 H^4} + \|z_t\|_{L^2 H^2} \\ & \leq e^{\nu/2} (A_0 + \nu^{-1/2} G) + \sqrt{2} \left( e^{\nu/2} (A_0 + \nu^{-1/2} G) + \nu^{-1/2} A_0 + \nu^{-1} G \right) + \nu^{1/2} A_0 + 2G \\ & \leq L_\nu (A_0 + G). \end{aligned}$$

The last inequality follows from

$$\nu^{1/2} \leq \frac{1+\nu}{2}, \quad \nu^{-1/2} \leq \frac{1+\nu^{-1}}{2}, \quad e^{\nu/2} \leq e^{(\nu+\nu^{-1})/2}.$$

Fourier truncation and passage to the limit justify the preceding linear estimates for the stated data class.

For  $v, z \in H_\sigma^3$ , Lemmas 2.4 and 2.5 give

$$\|\mathbb{P} \nabla \cdot (v \otimes z)\|_{H^2} \leq C_q \|v\|_{H^3} \|z\|_{H^3}, \quad (32)$$

$$\|\mathbb{P} \nabla \cdot (v \otimes v - z \otimes z)\|_{H^2} \leq C_q (\|v\|_{H^3} + \|z\|_{H^3}) \|v - z\|_{H^3}. \quad (33)$$

Indeed,

$$\|\nabla \cdot (v \otimes z)\|_{H^2}^2 \leq 3 \sum_{i,j=1}^3 \|v_i z_j\|_{H^3}^2 \leq 3(C_3^{\text{alg}})^2 \|v\|_{H^3}^2 \|z\|_{H^3}^2.$$

Define

$$X_\delta := \{v \in C([0, \delta]; H_\sigma^3) \cap L^2(0, \delta; H_\sigma^4) : v_t \in L^2(0, \delta; H_\sigma^2)\},$$

$$\|v\|_{X_\delta} := \|v\|_{CH^3} + \|v\|_{L^2 H^4} + \|v_t\|_{L^2 H^2}, \quad X_\delta(a) := \{v \in X_\delta : v(0) = a\}.$$

$X_\delta$  is a Banach space with this norm, and  $X_\delta(a)$  is closed. For  $A_0 > 0$ , set  $R := 2L_\nu A_0$  and

$$\mathbb{B}_R := \{v \in X_\delta(a) : \|v\|_{X_\delta} \leq R\}.$$

The set  $\mathbb{B}_R$  is nonempty: the zero-forcing heat trajectory  $v^{\text{lin}}(t) = e^{\nu t \Delta} a$  belongs to  $X_\delta(a)$ , and the linear estimate with  $F = 0$  gives  $\|v^{\text{lin}}\|_{X_\delta} \leq L_\nu A_0 < R$ . For  $v \in \mathbb{B}_R$ , define

$$\Phi(v)(t) := e^{\nu t \Delta} a - \int_0^t e^{\nu(t-s)\Delta} \mathbb{P} \nabla \cdot (v \otimes v)(s) ds.$$

If  $\delta \leq \delta_\nu(a)$ , then

$$\delta^{1/2} \leq \frac{1}{8C_q L_\nu^2 (1 + A_0)}.$$

The linear estimate and (32) imply

$$\begin{aligned} \|\Phi(v)\|_{X_\delta} &\leq L_\nu (A_0 + C_q \delta^{1/2} R^2) \\ &\leq L_\nu A_0 + \frac{L_\nu A_0^2}{2(1 + A_0)} < R. \end{aligned}$$

For  $v, z \in \mathbb{B}_R$ ,

$$\begin{aligned} \|\Phi(v) - \Phi(z)\|_{X_\delta} &\leq L_\nu C_q \delta^{1/2} (\|v\|_{CH^3} + \|z\|_{CH^3}) \|v - z\|_{CH^3} \\ &\leq \frac{A_0}{2(1 + A_0)} \|v - z\|_{X_\delta} \leq \frac{1}{2} \|v - z\|_{X_\delta}. \end{aligned}$$

Thus  $\Phi$  has a unique fixed point in  $\mathbb{B}_R$ . If  $A_0 = 0$ , the zero function is the fixed point. Let  $b > 0$  and let

$$(v, q_v), (z, q_z) \in \mathcal{M}_\nu^3(0, b; a).$$

Set

$$M_b := \|v\|_{C([0, b]; H^3)} + \|z\|_{C([0, b]; H^3)}.$$

On every subinterval  $I \subset [0, b]$  of length  $h \leq 1$  on which the two velocities have the same left trace, their shifted Duhamel identities and the zero-data linear estimate give

$$\|v - z\|_{C(I; H^3)} \leq L_\nu C_q h^{1/2} M_b \|v - z\|_{C(I; H^3)}.$$

A finite partition with  $L_\nu C_q h^{1/2} M_b < 1$  proves  $v = z$  on  $[0, b]$ . The pressure formula gives  $q_v = q_z$ . This proves the at-most-one assertion for the full shifted mild class. For the fixed point, (32) places the forcing in  $L^2(0, \delta; H^2)$ ; the linear estimates give

$$w \in L^2(0, \delta; H^4), \quad w_t \in L^2(0, \delta; H^2).$$

Lemma 2.5 gives the normalized pressure  $q = R_i R_j (w_i w_j)$ .

Fix  $m \geq 3$  and assume  $a \in H_\sigma^m$ . Let  $S_k$  be Fourier restriction to  $\{|\xi| \leq k\}$ , set  $a_k = S_k a$ , and replace the fixed-point map above by

$$\Phi_k(v)(t) := e^{\nu t \Delta} a_k - \int_0^t e^{\nu(t-s)\Delta} S_k \mathbb{P} \nabla \cdot (v \otimes v)(s) \, ds.$$

The projection  $S_k$  is self-adjoint, contractive on every Sobolev space, and commutes with the heat flow,  $\mathbb{P}$ , and spatial derivatives. The same estimates that gave  $\Phi$  its contraction factor at most  $1/2$  apply to  $\Phi_k$  on the closed radius- $R$  ball in  $X_\delta(a_k)$ . Its fixed point  $w^{(k)} = S_k w^{(k)}$  consequently exists on  $[0, \delta]$  and satisfies

$$\sup_k \|w^{(k)}\|_{X_\delta} \leq R. \quad (34)$$

Because its Fourier support is bounded,  $w^{(k)}$  belongs to every spatial Sobolev space. Self-adjointness of  $S_k$  removes the cutoff from the energy pairing, so Lemma 2.8 gives

$$\frac{1}{2} \frac{d}{dt} \|w^{(k)}\|_{H^m}^2 + \nu \|\nabla w^{(k)}\|_{H^m}^2 \leq K_m \|\nabla w^{(k)}\|_{L^\infty} \|w^{(k)}\|_{H^m}^2.$$

Set

$$C_{\nabla,3} := (2\pi)^{-3/2} \left( \int_{\mathbb{R}^3} |\xi|^2 \langle \xi \rangle^{-6} \, d\xi \right)^{1/2}.$$

Fourier inversion and Cauchy–Schwarz give  $\|\nabla v\|_\infty \leq C_{\nabla,3} \|v\|_{H^3}$ . From (34),

$$\int_0^\delta \|\nabla w^{(k)}(s)\|_\infty \, ds \leq C_{\nabla,3} \delta R.$$

Put

$$F_k(t) := \|w^{(k)}(t)\|_{H^m}^2 + 2\nu \int_0^t \|\nabla w^{(k)}(s)\|_{H^m}^2 \, ds.$$

The differential inequality above gives

$$F_k(t) \leq \|a\|_{H^m}^2 + 2K_m \int_0^t \|\nabla w^{(k)}(s)\|_\infty F_k(s) \, ds.$$

Gronwall's inequality now gives the uniform bound

$$\sup_{0 \leq t \leq \delta} \left( \|w^{(k)}(t)\|_{H^m}^2 + 2\nu \int_0^t \|\nabla w^{(k)}(s)\|_{H^m}^2 \, ds \right) \leq e^{2K_m C_{\nabla,3} \delta R} \|a\|_{H^m}^2. \quad (35)$$

Since  $\|f\|_{H^{m+1}}^2 = \|f\|_{H^m}^2 + \|\nabla f\|_{H^m}^2$ , this also bounds  $w^{(k)}$  in  $L^2 H^{m+1}$ . The cutoff equations and Lemma 2.4 give, with a constant depending only on  $m$ ,

$$\|\partial_t w^{(k)}\|_{L^2 H^{m-1}} \leq \nu \|w^{(k)}\|_{L^2 H^{m+1}} + \sqrt{3} C_m^{\text{alg}} \delta^{1/2} \|w^{(k)}\|_{L^\infty H^m}^2,$$

so the time derivatives are uniformly bounded in  $L^2 H^{m-1}$ .

It remains to identify the higher-order weak limit with the fixed point already constructed in  $X_\delta(a)$ . The contraction estimate gives

$$\|w^{(k)} - w\|_{X_\delta} \leq 2 \|\Phi_k(w) - \Phi(w)\|_{X_\delta}.$$

The linear bound proved above controls the right-hand side by a constant times

$$\|(S_k - I)a\|_{H^3} + \|(S_k - I)\mathbb{P}\nabla \cdot (w \otimes w)\|_{L^2 H^2},$$

which tends to zero. Hence  $w^{(k)} \rightarrow w$  in  $X_\delta$ . Weak compactness in the three uniformly bounded higher-order spaces and uniqueness of the distributional limit place

$$w \in L^\infty(0, \delta; H_\sigma^m) \cap L^2(0, \delta; H_\sigma^{m+1}), \quad w_t \in L^2(0, \delta; H_\sigma^{m-1}).$$

Lemma 2.7 supplies the representative  $w \in C([0, \delta]; H_\sigma^m)$ ; its trace at zero is  $a$ , since it is the same distributional trace as the already constructed  $CH^3$  solution. Assume now that  $a \in H_\sigma^\infty$ . Set  $W_0 = w$  and define recursively

$$Q_n = R_i R_j \sum_{a=0}^n \binom{n}{a} W_{a,i} W_{n-a,j},$$

$$W_{n+1} = \nu \Delta W_n - \sum_{a=0}^n \binom{n}{a} (W_a \cdot \nabla) W_{n-a} - \nabla Q_n.$$

If  $W_0, \dots, W_n \in CH^{m+2}$ , then Lemmas 2.4 and 2.5 give

$$Q_n \in CH^{m+1}, \quad W_{n+1} \in CH^m.$$

Since  $m$  is arbitrary, induction gives  $W_n, Q_n \in CH^m$  for every  $n, m \in \mathbb{N}_0$ . The equation first gives  $W_0 \in C^1 H^m$  and  $\partial_t W_0 = W_1$ . If  $W_a = \partial_t^a w$  for  $0 \leq a \leq n$ , distributional differentiation of  $q = R_i R_j (w_i w_j)$  and Lemma 2.6 give  $\partial_t^n q = Q_n$ . Differentiation of the momentum equation then gives  $\partial_t W_n = W_{n+1}$ . Induction gives

$$W_n \in C^1 H^m, \quad \partial_t W_n = W_{n+1}, \quad W_n = \partial_t^n w, \quad Q_n = \partial_t^n q.$$

Time translation proves the result for every  $s \geq 0$ . Lemma 2.10 and the at-most-one assertion make all nonempty classes below  $\tau_{\nu,3}^{\max}(s, a)$  compatible; their union is the maximal shifted pressure-normalized  $H^3$  mild history.  $\square$

### 3 The pressure-complete setting and mixed jet

**Proposition 3.1** (Pressure normalization). *Let  $u, p$  be smooth on  $I \times \mathbb{R}^3$ , let  $\nabla \cdot u = 0$ , and suppose*

$$u(t) \in H_\sigma^1(\mathbb{R}^3), \quad p(t) \in H^0(\mathbb{R}^3) \quad (t \in I),$$

and

$$u_t + (u \cdot \nabla)u + \nabla p = \nu \Delta u. \tag{36}$$

Then

$$-\Delta p = \partial_i \partial_j (u_i u_j), \tag{37}$$

$$p = R_i R_j (u_i u_j), \tag{38}$$

$$(u \cdot \nabla)u + \nabla p = \mathbb{P}\nabla \cdot (u \otimes u). \tag{39}$$

The condition  $p(t) \in H^0(\mathbb{R}^3)$  fixes the pressure uniquely.

*Proof.* Incompressibility gives

$$(u \cdot \nabla)u_i = \partial_j(u_j u_i).$$

Taking the divergence of (36) cancels  $\nabla \cdot u_t$  and  $\nabla \cdot \Delta u$ , which yields (37). Lemma 2.5, applied with  $m = 0$ , gives  $q := R_i R_j(u_i u_j) \in H^0$ , the same Poisson identity, and

$$(u \cdot \nabla)u + \nabla q = \mathbb{P} \nabla \cdot (u \otimes u).$$

If  $p_1, p_2 \in H^0$  satisfy (37), then  $h = p_1 - p_2$  satisfies  $|\xi|^2 \widehat{h} = 0$ . Since  $\widehat{h} \in L^2$  and is supported at  $\{0\}$ , it vanishes. Hence  $p = q$ , which proves (38) and (39).  $\square$

**Definition 3.2** (Maximal pressure-complete classical history). Fix  $\nu > 0$  and  $u_0 \in H_\sigma^\infty(\mathbb{R}^3)$ . A pressure-complete classical history generated by  $(u_0, \nu)$  is a triple

$$(T_*, u, p), \quad 0 < T_* \leq \infty,$$

such that, for every  $0 < T < T_*$  and  $n, m \in \mathbb{N}_0$ ,

$$\partial_t^n u \in C([0, T]; H_\sigma^m(\mathbb{R}^3)),$$

$$\partial_t^n p \in C([0, T]; H^m(\mathbb{R}^3)),$$

and

$$u_t + (u \cdot \nabla)u + \nabla p = \nu \Delta u, \tag{40}$$

$$\nabla \cdot u = 0, \tag{41}$$

$$u(0) = u_0, \tag{42}$$

$$p(t) = R_i R_j(u_i(t) u_j(t)). \tag{43}$$

The history is maximal if no pressure-complete classical history  $(\widetilde{T}, \widetilde{u}, \widetilde{p})$  generated by  $(u_0, \nu)$  satisfies

$$\widetilde{T} > T_*, \quad (\widetilde{u}, \widetilde{p})|_{[0, T_*]} = (u, p).$$

**Definition 3.3** (Temporal and mixed velocity–pressure jets). Let  $(T_*, u, p)$  be a maximal pressure-complete classical history. For  $n \in \mathbb{N}_0$  and  $\alpha \in \mathbb{N}_0^3$ , define

$$V_n := \partial_t^n u, \quad Q_n := \partial_t^n p, \quad J_{n, \alpha} := \partial_t^n \partial_x^\alpha u, \quad \Pi_{n, \alpha} := \partial_t^n \partial_x^\alpha p.$$

For  $\beta \leq \alpha$ , define

$$\binom{\alpha}{\beta} := \prod_{\ell=1}^3 \binom{\alpha_\ell}{\beta_\ell}, \quad |\alpha| := \alpha_1 + \alpha_2 + \alpha_3.$$

**Proposition 3.4** (Pressure-complete binomial recurrences). *For every  $n \in \mathbb{N}_0$  and  $\alpha \in \mathbb{N}_0^3$ ,*

$$Q_n = R_i R_j \sum_{a=0}^n \binom{n}{a} V_{a,i} V_{n-a,j}, \quad (44)$$

$$V_{n+1} = \nu \Delta V_n - \sum_{a=0}^n \binom{n}{a} (V_a \cdot \nabla) V_{n-a} - \nabla Q_n, \quad (45)$$

$$\Pi_{n,\alpha} = R_i R_j \sum_{a=0}^n \sum_{\beta \leq \alpha} \binom{n}{a} \binom{\alpha}{\beta} J_{a,\beta,i} J_{n-a,\alpha-\beta,j}, \quad (46)$$

$$J_{n+1,\alpha} + \sum_{a=0}^n \sum_{\beta \leq \alpha} \binom{n}{a} \binom{\alpha}{\beta} (J_{a,\beta} \cdot \nabla) J_{n-a,\alpha-\beta} + \nabla \Pi_{n,\alpha} = \nu \Delta J_{n,\alpha}, \quad (47)$$

$$\nabla \cdot J_{n,\alpha} = 0. \quad (48)$$

*Proof.* Apply  $\partial_t^n$  to (43) and use Lemma 2.6 with  $\alpha = 0$  to obtain (44). Apply  $\partial_t^n$  to (40) to obtain (45). Apply  $\partial_x^\alpha$  to both identities, use Lemma 2.6, and commute the derivatives with  $R_i R_j$  to obtain (46) and (47). Differentiation of (41) gives (48).  $\square$

**Definition 3.5** (Canonical pressure constant). For vector fields  $f, g$ , set

$$\mathcal{Q}(f, g) := \sum_{i,j=1}^3 R_i R_j (f_i g_j).$$

For  $m \in \mathbb{N}_0$ , define

$$C_m^p := \sup_{f,g \in H^{m+1}(\mathbb{R}^3; \mathbb{R}^3) \setminus \{0\}} \frac{\|\mathcal{Q}(f, g)\|_{H^m} + \|\nabla \mathcal{Q}(f, g)\|_{H^{m-1}}}{\|f\|_{H^{m+1}} \|g\|_{H^{m+1}}}.$$

**Lemma 3.6.** *For every  $m \in \mathbb{N}_0$ ,*

$$C_m^p \leq 2C_m^{\text{alg}}.$$

*Proof.* Regard  $(f_i g_j)_{i,j}$  as one tensor. The contraction  $F \mapsto R_i R_j F_{ij}$  has Fourier-multiplier norm one from the tensor  $H^m$  norm to scalar  $H^m$ . The same product lemma and  $\|\nabla h\|_{H^{m-1}} \leq \|h\|_{H^m}$  give

$$\|\mathcal{Q}(f, g)\|_{H^m} + \|\nabla \mathcal{Q}(f, g)\|_{H^{m-1}} \leq 2C_m^{\text{alg}} \|f\|_{H^{m+1}} \|g\|_{H^{m+1}}.$$

$\square$

**Lemma 3.7** (Initial pressure-complete jet). *Set*

$$V_n^0 := V_n(0), \quad Q_n^0 := Q_n(0), \quad J_{n,\alpha}^0 := J_{n,\alpha}(0), \quad \Pi_{n,\alpha}^0 := \Pi_{n,\alpha}(0).$$

*Then*

$$V_0^0 = u_0, \quad (49)$$

$$Q_n^0 = R_i R_j \sum_{a=0}^n \binom{n}{a} V_{a,i}^0 V_{n-a,j}^0, \quad (50)$$

$$V_{n+1}^0 = \nu \Delta V_n^0 - \sum_{a=0}^n \binom{n}{a} (V_a^0 \cdot \nabla) V_{n-a}^0 - \nabla Q_n^0. \quad (51)$$

For every  $n, m \in \mathbb{N}_0$  and  $\alpha \in \mathbb{N}_0^3$ ,

$$V_n^0, Q_n^0, J_{n,\alpha}^0, \Pi_{n,\alpha}^0 \in H^m(\mathbb{R}^3).$$

*Proof.* Equations (49)–(51) are (44) and (45) at  $t = 0$ . Assume  $V_0^0, \dots, V_n^0 \in \bigcap_{m \geq 0} H^m$ . Lemmas 2.4 and 2.5 give

$$Q_n^0 \in \bigcap_{m \geq 0} H^m, \quad V_{n+1}^0 \in \bigcap_{m \geq 0} H^m.$$

Induction starts from  $u_0 \in H_\sigma^\infty$ . Spatial differentiation and (46) give the claims for  $J_{n,\alpha}^0$  and  $\Pi_{n,\alpha}^0$ .  $\square$

**Theorem 3.8** (Finite-row pressure estimates). *Let  $I = (0, T) \subset (0, T_*)$ , let  $n, M \in \mathbb{N}_0$ , and suppose*

$$V_a \in L^2(I; H^{M+1}(\mathbb{R}^3)) \quad (0 \leq a \leq n).$$

*Then*

$$\begin{aligned} & \|Q_n\|_{L^1(I; H^M)} + \|\nabla Q_n\|_{L^1(I; H^{M-1})} \\ & \leq C_M^p \sum_{a=0}^n \binom{n}{a} \|V_a\|_{L^2(I; H^{M+1})} \|V_{n-a}\|_{L^2(I; H^{M+1})}. \end{aligned} \quad (52)$$

*If*

$$\mathcal{V}_{n,M}(I)^2 := \sum_{a=0}^n \|V_a\|_{L^2(I; H^{M+1})}^2,$$

*then*

$$\|Q_n\|_{L^1(I; H^M)} + \|\nabla Q_n\|_{L^1(I; H^{M-1})} \leq C_M^p 2^n \mathcal{V}_{n,M}(I)^2. \quad (53)$$

For  $\alpha \in \mathbb{N}_0^3$ , assume

$$J_{a,\beta} \in L^2(I; H^{M+1}(\mathbb{R}^3)) \quad (0 \leq a \leq n, \beta \leq \alpha).$$

*Then*

$$\begin{aligned} & \|\Pi_{n,\alpha}\|_{L^1(I; H^M)} + \|\nabla \Pi_{n,\alpha}\|_{L^1(I; H^{M-1})} \\ & \leq C_M^p \sum_{a=0}^n \sum_{\beta \leq \alpha} \binom{n}{a} \binom{\alpha}{\beta} \|J_{a,\beta}\|_{L^2(I; H^{M+1})} \|J_{n-a,\alpha-\beta}\|_{L^2(I; H^{M+1})}. \end{aligned} \quad (54)$$

*If*

$$\mathcal{J}_{n,\alpha,M}(I)^2 := \sum_{a=0}^n \sum_{\beta \leq \alpha} \|J_{a,\beta}\|_{L^2(I; H^{M+1})}^2,$$

*then*

$$\|\Pi_{n,\alpha}\|_{L^1(I; H^M)} + \|\nabla \Pi_{n,\alpha}\|_{L^1(I; H^{M-1})} \leq C_M^p 2^{n+|\alpha|} \mathcal{J}_{n,\alpha,M}(I)^2. \quad (55)$$

*Proof.* Lemmas 2.4 and 2.5 give

$$\|R_i R_j(f_i g_j)\|_{H^M} + \|\nabla R_i R_j(f_i g_j)\|_{H^{M-1}} \leq C_M^p \|f\|_{H^{M+1}} \|g\|_{H^{M+1}}. \quad (56)$$

Equations (44) and (56) give, for almost every  $t \in I$ ,

$$\|Q_n(t)\|_{H^M} + \|\nabla Q_n(t)\|_{H^{M-1}} \leq C_M^p \sum_{a=0}^n \binom{n}{a} \|V_a(t)\|_{H^{M+1}} \|V_{n-a}(t)\|_{H^{M+1}}.$$

Integration and Cauchy–Schwarz prove (52). For nonnegative  $x_a$ ,

$$\sum_{a=0}^n \binom{n}{a} x_a x_{n-a} \leq \sum_{a=0}^n \binom{n}{a} x_a^2 \leq 2^n \sum_{a=0}^n x_a^2.$$

This proves (53).

Apply (56) to each summand in (46). Integration and Cauchy–Schwarz give (54). The involution

$$(a, \beta) \mapsto (n - a, \alpha - \beta)$$

preserves the weights  $\binom{n}{a} \binom{\alpha}{\beta}$ , and

$$\sum_{a=0}^n \sum_{\beta \leq \alpha} \binom{n}{a} \binom{\alpha}{\beta} = 2^{n+|\alpha|}.$$

In particular, every individual weight is at most  $2^{n+|\alpha|}$ . The inequality  $2xy \leq x^2 + y^2$  proves (55).  $\square$

## 4 Datum-generated compatible rectangles

**Proposition 4.1** (Kinetic-energy identity). *For every  $0 \leq t < T_*$ ,*

$$\frac{1}{2} \|u(t)\|_2^2 + \nu \int_0^t \|\nabla u(s)\|_2^2 \, ds = \frac{1}{2} \|u_0\|_2^2. \quad (57)$$

Consequently, for  $0 < T < T_*$ ,

$$\|u\|_{L^\infty(0,T;L^2)} \leq \|u_0\|_2, \quad (58)$$

$$\|u\|_{L^2(0,T;H^1)}^2 \leq \left(T + \frac{1}{2\nu}\right) \|u_0\|_2^2. \quad (59)$$

*Proof.* Take the  $L^2$  pairing of the momentum equation with  $u$ . Integration by parts and  $\nabla \cdot u = 0$  give

$$\langle (u \cdot \nabla)u, u \rangle_{L^2} = 0, \quad \langle \nabla p, u \rangle_{L^2} = 0, \quad \langle \Delta u, u \rangle_{L^2} = -\|\nabla u\|_2^2.$$

Integration in time proves (57). The two estimates follow by discarding the nonnegative dissipation and adding  $\int_0^T \|u(s)\|_2^2 \, ds$ .  $\square$

**Definition 4.2** (Completed height rows). For  $r \in \mathbb{N}_0$  and  $s \geq 0$ , set

$$E_{r,s}(t) := \frac{1}{2} \|\Lambda^s V_r(t)\|_2^2, \quad (60)$$

$$\mathcal{P}_{r,s}(t) := -\operatorname{Re} \left\langle \Lambda^s V_r(t), \Lambda^s \left( \sum_{a=0}^r \binom{r}{a} (V_a \cdot \nabla) V_{r-a} + \nabla Q_r \right) (t) \right\rangle_{L^2}. \quad (61)$$

For the base row write  $E_s := E_{0,s}$ ,  $\mathcal{P}_s := \mathcal{P}_{0,s}$ , and  $H := E_{1/2}$ .

**Proposition 4.3** (Completed height recurrence in every temporal row). *For every strict classical interval  $(0, T)$ , every  $r \in \mathbb{N}_0$ , and every  $s \geq 0$ ,*

$$\partial_t E_{r,s} = \mathcal{P}_{r,s} - 2\nu E_{r,s+1}. \quad (62)$$

For every  $k \in \mathbb{N}$ ,

$$\partial_t^k E_{r,1/2} = (-2\nu)^k E_{r,k+1/2} + \sum_{j=0}^{k-1} (-2\nu)^j \partial_t^{k-1-j} \mathcal{P}_{r,j+1/2}. \quad (63)$$

In the base row  $r = 0$ ,

$$H^{(k)} = (-2\nu)^k E_{k+1/2} + \sum_{j=0}^{k-1} (-2\nu)^j \partial_t^{k-1-j} \mathcal{P}_{j+1/2}, \quad (64)$$

and the highest spatial energy is the exact coordinate

$$E_{k+1/2} = \frac{H^{(k)} - \sum_{j=0}^{k-1} (-2\nu)^j \partial_t^{k-1-j} \mathcal{P}_{j+1/2}}{(-2\nu)^k}. \quad (65)$$

*Proof.* Pair the temporal recurrence for  $V_r$  with  $\Lambda^{2s} V_r$ . The pressure-complete nonlinear sum is exactly  $\mathcal{P}_{r,s}$ , while

$$\nu \operatorname{Re} \langle \Lambda^s V_r, \Lambda^s \Delta V_r \rangle_{L^2} = -\nu \|\Lambda^{s+1} V_r\|_2^2 = -2\nu E_{r,s+1}.$$

This proves (62). Induction on  $k$  gives the stronger identity

$$\partial_t^k E_{r,s} = (-2\nu)^k E_{r,s+k} + \sum_{j=0}^{k-1} (-2\nu)^j \partial_t^{k-1-j} \mathcal{P}_{r,s+j}. \quad (66)$$

Indeed, differentiation of the order- $k$  identity followed by  $\partial_t E_{r,s+k} = \mathcal{P}_{r,s+k} - 2\nu E_{r,s+k+1}$  produces the order- $k+1$  identity. Taking  $s = 1/2$  proves (63). Taking  $r = 0$  gives (64); since  $\nu > 0$ , division by  $(-2\nu)^k$  gives (65).  $\square$

**Definition 4.4** (Completed critical-height coordinates). Set

$$R_0 := H = E_{1/2}, \quad R_k := \frac{H^{(k)} - \sum_{j=0}^{k-1} (-2\nu)^j \partial_t^{k-1-j} \mathcal{P}_{j+1/2}}{(-2\nu)^k} \quad (k \in \mathbb{N}). \quad (67)$$

Proposition 4.3 gives the exact identity

$$R_k = E_{k+1/2} = \frac{1}{2} \left\| \Lambda^{k+1/2} u \right\|_2^2 \quad (k \in \mathbb{N}_0). \quad (68)$$

**Definition 4.5** (Finite mixed rectangle). Let  $0 < T < T_*$  and  $N, M \in \mathbb{N}_0$ . Define

$$\mathfrak{R}_{N,M}(u; T) := \sum_{n=0}^N \|V_n\|_{L^2(0,T;H^{M+1})}^2 + \sum_{n=0}^N \|V_{n+1}\|_{L^2(0,T;H^{M-1})}^2, \quad (69)$$

$$\mathfrak{Q}_{N,M}(p; T) := \sum_{n=0}^N \left( \|Q_n\|_{L^1(0,T;H^M)} + \|\nabla Q_n\|_{L^1(0,T;H^{M-1})} \right). \quad (70)$$

The pressure-complete record is

$$\mathcal{R}_{N,M}[u_0; T] := ((V_n|_{(0,T)})_{n=0}^N, (V_{n+1}|_{(0,T)})_{n=0}^N; (Q_n|_{(0,T)})_{n=0}^N),$$

with the pressure-gradient coordinate retained through (70) and with every row constrained by Proposition 3.4.

**Theorem 4.6** (Datum-generated whole-terminal rectangles). *Let  $\nu > 0$ ,  $u_0 \in H_\sigma^\infty(\mathbb{R}^3)$ , and let  $(T_*, u, p)$  be the maximal pressure-complete classical history generated by  $(u_0, \nu)$ . If  $T_* < \infty$ , then, for every  $N, M \in \mathbb{N}_0$ ,*

$$\mathfrak{R}_{N,M}^*[u_0, \nu, T_*] := \sup_{0 < T < T_*} \mathfrak{R}_{N,M}(u; T) < \infty. \quad (71)$$

*This assertion is coordinatewise in  $(N, M)$ ; no bound uniform in all derivative orders is asserted. The quantity  $\mathfrak{R}_{N,M}^*$  is the exact supremum for this history; for general data the theorem does not replace it by a formula involving only finitely many norms of  $u_0$ . Equivalently, for every  $n, m \in \mathbb{N}_0$ ,*

$$V_n \in L^2(0, T_*; H^{m+1}), \quad V_{n+1} \in L^2(0, T_*; H^{m-1}). \quad (72)$$

*Every coordinate in (72) is the corresponding distributional derivative of the single maximal history.*

The theorem is displayed in rectangle form, but its proof order is

$$u_0 \longrightarrow \text{pressure-complete mixed jet} \longrightarrow \text{whole-terminal adjacent rows} \longrightarrow \text{finite rectangles}.$$

The middle arrow is the fixed datum-side payment: the rows in (72) are the actual derivatives of the one maximal history and have the stated whole-terminal integrability. The last arrow is finite summation. The next proposition writes out that final step, proves the converse by monotone exhaustion, and realizes the cutoff norms on  $(0, T_*)$ .

**Proposition 4.7** (Equivalence and whole-terminal realization). *For the fixed maximal history, the family of bounds (71) is equivalent to the adjacent-row statement (72). Whenever either formulation holds, for every  $N, M \in \mathbb{N}_0$ ,*

$$\sum_{n=0}^N \left( \|V_n\|_{L^2(0, T_*; H^{M+1})}^2 + \|V_{n+1}\|_{L^2(0, T_*; H^{M-1})}^2 \right) = \lim_{T \uparrow T_*} \mathfrak{R}_{N,M}(u; T) = \mathfrak{R}_{N,M}^*[u_0, \nu, T_*]. \quad (73)$$

*Every row is the corresponding distributional derivative of the single pressure-normalized maximal history.*

*Proof.* Assume first (72). For fixed  $N, M$ , set

$$C_{N,M} := \sum_{n=0}^N \left( \|V_n\|_{L^2(0, T_*; H^{M+1})}^2 + \|V_{n+1}\|_{L^2(0, T_*; H^{M-1})}^2 \right).$$

The sum is finite because it contains finitely many instances of (72). Restriction from  $(0, T_*)$  to  $(0, T)$  gives

$$\mathfrak{R}_{N,M}(u; T) \leq C_{N,M} \quad (0 < T < T_*),$$

and hence (71).

Conversely, assume (71). Every term in  $\mathfrak{R}_{N,M}(u; T)$  is the integral of a nonnegative function. Monotone exhaustion of  $(0, T_*)$  gives (73). Taking  $N = n$  and  $M = m$  yields (72) for each finite  $(n, m)$ . Proposition 3.4 identifies every entry with the corresponding distributional derivative of the fixed pressure-normalized history.  $\square$

**Corollary 4.8** (Whole-terminal completion of every critical height). *Under the hypotheses of Theorem 4.6, every completed critical coordinate belongs to  $L^1(0, T_*)$ , and*

$$\mathcal{H}_k[u_0, \nu, T_*] := \|R_k\|_{L^1(0, T_*)} = \frac{1}{2} \left\| \Lambda^{k+1/2} u \right\|_{L^2(0, T_*; L^2)}^2 \leq \frac{1}{2} \mathfrak{R}_{0,k}^*[u_0, \nu, T_*] \quad (k \in \mathbb{N}_0). \quad (74)$$

For every  $m \in \mathbb{N}_0$ , the same row yields the spatial-column projection

$$\|u\|_{L^2(0, T_*; H^m)}^2 \leq 2^m \left( T_* \|u_0\|_2^2 + \mathfrak{R}_{0,m}^*[u_0, \nu, T_*] \right). \quad (75)$$

Thus

$$(R_k)_{k \in \mathbb{N}_0} \in \prod_{k \in \mathbb{N}_0} L^1(0, T_*), \quad u \in \bigcap_{m \in \mathbb{N}_0} L^2(0, T_*; H^m(\mathbb{R}^3)). \quad (76)$$

*Proof.* Take  $N = 0$  and  $M = k$  in (73). Since  $|\xi|^{2k+1} \leq \langle \xi \rangle^{2k+2}$ ,

$$\begin{aligned} 2 \|R_k\|_{L^1(0, T_*)} &= \left\| \Lambda^{k+1/2} u \right\|_{L^2(0, T_*; L^2)}^2 \\ &\leq \|u\|_{L^2(0, T_*; H^{k+1})}^2 \leq \mathfrak{R}_{0,k}^*[u_0, \nu, T_*], \end{aligned}$$

which proves (74). For  $|\xi| \leq 1$ ,  $\langle \xi \rangle^{2m} \leq 2^m$ ; for  $|\xi| > 1$ ,  $\langle \xi \rangle^{2m} \leq 2^m |\xi|^{2m+1}$ . Hence

$$\|f\|_{H^m}^2 \leq 2^m \left( \|f\|_2^2 + \left\| \Lambda^{m+1/2} f \right\|_2^2 \right).$$

Integrating this inequality, using (58) and (74), proves (75). Since  $k$  and  $m$  are arbitrary finite indices, (76) follows.  $\square$

**Proposition 4.9** (Pressure completion). *Under the hypotheses of Theorem 4.6, for every  $N, M \in \mathbb{N}_0$ ,*

$$\mathfrak{Q}_{N,M}^*[u_0, \nu, T_*] := \sup_{0 < T < T_*} \mathfrak{Q}_{N,M}(p; T) \leq C_M^p (2^{N+1} - 1) \mathfrak{R}_{N,M}^*[u_0, \nu, T_*]. \quad (77)$$

For every  $0 \leq n \leq N$ , the identities

$$Q_n = R_i R_j \sum_{a=0}^n \binom{n}{a} V_{a,i} V_{n-a,j}, \quad (78)$$

$$V_{n+1} = \nu \Delta V_n - \sum_{a=0}^n \binom{n}{a} (V_a \cdot \nabla) V_{n-a} - \nabla Q_n \quad (79)$$

hold respectively in  $L^1(0, T_*; H^M)$  and  $L^1(0, T_*; H^{M-1})$ , including when  $M = 0$ .

*Proof.* For every  $0 < T < T_*$ , Theorem 3.8 gives

$$\|Q_n\|_{L^1(0, T; H^M)} + \|\nabla Q_n\|_{L^1(0, T; H^{M-1})} \leq C_M^p 2^n \mathfrak{R}_{N,M}^*[u_0, \nu, T_*].$$

Sum over  $0 \leq n \leq N$ , then let  $T \uparrow T_*$ . Monotone convergence in the nonnegative norm integrals proves (77). Since  $\nabla \cdot V_a = 0$ ,

$$(V_a \cdot \nabla) V_{n-a} = \nabla \cdot (V_a \otimes V_{n-a}).$$

Lemma 2.4 and Cauchy–Schwarz in time give

$$\|V_a \otimes V_{n-a}\|_{L^1(0,T_*;H^M)} \leq C_M^{\text{alg}} \|V_a\|_{L^2(0,T_*;H^{M+1})} \|V_{n-a}\|_{L^2(0,T_*;H^{M+1})}.$$

Thus every transport term belongs to  $L^1(0, T_*; H^{M-1})$ , also for  $M = 0$ . Moreover,

$$\|\Delta V_n\|_{L^1 H^{M-1}} \leq T_*^{1/2} \|V_n\|_{L^2 H^{M+1}}.$$

Theorem 4.6 places  $V_{n+1}$  in  $L^2(0, T_*; H^{M-1})$ , hence in  $L^1(0, T_*; H^{M-1})$ . The strict-interval identities are identities of distributions; their whole-terminal representatives belong to the stated spaces. This proves the two identities on  $(0, T_*)$ .  $\square$

**Definition 4.10** (Finite-window spaces). For  $N, M \in \mathbb{N}_0$ , set

$$\mathcal{X}_{N,M} := \prod_{n=0}^N L^2(0, T_*; H^{M+1}) \times \prod_{n=0}^N L^2(0, T_*; H^{M-1}) \times \prod_{n=0}^N L^1(0, T_*; H^M).$$

The whole-terminal rectangle is the element

$$\mathcal{R}_{N,M}[u_0; T_*] := ((V_n)_{n=0}^N, (V_{n+1})_{n=0}^N; (Q_n)_{n=0}^N) \in \mathcal{X}_{N,M}. \quad (80)$$

If  $N' \leq N$  and  $M' \leq M$ , let

$$\rho_{N',M'}^{N,M} : \mathcal{X}_{N,M} \longrightarrow \mathcal{X}_{N',M'}$$

discard the temporal coordinates with index larger than  $N'$  and apply

$$H^{M+1} \hookrightarrow H^{M'+1}, \quad H^{M-1} \hookrightarrow H^{M'-1}, \quad H^M \hookrightarrow H^{M'}$$

to the retained coordinates.

**Proposition 4.11** (Restriction compatibility). For  $N' \leq N$  and  $M' \leq M$ ,

$$\rho_{N',M'}^{N,M} \mathcal{R}_{N,M}[u_0; T_*] = \mathcal{R}_{N',M'}[u_0; T_*]. \quad (81)$$

If  $(N'', M'') \leq (N', M') \leq (N, M)$ , then

$$\rho_{N'',M''}^{N',M'} \rho_{N',M'}^{N,M} = \rho_{N'',M''}^{N,M}.$$

Thus the spaces  $\mathcal{X}_{N,M}$  and the maps  $\rho_{N',M'}^{N,M}$  form an inverse system, and the datum-generated rectangles form a compatible family in it.

*Proof.* All retained entries are the same distributions  $\partial_t^n u$ ,  $\partial_t^{n+1} u$ , and  $\partial_t^n p$ . The Sobolev embeddings preserve those distributions. Deletion of rows and composition of canonical embeddings are associative.  $\square$

## 5 Compatible intersection, endpoint jet, and restart

**Definition 5.1** (Compatible intersection). Let

$$\mathcal{X}_\infty := \varprojlim_{N,M} \mathcal{X}_{N,M} = \left\{ (X_{N,M})_{N,M} : \begin{array}{l} X_{N,M} \in \mathcal{X}_{N,M}, \\ \rho_{N',M'}^{N,M} X_{N,M} = X_{N',M'} \quad \text{whenever } N' \leq N, M' \leq M \end{array} \right\}. \quad (82)$$

The datum-specific family is

$$\mathcal{I}[u_0] := (\mathcal{R}_{N,M}[u_0; T_*])_{N,M \in \mathbb{N}_0}. \quad (83)$$

**Theorem 5.2** (Compatible assembly and whole-terminal mixed regularity). *Let  $\nu > 0$ ,  $u_0 \in H_\sigma^\infty(\mathbb{R}^3)$ , and let  $(T_*, u, p)$  be the maximal pressure-complete classical history generated by  $(u_0, \nu)$ . If  $T_* < \infty$ , then*

$$\mathcal{I}[u_0] = (\mathcal{R}_{N,M}[u_0; T_*])_{N,M \in \mathbb{N}_0} \in \mathcal{X}_\infty \quad (84)$$

*and all of its finite coordinates are the restrictions of the same pressure-complete history. Its coordinates  $(V_n, Q_n)_{n \in \mathbb{N}_0}$  satisfy*

$$V_n \in \bigcap_{m \in \mathbb{N}_0} L^2(0, T_*; H_\sigma^m(\mathbb{R}^3)), \quad (85)$$

$$Q_n \in \bigcap_{m \in \mathbb{N}_0} L^1(0, T_*; H^m(\mathbb{R}^3)), \quad (86)$$

*and every finite restriction is the corresponding rectangle  $\mathcal{R}_{N,M}[u_0; T_*]$ . Its velocity realization satisfies*

$$u \in \bigcap_{r,m \in \mathbb{N}_0} H^r(0, T_*; H^m(\mathbb{R}^3)), \quad (87)$$

*and, for every  $r, m \in \mathbb{N}_0$ ,*

$$\sum_{n=0}^r \|\partial_t^n u\|_{L^2(0, T_*; H^m)}^2 \leq \mathfrak{R}_{r,m}^*[u_0, \nu, T_*]. \quad (88)$$

*Proof.* Theorem 4.6 places every velocity coordinate of  $\mathcal{R}_{N,M}[u_0; T_*]$  in its whole-terminal adjacent spaces. Proposition 4.9 places every retained pressure coordinate in its whole-terminal pressure space. Proposition 4.11 gives

$$\rho_{N',M'}^{N,M} \mathcal{R}_{N,M}[u_0; T_*] = \mathcal{R}_{N',M'}[u_0; T_*] \quad (N' \leq N, M' \leq M),$$

so (84) belongs to  $\mathcal{X}_\infty$ .

Fix  $n$ . Every rectangle with  $N \geq n$  contains the same distributions  $\partial_t^n u$  and  $\partial_t^n p$ . Increasing  $M$ , using the canonical Sobolev embeddings, gives (85) and (86). Thus each finite rectangle reads the same distributional coordinate in its own Sobolev topology.

Fix finite  $r, m$  and choose  $N \geq r, M \geq m$ . The upper rectangle edges give

$$\partial_t^n u \in L^2(0, T_*; H^{M+1}) \hookrightarrow L^2(0, T_*; H^m) \quad (0 \leq n \leq r).$$

They are the distributional time derivatives of the same  $u$ , so  $u \in H^r(0, T_*; H^m)$ . Since  $(r, m)$  was arbitrary, (87) follows. Summing the retained coordinates and using Sobolev nesting gives (88).  $\square$

**Proposition 5.3** (Complete spatial row and common endpoint). *Under the hypotheses of Theorem 5.2, for every  $m \in \mathbb{N}_0$ ,*

$$u \in W^{1,1}(0, T_*; H_\sigma^m(\mathbb{R}^3)) \hookrightarrow C([0, T_*]; H_\sigma^m(\mathbb{R}^3)). \quad (89)$$

*The endpoint values supplied at the different Sobolev orders are the images of one common state*

$$u_* := u(T_*) \in \bigcap_{m \in \mathbb{N}_0} H_\sigma^m(\mathbb{R}^3) = H_\sigma^\infty(\mathbb{R}^3). \quad (90)$$

*Proof.* Fix  $m$ . The zeroth temporal coordinate of (87) gives

$$u \in L^2(0, T_*; H^{m+2}).$$

The pressure-completed nonlinearity satisfies

$$\|\mathbb{P}\nabla \cdot (u \otimes u)\|_{H^m} \leq C_m \|u\|_{H^{m+2}}^2$$

by Lemmas 2.4 and 2.5. It belongs to  $L^1(0, T_*; H^m)$ . The same spatial row and  $T_* < \infty$  give  $\Delta u \in L^1(0, T_*; H^m)$ , and the projected equation

$$u_t = \nu \Delta u - \mathbb{P}\nabla \cdot (u \otimes u)$$

places  $u_t$  in  $L^1(0, T_*; H^m)$ . Also  $u \in L^1(0, T_*; H^m)$ , so the Banach-valued fundamental theorem gives (89).

If  $k \geq m$ , the  $H^k$ - and  $H^m$ -continuous representatives are the same preterminal function after the embedding  $H^k \hookrightarrow H^m$ . Uniqueness of continuous representatives makes their endpoint values agree under that embedding. These compatible values define the single state in (90). Continuity of divergence from  $H^{m+1}$  to  $H^m$  gives  $\nabla \cdot u_* = 0$ .  $\square$

**Corollary 5.4** (Whole-terminal pressure-complete mixed jet). *Under the hypotheses of Theorem 5.2, for every  $r, m, n \in \mathbb{N}_0$  and  $\alpha \in \mathbb{N}_0^3$ ,*

$$u \in H^r(0, T_*; H_\sigma^m(\mathbb{R}^3)), \quad (91)$$

$$J_{n,\alpha} := \partial_x^\alpha V_n \in L^2(0, T_*; H^{m+1}(\mathbb{R}^3)), \quad (92)$$

$$J_{n+1,\alpha} \in L^2(0, T_*; H^{m-1}(\mathbb{R}^3)), \quad (93)$$

$$\Pi_{n,\alpha} := \partial_x^\alpha Q_n \in L^1(0, T_*; H^m(\mathbb{R}^3)). \quad (94)$$

The following finite bounds hold:

$$\sum_{j=0}^r \|V_j\|_{L^2(0, T_*; H^m)}^2 \leq \mathfrak{R}_{r,m}^*[u_0, \nu, T_*], \quad (95)$$

$$\|J_{n,\alpha}\|_{L^2(0, T_*; H^{m+1})}^2 + \|J_{n+1,\alpha}\|_{L^2(0, T_*; H^{m-1})}^2 \leq \mathfrak{R}_{n,m+|\alpha|}^*[u_0, \nu, T_*], \quad (96)$$

$$\|\Pi_{n,\alpha}\|_{L^1(0, T_*; H^m)} \leq C_{m+|\alpha|}^p (2^{n+1} - 1) \mathfrak{R}_{n,m+|\alpha|}^*[u_0, \nu, T_*]. \quad (97)$$

Every mixed row satisfies the recurrences in Proposition 3.4 on  $(0, T_*)$ .

*Proof.* Equation (91) and (95) follow from Theorem 5.2. The Fourier multiplier bound

$$\|\partial_x^\alpha f\|_{H^s} \leq \|f\|_{H^{s+|\alpha|}}$$

and the rectangle  $(n, m + |\alpha|)$  give (92)–(96). Apply the same multiplier bound to the pressure completion in Proposition 4.9 to obtain (94) and (97). Proposition 3.4 supplies the stated identities.  $\square$

**Theorem 5.5** (Compatible whole-terminal Sobolev traces). *Assume the hypotheses of Theorem 5.2. For  $N, m \in \mathbb{N}_0$ , define*

$$\mathfrak{E}_{N,m} := (1 + T_*^{-1})^{1/2} (\mathfrak{R}_{N,m}^*[u_0, \nu, T_*])^{1/2}. \quad (98)$$

*If  $0 \leq n \leq N$  and  $\alpha \in \mathbb{N}_0^3$ , then*

$$V_n \in C([0, T_*]; H_\sigma^m(\mathbb{R}^3)), \quad \sup_{0 \leq t \leq T_*} \|V_n(t)\|_{H^m} \leq \mathfrak{E}_{N,m}, \quad (99)$$

$$J_{n,\alpha} \in C([0, T_*]; H^m(\mathbb{R}^3)), \quad \sup_{0 \leq t \leq T_*} \|J_{n,\alpha}(t)\|_{H^m} \leq \mathfrak{E}_{N,m+|\alpha|}. \quad (100)$$

*There are unique values*

$$V_n^* := \lim_{t \uparrow T_*} V_n(t) \in H^\infty(\mathbb{R}^3), \quad (101)$$

$$J_{n,\alpha}^* := \lim_{t \uparrow T_*} J_{n,\alpha}(t) = \partial_x^\alpha V_n^* \in H^\infty(\mathbb{R}^3), \quad (102)$$

*such that, for  $k \geq m$ , the image of the  $H^k$  trace under  $H^k \hookrightarrow H^m$  is the  $H^m$  trace. Quantitatively,*

$$\|V_n^*\|_{H^m} \leq \mathfrak{E}_{n,m}, \quad (103)$$

$$\|V_n(t) - V_n^*\|_{H^m} \leq (T_* - t) \mathfrak{E}_{n+1,m}, \quad 0 \leq t \leq T_*, \quad (104)$$

$$\|J_{n,\alpha}^*\|_{H^m} \leq \mathfrak{E}_{n,m+|\alpha|}, \quad (105)$$

$$\|J_{n,\alpha}(t) - J_{n,\alpha}^*\|_{H^m} \leq (T_* - t) \mathfrak{E}_{n+1,m+|\alpha|}. \quad (106)$$

*In particular, the zeroth trace agrees with the common endpoint in Proposition 5.3, and*

$$u_* = V_0^* \in H_\sigma^\infty(\mathbb{R}^3), \quad u \in \bigcap_{a,m \in \mathbb{N}_0} C^a([0, T_*]; H_\sigma^m(\mathbb{R}^3)), \quad \partial_t^a u(T_*) = V_a^*. \quad (107)$$

*Proof.* For  $0 \leq n \leq N$ , Theorem 4.6 gives

$$V_n \in L^2(0, T_*; H^{m+1}), \quad \partial_t V_n = V_{n+1} \in L^2(0, T_*; H^{m-1}).$$

Lemma 2.7 gives the first membership in (99) and

$$\begin{aligned} \sup_{0 \leq t \leq T_*} \|V_n(t)\|_{H^m}^2 &\leq T_*^{-1} \|V_n\|_{L^2 H^m}^2 + 2 \|V_n\|_{L^2 H^{m+1}} \|V_{n+1}\|_{L^2 H^{m-1}} \\ &\leq (1 + T_*^{-1}) \mathfrak{R}_{N,m}^*[u_0, \nu, T_*]. \end{aligned}$$

This proves the bound in (99). Apply the same lemma to  $J_{n,\alpha}$ , using (96), to prove (100).

For  $k \geq m$ , the  $H^k$ -continuous and  $H^m$ -continuous representatives equal the same preterminal distribution. After applying  $H^k \hookrightarrow H^m$ , uniqueness of continuous representatives makes them equal on  $[0, T_*]$ . Their endpoint values are compatible and define (101)–(102). Spatial differentiation is continuous between the corresponding finite Sobolev spaces, so  $J_{n,\alpha}^* = \partial_x^\alpha V_n^*$ .

Corollary 5.4 gives  $V_n, V_{n+1} \in L^1(0, T_*; H^m)$  and  $\partial_t V_n = V_{n+1}$  in distributions. The Banach-valued fundamental identity gives

$$V_n(t) - V_n(s) = \int_s^t V_{n+1}(\tau) d\tau \quad \text{in } H^m, \quad 0 \leq s \leq t \leq T_*. \quad (108)$$

The continuous representative of  $V_{n+1}$  is bounded by  $\mathfrak{E}_{n+1,m}$ . Set  $t = T_*$  in (108) to prove (104); spatial differentiation gives (106). The remaining size estimates follow from (99)–(100).

The divergence map  $H^{m+1} \rightarrow H^m$  is continuous, so  $\nabla \cdot u_* = 0$ . For each  $a, m$ , the curves  $V_0, \dots, V_{a+1}$  are continuous in  $H^m$ , and (108) identifies the derivative of  $V_j$  with  $V_{j+1}$ . Induction in  $j$  proves (107).  $\square$

**Theorem 5.6** (Pressure-complete endpoint jet). *Assume the hypotheses of Theorem 5.2. For  $n \in \mathbb{N}_0$ , define*

$$Q_n^* := R_i R_j \sum_{a=0}^n \binom{n}{a} V_{a,i}^* V_{n-a,j}^*. \quad (109)$$

*Then  $Q_n^* \in H^\infty(\mathbb{R}^3)$ , and, for every  $s \in \mathbb{N}_0$ ,*

$$\|Q_n^*\|_{H^s} \leq 9C_s^{\text{alg}} 2^n \mathfrak{E}_{n,s+1}^2, \quad (110)$$

$$\|Q_n(t) - Q_n^*\|_{H^s} \leq 9C_s^{\text{alg}} 2^{n+1} (T_* - t) \mathfrak{E}_{n+1,s+1}^2. \quad (111)$$

*For  $\alpha \in \mathbb{N}_0^3$ ,*

$$\Pi_{n,\alpha}^* := \partial_x^\alpha Q_n^* = \lim_{t \uparrow T_*} \Pi_{n,\alpha}(t) \quad \text{in } H^m(\mathbb{R}^3) \text{ for every } m \in \mathbb{N}_0. \quad (112)$$

*More precisely, for every  $m \in \mathbb{N}_0$ ,*

$$\|\Pi_{n,\alpha}^*\|_{H^m} \leq 9C_{m+|\alpha|}^{\text{alg}} 2^n \mathfrak{E}_{n,m+|\alpha|+1}^2, \quad (113)$$

$$\|\Pi_{n,\alpha}(t) - \Pi_{n,\alpha}^*\|_{H^m} \leq 9C_{m+|\alpha|}^{\text{alg}} 2^{n+1} (T_* - t) \mathfrak{E}_{n+1,m+|\alpha|+1}^2. \quad (114)$$

*The endpoint rows satisfy, for every  $n, q \in \mathbb{N}_0$ ,*

$$V_{n+1}^* = \nu \Delta V_n^* - \sum_{a=0}^n \binom{n}{a} (V_a^* \cdot \nabla) V_{n-a}^* - \nabla Q_n^* \quad \text{in } H^q(\mathbb{R}^3). \quad (115)$$

*Among families rooted at  $u_*$  that satisfy (109) and (115) in every finite Sobolev space,  $(V_n^*, Q_n^*)_{n \in \mathbb{N}_0}$  is the unique pressure-normalized endpoint jet. Moreover,*

$$p \in \bigcap_{a,m \in \mathbb{N}_0} C^a([0, T_*]; H^m(\mathbb{R}^3)), \quad \partial_t^a p(T_*) = Q_a^*. \quad (116)$$

*Proof.* Lemmas 2.4 and 2.5, followed by (99), give

$$\begin{aligned} \|Q_n^*\|_{H^s} &\leq 9C_s^{\text{alg}} \sum_{a=0}^n \binom{n}{a} \|V_a^*\|_{H^{s+1}} \|V_{n-a}^*\|_{H^{s+1}} \\ &\leq 9C_s^{\text{alg}} 2^n \mathfrak{E}_{n,s+1}^2. \end{aligned}$$

For each product in  $Q_n(t) - Q_n^*$ , insert  $V_a^* V_{n-a}(t)$ , use (104), and bound every retained factor by  $\mathfrak{E}_{n+1,s+1}$ . The two product differences and  $\sum_{a=0}^n \binom{n}{a} = 2^n$  prove (111). The Fourier multiplier inequality

$$\|\partial_x^\alpha f\|_{H^m} \leq \|f\|_{H^{m+|\alpha|}}$$

applied to (110) and (111) proves (113) and (114). The latter tends to zero as  $t \uparrow T_*$ , which proves (112) in every  $H^m$ .

Fix  $n, q$ . The trace of  $V_n$  in  $H^{q+2}$  carries the diffusion term in Proposition 3.4 to  $H^q$  at  $T_*$ . The product estimate at order  $q+1$ , using velocity convergence in  $H^{q+2}$ , carries every transport term to  $H^q$ . Equation (111) at order  $q+1$  carries the pressure gradient to  $H^q$ . The left side converges to  $V_{n+1}^*$  in  $H^q$ , which proves (115).

The value  $V_0^* = u_*$  fixes  $Q_0^*$ , and the velocity recurrence fixes  $V_1^*$ . Inductively,  $V_0^*, \dots, V_n^*$  fix  $Q_n^*$ , which fixes  $V_{n+1}^*$ . This proves uniqueness within the displayed recurrence class. Extend each

$Q_n$  to  $T_*$  by  $Q_n(T_*) := Q_n^*$ . On  $(0, T_*)$ ,  $Q_n = \partial_t^n p$ , so  $\partial_t Q_n = Q_{n+1}$  in distributions. Both rows are continuous in every  $H^m$  by (111). Hence, for  $0 \leq s \leq t \leq T_*$ ,

$$Q_n(t) - Q_n(s) = \int_s^t Q_{n+1}(\tau) d\tau \quad \text{in } H^m(\mathbb{R}^3). \quad (117)$$

Indeed, the identity holds for  $t < T_*$ , and the endpoint bounds permit passage to  $t = T_*$ . Thus  $Q_n \in C^1([0, T_*]; H^m)$  with derivative  $Q_{n+1}$ . Induction gives (116).  $\square$

**Lemma 5.7** (Late-slice overlap uniqueness). *Assume the hypotheses of Theorem 5.2. Define*

$$\delta_0 := \min\{1, c_\nu(1 + \mathfrak{E}_{0,3})^{-2}\} > 0, \quad (118)$$

where  $c_\nu$  is the constant in Theorem 2.11. For every  $t_0 \in (\max\{0, T_* - \delta_0/2\}, T_*)$ , let

$$(w^{t_0}, q^{t_0}) \in \mathcal{M}_\nu^3(t_0, t_0 + \delta_0; u(t_0)), \quad (w^*, q^*) \in \mathcal{M}_\nu^3(T_*, T_* + \delta_0; u_*)$$

be the unique pairs supplied by Theorem 2.11. Then

$$(w^{t_0}, q^{t_0}) = (u, p) \quad \text{on } [t_0, T_*], \quad (119)$$

$$(w^{t_0}, q^{t_0}) = (w^*, q^*) \quad \text{on } [T_*, t_0 + \delta_0]. \quad (120)$$

*Proof.* Equation (99) gives

$$\|u(t_0)\|_{H^3} \leq \mathfrak{E}_{0,3}, \quad \|u_*\|_{H^3} \leq \mathfrak{E}_{0,3}.$$

Hence

$$\delta_0 \leq \delta_\nu(u(t_0)), \quad \delta_0 \leq \delta_\nu(u_*),$$

with  $\delta_\nu$  as in Theorem 2.11. Theorem 2.11 gives both pairs on the stated intervals. For each  $b < T_*$ , Lemma 2.10 places  $(u, p)|_{[t_0, b]}$  and  $(w^{t_0}, q^{t_0})|_{[t_0, b]}$  in  $\mathcal{M}_\nu^3(t_0, b; u(t_0))$ . Uniqueness in that class gives (119). Continuity in  $H^3$  gives

$$w^{t_0}(T_*) = \lim_{t \uparrow T_*} u(t) = u_*.$$

Lemma 2.10 places the restrictions of  $(w^{t_0}, q^{t_0})$  and  $(w^*, q^*)$  to  $[T_*, t_0 + \delta_0]$  in  $\mathcal{M}_\nu^3(T_*, t_0 + \delta_0; u_*)$ . Uniqueness in that class gives (120). The right overlap has positive length because  $t_0 + \delta_0 > T_* + \delta_0/2$ .  $\square$

**Theorem 5.8** (Same-history endpoint restart). *Assume the hypotheses of Theorem 5.2. There is a pressure-normalized classical solution  $(\tilde{u}, \tilde{p})$  on  $[0, T_* + \delta_0/2] \times \mathbb{R}^3$  such that*

$$(\tilde{u}, \tilde{p})|_{[0, T_*] \times \mathbb{R}^3} = (u, p), \quad (121)$$

$$\tilde{u}, \tilde{p} \in C^\infty([0, T_* + \delta_0/2] \times \mathbb{R}^3), \quad (122)$$

$$\partial_t^n \tilde{u}(T_*) = V_n^*, \quad \partial_t^n \tilde{p}(T_*) = Q_n^* \quad (n \in \mathbb{N}_0). \quad (123)$$

If  $0 < \eta \leq \delta_0/2$  and  $(\hat{u}, \hat{p})$  is another pressure-normalized classical extension satisfying

$$(\hat{u}, \hat{p})|_{[0, T_*]} = (u, p), \quad (\hat{u}, \hat{p})|_{[T_*, T_* + \eta]} \in \mathcal{M}_\nu^3(T_*, T_* + \eta; u_*), \quad (124)$$

then

$$(\hat{u}, \hat{p}) = (\tilde{u}, \tilde{p}) \quad \text{on } [0, T_* + \eta]. \quad (125)$$

*Proof.* Let  $(w^*, q^*)$  be the endpoint solution from Lemma 5.7, and set

$$W_n^+ := \partial_t^n w^*(T_*), \quad P_n^+ := \partial_t^n q^*(T_*).$$

Theorem 2.11 makes  $(w^*, q^*)$  smooth and pressure-normalized. Differentiating its equations exactly as in the proof of Proposition 3.4 gives

$$\begin{aligned} P_n^+ &= R_i R_j \sum_{a=0}^n \binom{n}{a} W_{a,i}^+ W_{n-a,j}^+, \\ W_{n+1}^+ &= \nu \Delta W_n^+ - \sum_{a=0}^n \binom{n}{a} (W_a^+ \cdot \nabla) W_{n-a}^+ - \nabla P_n^+. \end{aligned}$$

Since  $W_0^+ = u_* = V_0^*$ , induction with (109) and (115) gives

$$W_n^+ = V_n^*, \quad P_n^+ = Q_n^*, \quad n \in \mathbb{N}_0. \quad (126)$$

Define

$$(\tilde{u}, \tilde{p})(t) := \begin{cases} (u, p)(t), & 0 \leq t < T_*, \\ (w^*, q^*)(t), & T_* \leq t \leq T_* + \delta_0/2. \end{cases}$$

Equations (107), (116), and (126) match every one-sided time derivative in every finite Sobolev space. Sobolev embedding gives (122), and the pressure-complete equation holds at  $T_*$  by (115). Lemma 5.7 identifies the joined pair near  $T_*$  with the single local history  $(w^{t_0}, q^{t_0})$ , so the gluing is one local classical history across the endpoint; the piecewise definition gives (121). Any pair in (124) has the same datum  $u_*$  on the right of  $T_*$  and belongs to the same shifted mild class as  $(w^*, q^*)$ . The at-most-one assertion in Theorem 2.11 proves (125).  $\square$

**Corollary 5.9** (Maximal-time contradiction). *The maximal pressure-complete classical history generated by  $(u_0, \nu)$  cannot have  $T_* < \infty$ .*

*Proof.* If  $T_* < \infty$ , Theorem 4.6 supplies the whole-terminal rectangles. The chain from Theorem 5.2 through Theorem 5.8 produces a pressure-normalized classical extension to  $[0, T_* + \delta_0/2]$ , where  $\delta_0 > 0$ . This contradicts maximality in Definition 3.2.  $\square$

**Theorem 5.10** (Global smoothness on  $\mathbb{R}^3$ ). *For every  $\nu > 0$  and every  $u_0 \in H_\sigma^\infty(\mathbb{R}^3)$ , there exists a pressure-normalized pair*

$$(u, p) \in C^\infty([0, \infty) \times \mathbb{R}^3; \mathbb{R}^3 \times \mathbb{R}) \quad (127)$$

*satisfying*

$$\partial_t u + (u \cdot \nabla) u + \nabla p = \nu \Delta u, \quad (128)$$

$$\nabla \cdot u = 0, \quad u(0) = u_0, \quad (129)$$

$$p(t) = R_i R_j (u_i(t) u_j(t)), \quad \lim_{|x| \rightarrow \infty} p(t, x) = 0, \quad (130)$$

$$\frac{1}{2} \|u(t)\|_{L^2}^2 + \nu \int_0^t \|\nabla u(s)\|_{L^2}^2 ds = \frac{1}{2} \|u_0\|_{L^2}^2, \quad t \geq 0. \quad (131)$$

*In particular,*

$$\sup_{t \geq 0} \|u(t)\|_{L^2(\mathbb{R}^3)}^2 \leq \|u_0\|_{L^2(\mathbb{R}^3)}^2 < \infty. \quad (132)$$

*For every finite  $T > 0$ , its restriction is the unique member of  $\mathcal{M}_\nu^3(0, T; u_0)$ .*

*Proof.* Let

$$\mathcal{T} := \left\{ \tau > 0 : \begin{array}{l} \text{there exists a pressure-complete classical history} \\ (\tau, u^\tau, p^\tau) \text{ generated by } (u_0, \nu) \end{array} \right\},$$

and put

$$T_* := \sup \mathcal{T}.$$

Theorem 2.11, applied to  $u_0 \in H_\sigma^\infty$ , supplies a shifted mild solution whose persistence statement gives every finite velocity and pressure row. Sobolev embedding and the displayed recurrences make this pair a pressure-complete classical history, so  $\mathcal{T}$  is nonempty. Lemma 2.10 admits each strict classical restriction to the shifted mild class, whose at-most-one property identifies any two such solutions on their common interval; their union defines a unique solution on  $[0, T_*)$ . Since  $u_0 \in H_\sigma^\infty$ , the persistence conclusion of Theorem 2.11, applied on overlapping local intervals, places every strict restriction in the class of Definition 3.2. An extension beyond  $T_*$  would restrict to an interval  $[0, \tau]$  with  $\tau > T_*$ , contrary to the definition of the supremum. Thus  $(T_*, u, p)$  is maximal. Corollary 5.9 gives  $T_* = \infty$ .

Fix  $T < \infty$ . Persistence in Theorem 2.11, iterated along the unique history, retains every finite spatial Sobolev order on  $[0, T]$ . Proposition 3.4 retains every finite temporal row. Lemma 2.2 gives (127); arbitrariness of  $T$  gives the global class. The pressure belongs to  $H^s(\mathbb{R}^3)$  for every  $s$ , and the Fourier inversion proof of Lemma 2.2 places its continuous representative in  $C_0(\mathbb{R}^3)$ , which proves the limit in (130). Lemma 2.10 places  $(u, p)|_{[0, T]}$  in  $\mathcal{M}_\nu^3(0, T; u_0)$ . If  $(\tilde{u}, \tilde{p}) \in \mathcal{M}_\nu^3(0, T; u_0)$ , the at-most-one assertion in Theorem 2.11 gives  $(\tilde{u}, \tilde{p}) = (u, p)$  on  $[0, T]$ . This proves the stated uniqueness.

Proposition 4.1 holds on every finite interval. Since  $T$  was arbitrary, it gives (131) for every  $t \geq 0$ ; (132) follows by discarding the nonnegative dissipation term.  $\square$

## 6 Matrix forms of the completed intersection

The compatible intersection, its endpoint jet, and the same-history restart have now been established. The matrices in this section reorganize those completed rows. The whole-terminal bounds and their compatibility have already been proved before any matrix is introduced.

**Proposition 6.1** (Mixed-index transport matrix and pressure completion). *Let*

$$\Gamma := \mathbb{N}_0 \times \mathbb{N}_0^3, \quad \kappa = (n, \alpha), \quad \lambda = (a, \beta),$$

and write

$$\lambda \preceq \kappa \iff a \leq n \text{ and } \beta \leq \alpha, \quad \kappa - \lambda := (n - a, \alpha - \beta), \quad \binom{\kappa}{\lambda} := \binom{n}{a} \binom{\alpha}{\beta}.$$

Set

$$J_\kappa := J_{n, \alpha}, \quad \Pi_\kappa := \Pi_{n, \alpha}, \quad \mathbf{J} := (J_\lambda)_{\lambda \in \Gamma}.$$

The block operator

$$\mathbb{L}(J)_{\kappa, \lambda} := \begin{cases} \binom{\kappa}{\lambda} \mathbb{P}[(J_{\kappa - \lambda} \cdot \nabla) \cdot], & \lambda \preceq \kappa, \\ 0, & \lambda \not\preceq \kappa, \end{cases} \quad (133)$$

has the finite block-row action

$$[\mathbb{L}(J)\mathbf{J}]_\kappa = \sum_{\lambda \preceq \kappa} \binom{\kappa}{\lambda} \mathbb{P}[(J_\lambda \cdot \nabla) J_{\kappa-\lambda}]. \quad (134)$$

Consequently the complete mixed row is the pair

$$J_{n+1,\alpha} + [\mathbb{L}(J)\mathbf{J}]_\kappa = \nu \Delta J_\kappa, \quad (135)$$

$$\Pi_\kappa = R_i R_j \sum_{\lambda \preceq \kappa} \binom{\kappa}{\lambda} J_{\lambda,i} J_{\kappa-\lambda,j}, \quad (136)$$

while the unprojected velocity identity retains  $\nabla \Pi_\kappa$  exactly as in (47).

Order  $\Gamma$  by increasing  $n + |\alpha|$ , with any fixed ordering among indices of equal depth. Then  $\mathbb{L}(J)$  is block lower triangular. For

$$\kappa! := n! \alpha!, \quad \alpha! := \alpha_1! \alpha_2! \alpha_3!, \quad \widehat{J}_\kappa := \frac{J_\kappa}{\kappa!}, \quad \widehat{\Pi}_\kappa := \frac{\Pi_\kappa}{\kappa!},$$

let  $D$  be the diagonal matrix with  $D_{\kappa,\kappa} = \kappa!$ . On every finite downward-closed truncation of  $\Gamma$ ,

$$D^{-1} \mathbb{L}(D\widehat{J})D = \widehat{\mathbb{L}}(\widehat{J}), \quad \widehat{\mathbb{L}}(\widehat{J})_{\kappa,\lambda} := \begin{cases} \mathbb{P}[(\widehat{J}_{\kappa-\lambda} \cdot \nabla) \cdot], & \lambda \preceq \kappa, \\ 0, & \lambda \not\preceq \kappa. \end{cases} \quad (137)$$

Thus the factorial-normalized pressure-complete recurrence is

$$(n+1)\widehat{J}_{n+1,\alpha} + \sum_{\lambda \preceq \kappa} (\widehat{J}_\lambda \cdot \nabla) \widehat{J}_{\kappa-\lambda} + \nabla \widehat{\Pi}_\kappa = \nu \Delta \widehat{J}_\kappa, \quad (138)$$

$$\widehat{\Pi}_\kappa = R_i R_j \sum_{\lambda \preceq \kappa} \widehat{J}_{\lambda,i} \widehat{J}_{\kappa-\lambda,j}. \quad (139)$$

The normalized transport blocks depend on the difference  $\kappa - \lambda$ ; hence (137) is the block lower-triangular convolution form of the mixed transport array.

*Proof.* For  $\lambda \preceq \kappa$ , change the summation index from  $\lambda$  to  $\kappa - \lambda$  and use  $\binom{\kappa}{\kappa-\lambda} = \binom{\kappa}{\lambda}$ . This gives (134). Apply  $\mathbb{P}$  to (47). The identities

$$\mathbb{P}J_{n+1,\alpha} = J_{n+1,\alpha}, \quad \mathbb{P}\Delta J_\kappa = \Delta J_\kappa, \quad \mathbb{P}\nabla \Pi_\kappa = 0$$

follow from (48) and the Fourier symbols. They prove (135); equation (136) is (46) in mixed-index notation. The pressure formula also shows that projection of the velocity row loses no pressure coordinate.

If  $\lambda \preceq \kappa$ , then  $a + |\beta| \leq n + |\alpha|$ , which proves lower triangularity in the stated ordering. Finally,

$$\binom{\kappa}{\lambda} \lambda! (\kappa - \lambda)! = \kappa!.$$

Dividing the mixed velocity and pressure rows by  $\kappa!$  proves (138) and (139). The same factorial cancellation in each operator block proves (137). Every row has  $(n+1) \prod_{j=1}^3 (\alpha_j + 1)$  supported blocks, so all displayed row sums are finite.  $\square$

**Proposition 6.2** (Sobolev rectangle matrix and adjacent stencil). *For  $0 < T < T_*$ ,  $N, M \in \mathbb{N}_0$ ,  $0 \leq n \leq N$ , and  $m \in \mathbb{Z}$ , set*

$$G_{n,m}(T) := \|V_n\|_{L^2(0,T;H^m)},$$

and set

$$P_{n,M}(T) := \|Q_n\|_{L^1(0,T;H^M)} + \|\nabla Q_n\|_{L^1(0,T;H^{M-1})}.$$

The object matrix of the finite pressure-complete rectangle is

$$\mathbf{R}_{N,M}[u_0; T] := \begin{pmatrix} V_0|_{(0,T)} & V_1|_{(0,T)} & Q_0|_{(0,T)} \\ V_1|_{(0,T)} & V_2|_{(0,T)} & Q_1|_{(0,T)} \\ \vdots & \vdots & \vdots \\ V_N|_{(0,T)} & V_{N+1}|_{(0,T)} & Q_N|_{(0,T)} \end{pmatrix}. \quad (140)$$

The corresponding matrix of the three column norms is

$$\mathbf{C}_{N,M}(T) := \begin{pmatrix} G_{0,M+1}(T) & G_{1,M-1}(T) & P_{0,M}(T) \\ G_{1,M+1}(T) & G_{2,M-1}(T) & P_{1,M}(T) \\ \vdots & \vdots & \vdots \\ G_{N,M+1}(T) & G_{N+1,M-1}(T) & P_{N,M}(T) \end{pmatrix}. \quad (141)$$

If  $\mathbf{G}_{N,M}(T)$  denotes its first two columns, then

$$\|\mathbf{G}_{N,M}(T)\|_{\mathbf{F}}^2 = \mathfrak{R}_{N,M}(u; T), \quad \sum_{n=0}^N P_{n,M}(T) = \mathfrak{Q}_{N,M}(p; T). \quad (142)$$

For each  $0 \leq n \leq N$ , the first two entries in row  $n$  form the adjacent Hilbert-scale stencil

$$\left. \begin{array}{l} (n, M+1) : \quad V_n \in L^2(0, T; H^{M+1}), \\ (n+1, M-1) : \quad V_{n+1} = \partial_t V_n \in L^2(0, T; H^{M-1}) \end{array} \right\} \longrightarrow (n, M) : \quad V_n \in C([0, T]; H^M). \quad (143)$$

For every  $m \in \mathbb{N}_0$ , the Sobolev matrix measures the mixed-index array through the exact identity

$$G_{n,m}(T)^2 = \sum_{\substack{\alpha \in \mathbb{N}_0^3 \\ |\alpha| \leq m}} \frac{m!}{(m-|\alpha|)! \alpha!} \|J_{n,\alpha}\|_{L^2((0,T) \times \mathbb{R}^3)}^2. \quad (144)$$

The pressure entry in row  $n$  is  $Q_n = \Pi_{n,0}$ , and its spatial rows  $\partial_x^\alpha Q_n = \Pi_{n,\alpha}$  are fixed by (136).

*Proof.* The object matrix is the tuple in Definition 4.5, arranged by temporal row. The two identities in (142) are (69) and (70). Lemma 2.7 gives (143). Finally, Plancherel and

$$(1 + |\xi|^2)^m = \sum_{\substack{\alpha \in \mathbb{N}_0^3 \\ |\alpha| \leq m}} \frac{m!}{(m-|\alpha|)! \alpha!} \xi^{2\alpha}$$

give (144). The pressure claims follow from the definitions of  $Q_n, \Pi_{n,\alpha}$  and (136).  $\square$

**Proposition 6.3** (Whole-terminal closed estimate matrix). *Under the hypotheses of Theorem 4.6, fix  $N, M \in \mathbb{N}_0$  and set*

$$\begin{aligned} a_{n,M} &:= \|V_n\|_{L^2(0,T_*;H^{M+1})}, \\ b_{n,M} &:= \|V_{n+1}\|_{L^2(0,T_*;H^{M-1})}, \quad 0 \leq n \leq N. \end{aligned} \quad (145)$$

Define entrywise order on nonnegative matrices by  $A \preceq_{\text{ent}} B$  when  $A_{ij} \leq B_{ij}$  for every entry. Then, for every  $0 < T < T_*$ ,

$$\mathbf{G}_{N,M}(T) \preceq_{\text{ent}} \mathbf{G}_{N,M}^* := \begin{pmatrix} a_{0,M} & b_{0,M} \\ a_{1,M} & b_{1,M} \\ \vdots & \vdots \\ a_{N,M} & b_{N,M} \end{pmatrix}. \quad (146)$$

Moreover,

$$\begin{aligned} \mathfrak{R}_{N,M}(u; T) &= \|\mathbf{G}_{N,M}(T)\|_{\text{F}}^2 \\ &\leq \|\mathbf{G}_{N,M}^*\|_{\text{F}}^2 = \sum_{n=0}^N (a_{n,M}^2 + b_{n,M}^2) = \mathfrak{R}_{N,M}^*[u_0, \nu, T_*] < \infty. \end{aligned} \quad (147)$$

*Proof.* Proposition 4.7 makes every  $a_{n,M}$  and  $b_{n,M}$  finite. Restriction from  $(0, T_*)$  to  $(0, T)$  decreases each nonnegative coordinate norm and gives (146). Taking Frobenius norms and using (142) together with (73) proves (147).  $\square$

## 7 Quantitative consequences of the compatible rectangles

### 7.1 Completion in the global $H^1$ mild class

The construction begins with every datum in  $H_\sigma^\infty(\mathbb{R}^3)$ , not only with Schwartz data. Parabolic smoothing now transfers that result to the full strong  $H^1$  class.

**Theorem 7.1** (Global  $H^1$  completion). *Let  $\nu > 0$  and  $a \in H_\sigma^1(\mathbb{R}^3)$ . There is a unique pressure-normalized mild solution*

$$u^a \in C([0, \infty); H_\sigma^1) \cap L_{\text{loc}}^2([0, \infty); H_\sigma^2), \quad u^a \in C^\infty((0, \infty) \times \mathbb{R}^3), \quad (148)$$

*of the three-dimensional Navier–Stokes equations with datum  $a$  and viscosity  $\nu$ . Its pressure is*

$$p^a = R_i R_j (u_i^a u_j^a)$$

*in distributions. The solution obeys the kinetic-energy identity on every finite interval.*

*Proof.* Write

$$Y(t) := \|\nabla u(t)\|_2^2, \quad Z(t) := \|\Delta u(t)\|_2^2, \quad c_1 := \frac{27}{16} S(6)^6.$$

For a smooth solenoidal approximation, the  $L^2$  pairing of the projected equation with  $-\Delta u$  gives

$$\frac{1}{2} Y'(t) + \nu Z(t) = \text{Re} \langle (u \cdot \nabla) u, \Delta u \rangle_{L^2}. \quad (149)$$

The explicit three-dimensional Sobolev inequality and interpolation give

$$\|u\|_6 \leq S(6)Y^{1/2}, \quad \|\nabla u\|_3 \leq S(6)^{1/2}Y^{1/4}Z^{1/4}.$$

Consequently,

$$|\langle (u \cdot \nabla)u, \Delta u \rangle_{L^2}| \leq S(6)^{3/2}Y^{3/4}Z^{3/4} \leq \frac{\nu}{2}Z + \frac{27S(6)^6}{32\nu^3}Y^3,$$

and hence

$$Y'(t) + \nu Z(t) \leq c_1 \nu^{-3} Y(t)^3. \quad (150)$$

Comparison with  $y' = c_1 \nu^{-3} y^3$  shows that

$$Y(t) \leq \frac{Y(0)}{(1 - 2c_1 \nu^{-3} Y(0)^2 t)^{1/2}} \quad (151)$$

while the denominator is positive.

The estimate holds on the explicit lifespan

$$\begin{aligned} \tau_\nu(a) &:= \min \left\{ 1, \frac{\nu^3}{4c_1(1 + \|\nabla a\|_2^2)^2} \right\}, \\ \sup_{0 \leq t \leq \tau_\nu(a)} Y(t) &\leq \sqrt{2}Y(0), \quad \int_0^{\tau_\nu(a)} Z(t) dt < \infty. \end{aligned} \quad (152)$$

Let  $\mathcal{J}_k$  be the orthogonal Fourier projection to  $\{|\xi| \leq k\}$ . On the range of  $\mathcal{J}_k$ , the regularized problem

$$\partial_t u^{(k)} - \nu \Delta u^{(k)} = -\mathcal{J}_k \mathbb{P} \nabla \cdot (u^{(k)} \otimes u^{(k)}), \quad u^{(k)}(0) = \mathcal{J}_k a, \quad (153)$$

is a Banach-space ordinary differential equation in  $H_\sigma^1$ . Indeed, if

$$\mathcal{N}_k(z) := \mathcal{J}_k \mathbb{P} \nabla \cdot (z \otimes z),$$

then the Bernstein inequalities on the range of  $\mathcal{J}_k$  give

$$\|\mathcal{N}_k(z) - \mathcal{N}_k(y)\|_{H^1} \leq C_k (\|z\|_{H^1} + \|y\|_{H^1}) \|z - y\|_{H^1}, \quad C_k < \infty. \quad (154)$$

Picard's theorem supplies a maximal trajectory. The projection is self-adjoint and commutes with  $\mathbb{P}$ ,  $\nabla$ , and  $\Delta$ , so the kinetic identity holds along that trajectory. Since

$$\|z\|_{H^1}^2 \leq (1 + k^2) \|z\|_2^2 \quad (z = \mathcal{J}_k z), \quad (155)$$

the kinetic identity and the Banach-space continuation criterion extend the trajectory for all positive times. Estimate (150) holds for  $u^{(k)}$ , with constants independent of  $k$ . Equations (151)–(152) give

$$\sup_k \left( \|u^{(k)}\|_{L^\infty(0, \tau_\nu(a); H^1)} + \|u^{(k)}\|_{L^2(0, \tau_\nu(a); H^2)} \right) < \infty. \quad (156)$$

The nonlinear estimate

$$\|\mathbb{P} \nabla \cdot (u \otimes u)\|_2 \leq \|u\|_6 \|\nabla u\|_3 \leq S(6)^{3/2}Y^{3/4}Z^{1/4},$$

shows that the regularized nonlinear terms are bounded in  $L^2(0, \tau_\nu(a); L^2)$ , uniformly in  $k$ .

The equations bound  $\partial_t u^{(k)}$  in the same space. Weak compactness and the Aubin–Lions–Simon lemma [1, 13] on every ball give, after diagonal extraction,

$$\begin{aligned} u^{(k)} &\overset{*}{\rightharpoonup} u && \text{in } L^\infty(0, \tau_\nu(a); H^1), \\ u^{(k)} &\rightharpoonup u && \text{in } L^2(0, \tau_\nu(a); H^2), \\ u^{(k)} &\longrightarrow u && \text{in } L^2(0, \tau_\nu(a); H^1(B_R)) \quad (R < \infty). \end{aligned}$$

The last convergence passes the quadratic term in distributions; lower semicontinuity preserves (156). The displayed nonlinear estimate now gives  $\mathbb{P}\nabla \cdot (u \otimes u) \in L^2 L^2$ , and the equation gives  $u_t \in L^2(0, \tau_\nu(a); L^2)$ . For every  $\phi \in C_c^\infty(\mathbb{R}^3; \mathbb{R}^3)$  with  $\nabla \cdot \phi = 0$  and every  $\eta \in C_c^\infty([0, \tau_\nu(a)))$ , passage to the limit in the time-integrated regularized equation gives

$$\begin{aligned} - \int_0^{\tau_\nu(a)} \langle u, \phi \rangle_{L^2} \eta' dt + \nu \int_0^{\tau_\nu(a)} \langle \nabla u, \nabla \phi \rangle_{L^2} \eta dt - \int_0^{\tau_\nu(a)} \langle u \otimes u, \nabla \phi \rangle_{L^2} \eta dt \\ = \langle a, \phi \rangle_{L^2} \eta(0). \end{aligned} \quad (157)$$

The right-hand side fixes the weak initial trace as  $a$ .

The Lions–Magenes theorem [11] for  $H_\sigma^2 \hookrightarrow H_\sigma^1 \hookrightarrow L_\sigma^2$  gives  $u \in C([0, \tau_\nu(a)]; H_\sigma^1)$ ; its continuous trace at zero is the trace in (157), hence  $u(0) = a$ . Variation of constants gives the mild equation, and  $p = R_i R_j (u_i u_j)$  supplies its normalized pressure. Since  $u_t \in L^2 L^2$  and  $u \in L^2 H^2$ , pairing the limiting equation with  $u$  is legitimate; incompressibility cancels the transport term and proves the kinetic-energy identity.

If  $u, v$  are two such solutions and  $w = u - v$ , incompressibility cancels the  $(u \cdot \nabla)w$  pairing and gives

$$\frac{1}{2} \frac{d}{dt} \|w\|_2^2 + \nu \|\nabla w\|_2^2 = -\operatorname{Re} \langle (w \cdot \nabla)v, w \rangle_{L^2}.$$

The same Sobolev constant yields

$$|\langle (w \cdot \nabla)v, w \rangle_{L^2}| \leq S(6)^{3/2} \|\nabla v\|_2 \|w\|_2^{1/2} \|\nabla w\|_2^{3/2} \leq \frac{\nu}{2} \|\nabla w\|_2^2 + \frac{27S(6)^6}{32\nu^3} \|\nabla v\|_2^4 \|w\|_2^2.$$

Gronwall’s inequality proves uniqueness. The same calculation one derivative higher also gives the finite-interval stability needed below. Indeed, pairing the difference equation with  $-\Delta w$ , integrating the  $(v \cdot \nabla)w$  term once in space, and applying Young’s inequality gives

$$\frac{d}{dt} \|\nabla w\|_2^2 + \nu \|\Delta w\|_2^2 \leq \frac{C_{\text{st}}}{\nu} \left( \|\nabla u\|_3^2 + \|\nabla v\|_3^2 \right) \|\nabla w\|_2^2. \quad (158)$$

The coefficient is integrable because

$$\|\nabla u\|_3^2 \leq S(6) \|\nabla u\|_2 \|\Delta u\|_2,$$

and likewise for  $v$ . The  $L^2$  difference estimate and (158) show continuous dependence in  $C([0, S]; H^1)$  on every finite interval  $[0, S]$ .

Let  $c_0 > 0$  be the absolute smallness constant in the full-space  $H^1$  local existence theorem, Theorem 5.4(ii) of [14], written at viscosity one as  $\|b\|_{H^1}^4 h \leq c_0$ . After viscosity rescaling, its condition is

$$h \|b\|_{H^1}^4 \leq c_0 \nu^3. \quad (159)$$

Fix  $0 < \varepsilon < S \leq \tau_\nu(a)$ , and put  $K_S := 1 + \sup_{0 \leq t \leq S} \|u(t)\|_{H^1}$ . Choose

$$0 < h \leq \min \{ \varepsilon, c_0 \nu^3 K_S^{-4} \}.$$

The intervals  $[0, h]$  and  $[t - h, t]$ ,  $h \leq t \leq S$ , obey (159). The quantitative positive-time regularity estimate in Lemma 5.5 of [14], after viscosity rescaling with interior time  $h/2$ , and the heat-semigroup/product iteration in its proof give, for every  $m \in \mathbb{N}$ ,

$$\sup_{\varepsilon \leq t \leq S} \|u(t)\|_{H^m} \leq C_{m, \varepsilon, S}(\nu, K_S) < \infty. \quad (160)$$

Apply the mild formula from time  $\varepsilon/2$ . The bounds (160), the heat multiplier, and Lemma 2.4 show that  $u \in C([\varepsilon, S]; H^m)$  for every  $m$ . The identities

$$p = R_i R_j (u_i u_j), \quad u_t = \nu \Delta u - \mathbb{P} \nabla \cdot (u \otimes u) \quad (161)$$

then give  $p \in C([\varepsilon, S]; H^m)$  and  $u_t \in C([\varepsilon, S]; H^m)$  for every  $m$ . Differentiating (161) and the pressure identity, and applying the same product estimate, proves inductively that all temporal derivatives of  $u$  and  $p$  belong to every spatial Sobolev space continuously on  $[\varepsilon, S]$ . Sobolev embedding proves  $(u, p) \in C^\infty((0, \tau_\nu(a)] \times \mathbb{R}^3)$ .

Choose  $0 < t_0 < \tau_\nu(a)$ . Equation (160) places  $u(t_0)$  in  $H_\sigma^\infty$ . Theorem 5.10, with time origin  $t_0$ , supplies a global smooth history from that datum. The uniqueness estimate above identifies it with the local history on their overlap, so the two histories glue to (148). The same estimate proves uniqueness in the full displayed class. The kinetic identity on the first interval and the classical identity after  $t_0$  agree at the common trace and prove the final assertion.  $\square$

## 7.2 Canonical coordinates and finite majorants

The compatible rectangles carry more information than the endpoint argument uses. We first name their finite coordinates and close the estimates that will feed the mixed rows, frequency tails, and continuation bounds below.

**Definition 7.2** (Rectangle and trace coordinates). Let  $u_0 \in H_\sigma^\infty(\mathbb{R}^3)$ ,  $\nu > 0$ , and let  $(u, p)$  be the datum-generated pressure-normalized solution. Let  $0 < T < \infty$  satisfy either  $T < T_*$  or  $T = T_* < \infty$ , and set  $I_T = (0, T)$ ,

$$V_n := \partial_t^n u, \quad Q_n := \partial_t^n p, \quad n \in \mathbb{N}_0.$$

Define the monotone rectangle coordinate

$$\mathfrak{P}_{N, M}[u_0, \nu; T] := 1 + \sup_{0 < S < T} \mathfrak{R}_{N, M}(u; S), \quad (162)$$

and the trace constant

$$\mathfrak{T}_{N, M}[u_0, \nu; T] := (1 + T^{-1})^{1/2} \mathfrak{P}_{N, M}[u_0, \nu; T]^{1/2}. \quad (163)$$

All occurrences of  $\mathfrak{P}_{N, M}$  and  $\mathfrak{T}_{N, M}$  below carry the arguments  $[u_0, \nu; T]$ . The datum  $u_0$ , viscosity  $\nu$ , pressure-normalized solution  $(u, p)$ , horizon  $T$ , interval  $I_T$ , and row notation  $V_n, Q_n$  fixed here remain in force throughout the rest of this section. When  $T = T_* < \infty$ , the whole-terminal representatives are those of Proposition 4.7, and every endpoint value used below is the compatible trace supplied by Theorem 5.5.

**Lemma 7.3** (Rectangle-coordinate finiteness). *For every  $N, M \in \mathbb{N}_0$ ,*

$$1 \leq \mathfrak{P}_{N, M}[u_0, \nu; T] < \infty. \quad (164)$$

*Proof.* If  $T = T_* < \infty$ , equation (71) proves (164). If  $T < T_*$ , classical persistence on  $[0, T]$  gives

$$\sup_{0 \leq t \leq T} \|V_n(t)\|_{H^m} < \infty \quad (n, m \in \mathbb{N}_0),$$

and (69) proves (164).  $\square$

**Lemma 7.4** (Fixed analytic constants). *For  $r \in \mathbb{N}_0$ , set*

$$A_r := 2C_r^{\text{alg}}.$$

*Then*

$$\|R_i R_j(f_i g_j)\|_{H^r} + \|\mathbb{P}\nabla \cdot (f \otimes g)\|_{H^{r-1}} \leq A_r \|f\|_{H^{r+1}} \|g\|_{H^{r+1}}. \quad (165)$$

*The same constant satisfies*

$$\|R_i R_j(f_i g_j)\|_{H^r} + \|\nabla R_i R_j(f_i g_j)\|_{H^{r-1}} \leq A_r \|f\|_{H^{r+1}} \|g\|_{H^{r+1}}. \quad (166)$$

*For  $s, k \in \mathbb{N}_0$ ,  $2 \leq b \leq \infty$ , and  $s > k + \frac{3}{2} - \frac{3}{b}$ , define*

$$\nabla^k f := (\partial^\alpha f)_{|\alpha|=k}, \quad \|\nabla^k f\|_{L^b} := \left\| \left( \sum_{|\alpha|=k} |\partial^\alpha f|^2 \right)^{1/2} \right\|_{L^b},$$

*and set*

$$d_k := \#\{\alpha \in \mathbb{N}_0^3 : |\alpha| = k\}, \quad \frac{1}{q_b} := \frac{1}{2} - \frac{1}{b},$$

*with  $q_2 = \infty$ , and*

$$S_{s,k,b} := d_k^{1/2} (2\pi)^{-3(1/2-1/b)} \left\| |\xi|^k \langle \xi \rangle^{-s} \right\|_{L_\xi^{q_b}}. \quad (167)$$

*Then  $S_{s,k,b} < \infty$  and*

$$\|\nabla^k f\|_{L^b} \leq S_{s,k,b} \|f\|_{H^s}. \quad (168)$$

*For  $m \geq 2$ , set  $L_m := \max\{1, C_m^{\text{alg}}\}$ . Then*

$$\|fg\|_{H^m} \leq L_m \|f\|_{H^m} \|g\|_{H^m}, \quad (169)$$

$$\|e^{\nu t \Delta} \mathbb{P}\nabla \cdot F\|_{H^m} \leq L_m (\nu t)^{-1/2} \|F\|_{H^m}, \quad t > 0. \quad (170)$$

*Proof.* Lemmas 2.4 and 2.5, applied componentwise before summing squares, give

$$\begin{aligned} \|R_i R_j(f_i g_j)\|_{H^r} &\leq \|f \otimes g\|_{H^r} \leq C_r^{\text{alg}} \|f\|_{H^{r+1}} \|g\|_{H^{r+1}}, \\ \|\mathbb{P}\nabla \cdot (f \otimes g)\|_{H^{r-1}} &\leq \|f \otimes g\|_{H^r} \leq C_r^{\text{alg}} \|f\|_{H^{r+1}} \|g\|_{H^{r+1}}. \end{aligned}$$

Moreover,

$$\|\nabla R_i R_j(f_i g_j)\|_{H^{r-1}} \leq \|R_i R_j(f_i g_j)\|_{H^r}.$$

This proves (165) and (166).

Let  $b' = b/(b-1)$ . Interpolation between Fourier inversion  $L_\xi^1 \rightarrow L_x^\infty$ , with norm  $(2\pi)^{-3/2}$ , and Plancherel gives

$$\|\mathcal{F}^{-1}h\|_{L_x^b} \leq (2\pi)^{-3(1/2-1/b)} \|h\|_{L_\xi^{b'}}.$$

For  $|\alpha| = k$ , Hölder's inequality with  $1/b' = 1/2 + 1/q_b$  gives

$$\|\partial^\alpha f\|_{L^b} \leq (2\pi)^{-3(1/2-1/b)} \left\| |\xi|^k \langle \xi \rangle^{-s} \right\|_{L_\xi^{q_b}} \|f\|_{H^s}.$$

The weight belongs to  $L^{q_b}$  under the displayed strict inequality for  $s$ ; for  $b = 2$ , its  $L^\infty$ -norm is finite because  $s > k$ . Summation over  $|\alpha| = k$  proves (168).

The first estimate in (169) is Lemma 2.4. For the heat estimate, the symbol of  $\mathbb{P}\nabla\cdot$ , acting from tensors to vectors, has norm at most  $|\xi|$ , while

$$\sup_{\rho \geq 0} \rho e^{-\nu t \rho^2} = (2e\nu t)^{-1/2}.$$

Plancherel gives

$$\|e^{\nu t \Delta} \mathbb{P}\nabla \cdot F\|_{H^m} \leq (2e\nu t)^{-1/2} \|F\|_{H^m} \leq L_m(\nu t)^{-1/2} \|F\|_{H^m}.$$

□

**Definition 7.5** (Canonical rectangle majorants). For  $N, M \in \mathbb{N}_0$ , define

$$\mathfrak{B}_{N,M}[u_0, \nu; T] := \inf \{B \geq 0 : \mathfrak{R}_{N,M}(u; S) \leq B \text{ for every } 0 < S < T\}. \quad (171)$$

Monotone convergence gives

$$\mathfrak{B}_{N,M}[u_0, \nu; T] = \begin{cases} \mathfrak{R}_{N,M}(u; T), & T < T_*, \\ \mathfrak{R}_{N,M}^*[u_0, \nu, T_*], & T = T_* < \infty. \end{cases} \quad (172)$$

The global theorem fixes one history for  $(u_0, \nu)$ . Thus  $\mathfrak{B}_{N,M}[u_0, \nu; T]$  is the exact minimal rectangle bound of that datum-generated history, not a supremum over a family of solutions and not an independently chosen constant. Equations (142) and (171) give the exact matrix identities

$$\mathfrak{B}_{N,M}[u_0, \nu; T] = \sup_{0 < S < T} \|\mathbf{G}_{N,M}(S)\|_{\mathbb{F}}^2, \quad (173)$$

$$\mathfrak{P}_{N,M}[u_0, \nu; T] = 1 + \sup_{0 < S < T} \|\mathbf{G}_{N,M}(S)\|_{\mathbb{F}}^2. \quad (174)$$

Consequently,

$$\mathfrak{P}_{N,M}[u_0, \nu; T] = 1 + \mathfrak{B}_{N,M}[u_0, \nu; T]. \quad (175)$$

For  $n, m \in \mathbb{N}_0$ , define the initial-jet majorants  $\mathfrak{D}_{n,m} = \mathfrak{D}_{n,m}[u_0, \nu]$  by

$$\mathfrak{D}_{0,m} := \|u_0\|_{H^m}, \quad (176)$$

$$\mathfrak{D}_{n+1,m} := \nu \mathfrak{D}_{n,m+2} + A_{m+1} \sum_{a=0}^n \binom{n}{a} \mathfrak{D}_{a,m+2} \mathfrak{D}_{n-a,m+2}. \quad (177)$$

Define the pointwise recursion  $\mathfrak{Y}_{n,m} = \mathfrak{Y}_{n,m}[u_0, \nu; T]$  by

$$\mathfrak{Y}_{0,0} := \|u_0\|_2, \quad (178)$$

$$\mathfrak{Y}_{0,m} := \min \left\{ (1 + T^{-1})^{1/2} \mathfrak{B}_{0,m}^{1/2} \|u_0\|_{H^m} + T^{1/2} \mathfrak{B}_{0,m+1}^{1/2} \right\}, \quad m \geq 1, \quad (179)$$

$$\mathfrak{Y}_{n+1,m} := \nu \mathfrak{Y}_{n,m+2} + A_{m+1} \sum_{a=0}^n \binom{n}{a} \mathfrak{Y}_{a,m+2} \mathfrak{Y}_{n-a,m+2}. \quad (180)$$

For  $m \in \{-1\} \cup \mathbb{N}_0$ , put

$$m_+ := \max\{m, 0\}, \quad \mu(m) := \max\{m-1, 0\},$$

and define

$$\tilde{\mathfrak{X}}_{n,m} := \min \left\{ T^{1/2} \mathfrak{Y}_{n,m_+}, \mathfrak{B}_{n,\mu(m)}^{1/2} \right\}. \quad (181)$$

Set

$$\mathfrak{X}_{0,0} := \min \left\{ \tilde{\mathfrak{X}}_{0,0}, T^{1/2} \|u_0\|_2 \right\}, \quad (182)$$

$$\mathfrak{X}_{0,1} := \min \left\{ \tilde{\mathfrak{X}}_{0,1}, \left( T + \frac{1}{2\nu} \right)^{1/2} \|u_0\|_2 \right\}, \quad (183)$$

$$\mathfrak{X}_{n,m} := \tilde{\mathfrak{X}}_{n,m} \quad \text{for all other } (n, m). \quad (184)$$

For  $n, m \in \mathbb{N}_0$ , define

$$\mathfrak{Z}_{n,m}[u_0, \nu; T] := \min \left\{ \mathfrak{Y}_{n,m}, (1 + T^{-1})^{1/2} \mathfrak{B}_{n,m}^{1/2}, \mathfrak{D}_{n,m} + T^{1/2} \mathfrak{X}_{n+1,m} \right\}. \quad (185)$$

All  $\mathfrak{B}, \mathfrak{D}, \mathfrak{X}, \mathfrak{Y}, \mathfrak{Z}$  in this section carry the arguments  $[u_0, \nu; T]$ , except that  $\mathfrak{D}$  is independent of  $T$ .

**Proposition 7.6** (Exact viscosity and horizon scaling). *Let*

$$a := \nu^{-1} u_0, \quad v(s, x) := \nu^{-1} u(s/\nu, x), \quad q(s, x) := \nu^{-2} p(s/\nu, x).$$

*Then  $(v, q)$  is the unit-viscosity solution generated by  $a$ . If*

$$\tilde{V}_n := \partial_s^n v, \quad \tilde{Q}_n := \partial_s^n q,$$

*then*

$$V_n(t) = \nu^{n+1} \tilde{V}_n(\nu t), \quad Q_n(t) = \nu^{n+2} \tilde{Q}_n(\nu t). \quad (186)$$

*Substitution of (186) into the defining norms gives*

$$\begin{aligned} \mathfrak{B}_{N,M}[u_0, \nu; T] &= \sum_{n=0}^N \left[ \nu^{2n+1} \left\| \tilde{V}_n \right\|_{L^2(0, \nu T; H^{M+1})}^2 \right. \\ &\quad \left. + \nu^{2n+3} \left\| \tilde{V}_{n+1} \right\|_{L^2(0, \nu T; H^{M-1})}^2 \right], \end{aligned} \quad (187)$$

$$\begin{aligned} \|Q_n\|_{L^1(0, T; H^m)} + \|\nabla Q_n\|_{L^1(0, T; H^{m-1})} &= \nu^{n+1} \left( \left\| \tilde{Q}_n \right\|_{L^1(0, \nu T; H^m)} \right. \\ &\quad \left. + \left\| \nabla \tilde{Q}_n \right\|_{L^1(0, \nu T; H^{m-1})} \right), \end{aligned} \quad (188)$$

$$\sup_{0 \leq t \leq T} \|Q_n(t)\|_{H^m} = \nu^{n+2} \sup_{0 \leq s \leq \nu T} \left\| \tilde{Q}_n(s) \right\|_{H^m}. \quad (189)$$

The initial-jet recursion obeys

$$\mathfrak{D}_{n,m}[u_0, \nu] = \nu^{n+1} \mathfrak{D}_{n,m}[u_0/\nu, 1]. \quad (190)$$

Write

$$E_\sigma^u(t) := \frac{1}{2} \|\Lambda^\sigma u(t)\|_2^2, \quad E_\sigma^v(s) := \frac{1}{2} \|\Lambda^\sigma v(s)\|_2^2.$$

For every  $j \in \mathbb{N}_0$ , the critical-height rows obey

$$E_{j+1/2}^u(t) = \nu^2 E_{j+1/2}^v(\nu t), \quad (191)$$

$$\int_0^T E_{j+1/2}^u(t) dt = \nu \int_0^{\nu T} E_{j+1/2}^v(s) ds. \quad (192)$$

For every  $b \in H_\sigma^3(\mathbb{R}^3)$ , the maximal elapsed lifespan in the shifted mild class has the exact covariance

$$\tau_{\nu,3}^{\max}(s, b) = \nu^{-1} \tau_{1,3}^{\max}(0, b/\nu). \quad (193)$$

*Proof.* The change of variables gives  $u(t) = \nu v(\nu t)$  and  $p(t) = \nu^2 q(\nu t)$ . Differentiation proves (186); the substitution  $s = \nu t$  proves (187)–(189). Equation (190) follows by induction in (176)–(177). The amplitude relation  $u(t) = \nu v(\nu t)$  proves (191), and the substitution  $s = \nu t$  proves (192). For  $(w, \pi) \in \mathcal{M}_\nu^3(s, s + \delta; b)$ , set

$$\mathcal{S}_\nu(w, \pi)(\sigma) := (\nu^{-1} w(s + \sigma/\nu), \nu^{-2} \pi(s + \sigma/\nu)), \quad 0 \leq \sigma \leq \nu \delta.$$

Time translation and the same change of variables show that  $\mathcal{S}_\nu$  is a bijection from  $\mathcal{M}_\nu^3(s, s + \delta; b)$  onto  $\mathcal{M}_1^3(0, \nu \delta; b/\nu)$ . Its inverse is the corresponding amplitude and time rescaling, which proves (193).  $\square$

**Proposition 7.7** (Height-coordinate re-expression of the finite rows). *For  $m \in \mathbb{N}_0$ , define*

$$\mathfrak{H}_m[u_0, \nu; T] := \frac{1}{2} \left\| \Lambda^{m+1/2} u \right\|_{L^2(I_T; L^2)}^2, \quad (194)$$

$$\mathfrak{S}_m[u_0, \nu; T]^2 := 2^m \left( T \|u_0\|_2^2 + 2 \mathfrak{H}_m[u_0, \nu; T] \right). \quad (195)$$

Then

$$2 \mathfrak{H}_m[u_0, \nu; T] \leq \mathfrak{B}_{0,m}[u_0, \nu; T], \quad \|u\|_{L^2(I_T; H^m)} \leq \mathfrak{S}_m[u_0, \nu; T]. \quad (196)$$

Define recursively, for  $n, m \in \mathbb{N}_0$ ,

$$\mathfrak{A}_{0,m}^v := T^{-1/2} \mathfrak{S}_m + \nu T^{1/2} \mathfrak{S}_{m+2} + A_{m+1} \mathfrak{S}_{m+2}^2, \quad (197)$$

$$\mathfrak{A}_{n,m}^p := A_m \sum_{a=0}^n \binom{n}{a} \mathfrak{A}_{a,m+1}^v \mathfrak{A}_{n-a,m+1}^v, \quad (198)$$

$$\mathfrak{A}_{n+1,m}^v := \nu \mathfrak{A}_{n,m+2}^v + A_{m+1} \sum_{a=0}^n \binom{n}{a} \mathfrak{A}_{a,m+2}^v \mathfrak{A}_{n-a,m+2}^v. \quad (199)$$

Every finite quantity in this recursion is finite, and

$$\sup_{0 \leq t \leq T} \|V_n(t)\|_{H^m} \leq \mathfrak{A}_{n,m}^v, \quad (200)$$

$$\sup_{0 \leq t \leq T} (\|Q_n(t)\|_{H^m} + \|\nabla Q_n(t)\|_{H^{m-1}}) \leq \mathfrak{A}_{n,m}^p. \quad (201)$$

Let  $M_- := \max\{M - 1, 0\}$  and set

$$\widehat{\mathfrak{B}}_{N,M}^h[u_0, \nu; T] := T \sum_{n=0}^N \left( (\mathfrak{A}_{n,M+1}^v)^2 + (\mathfrak{A}_{n+1,M_-}^v)^2 \right), \quad (202)$$

$$\widehat{\mathfrak{Q}}_{N,M}^h[u_0, \nu; T] := T \sum_{n=0}^N \mathfrak{A}_{n,M}^p. \quad (203)$$

Then

$$\mathfrak{B}_{N,M}[u_0, \nu; T] \leq \widehat{\mathfrak{B}}_{N,M}^h[u_0, \nu; T], \quad (204)$$

$$\mathfrak{Q}_{N,M}(p; T) \leq \widehat{\mathfrak{Q}}_{N,M}^h[u_0, \nu; T]. \quad (205)$$

For fixed  $N, M$ , the right-hand sides use only

$$\nu, \quad T, \quad \|u_0\|_2, \quad N, \quad M, \quad \{\mathfrak{H}_q[u_0, \nu; T] : 0 \leq q \leq M + 2N + 4\}, \quad \{A_q : 0 \leq q \leq M + 2N + 4\}. \quad (206)$$

The quantities  $\mathfrak{H}_q$  are the exact completed-height coordinates of the fixed datum-generated history. Equations (196) and (204) express the same finite rows in the rectangle and height coordinates, respectively. This is a change of coordinates, not an independent estimate of either family, and it is not used to prove the whole-terminal rectangle theorem.

*Proof.* The first estimate in (196) follows from the upper row of  $\mathfrak{B}_{0,m}$ . The Fourier inequality

$$(1 + |\xi|^2)^m \leq 2^m (1 + |\xi|^{2m+1})$$

and (58) give the second estimate.

The projected equation and (165) give

$$\|u_t\|_{L^1(I_T; H^m)} \leq \nu T^{1/2} \mathfrak{S}_{m+2} + A_{m+1} \mathfrak{S}_{m+2}^2.$$

Choose a Lebesgue time  $t_m \in I_T$  satisfying

$$\|u(t_m)\|_{H^m} \leq T^{-1/2} \|u\|_{L^2(I_T; H^m)}.$$

The Banach-valued fundamental identity gives (200) at  $n = 0$ . Assume that bound through row  $n$ . The pressure recurrence and (166) give (201). The projected temporal recurrence and (165) give the next velocity row through (199). Induction proves (200)–(201).

Integrating the pointwise bounds over  $I_T$  gives (204); when  $M = 0$ , use  $H^0 \hookrightarrow H^{-1}$  on the lower row. The same integration in (201) gives (205). Induction in (199) gives the finite dependency range in (206).  $\square$

**Lemma 7.8** (Closed velocity majorants). *For every  $n, m \in \mathbb{N}_0$ ,*

$$\|V_n(0)\|_{H^m} \leq \mathfrak{D}_{n,m}, \quad (207)$$

$$\|V_n\|_{L^2(I_T; H^m)} \leq \mathfrak{X}_{n,m}, \quad (208)$$

$$\sup_{0 \leq t \leq T} \|V_n(t)\|_{H^m} \leq \mathfrak{Z}_{n,m}. \quad (209)$$

For  $m = -1$ , equation (208) remains valid. Every right-hand side is finite and depends on  $(u_0, \nu, n, m, T)$  through the finite rectangle-matrix suprema in (173).

*Proof.* Applying  $\mathbb{P}$  to the temporal recurrence gives

$$V_{n+1} = \nu \Delta V_n - \mathbb{P} \nabla \cdot \sum_{a=0}^n \binom{n}{a} (V_a \otimes V_{n-a}). \quad (210)$$

Equation (165) at order  $m+1$ , evaluated at  $t=0$ , proves (207) by induction through (177). The same estimate, taken pointwise on  $[0, T]$ , proves by induction that

$$\sup_{0 \leq t \leq T} \|V_n(t)\|_{H^m} \leq \mathfrak{Y}_{n,m}. \quad (211)$$

The base case  $m=0$  is (58). For  $m \geq 1$ , both terms in (179) are valid: the first is the adjacent trace bound, and the second follows from

$$u(t) = u_0 + \int_0^t V_1(s) \, ds \quad \text{in } H^m.$$

The rectangle with middle order  $\mu(m)$ , Sobolev nesting, and (211) give respectively

$$\|V_n\|_{L^2(I_T; H^m)} \leq \mathfrak{B}_{n, \mu(m)}^{1/2}, \quad \|V_n\|_{L^2(I_T; H^m)} \leq T^{1/2} \mathfrak{Y}_{n, m+}.$$

This proves (208) through (181); the two exceptional improvements are (58) and (59). The adjacent trace lemma gives

$$\sup_{0 \leq t \leq T} \|V_n(t)\|_{H^m} \leq (1 + T^{-1})^{1/2} \mathfrak{B}_{n, m}^{1/2},$$

while the Banach-valued fundamental identity and (207) give

$$\sup_{0 \leq t \leq T} \|V_n(t)\|_{H^m} \leq \mathfrak{D}_{n, m} + T^{1/2} \mathfrak{X}_{n+1, m}.$$

Taking the minimum of these bounds and (211) proves (209).  $\square$

**Definition 7.9** (Closed pressure majorants). For  $n, m \in \mathbb{N}_0$ , define

$$\mathfrak{Q}_{n, m}^{(1)} := A_m \sum_{a=0}^n \binom{n}{a} \mathfrak{X}_{a, m+1} \mathfrak{X}_{n-a, m+1}, \quad (212)$$

$$\mathfrak{Q}_{n, m}^{(2)} := A_m \sum_{a=0}^n \binom{n}{a} \min \{ \mathfrak{Z}_{a, m+1} \mathfrak{X}_{n-a, m+1}, \mathfrak{X}_{a, m+1} \mathfrak{Z}_{n-a, m+1} \}, \quad (213)$$

$$\mathfrak{Q}_{n, m}^{(\infty)} := A_m \sum_{a=0}^n \binom{n}{a} \mathfrak{Z}_{a, m+1} \mathfrak{Z}_{n-a, m+1}, \quad (214)$$

$$\mathfrak{Q}_{n, m}^{(\text{mod})} := A_m \sum_{a=0}^n \binom{n}{a} (\mathfrak{X}_{a+1, m+1} \mathfrak{Z}_{n-a, m+1} + \mathfrak{Z}_{a, m+1} \mathfrak{X}_{n-a+1, m+1}). \quad (215)$$

These quantities carry the arguments  $[u_0, \nu; T]$  through  $\mathfrak{X}$  and  $\mathfrak{Z}$ .

### 7.3 Mixed velocity and pressure rows

The canonical majorants now turn each finite rectangle into bounds for the actual differentiated velocity and pressure rows, including their traces and time moduli.

**Theorem 7.10** (Quantitative mixed-jet rows and traces). *Let  $N, s \in \mathbb{N}_0$ , let  $0 \leq n \leq N$ , and let  $\alpha \in \mathbb{N}_0^3$ . Put*

$$J_{n,\alpha} := \partial_x^\alpha V_n.$$

*Then*

$$\|J_{n,\alpha}\|_{L^2(I_T; H^{s+1})}^2 + \|J_{n+1,\alpha}\|_{L^2(I_T; H^{s-1})}^2 \leq \mathfrak{P}_{N,s+|\alpha|}, \quad (216)$$

$$\sup_{0 \leq t \leq T} \|J_{n,\alpha}(t)\|_{H^s} \leq \mathfrak{T}_{N,s+|\alpha|}. \quad (217)$$

*There is a unique trace  $J_{n,\alpha}^T \in H^s$  at  $t = T$ , and for  $0 \leq r \leq t \leq T$ ,*

$$\|J_{n,\alpha}(t) - J_{n,\alpha}(r)\|_{H^{s-1}} \leq (t-r)^{1/2} \mathfrak{P}_{N,s+|\alpha|}^{1/2}, \quad (218)$$

$$\|J_{n,\alpha}(t) - J_{n,\alpha}(r)\|_{H^s} \leq (t-r)^{1/2} \mathfrak{P}_{N,s+|\alpha|+1}^{1/2}, \quad (219)$$

$$\|J_{n,\alpha}^T\|_{H^s} \leq \mathfrak{T}_{N,s+|\alpha|}. \quad (220)$$

*If  $0 \leq \theta \leq 1$ , then*

$$\|J_{n,\alpha}(t) - J_{n,\alpha}(r)\|_{H^{s-\theta}} \leq 2^{1-\theta} \mathfrak{T}_{N,s+|\alpha|}^{1-\theta} \mathfrak{P}_{N,s+|\alpha|}^{\theta/2} (t-r)^{\theta/2}. \quad (221)$$

*For  $1 \leq a \leq \infty$ ,  $2 \leq b \leq \infty$ ,  $k, q \in \mathbb{N}_0$ , and  $q > k + \frac{3}{2} - \frac{3}{b}$ ,*

$$\left\| \nabla^k J_{n,\alpha} \right\|_{L^a(I_T; L^b)} \leq S_{q,k,b} T^{1/a} \mathfrak{T}_{N,q+|\alpha|}, \quad (222)$$

$$\left\| \nabla^k J_{n,\alpha} \right\|_{L^2(I_T; L^b)} \leq S_{q,k,b} \mathfrak{P}_{N,q+|\alpha|}^{1/2}, \quad (223)$$

*where  $T^{1/\infty} = 1$ .*

*Proof.* Take the rectangle with middle order  $M = s + |\alpha|$ . The Fourier multiplier inequality

$$\|\partial_x^\alpha f\|_{H^r} \leq \|f\|_{H^{r+|\alpha|}}$$

and (162) give (216). The adjacent-row trace estimate gives

$$\begin{aligned} \sup_{0 \leq t \leq T} \|J_{n,\alpha}(t)\|_{H^s}^2 &\leq T^{-1} \|J_{n,\alpha}\|_{L^2(I_T; H^s)}^2 \\ &\quad + 2 \|J_{n,\alpha}\|_{L^2(I_T; H^{s+1})} \|J_{n+1,\alpha}\|_{L^2(I_T; H^{s-1})} \\ &\leq (T^{-1} + 1) \mathfrak{P}_{N,s+|\alpha|}, \end{aligned}$$

which proves (217) and (220).

The Bochner identity

$$J_{n,\alpha}(t) - J_{n,\alpha}(r) = \int_r^t J_{n+1,\alpha}(\tau) d\tau$$

in  $H^{s-1}$ , followed by Cauchy–Schwarz, proves (218). Applying the same argument to the rectangle with middle order  $s + |\alpha| + 1$  proves (219). Interpolation between  $H^{s-1}$  and  $H^s$ , together with  $\|J(t) - J(r)\|_{H^s} \leq 2 \mathfrak{T}_{N,s+|\alpha|}$ , proves (221). Sobolev embedding and (217) prove (222); Sobolev embedding and (216) prove (223).  $\square$

**Theorem 7.11** (Quantitative pressure rows). *Let  $N, s \in \mathbb{N}_0$ ,  $0 \leq n \leq N$ , and  $\alpha \in \mathbb{N}_0^3$ . Set*

$$\Pi_{n,\alpha} := \partial_x^\alpha Q_n, \quad r := s + |\alpha|.$$

*Then*

$$\|\Pi_{n,\alpha}\|_{L^1(I_T; H^s)} + \|\nabla \Pi_{n,\alpha}\|_{L^1(I_T; H^{s-1})} \leq 2A_r 2^n \mathfrak{P}_{N,r}, \quad (224)$$

$$\|\Pi_{n,\alpha}\|_{L^2(I_T; H^s)} \leq A_r 2^n \mathfrak{T}_{N,r+1} \mathfrak{P}_{N,r}^{1/2}, \quad (225)$$

$$\sup_{0 \leq t \leq T} \|\Pi_{n,\alpha}(t)\|_{H^s} \leq A_r 2^n \mathfrak{T}_{N,r+1}^2. \quad (226)$$

*There is a unique endpoint pressure row  $\Pi_{n,\alpha}^T \in H^s$ , and*

$$\|\Pi_{n,\alpha}^T\|_{H^s} \leq A_r 2^n \mathfrak{T}_{N,r+1}^2, \quad (227)$$

$$\|\Pi_{n,\alpha}(t) - \Pi_{n,\alpha}(r_0)\|_{H^s} \leq 2^{n+1} A_r \mathfrak{T}_{N,r+1} \mathfrak{P}_{N,r+2}^{1/2} |t - r_0|^{1/2} \quad (228)$$

*for  $0 \leq r_0 \leq t \leq T$ . If  $1 \leq a \leq \infty$ ,  $2 \leq b \leq \infty$ ,  $k, q \in \mathbb{N}_0$ , and  $q > k + \frac{3}{2} - \frac{3}{b}$ , then*

$$\left\| \nabla^k \Pi_{n,\alpha} \right\|_{L^a(I_T; L^b)} \leq S_{q,k,b} T^{1/a} A_{q+|\alpha|} 2^n \mathfrak{T}_{N,q+|\alpha|+1}^2. \quad (229)$$

*Proof.* The normalized pressure recurrence is

$$Q_n = R_i R_j \sum_{a=0}^n \binom{n}{a} V_{a,i} V_{n-a,j}. \quad (230)$$

Equation (165), Cauchy–Schwarz in time, and  $\sum_{a=0}^n \binom{n}{a} = 2^n$  give

$$\|Q_n\|_{L^1(I_T; H^r)} \leq A_r 2^n \mathfrak{P}_{N,r}. \quad (231)$$

The gradient estimate in (224) follows from  $\|\nabla f\|_{H^{r-1}} \leq \|f\|_{H^r}$ . Using one factor in  $L_t^\infty H_x^{r+1}$  and the other in  $L_t^2 H_x^{r+1}$  proves (225). Using both factors in  $L_t^\infty H_x^{r+1}$  proves (226) and (227).

Subtract (230) at times  $t$  and  $r_0$ , split each product difference into two terms, apply (219) in  $H^{r+1}$ , and use (217). The two product differences and the binomial sum give (228). Equation (229) follows from (226) and Sobolev embedding.  $\square$

**Proposition 7.12** (Matrix form of the closed rectangle estimates). *For nonnegative matrices of the same size, write  $A \preceq_{\text{ent}} B$  when every entry of  $A$  is at most the corresponding entry of  $B$ . When  $T = T_* < \infty$ , interpret the closed rectangle matrix and the two scalar rectangle coordinates by*

$$\begin{aligned} \mathbf{C}_{N,M}(T_*) &:= \lim_{S \uparrow T_*} \mathbf{C}_{N,M}(S), \\ \mathfrak{R}_{N,M}(u; T_*) &:= \lim_{S \uparrow T_*} \mathfrak{R}_{N,M}(u; S), \\ \mathfrak{Q}_{N,M}(p; T_*) &:= \lim_{S \uparrow T_*} \mathfrak{Q}_{N,M}(p; S). \end{aligned}$$

*All three limits exist in the extended nonnegative reals by monotonicity. Theorem 4.6 and Proposition 4.9 make the scalar limits finite, and the estimates below make every matrix entry finite. Then*

$$\mathbf{C}_{N,M}(T) \preceq_{\text{ent}} \left( \mathfrak{X}_{n,M+1} \quad \mathfrak{X}_{n+1,M-1} \quad \mathfrak{Q}_{n,M}^{(1)} \right)_{n=0}^N. \quad (232)$$

Consequently,

$$\mathfrak{R}_{N,M}(u; T) \leq \sum_{n=0}^N (\mathfrak{X}_{n,M+1}^2 + \mathfrak{X}_{n+1,M-1}^2), \quad (233)$$

$$\mathfrak{Q}_{N,M}(p; T) \leq \sum_{n=0}^N \mathfrak{Q}_{n,M}^{(1)}. \quad (234)$$

*Proof.* The first two columns of (232) are (208). The pressure recurrence (230), the product estimate (166), and Cauchy–Schwarz in time give the third column through (212). Take the Frobenius norm of the first two columns and sum the third column, using (142).  $\square$

## 7.4 Frequency and critical-scale consequences

**Proposition 7.13** (Velocity and pressure frequency tails). *Let  $\mathbf{P}_{\geq K}$  be the Fourier multiplier of  $\mathbf{1}_{\{|\xi| \geq K\}}$ , where  $K \geq 1$ . Let  $r_1 \in \mathbb{N}$ ,  $s \in \mathbb{N}_0$ ,  $0 \leq n \leq N$ , and  $\alpha \in \mathbb{N}_0^3$ . Then*

$$\|\mathbf{P}_{\geq K} J_{n,\alpha}\|_{L^2(I_T; H^s)} \leq K^{-r_1} \mathfrak{P}_{N,s+r_1+|\alpha|}^{1/2}, \quad (235)$$

$$\sup_{0 \leq t \leq T} \|\mathbf{P}_{\geq K} J_{n,\alpha}(t)\|_{H^s} \leq K^{-r_1} \mathfrak{T}_{N,s+r_1+|\alpha|}, \quad (236)$$

$$\|\mathbf{P}_{\geq K} \Pi_{n,\alpha}\|_{L^1(I_T; H^s)} \leq K^{-r_1} A_{s+r_1+|\alpha|} 2^n \mathfrak{P}_{N,s+r_1+|\alpha|}, \quad (237)$$

$$\sup_{0 \leq t \leq T} \|\mathbf{P}_{\geq K} \Pi_{n,\alpha}(t)\|_{H^s} \leq K^{-r_1} A_{s+r_1+|\alpha|} 2^n \mathfrak{T}_{N,s+r_1+|\alpha|+1}^2. \quad (238)$$

The pointwise estimates (236) and (238) hold at  $t = T$ .

*Proof.* For every  $f \in H^{s+r_1}$ ,

$$\|\mathbf{P}_{\geq K} f\|_{H^s} \leq K^{-r_1} \|f\|_{H^{s+r_1}}.$$

Apply this inequality to (216) and (217) at spatial order  $s + r_1$  to obtain the two velocity bounds. Since Fourier multipliers commute,

$$\|\mathbf{P}_{\geq K} \Pi_{n,\alpha}\|_{H^s} \leq K^{-r_1} \|Q_n\|_{H^{s+r_1+|\alpha|}}.$$

Integration in time followed by (231) at order  $s + r_1 + |\alpha|$  gives (237). Applying the tail inequality directly to  $\Pi_{n,\alpha}$ , followed by (226) with  $s$  replaced by  $s + r_1$ , gives (238).  $\square$

**Proposition 7.14** (Fixed weighted dyadic shell sums). *Let  $(\Delta_j)_{j \geq -1}$  be a fixed inhomogeneous Littlewood–Paley decomposition on  $\mathbb{R}^3$ . Let  $\sigma \in \mathbb{R}$ ,  $k \in \mathbb{N}_0$ , and choose  $M \in \mathbb{N}_0$  such that*

$$M > \sigma + k + \frac{3}{2}. \quad (239)$$

Then

$$\int_0^T \sum_{j \geq -1} 2^{\sigma j} \left\| \Delta_j \nabla^k u(t) \right\|_{L^\infty} dt \leq C_{\Delta, \sigma, k, M} T^{1/2} \mathfrak{X}_{0,M}. \quad (240)$$

In particular, with  $\omega = \nabla \times u$ ,

$$\sum_{j \geq -1} \int_0^T \left\| \Delta_j \omega(t) \right\|_{L^\infty} dt \leq C_\Delta T^{1/2} \mathfrak{X}_{0,3} < \infty. \quad (241)$$

Each estimate uses one fixed finite Sobolev coordinate; no summation over Sobolev orders is asserted.

*Proof.* Bernstein's inequality on a shell gives, for  $j \geq 0$ ,

$$2^{\sigma j} \left\| \Delta_j \nabla^k f \right\|_{L^\infty} \leq C_\Delta 2^{(\sigma+k+3/2)j} \left\| \Delta_j f \right\|_2.$$

Put  $d := M - \sigma - k - 3/2 > 0$ . Cauchy–Schwarz in  $j$  yields

$$\begin{aligned} \sum_{j \geq 0} 2^{\sigma j} \left\| \Delta_j \nabla^k f \right\|_{L^\infty} &\leq C_\Delta \left( \sum_{j \geq 0} 2^{-2dj} \right)^{1/2} \left( \sum_{j \geq 0} 2^{2Mj} \left\| \Delta_j f \right\|_2^2 \right)^{1/2} \\ &\leq C_{\Delta, \sigma, k, M} \|f\|_{H^M}. \end{aligned}$$

The single low block  $j = -1$  obeys the same final bound. The last inequality follows from Plancherel's theorem and the finite overlap of the dyadic multipliers. Integrating in time, applying Cauchy–Schwarz, and using (208) prove (240). Since curl commutes with every  $\Delta_j$ , the choice  $(\sigma, k, M) = (0, 1, 3)$  gives (241).  $\square$

**Proposition 7.15** (Critical-height bounds). *Define*

$$E_\sigma(t) := \frac{1}{2} \left\| \Lambda^\sigma u(t) \right\|_2^2, \quad H(t) := E_{1/2}(t),$$

and

$$E_{j+1/2, \geq K}(t) := \frac{1}{2} \int_{|\xi| \geq K} |\xi|^{2j+1} |\widehat{u}(t, \xi)|^2 d\xi.$$

For  $j \in \mathbb{N}_0$ ,

$$\sup_{0 \leq t \leq T} E_{j+1/2}(t) \leq \frac{1}{2} \mathfrak{T}_{0, j+1}^2, \quad (242)$$

$$\int_0^T E_{j+1/2}(t) dt \leq \frac{1}{2} \mathfrak{P}_{0, j}. \quad (243)$$

If  $m > j + \frac{1}{2}$  is an integer and  $K \geq 1$ , then

$$\sup_{0 \leq t \leq T} E_{j+1/2, \geq K}(t) \leq \frac{1}{2} K^{2j+1-2m} \mathfrak{T}_{0, m}^2, \quad (244)$$

$$\int_0^T E_{j+1/2, \geq K}(t) dt \leq \frac{1}{2} K^{2j+1-2m} \mathfrak{P}_{0, m-1}. \quad (245)$$

For  $k \in \mathbb{N}$ ,

$$\left\| H^{(k)} \right\|_{L^1(0, T)} \leq 2^{k-1} \mathfrak{P}_{k, 0}. \quad (246)$$

In particular,

$$\left\| H' \right\|_{L^1(0, T)} \leq \frac{1}{2} \mathfrak{P}_{0, 2}, \quad (247)$$

$$\int_0^T \left\| \Lambda^{3/2} u(t) \right\|_2^2 dt \leq \mathfrak{P}_{0, 1}, \quad (248)$$

$$\left\| \mathcal{P}_{1/2} \right\|_{L^1(0, T)} \leq \frac{1}{2} \mathfrak{P}_{0, 2} + \nu \mathfrak{P}_{0, 1}, \quad (249)$$

$$|H(T) - H(t)| \leq \mathfrak{T}_{0, 1} \mathfrak{P}_{0, 2}^{1/2} (T - t)^{1/2}, \quad 0 \leq t \leq T, \quad (250)$$

where  $\mathcal{P}_{1/2} = H' + \nu \left\| \Lambda^{3/2} u \right\|_2^2$ .

*Proof.* Equations (242) and (243) follow from  $\dot{H}^{j+1/2} \hookleftarrow H^{j+1}$ , (217), and (216). On  $|\xi| \geq K$ ,

$$|\xi|^{2j+1} \leq K^{2j+1-2m} |\xi|^{2m},$$

which proves (244) and (245).

The identity

$$H^{(k)}(t) = \frac{1}{2} \sum_{a=0}^k \binom{k}{a} \operatorname{Re} \left\langle \Lambda^{1/2} V_a(t), \Lambda^{1/2} V_{k-a}(t) \right\rangle$$

and Cauchy–Schwarz in time give (246). At  $k = 1$ , the two equal terms give

$$\|H'\|_{L^1(0,T)} \leq \|u_t\|_{L^2(I_T; \dot{H}^{1/2})} \|u\|_{L^2(I_T; \dot{H}^{1/2})} \leq \frac{1}{2} \mathfrak{P}_{0,2}.$$

Equation (248) follows from the upper velocity row of the  $(0, 1)$  rectangle. The identity defining  $\mathcal{P}_{1/2}$  gives (249). Finally,

$$\|u(T) - u(t)\|_{\dot{H}^{1/2}} \leq \|u(T) - u(t)\|_{H^1} \leq (T-t)^{1/2} \|u_t\|_{L^2(t,T; H^1)} \leq (T-t)^{1/2} \mathfrak{P}_{0,2}^{1/2},$$

and

$$|H(T) - H(t)| \leq \frac{1}{2} (\|u(T)\|_{\dot{H}^{1/2}} + \|u(t)\|_{\dot{H}^{1/2}}) \|u(T) - u(t)\|_{\dot{H}^{1/2}}.$$

Equation (217) bounds each retained velocity factor, so these two estimates prove (250).  $\square$

Taking  $j = 0$  and any integer  $m \geq 1$  in (244) gives the uniform tail limit

$$\lim_{K \rightarrow \infty} \sup_{0 \leq t \leq T} E_{1/2, \geq K}(t) = 0. \quad (251)$$

## 7.5 Continuation and quantitative restart

**Proposition 7.16** (Continuation norms). *The finite-horizon solution satisfies*

$$\|u\|_{L^2(I_T; L^\infty)} \leq S_{2,0,\infty} \mathfrak{P}_{0,1}^{1/2}, \quad (252)$$

$$\int_0^T \|\nabla u(t)\|_{L^\infty} dt \leq S_{3,1,\infty} T^{1/2} \mathfrak{P}_{0,2}^{1/2}, \quad (253)$$

$$\int_0^T \|\nabla \times u(t)\|_{L^\infty} dt \leq 2S_{3,1,\infty} T^{1/2} \mathfrak{P}_{0,2}^{1/2}. \quad (254)$$

For every integer  $m \geq 3$ , let  $K_m$  be the explicit constant in (14). Then

$$|\operatorname{Re} \langle A^m \mathbb{P} \nabla \cdot (u \otimes u), A^m u \rangle_{L^2}| \leq K_m \|\nabla u\|_{L^\infty} \|u\|_{H^m}^2. \quad (255)$$

$$\sup_{0 \leq t \leq T} \|u(t)\|_{H^m}^2 \leq \|u_0\|_{H^m}^2 \exp \left( 2K_m S_{3,1,\infty} T^{1/2} \mathfrak{P}_{0,2}^{1/2} \right). \quad (256)$$

*Proof.* Sobolev embedding and the upper rows of the  $(0, 1)$  and  $(0, 2)$  rectangles give (252) and (253); the pointwise bound  $\|\nabla \times u\|_\infty \leq 2 \|\nabla u\|_\infty$  gives (254). The  $H^m$  energy identity and (255) give

$$\frac{1}{2} \frac{d}{dt} \|u\|_{H^m}^2 + \nu \|\nabla u\|_{H^m}^2 \leq K_m \|\nabla u\|_{L^\infty} \|u\|_{H^m}^2,$$

and hence

$$\frac{d}{dt} \|u\|_{H^m}^2 \leq 2K_m \|\nabla u\|_{L^\infty} \|u\|_{H^m}^2.$$

Gronwall's inequality and (253) prove (256).  $\square$

**Theorem 7.17** (Explicit endpoint-restart bound). *Assume  $T = T_* < \infty$ , and set  $u_T := u_*$ . Define the maximal restart lifespan inside the shifted pressure-normalized  $H^3$  mild class by*

$$\begin{aligned}\delta_T^{\max} &:= \tau_{\nu,3}^{\max}(T, u_T) \\ &= \sup \{ \delta > 0 : \mathcal{M}_\nu^3(T, T + \delta; u_T) \neq \emptyset \} \in (0, \infty].\end{aligned}\tag{257}$$

Define

$$\delta_{\text{qc}}[u_0, \nu; T] := \min \left\{ 1, c_\nu (1 + \mathfrak{Z}_{0,3}[u_0, \nu; T])^{-2} \right\}.\tag{258}$$

The class  $\mathcal{M}_\nu^3(T, T + \delta_{\text{qc}}; u_T)$  contains exactly one pair  $(w, q)$ . It is the same-history continuation constructed in Theorem 5.8, and

$$\delta_T^{\max} \geq \delta_{\text{qc}}[u_0, \nu; T] > 0.\tag{259}$$

At the base restart order,

$$\begin{aligned}\|w\|_{C([T, T+\delta_{\text{qc}}]; H^3)} + \|w\|_{L^2(T, T+\delta_{\text{qc}}; H^4)} + \|w_t\|_{L^2(T, T+\delta_{\text{qc}}; H^2)} \\ \leq 2L_\nu \mathfrak{Z}_{0,3}[u_0, \nu; T].\end{aligned}\tag{260}$$

Set

$$\mathfrak{A}_T := 2S_{3,1,\infty} L_\nu \delta_{\text{qc}} \mathfrak{Z}_{0,3}[u_0, \nu; T].\tag{261}$$

For every integer  $m \geq 3$ ,

$$\sup_{T \leq t \leq T+\delta_{\text{qc}}} \|w(t)\|_{H^m}^2 \leq \mathfrak{Z}_{0,m}[u_0, \nu; T]^2 \exp(2K_m \mathfrak{A}_T).\tag{262}$$

*Proof.* Equations (103) and (209) give

$$\|u_T\|_{H^3} \leq \mathfrak{Z}_{0,3}.$$

Consequently,

$$\delta_{\text{qc}}[u_0, \nu; T] \leq \min \{ 1, c_\nu (1 + \|u_T\|_{H^3})^{-2} \}.$$

Theorem 2.11 gives the unique member of  $\mathcal{M}_\nu^3(T, T + \delta_{\text{qc}}; u_T)$ . The fixed-point ball in its proof gives (260). Hence

$$\int_T^{T+\delta_{\text{qc}}} \|\nabla w(t)\|_\infty dt \leq S_{3,1,\infty} \delta_{\text{qc}} \|w\|_{C([T, T+\delta_{\text{qc}}]; H^3)} \leq \mathfrak{A}_T.$$

The  $H^m$  energy estimate in Proposition 7.16, now on the restart interval, and  $\|u_T\|_{H^m} \leq \mathfrak{Z}_{0,m}$  prove (262). Theorem 5.8 identifies this unique local pair with the continuation of the preterminal history, and its membership on the displayed interval proves (259).  $\square$

## 8 Three-dimensional applications

### 8.1 Explicit closure for small critical data

The construction below gives an explicit three-dimensional  $\dot{H}^{1/2}$  realization of the critical-data mechanism introduced by Fujita and Kato [5]; every constant used here is derived in the displayed estimates.

Use the explicit homogeneous Sobolev constants

$$S_3 := S(3), \quad C_{\text{src}} := S_3^3 = \frac{1}{\sqrt{2}\pi}.$$

The same constants hold for vector fields. Indeed, apply the scalar inequality to the Euclidean length  $|F|$ : one has  $|\nabla|F|| \leq |\nabla F|$ , while the difference-quotient representation of  $\dot{H}^{1/2}$  and  $||F(x)| - |F(y)|| \leq |F(x) - F(y)|$  give  $||F||_{\dot{H}^{1/2}} \leq \|F\|_{\dot{H}^{1/2}}$ .

**Lemma 8.1** (Critical action identity). *Let  $(u, p)$  be a smooth pressure-normalized solution on a time interval  $I \subset [0, T_*)$ , and set*

$$\mathcal{B}(u) := \mathbb{P}\nabla \cdot (u \otimes u).$$

*Then, pointwise on  $I$ ,*

$$\nu \frac{d}{dt} \|u\|_{\dot{H}^{1/2}}^2 + \nu^2 \|u\|_{\dot{H}^{3/2}}^2 + \|u_t\|_{\dot{H}^{-1/2}}^2 = \|\mathcal{B}(u)\|_{\dot{H}^{-1/2}}^2. \quad (263)$$

*Proof.* The projected equation is

$$u_t = -\nu\Lambda^2 u - \mathcal{B}(u).$$

Put

$$g := \Lambda^{-1/2} \mathcal{B}(u), \quad w := \Lambda^{-1/2} u_t = -\nu\Lambda^{3/2} u - g.$$

Pairing the projected equation with  $\Lambda u$  gives

$$\frac{d}{dt} \|u\|_{\dot{H}^{1/2}}^2 = -2\nu \|u\|_{\dot{H}^{3/2}}^2 - 2 \operatorname{Re} \left\langle g, \Lambda^{3/2} u \right\rangle_2.$$

Multiply this equality by  $\nu$  and expand  $\|w\|_2^2 = \|\nu\Lambda^{3/2} u + g\|_2^2$ . The cross terms cancel and give (263).  $\square$

For  $m \in \mathbb{N}$ , define the explicit homogeneous commutator constant

$$\mathsf{K}_m^{\text{hom}} := 3^m \sum_{\ell=1}^m \binom{m}{\ell} \mathsf{S}(\mathfrak{p}_{m,\ell}) \mathsf{S}(\mathfrak{q}_{m,\ell}), \quad (264)$$

where  $(\mathfrak{p}_{1,1}, \mathfrak{q}_{1,1}) = (3, 6)$ , and, for  $m \geq 2$ ,

$$\vartheta_{m,\ell} := \frac{\ell-1}{m-1}, \quad \frac{1}{\mathfrak{p}_{m,\ell}} := \frac{1}{3} - \frac{\vartheta_{m,\ell}}{6}, \quad \frac{1}{\mathfrak{q}_{m,\ell}} := \frac{1}{6} + \frac{\vartheta_{m,\ell}}{6}.$$

The corresponding estimate is

$$|\operatorname{Re} \langle \Lambda^m \mathbb{P}\nabla \cdot (z \otimes z), \Lambda^m z \rangle_{L^2}| \leq \mathsf{K}_m^{\text{hom}} \left\| \Lambda^{3/2} z \right\|_2 \|\Lambda^m z\|_2 \|\Lambda^{m+1} z\|_2 \quad (265)$$

for every  $z \in \mathcal{S}_\sigma(\mathbb{R}^3)$ .

**Lemma 8.2** (Explicit homogeneous critical commutator). *For every  $m \in \mathbb{N}$ ,  $\mathsf{K}_m^{\text{hom}} < \infty$ , and (265) holds with the displayed constant. Moreover, (265) extends to every divergence-free  $z \in H^{m+1}(\mathbb{R}^3) \cap \dot{H}^{3/2}(\mathbb{R}^3)$ .*

*Proof.* The multinomial identity gives the exact pairing

$$\langle \Lambda^m f, \Lambda^m g \rangle_{L^2} = \sum_{|\alpha|=m} \frac{m!}{\alpha!} \langle \partial^\alpha f, \partial^\alpha g \rangle_{L^2}.$$

After differentiating  $(z \cdot \nabla)z$ , the undifferentiated transport term has zero pairing by incompressibility. A remaining term has the form

$$\left\langle (\partial^\beta z \cdot \nabla) \partial^{\alpha-\beta} z, \partial^\alpha z \right\rangle_{L^2}, \quad |\alpha| = m, \quad 0 < \beta \leq \alpha.$$

Put  $\ell = |\beta|$ . For  $m \geq 2$ , Hölder's inequality, Lemma 2.3, and Fourier log-convexity give

$$\begin{aligned} & \left\| \partial^\beta z \right\|_{\mathfrak{p}_{m,\ell}} \left\| \nabla \partial^{\alpha-\beta} z \right\|_{\mathfrak{q}_{m,\ell}} \\ & \leq \mathbf{S}(\mathfrak{p}_{m,\ell}) \mathbf{S}(\mathfrak{q}_{m,\ell}) \|z\|_{\dot{H}^{3/2}} \|z\|_{\dot{H}^{m+1}}. \end{aligned}$$

Indeed, the two differentiated Sobolev orders are the complementary interpolants

$$\begin{aligned} \ell + s(\mathfrak{p}_{m,\ell}) &= (1 - \vartheta_{m,\ell}) \frac{3}{2} + \vartheta_{m,\ell}(m+1), \\ m - \ell + 1 + s(\mathfrak{q}_{m,\ell}) &= \vartheta_{m,\ell} \frac{3}{2} + (1 - \vartheta_{m,\ell})(m+1). \end{aligned}$$

For  $m = 1$ , the same bound is the direct  $L^3$ - $L^6$  estimate. Finally,

$$\sum_{|\alpha|=m} \frac{m!}{\alpha!} = 3^m, \quad \sum_{\substack{\beta \leq \alpha \\ |\beta|=\ell}} \binom{\alpha}{\beta} = \binom{m}{\ell}.$$

Summation gives exactly (264) and proves (265). Solenoidal Fourier truncation and convergence in  $H^{m+1} \cap \dot{H}^{3/2}$  give the asserted extension.  $\square$

**Theorem 8.3** (Explicit small-critical-data global regularity). *Let  $\nu > 0$  and  $a \in \mathcal{S}_\sigma(\mathbb{R}^3)$ . Set*

$$a_c := \left\| \Lambda^{1/2} a \right\|_2, \quad \delta_0 := \nu^2 - C_{\text{src}}^2 a_c^2.$$

*If*

$$a_c < \frac{\nu}{C_{\text{src}}} = \sqrt{2} \pi \nu, \tag{266}$$

*then  $a$  generates a unique global pressure-normalized smooth solution*

$$(u, p) \in C^\infty([0, \infty) \times \mathbb{R}^3; \mathbb{R}^3 \times \mathbb{R}).$$

*For every finite  $T > 0$ , the solution is the unique member of  $\mathcal{M}_\nu^3(0, T; a)$ . For every  $t \geq 0$ ,*

$$\nu \left\| \Lambda^{1/2} u(t) \right\|_2^2 + \delta_0 \int_0^t \left\| \Lambda^{3/2} u(s) \right\|_2^2 ds + \int_0^t \|u_t(s)\|_{\dot{H}^{-1/2}}^2 ds \leq \nu a_c^2, \quad \delta_0 > 0. \tag{267}$$

*The critical nonlinear source and the first temporal response satisfy*

$$\int_0^\infty \|\mathbb{P} \nabla \cdot (u \otimes u)(t)\|_{\dot{H}^{-1/2}}^2 dt \leq \frac{C_{\text{src}}^2 \nu a_c^4}{\delta_0} = \frac{\nu a_c^4}{2\pi^2 \nu^2 - a_c^2}, \tag{268}$$

$$\nu^2 \int_0^\infty \left\| \Lambda^{3/2} u(t) \right\|_2^2 dt + \int_0^\infty \|u_t(t)\|_{\dot{H}^{-1/2}}^2 dt \leq \nu a_c^2 + \frac{C_{\text{src}}^2 \nu a_c^4}{\delta_0}. \tag{269}$$

*For every  $m \in \mathbb{N}$ , define*

$$\Gamma_m^{\text{sm}} := \frac{(\mathbf{K}_m^{\text{hom}})^2 a_c^2}{\delta_0}.$$

Then

$$\begin{aligned} \|\Lambda^m u(t)\|_2^2 + \nu \int_0^t \|\Lambda^{m+1} u(s)\|_2^2 \, ds \\ \leq \|\Lambda^m a\|_2^2 \exp(\Gamma_m^{\text{sm}}). \end{aligned} \quad (270)$$

In particular,

$$\sup_{t \geq 0} \|u(t)\|_{H^m}^2 \leq 2^{m-1} \left[ \|a\|_2^2 + \|\Lambda^m a\|_2^2 \exp(\Gamma_m^{\text{sm}}) \right]. \quad (271)$$

*Proof.* Iterate Theorem 2.11 and join overlapping intervals by its at-most-one assertion for the shifted mild class. The union is the maximal shifted pressure-normalized  $H^3$  mild history on  $[0, T_*)$ , and persistence retains  $H^\infty$  on every compact subinterval. Put

$$A(t) := \left\| \Lambda^{1/2} u(t) \right\|_2, \quad B(t) := \left\| \Lambda^{3/2} u(t) \right\|_2.$$

Dual Sobolev, Hölder, and the unit Fourier-multiplier norm of  $\mathbb{P}$  give the complete source-constant chain

$$\begin{aligned} \|\mathbb{P} \nabla \cdot (u \otimes u)\|_{\dot{H}^{-1/2}} &\leq S_3 \|(u \cdot \nabla) u\|_{L^{3/2}} \\ &\leq S_3 \|u\|_{L^3} \|\nabla u\|_{L^3} \\ &\leq S_3^3 \left\| \Lambda^{1/2} u \right\|_2 \left\| \Lambda^{3/2} u \right\|_2 = C_{\text{src}} A(t) B(t). \end{aligned} \quad (272)$$

Combining (272) with the exact identity (263) gives, on every classical subinterval,

$$\nu \frac{d}{dt} A(t)^2 + (\nu^2 - C_{\text{src}}^2 A(t)^2) B(t)^2 + \|u_t(t)\|_{\dot{H}^{-1/2}}^2 \leq 0.$$

Continuity and a first-exit argument give  $A(t) \leq a_c$  throughout  $[0, T_*)$ . Integration and the definition of  $\delta_0$  prove (267). In particular,

$$\int_0^t B(s)^2 \, ds \leq \frac{\nu a_c^2}{\delta_0}.$$

For every  $t < T_*$ , the same source estimate and the integrated identity (263) prove (268) and (269) with  $\infty$  replaced by  $t$ . After the continuation argument below proves  $T_* = \infty$ , monotone convergence gives the stated forms.

At order  $m$ , Lemma 8.2 and Young's inequality yield

$$\frac{d}{dt} \|\Lambda^m u\|_2^2 + \nu \|\Lambda^{m+1} u\|_2^2 \leq \frac{(\mathbf{K}_m^{\text{hom}})^2}{\nu} B(t)^2 \|\Lambda^m u\|_2^2.$$

Multiplication by the corresponding integrating factor and

$$\int_0^t B(s)^2 \, ds \leq \frac{\nu a_c^2}{\delta_0}$$

prove (270). The Fourier inequality

$$(1 + |\xi|^2)^m \leq 2^{m-1} (1 + |\xi|^{2m})$$

and the kinetic-energy identity prove (271).

Let  $M_3$  be the square root of the right-hand side of (271) with  $m = 3$ , and set

$$\delta_{\text{crit}} := \min\{1, c_\nu(1 + M_3)^{-2}\} > 0.$$

Every datum  $u(t_0)$ ,  $t_0 < T_*$ , has  $H^3$ -norm at most  $M_3$ . Theorem 2.11 restarts the solution for at least  $\delta_{\text{crit}}$  units of time. If  $T_* < \infty$ , choose  $t_0 < T_*$  with  $T_* - t_0 < \delta_{\text{crit}}/2$ . Both restrictions belong to  $\mathcal{M}_\nu^3(t_0, b; u(t_0))$  for every  $b < T_*$ , so uniqueness on each such closed subinterval identifies them throughout  $[t_0, T_*)$ . The restarted member then extends the original history beyond  $T_*$ , contradicting maximality. Hence  $T_* = \infty$ , and persistence gives the asserted smoothness. The same local uniqueness on each finite interval proves the stated global uniqueness class.  $\square$

For  $m \geq 1$ , define

$$U_m^{\text{sm}} := 2^{(m-1)/2} \left[ \|a\|_2^2 + \|\Lambda^m a\|_2^2 \exp(\Gamma_m^{\text{sm}}) \right]^{1/2}, \quad U_0^{\text{sm}} := \|a\|_2, \quad U_{-1}^{\text{sm}} := U_0^{\text{sm}}.$$

For  $n \in \mathbb{N}_0$  and  $m \geq -1$ , define the finite-order recursion

$$\mathbf{Y}_{0,m}^{\text{sm}} := U_m^{\text{sm}}, \quad (273)$$

$$\mathbf{Y}_{n+1,m}^{\text{sm}} := \nu \mathbf{Y}_{n,m+2}^{\text{sm}} + A_{m+1} \sum_{a'=0}^n \binom{n}{a'} \mathbf{Y}_{a',m+2}^{\text{sm}} \mathbf{Y}_{n-a',m+2}^{\text{sm}}. \quad (274)$$

**Proposition 8.4** (Explicit small-data pressure-complete rectangles). *Under the hypotheses of Theorem 8.3, for every  $n \in \mathbb{N}_0$  and  $m \geq -1$ ,*

$$\sup_{t \geq 0} \|V_n(t)\|_{H^m} \leq \mathbf{Y}_{n,m}^{\text{sm}}. \quad (275)$$

For every  $T < \infty$  and  $N, M \in \mathbb{N}_0$ ,

$$\mathfrak{R}_{N,M}(u; T) \leq T \sum_{n=0}^N [(\mathbf{Y}_{n,M+1}^{\text{sm}})^2 + (\mathbf{Y}_{n+1,M-1}^{\text{sm}})^2], \quad (276)$$

$$\mathfrak{Q}_{N,M}(p; T) \leq T A_M \sum_{n=0}^N \sum_{a'=0}^n \binom{n}{a'} \mathbf{Y}_{a',M+1}^{\text{sm}} \mathbf{Y}_{n-a',M+1}^{\text{sm}}. \quad (277)$$

In the entrywise order from Proposition 7.12, these bounds have the matrix form

$$\mathbf{C}_{N,M}(T) \preceq_{\text{ent}} \begin{pmatrix} T^{1/2} \mathbf{Y}_{n,M+1}^{\text{sm}} & T^{1/2} \mathbf{Y}_{n+1,M-1}^{\text{sm}} & T A_M \sum_{a'=0}^n \binom{n}{a'} \mathbf{Y}_{a',M+1}^{\text{sm}} \mathbf{Y}_{n-a',M+1}^{\text{sm}} \end{pmatrix}_{n=0}^N. \quad (278)$$

Every right-hand side is an explicit finite function of  $\nu, T, N, M$  and the datum norms

$$\|a\|_2, \quad \left\| \Lambda^{1/2} a \right\|_2, \quad \{ \|\Lambda^r a\|_2 : 1 \leq r \leq M + 2N + 1 \}. \quad (279)$$

Indeed, induction in  $n$  shows that  $\mathbf{Y}_{n,m}^{\text{sm}}$  uses only the entries  $U_j^{\text{sm}}$  with  $-1 \leq j \leq m + 2n$ ; both velocity columns of (278) consequently stop at  $j = M + 2N + 1$ , and its pressure column stops no later.

*Proof.* Equation (271) proves the base (275) for  $m \geq 1$ ; the kinetic-energy identity proves it for  $m = 0$ , and  $H^0 \hookrightarrow H^{-1}$  proves it for  $m = -1$ . Apply  $\mathbb{P}$  to the temporal recurrence. This is the  $\alpha = 0$  temporal block-row of (135); its factorial conjugate is (138). Equation (165) at order  $m + 1$  and induction give (275) through (274); for  $m = -1$ , use the order-zero product estimate and  $H^0 \hookrightarrow H^{-1}$ . Integration in time gives (276). The pressure recurrence and (166) at order  $M$ , followed by integration and summation in  $n$ , give (277). The same rowwise bounds give (278); taking the Frobenius norm of its first two columns and summing its pressure column recovers (276) and (277).  $\square$

## 8.2 Low-rectangle and scale-local bounds

The following results use distinct outputs of the compatible rectangles: mixed space–time embeddings, pressure completion, and integrable Lipschitz action.

**Proposition 8.5** (Three low-rectangle regularity classes). *Let  $u$  be the global solution generated by Theorem 5.10, and let  $0 < T < \infty$ . Then*

$$\|u\|_{L^\infty(0,T;L^3)} \leq S_{1,0,3}\mathfrak{Z}_{0,1}[u_0, \nu; T], \quad (280)$$

$$\|u\|_{L^2(0,T;L^\infty)} \leq S_{2,0,\infty}\mathfrak{X}_{0,2}[u_0, \nu; T], \quad (281)$$

$$\|\nabla u\|_{L^2(0,T;L^3)} \leq S_{2,1,3}\mathfrak{X}_{0,2}[u_0, \nu; T]. \quad (282)$$

*In particular, each of the three norms is finite on every finite horizon, with the stated datum, viscosity, and horizon dependence.*

*Proof.* Use (324) with  $(n, \alpha, a, b, q, k) = (0, 0, \infty, 3, 1, 0)$  to obtain (280). Use (325) with  $(n, \alpha, b, q, k) = (0, 0, \infty, 2, 0)$  and  $(0, 0, 3, 2, 1)$  to obtain (281) and (282). The strict embedding conditions are, respectively,  $1 > 1/2$ ,  $2 > 3/2$ , and  $2 > 3/2$ .  $\square$

**Corollary 8.6** (Uniform extinction of parabolic scale content). *Let  $\nu > 0$ , let  $u_0 \in H_\sigma^\infty(\mathbb{R}^3)$ , and let  $(u, p)$  be the pressure-normalized global pair supplied by Theorem 5.10. This is a post-regularity consequence for the Caffarelli–Kohn–Nirenberg scale-normalized quantity [2]; it is not used as an  $\varepsilon$ -regularity input or continuation criterion. For  $0 < T < \infty$ , define*

$$\begin{aligned} U_T &:= S_{2,0,\infty}\mathfrak{Z}_{0,2}[u_0, \nu; T], \\ P_T &:= S_{2,0,\infty}\mathfrak{Q}_{0,2}^{(\infty)}[u_0, \nu; T]. \end{aligned} \quad (283)$$

*Let  $z_0 = (t_0, x_0)$ ,  $\rho > 0$ , and suppose that*

$$Q_\rho(z_0) := (t_0 - \rho^2, t_0) \times B_\rho(x_0) \subset (0, T) \times \mathbb{R}^3.$$

*Then the scale-normalized velocity–pressure content satisfies*

$$\rho^{-2} \iint_{Q_\rho(z_0)} (|u|^3 + |p|^{3/2}) \, dx \, dt \leq \frac{4\pi}{3} \rho^3 (U_T^3 + P_T^{3/2}). \quad (284)$$

*Consequently,*

$$\lim_{\rho \downarrow 0} \sup_{\{z_0: Q_\rho(z_0) \subset (0, T) \times \mathbb{R}^3\}} \rho^{-2} \iint_{Q_\rho(z_0)} (|u|^3 + |p|^{3/2}) \, dx \, dt = 0. \quad (285)$$

*Proof.* Equation (319), followed by (168) with  $(s, k, b) = (2, 0, \infty)$ , gives

$$\|u\|_{L^\infty((0,T) \times \mathbb{R}^3)} \leq U_T.$$

Equations (328) and (168) give

$$\|p\|_{L^\infty((0,T) \times \mathbb{R}^3)} \leq P_T.$$

Since  $|B_\rho| = \frac{4\pi}{3} \rho^3$  and the time length of  $Q_\rho$  is  $\rho^2$ ,

$$\iint_{Q_\rho(z_0)} (|u|^3 + |p|^{3/2}) \, dx \, dt \leq \frac{4\pi}{3} \rho^5 (U_T^3 + P_T^{3/2}).$$

Division by  $\rho^2$  proves (284); its right-hand side is independent of the center and tends to zero with  $\rho$ , which proves (285).  $\square$

### 8.3 Vorticity and material flow

**Theorem 8.7** (Global vorticity amplification and dissipation). *Let  $\nu > 0$ , let  $u_0 \in H_\sigma^\infty(\mathbb{R}^3)$ , and let  $(u, p)$  be the pressure-normalized global pair supplied by Theorem 5.10. Put*

$$\omega := \nabla \times u, \quad \omega_0 := \nabla \times u_0, \quad L_T^\omega := S_{3,1,\infty} T^{1/2} \mathfrak{X}_{0,3}[u_0, \nu; T].$$

For every  $T < \infty$  and  $2 \leq r \leq \infty$ ,

$$\sup_{0 \leq t \leq T} \|\omega(t)\|_{L^r} \leq e^{L_T^\omega} \|\omega_0\|_{L^r}, \quad (286)$$

$$\int_0^T \|(\omega \cdot \nabla)u(t)\|_{L^r} dt \leq L_T^\omega e^{L_T^\omega} \|\omega_0\|_{L^r}. \quad (287)$$

If  $2 \leq r < \infty$ , then

$$\int_0^T \left\| \nabla |\omega(t)|^{r/2} \right\|_2^2 dt \leq \frac{r}{4\nu(r-1)} e^{rL_T^\omega} \|\omega_0\|_{L^r}^r. \quad (288)$$

*Proof.* Taking the curl of the momentum equation gives the three-dimensional vorticity equation

$$\partial_t \omega + (u \cdot \nabla) \omega = (\omega \cdot \nabla) u + \nu \Delta \omega. \quad (289)$$

Fix  $2 \leq r < \infty$ . Pair (289) with  $|\omega|^{r-2} \omega$ . To justify this power test, choose  $\chi \in C_c^\infty(\mathbb{R}^3)$ , with  $0 \leq \chi \leq 1$ ,  $\chi = 1$  on  $B_1$ , and  $\chi = 0$  outside  $B_2$ , put  $\chi_R(x) = \chi(x/R)$ , and first test against

$$\chi_R \Phi_\varepsilon(\omega), \quad \Phi_\varepsilon(z) := (|z|^2 + \varepsilon)^{(r-2)/2} z.$$

Define the primitive

$$F_\varepsilon(z) := \frac{(|z|^2 + \varepsilon)^{r/2} - \varepsilon^{r/2}}{r}. \quad (290)$$

Then  $\nabla_z F_\varepsilon = \Phi_\varepsilon$ , and incompressibility gives

$$\int_{\mathbb{R}^3} \chi_R (u \cdot \nabla) \omega \cdot \Phi_\varepsilon(\omega) dx = - \int_{\mathbb{R}^3} F_\varepsilon(\omega) u \cdot \nabla \chi_R dx. \quad (291)$$

With  $A_R := B_{2R} \setminus B_R$ ,

$$\left| \int F_\varepsilon(\omega) u \cdot \nabla \chi_R dx \right| \leq \frac{C_r}{R} \|u\|_\infty \left( \|\omega\|_{L^r(A_R)}^r + \varepsilon^{(r-2)/2} \|\omega\|_{L^2(A_R)}^2 \right), \quad (292)$$

$$\left| \int \nabla \chi_R \cdot \nabla \omega \Phi_\varepsilon(\omega) dx \right| \leq \frac{C_r}{R} \|\nabla \omega\|_{L^2(A_R)} \left( \|\omega\|_{L^{2r-2}(A_R)}^{r-1} + \varepsilon^{(r-2)/2} \|\omega\|_{L^2(A_R)} \right). \quad (293)$$

All norms on the right are finite because  $u, \omega$  belong to every spatial Sobolev space. Their annular tails vanish as  $R \rightarrow \infty$ , so (291) converges to the transport cancellation and the diffusion cutoff term vanishes.

The remaining diffusion form is nonnegative:

$$\begin{aligned} \nabla \omega : D \Phi_\varepsilon(\omega) \nabla \omega &= (|\omega|^2 + \varepsilon)^{(r-2)/2} |\nabla \omega|^2 \\ &\quad + (r-2)(|\omega|^2 + \varepsilon)^{(r-4)/2} \sum_{j=1}^3 (\omega \cdot \partial_j \omega)^2. \end{aligned} \quad (294)$$

After the limit  $R \rightarrow \infty$ , dominated convergence applies to  $F_\varepsilon(\omega)$  and to the stretching term, while Fatou's lemma applies to (294) as  $\varepsilon \downarrow 0$ . This yields the unregularized chain rule and diffusion inequality used below. The stretching term satisfies

$$\left| \int_{\mathbb{R}^3} (\omega \cdot \nabla) u \cdot |\omega|^{r-2} \omega \, dx \right| \leq \|\nabla u\|_\infty \|\omega\|_r^r.$$

For the diffusion term, spatial integration by parts gives

$$\begin{aligned} - \int_{\mathbb{R}^3} \Delta \omega \cdot |\omega|^{r-2} \omega \, dx &\geq (r-1) \int_{\mathbb{R}^3} |\omega|^{r-2} |\nabla |\omega||^2 \, dx \\ &= \frac{4(r-1)}{r^2} \left\| \nabla |\omega|^{r/2} \right\|_2^2. \end{aligned}$$

Hence

$$\frac{1}{r} \frac{d}{dt} \|\omega\|_r^r + \frac{4\nu(r-1)}{r^2} \left\| \nabla |\omega|^{r/2} \right\|_2^2 \leq \|\nabla u\|_\infty \|\omega\|_r^r. \quad (295)$$

Combining (295) with the quantitative rectangle estimate (350) gives

$$\int_0^T \|\nabla u(t)\|_\infty \, dt \leq L_T^\omega.$$

Gronwall's inequality proves (286) for finite  $r$ . Multiplying (295) by  $\exp(-r \int_0^t \|\nabla u(s)\|_\infty \, ds)$  and integrating proves (288). Since  $\omega_0, \omega(t) \in L^2 \cap L^\infty$ ,  $\|f\|_r \rightarrow \|f\|_\infty$  as  $r \rightarrow \infty$  for each such  $f$ . Consequently,

$$\sup_{0 \leq t \leq T} \|\omega(t)\|_\infty \leq \liminf_{r \rightarrow \infty} \sup_{0 \leq t \leq T} \|\omega(t)\|_r \leq e^{L_T^\omega} \lim_{r \rightarrow \infty} \|\omega_0\|_r = e^{L_T^\omega} \|\omega_0\|_\infty.$$

This is (286) at the endpoint. Finally,

$$\|(\omega \cdot \nabla) u(t)\|_r \leq \|\nabla u(t)\|_\infty \|\omega(t)\|_r;$$

integration and (286) give (287).  $\square$

**Theorem 8.8** (Quantitative volume-preserving material flow). *Let  $\nu > 0$ , let  $u_0 \in H_\sigma^\infty(\mathbb{R}^3)$ , and let  $(u, p)$  be the pressure-normalized global pair supplied by Theorem 5.10. For  $0 < T < \infty$ , put*

$$\begin{aligned} D_T &:= S_{2,0,\infty} T^{1/2} \mathfrak{X}_{0,2}[u_0, \nu; T], \\ L_T &:= S_{3,1,\infty} T^{1/2} \mathfrak{X}_{0,3}[u_0, \nu; T]. \end{aligned} \quad (296)$$

For every  $a \in \mathbb{R}^3$ , the problem

$$\partial_t X(t, a) = u(t, X(t, a)), \quad X(0, a) = a, \quad (297)$$

has a unique solution on  $[0, T]$ . For every  $t \in [0, T]$ , the map  $X(t, \cdot) : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  is a smooth volume-preserving diffeomorphism and

$$|X(t, a) - a| \leq D_T, \quad (298)$$

$$e^{-L_T} |a - b| \leq |X(t, a) - X(t, b)| \leq e^{L_T} |a - b|, \quad (299)$$

$$\det D_a X(t, a) = 1. \quad (300)$$

*Proof.* Equations (349) and (350) give

$$\begin{aligned} \int_0^T \|u(t)\|_\infty \, dt &\leq D_T, \\ \int_0^T \|\nabla u(t)\|_\infty \, dt &\leq L_T. \end{aligned}$$

Picard iteration on successive compact time intervals gives local existence and uniqueness for (297). The first integral bound prevents escape to spatial infinity and extends every trajectory to  $[0, T]$ ; it also proves (298). For two trajectories, the second integral bound and Gronwall's inequality give the upper bound in (299). Solving the same equation backward from time  $t$  produces the inverse map and gives the lower bound. Spatial differentiation of (297) and induction give smoothness of the map and its inverse. Finally, Jacobi's formula and  $\nabla \cdot u = 0$  give

$$\partial_t \det D_a X = (\nabla \cdot u)(t, X) \det D_a X = 0, \quad \det D_a X(0, a) = 1,$$

which proves (300).  $\square$

#### 8.4 Relative energy and the Leray–Hopf class

The weak class is stated in the whole-space form originating with Leray and Hopf [7, 6]. The argument is the classical weak–strong uniqueness mechanism associated with Serrin [12], written here against the pressure-complete smooth history and derived directly from the weak formulation and the energy inequality.

**Definition 8.9** (Three-dimensional Leray–Hopf solution). Let  $\nu > 0$ ,  $T > 0$ , and  $v_0 \in H_\sigma^0(\mathbb{R}^3)$ . A Leray–Hopf solution on  $[0, T]$  is a pair  $(v, q)$  such that

$$v \in C_w([0, T]; H_\sigma^0) \cap L^\infty(0, T; L_\sigma^2) \cap L^2(0, T; \dot{H}_\sigma^1), \quad q \in \mathcal{D}'((0, T) \times \mathbb{R}^3),$$

the equations

$$\partial_t v + (v \cdot \nabla)v + \nabla q = \nu \Delta v, \quad \nabla \cdot v = 0$$

hold in distributions,  $v(t) \rightarrow v_0$  strongly in  $L^2$  as  $t \downarrow 0$ , and

$$\frac{1}{2} \|v(t)\|_2^2 + \nu \int_0^t \|\nabla v(s)\|_2^2 \, ds \leq \frac{1}{2} \|v_0\|_2^2 \quad (301)$$

for every  $t \in [0, T]$ . Every divergence-free test field  $\phi \in C_c^\infty([0, T] \times \mathbb{R}^3; \mathbb{R}^3)$  satisfies

$$\int_0^T \left[ -\langle v, \partial_t \phi \rangle - \int_{\mathbb{R}^3} (v \otimes v) : \nabla \phi \, dx + \nu \langle \nabla v, \nabla \phi \rangle \right] dt = \langle v_0, \phi(0) \rangle. \quad (302)$$

A global Leray–Hopf solution at viscosity  $\nu$  is a pair on  $[0, \infty)$  whose restriction to every finite  $[0, T]$  is a Leray–Hopf solution in Definition 8.9, with the same datum and viscosity.

**Lemma 8.10** (Solenoidal cutoff density and time commutation). *Let  $\mathcal{A}$  be a finite subset of  $\{(r, p) : r \in \mathbb{N}_0, 2 \leq p < \infty\}$ . There are linear operators  $\mathcal{T}_k$ , independent of time, such that, for every  $f \in H_\sigma^\infty(\mathbb{R}^3)$ ,*

$$\mathcal{T}_k f \in C_{c,\sigma}^\infty(\mathbb{R}^3)$$

and, for every  $(r, p) \in \mathcal{A}$ ,

$$\sup_k \|\mathcal{T}_k f\|_{W^{r,p}} \leq C_{\mathcal{A}} \|f\|_{W^{r,p}}, \quad (303)$$

$$\|\mathcal{T}_k f - f\|_{W^{r,p}} \longrightarrow 0. \quad (304)$$

The convergence is uniform for  $f$  in a compact subset of the retained intersection of  $W^{r,p}$  spaces. If  $F \in C^\infty([0, T]; H_\sigma^\infty)$ , then

$$\partial_t^a (\mathcal{T}_k F) = \mathcal{T}_k (\partial_t^a F) \quad (a \in \mathbb{N}_0), \quad (305)$$

and the convergence in (304) is uniform in  $t$  for every fixed temporal derivative.

*Proof.* Choose  $\chi \in C_c^\infty(B_2)$  with  $\chi = 1$  on  $B_1$ , and put  $\chi_R(x) = \chi(x/R)$  and  $A_R = B_{2R} \setminus \overline{B_R}$ . For a smooth solenoidal  $f$ , set

$$g_R := \nabla \chi_R \cdot f.$$

It is supported in  $A_R$ , and

$$\int_{A_R} g_R \, dx = \int_{\mathbb{R}^3} \nabla \cdot (\chi_R f) \, dx = 0.$$

We now construct the required compact anti-divergence explicitly. Fix  $\rho \in C_c^\infty((-3, 3))$  with  $\int_{\mathbb{R}} \rho = 1$ . If  $h \in C_c^\infty(B_2)$  and  $\int h = 0$ , set

$$m_1(y_2, y_3) := \int_{\mathbb{R}} h(s, y_2, y_3) \, ds, \quad m_2(y_3) := \int_{\mathbb{R}} m_1(s, y_3) \, ds$$

and define  $\mathcal{G}h = (G_1, G_2, G_3)$  by

$$\begin{aligned} G_1(y) &:= \int_{-\infty}^{y_1} (h(s, y_2, y_3) - \rho(s)m_1(y_2, y_3)) \, ds, \\ G_2(y) &:= \rho(y_1) \int_{-\infty}^{y_2} (m_1(s, y_3) - \rho(s)m_2(y_3)) \, ds, \\ G_3(y) &:= \rho(y_1)\rho(y_2) \int_{-\infty}^{y_3} m_2(s) \, ds. \end{aligned}$$

The three integrands have zero total integral in their integration variables; for the last one this is  $\int h = 0$ . Thus  $\mathcal{G}h \in C_c^\infty((-3, 3)^3; \mathbb{R}^3)$ , and direct differentiation gives

$$\nabla \cdot \mathcal{G}h = h - \rho(y_1)m_1 + \rho(y_1)m_1 - \rho(y_1)\rho(y_2)m_2 + \rho(y_1)\rho(y_2)m_2 = h.$$

Integration over intervals of length at most six, Hölder's inequality, Fubini's theorem, and differentiation under the integral give, for  $a \in \mathbb{N}_0$  and  $1 < p < \infty$ ,

$$\|\mathcal{G}h\|_{W^{a,p}((-3,3)^3)} \leq C_{a,p,\rho} \|h\|_{W^{a,p}(B_2)}. \quad (306)$$

For  $g \in C_c^\infty(B_{2R})$  of mean zero, put

$$(\mathcal{G}_R g)(x) := R[\mathcal{G}(g(R \cdot))](x/R).$$

Then  $\nabla \cdot \mathcal{G}_R g = g$ , its support lies in  $(-3R, 3R)^3$ , and scaling (306) gives

$$\begin{aligned} \|\mathcal{G}_R g\|_{L^p} &\leq C_p R \|g\|_{L^p}, \\ \|\nabla^j \mathcal{G}_R g\|_{L^p} &\leq C_{j,p} \sum_{b=0}^j R^{1-j+b} \|\nabla^b g\|_{L^p}, \quad j \geq 1. \end{aligned} \quad (307)$$

Define

$$\mathcal{S}_R f := \chi_R f - \mathcal{G}_R g_R.$$

Then  $\mathcal{S}_R f \in C_{c,\sigma}^\infty(\mathbb{R}^3)$ , with support in  $(-3R, 3R)^3$ . Leibniz' rule gives, for  $0 \leq a \leq r-1$ ,

$$\|\nabla^a g_R\|_{L^p(A_R)} \leq C_a \sum_{b=0}^a R^{-1-a+b} \|\nabla^b f\|_{L^p(A_R)}.$$

Combining this estimate with (307) and the analogous cutoff estimate yields

$$\|\mathcal{S}_R f - f\|_{W^{r,p}} \leq C_{r,p} \sum_{b=0}^r (1 + R^{b-r}) \|\nabla^b f\|_{L^p(\mathbb{R}^3 \setminus B_R)}. \quad (308)$$

For  $r = 0$ , the same conclusion follows from (307) and  $\|g_R\|_p \leq CR^{-1} \|f\|_{L^p(A_R)}$ . The right side tends to zero as  $R \rightarrow \infty$ , and the same calculations give operator norms bounded independently of  $R$ .

Choose  $R_k \uparrow \infty$  and set  $\mathcal{T}_k := \mathcal{S}_{R_k}$ . Estimate (308) holds simultaneously for the finite family  $\mathcal{A}$ , while the preceding operator estimates give (303) simultaneously for that family. Strong convergence plus the uniform operator bound makes convergence uniform on compact subsets: cover a compact set by finitely many norm balls and use the bound on each remainder. Each  $\mathcal{T}_k$  is independent of time, so differentiation commutes with it. The ranges of  $\partial_t^\alpha F$  on  $[0, T]$  are compact in every retained space, which proves the final assertion.  $\square$

**Lemma 8.11** (Admissible strong-history test). *Let  $u$  be the global solution from Theorem 5.10, and let  $v$  be a Leray–Hopf solution on  $[0, T]$  at the same viscosity. Put  $w := v - u$ . Then, for every  $t \in [0, T]$ ,*

$$\begin{aligned} \langle v(t), u(t) \rangle &= \langle v_0, u_0 \rangle + \int_0^t \int_{\mathbb{R}^3} (w \cdot \nabla u) \cdot w \, dx \, ds \\ &\quad - 2\nu \int_0^t \langle \nabla v, \nabla u \rangle \, ds. \end{aligned} \quad (309)$$

*Proof.* The energy class and spatial interpolation give

$$v \in L^4(0, T; L^3), \quad \|v\|_{L^4 L^3}^4 \leq C \|v\|_{L^\infty L^2}^2 \|\nabla v\|_{L^2 L^2}^2,$$

so  $v \otimes v \in L^2(0, T; L^{3/2})$ .

Fix  $0 < t < T$ . Solenoidal density on  $\mathbb{R}^3$ , applied uniformly to the compact sets  $u([0, T])$  and  $u_t([0, T])$  in their finite Sobolev spaces, gives

$$u^{(k)} \in C^\infty([0, T]; C_{c,\sigma}^\infty(\mathbb{R}^3))$$

such that

$$u^{(k)} \longrightarrow u \quad \text{in} \quad W^{1,1}(0, T; L^2) \cap L^2(0, T; H^1) \cap L^2(0, T; W^{1,3}) \cap C([0, T]; L^2). \quad (310)$$

Apply Lemma 8.10 with

$$\mathcal{A} = \{(0, 2), (1, 2), (1, 3)\}$$

and set  $u^{(k)} = \mathcal{T}_k u$ . Its uniform compact-range conclusion for  $u$  and  $u_t$ , together with time commutation (305), gives (310).

For  $0 < h < T - t$ , choose  $\eta_h \in C_c^\infty([0, T])$  with  $0 \leq \eta_h \leq 1$ ,  $\eta_h = 1$  on  $[0, t]$ ,  $\eta_h = 0$  on  $[t + h, T]$ , and  $\eta_h' \leq 0$ . The field  $\eta_h u^{(k)}$  is an admissible test in (302). First let  $k \rightarrow \infty$ . The convergence in (310), together with  $v \otimes v \in L^2 L^{3/2}$ , permits passage in every term.

Choose the time cutoffs so that  $-\eta_h'$  is an approximate identity from the right at  $t$ , and let  $h \downarrow 0$ . Weak continuity of  $v$ , strong continuity of  $u$ , and the Lebesgue differentiation theorem give

$$\begin{aligned} \langle v(t), u(t) \rangle &= \langle v_0, u_0 \rangle + \int_0^t \langle v, u_t \rangle \, ds + \int_0^t \int_{\mathbb{R}^3} (v \otimes v) : \nabla u \, dx \, ds \\ &\quad - \nu \int_0^t \langle \nabla v, \nabla u \rangle \, ds. \end{aligned}$$

This identity first holds for almost every  $t$ . Its right-hand side is continuous in  $t$ , while  $t \mapsto \langle v(t), u(t) \rangle$  is continuous by weak continuity of  $v$  and norm continuity of  $u$ . Hence it holds for every  $0 \leq t \leq T$ , including  $t = T$  by passage from below. Using

$$u_t = -(u \cdot \nabla)u - \nabla p + \nu \Delta u$$

is admissible because  $v$  is divergence free. It gives

$$\int_0^t \langle v, u_t \rangle \, ds = - \int_0^t \int_{\mathbb{R}^3} v \cdot (u \cdot \nabla)u \, dx \, ds - \nu \int_0^t \langle \nabla v, \nabla u \rangle \, ds.$$

Finally,

$$\int_{\mathbb{R}^3} [(v \otimes v) : \nabla u - v \cdot (u \cdot \nabla)u] \, dx = \int_{\mathbb{R}^3} (w \cdot \nabla u) \cdot w \, dx,$$

because

$$\int_{\mathbb{R}^3} (w \cdot \nabla u) \cdot u \, dx = \frac{1}{2} \int_{\mathbb{R}^3} w \cdot \nabla |u|^2 \, dx = 0.$$

This proves (309).  $\square$

**Theorem 8.12** (Quantitative Leray–Hopf stability). *Let  $\nu > 0$  and  $u_0 \in H_\sigma^\infty(\mathbb{R}^3)$ , and let  $u$  be the global strong history from Theorem 5.10. Fix  $T > 0$ , and let  $v$  be any Leray–Hopf solution on  $[0, T]$  at the same viscosity  $\nu$ , with datum  $v_0 \in H_\sigma^0(\mathbb{R}^3)$ . Then, for  $0 \leq t \leq T$ ,*

$$\begin{aligned} &\frac{1}{2} \|v(t) - u(t)\|_2^2 + \nu \int_0^t \|\nabla(v - u)(s)\|_2^2 \, ds \\ &\leq \frac{1}{2} \|v_0 - u_0\|_2^2 \exp\left(2 \int_0^t \|\nabla u(s)\|_\infty \, ds\right) \end{aligned} \tag{311}$$

$$\leq \frac{1}{2} \|v_0 - u_0\|_2^2 \exp\left(2S_{3,1,\infty} T^{1/2} \mathfrak{P}_{0,2}[u_0, \nu; T]^{1/2}\right). \tag{312}$$

*The lower spatial rectangle also gives the viscosity-weighted estimate*

$$\begin{aligned} &\frac{1}{2} \|v(t) - u(t)\|_2^2 + \frac{\nu}{2} \int_0^t \|\nabla(v - u)(s)\|_2^2 \, ds \\ &\leq \frac{1}{2} \|v_0 - u_0\|_2^2 \exp\left(\frac{S_{2,0,\infty}^2}{\nu} \mathfrak{X}_{0,2}[u_0, \nu; T]^2\right). \end{aligned} \tag{313}$$

*Proof.* Combine (301), the strong energy equality (131), and (309). With  $w = v - u$ , this gives

$$\frac{1}{2} \|w(t)\|_2^2 + \nu \int_0^t \|\nabla w(s)\|_2^2 \, ds \leq \frac{1}{2} \|w(0)\|_2^2 - \int_0^t \int_{\mathbb{R}^3} (w \cdot \nabla u) \cdot w \, dx \, ds. \quad (314)$$

The integral term in (314) has absolute value at most

$$\|\nabla u(s)\|_\infty \|w(s)\|_2^2.$$

Gronwall's inequality bounds  $\|w(t)\|_2^2$ . Substitution of that bound back into (314) proves (311). Proposition 7.16 gives

$$\int_0^T \|\nabla u(t)\|_\infty \, dt \leq S_{3,1,\infty} T^{1/2} \mathfrak{P}_{0,2}[u_0, \nu; T]^{1/2},$$

which proves (312).

For the lower-rectangle estimate, integration by parts and  $\nabla \cdot w = 0$  give

$$\left| \int_{\mathbb{R}^3} (w \cdot \nabla u) \cdot w \, dx \right| = \left| \int_{\mathbb{R}^3} (w \cdot \nabla w) \cdot u \, dx \right| \leq \|u\|_\infty \|w\|_2 \|\nabla w\|_2.$$

Young's inequality in (314) gives

$$\frac{1}{2} \|w(t)\|_2^2 + \frac{\nu}{2} \int_0^t \|\nabla w(s)\|_2^2 \, ds \leq \frac{1}{2} \|w(0)\|_2^2 + \frac{1}{2\nu} \int_0^t \|u(s)\|_\infty^2 \|w(s)\|_2^2 \, ds.$$

Set

$$A(t) := \frac{1}{\nu} \int_0^t \|u(s)\|_\infty^2 \, ds.$$

Gronwall's inequality gives  $\|w(t)\|_2^2 \leq \|w(0)\|_2^2 e^{A(t)}$ . Substitution in the preceding inequality gives

$$\frac{1}{2} \|w(t)\|_2^2 + \frac{\nu}{2} \int_0^t \|\nabla w(s)\|_2^2 \, ds \leq \frac{1}{2} \|w(0)\|_2^2 e^{A(t)}.$$

Finally, (349) yields

$$A(t) \leq \frac{S_{2,0,\infty}^2}{\nu} \mathfrak{X}_{0,2}[u_0, \nu; T]^2,$$

which proves (313).  $\square$

**Corollary 8.13** (Explicit weak-class stability for small critical data). *Under the hypotheses of Theorem 8.3, let  $v$  be a global Leray–Hopf solution at the same viscosity  $\nu$ , with datum  $v_0 \in H_\sigma^0(\mathbb{R}^3)$ . Then*

$$\begin{aligned} & \frac{1}{2} \|v(t) - u(t)\|_2^2 + \nu \int_0^t \|\nabla(v - u)(s)\|_2^2 \, ds \\ & \leq \frac{1}{2} \|v_0 - a\|_2^2 \exp(2S_{3,1,\infty} t U_3^{\text{sm}}) \quad (t \geq 0). \end{aligned} \quad (315)$$

*Proof.* Equation (271) gives  $\sup_{s \geq 0} \|u(s)\|_{H^3} \leq U_3^{\text{sm}}$ . Hence

$$\int_0^t \|\nabla u(s)\|_\infty \, ds \leq S_{3,1,\infty} t U_3^{\text{sm}}.$$

The solution constructed in Theorem 8.3 and the global history from Theorem 5.10, with datum  $a$ , both restrict to members of  $\mathcal{M}_\nu^3(0, T; a)$  on every finite interval. The at-most-one assertion in Theorem 2.11 identifies the two pressure-normalized histories. Apply (311).  $\square$

**Corollary 8.14** (Uniqueness and energy equality in the Leray–Hopf class). *Let  $\nu > 0$  and  $u_0 \in H^\infty_\sigma(\mathbb{R}^3)$ . If  $(v, q)$  is any global three-dimensional Leray–Hopf solution at viscosity  $\nu$ , with datum  $u_0$ , then  $v$  agrees on every finite interval with the velocity of the global solution from Theorem 5.10. Thus  $v$  has the smooth representative*

$$u \in C^\infty([0, \infty) \times \mathbb{R}^3; \mathbb{R}^3),$$

and it satisfies

$$\frac{1}{2} \|u(t)\|_2^2 + \nu \int_0^t \|\nabla u(s)\|_2^2 \, ds = \frac{1}{2} \|u_0\|_2^2 \quad (t \geq 0).$$

Its distributional pressure has the form

$$q = p + c(t) \quad \text{in } \mathcal{D}'((0, \infty) \times \mathbb{R}^3), \quad (316)$$

where  $p$  is the pressure in (130) and  $c$  is a scalar distribution in time. The canonical Riesz representative  $R_i R_j(u_i u_j)$  equals  $p$ . If  $q$  is represented by a locally integrable function with  $q(t, \cdot) \in C_0(\mathbb{R}^3)$  for almost every  $t$ , then  $q = p$  almost everywhere.

*Proof.* Set  $v_0 = u_0$  in Theorem 8.12. The right-hand side of (312) is zero, so  $v = u$  in  $C_w([0, T]; L^2) \cap L^2(0, T; \dot{H}^1)$ . Since  $T$  is arbitrary, the identity holds globally. Smoothness and energy equality follow from (127) and (131). If  $q$  is a distributional pressure paired with this velocity and  $p$  is the pressure in (130), subtraction of the two momentum equations gives

$$\nabla(q - p) = 0 \quad \text{in } \mathcal{D}'((0, \infty) \times \mathbb{R}^3).$$

A distribution on a product domain whose spatial gradient vanishes is the pullback of a scalar distribution in time: testing first against spatial test functions of integral zero and then against one fixed test function of integral one gives (316). The Riesz formula for  $p$  proves the canonical-representative assertion. In the function-valued case,  $q(t, x) - p(t, x) = c(t)$  for almost every  $(t, x)$ ; both spatial slices tend to zero as  $|x| \rightarrow \infty$ , so  $c(t) = 0$  for almost every  $t$ .  $\square$

## 9 Conclusion

The proof follows one pressure-normalized velocity history from its initial datum to the endpoint of its maximal smooth interval. Every finite temporal depth and every finite Sobolev depth determine a whole-terminal rectangle, and the restriction identities assemble their shared entries without changing the history. The rectangle estimates then give  $u \in L^2(0, T_*; H^m)$  for every finite  $m$ . The equation upgrades the zeroth row to an  $H^m$ -valued endpoint, and the pressure formula together with the velocity recurrence recovers the complete endpoint jet. Local well-posedness restarts from that state, while uniqueness on the overlap keeps the restarted solution on the history already generated by the datum. If  $T_*$  were finite, this continuation would contradict maximality. Hence the maximal interval is  $[0, \infty)$ .

The same rectangles also quantify the critical-height rows, frequency tails, Lipschitz action, and relative-energy stability, with constants depending on the datum, viscosity, and horizon. In the small-critical regime, the scale-invariant norm closes the critical stock/action estimate, and the higher rectangles admit explicit formulas using the corresponding finite collection of higher datum norms.

The resulting global solution is smooth, satisfies the exact energy identity, and carries the decay-normalized pressure. Relative energy then identifies every Leray–Hopf solution with the same datum with this smooth velocity, and the pressure is unique up to the unavoidable time-dependent gauge unless the same decay normalization is imposed. The endpoint contradiction and these consequences all belong to the one fixed-viscosity evolution constructed from  $u_0$ .

## A Canonical coordinate catalogue

The estimates in Section 7 are organized by their role in the argument. For reference, the following theorem collects their canonical-coordinate forms in one place.

**Theorem A.1** (Canonical finite-horizon coordinate estimates). *Under the hypotheses and notation of Definition 7.2, let  $J_{n,\alpha} := \partial_x^\alpha V_n$  and  $\Pi_{n,\alpha} := \partial_x^\alpha Q_n$ , as in Theorems 7.10 and 7.11. Let  $n, s, k, q \in \mathbb{N}_0$ ,  $\alpha \in \mathbb{N}_0^3$ , and set  $r := s + |\alpha|$ . Then*

$$\|J_{n,\alpha}\|_{L^2(I_T; H^{s+1})} \leq \mathfrak{X}_{n,r+1}, \quad (317)$$

$$\|J_{n+1,\alpha}\|_{L^2(I_T; H^{s-1})} \leq \mathfrak{X}_{n+1,r-1}, \quad (318)$$

$$\sup_{0 \leq t \leq T} \|J_{n,\alpha}(t)\|_{H^s} \leq \mathfrak{Z}_{n,r}, \quad (319)$$

$$\|J_{n,\alpha}^T\|_{H^s} \leq \mathfrak{Z}_{n,r}, \quad (320)$$

$$\|J_{n,\alpha}(t) - J_{n,\alpha}(r_0)\|_{H^{s-1}} \leq \min \left\{ |t - r_0|^{1/2} \mathfrak{X}_{n+1,r-1}, |t - r_0| \mathfrak{Z}_{n+1,r} \right\}, \quad (321)$$

$$\|J_{n,\alpha}(t) - J_{n,\alpha}(r_0)\|_{H^s} \leq \min \left\{ |t - r_0|^{1/2} \mathfrak{X}_{n+1,r}, |t - r_0| \mathfrak{Z}_{n+1,r} \right\}, \quad (322)$$

for  $0 \leq r_0 \leq t \leq T$ .

For  $0 \leq \theta \leq 1$ ,

$$\|J_{n,\alpha}(t) - J_{n,\alpha}(r_0)\|_{H^{s-\theta}} \leq 2^{1-\theta} \mathfrak{Z}_{n,r}^{1-\theta} \mathfrak{X}_{n+1,r-1}^\theta |t - r_0|^{\theta/2}. \quad (323)$$

If  $1 \leq a \leq \infty$ ,  $2 \leq b \leq \infty$ , and  $q > k + \frac{3}{2} - \frac{3}{b}$ , then

$$\left\| \nabla^k J_{n,\alpha} \right\|_{L^a(I_T; L^b)} \leq S_{q,k,b} T^{1/a} \mathfrak{Z}_{n,q+|\alpha|}, \quad (324)$$

$$\left\| \nabla^k J_{n,\alpha} \right\|_{L^2(I_T; L^b)} \leq S_{q,k,b} \mathfrak{X}_{n,q+|\alpha|}, \quad (325)$$

where  $T^{1/\infty} = 1$ .

The pressure rows obey

$$\|\Pi_{n,\alpha}\|_{L^1(I_T; H^s)} + \|\nabla \Pi_{n,\alpha}\|_{L^1(I_T; H^{s-1})} \leq \mathfrak{Q}_{n,r}^{(1)}, \quad (326)$$

$$\|\Pi_{n,\alpha}\|_{L^2(I_T; H^s)} \leq \mathfrak{Q}_{n,r}^{(2)}, \quad (327)$$

$$\sup_{0 \leq t \leq T} \|\Pi_{n,\alpha}(t)\|_{H^s} \leq \mathfrak{Q}_{n,r}^{(\infty)}, \quad (328)$$

$$\|\Pi_{n,\alpha}^T\|_{H^s} \leq \mathfrak{Q}_{n,r}^{(\infty)}, \quad (329)$$

$$\|\Pi_{n,\alpha}(t) - \Pi_{n,\alpha}(r_0)\|_{H^s} \leq \min \left\{ |t - r_0|^{1/2} \mathfrak{Q}_{n,r}^{(\text{mod})}, |t - r_0| \mathfrak{Q}_{n+1,r}^{(\infty)} \right\}. \quad (330)$$

Under the preceding restrictions on  $a, b, k, q$ ,

$$\left\| \nabla^k \Pi_{n,\alpha} \right\|_{L^a(I_T; L^b)} \leq S_{q,k,b} T^{1/a} \mathfrak{Q}_{n,q+|\alpha|}^{(\infty)}. \quad (331)$$

Let  $r_1 \in \mathbb{N}$  and  $K \geq 1$ . Then

$$\|\mathbf{P}_{\geq K} J_{n,\alpha}\|_{L^2(I_T; H^s)} \leq K^{-r_1} \mathfrak{X}_{n,s+r_1+|\alpha|}, \quad (332)$$

$$\sup_{0 \leq t \leq T} \|\mathbf{P}_{\geq K} J_{n,\alpha}(t)\|_{H^s} \leq K^{-r_1} \mathfrak{Z}_{n,s+r_1+|\alpha|}, \quad (333)$$

$$\|\mathbf{P}_{\geq K} \Pi_{n,\alpha}\|_{L^1(I_T; H^s)} \leq K^{-r_1} \mathfrak{Q}_{n,s+r_1+|\alpha|}^{(1)}, \quad (334)$$

$$\sup_{0 \leq t \leq T} \|\mathbf{P}_{\geq K} \Pi_{n,\alpha}(t)\|_{H^s} \leq K^{-r_1} \mathfrak{Q}_{n,s+r_1+|\alpha|}^{(\infty)}. \quad (335)$$

For  $j \in \mathbb{N}_0$ ,

$$\sup_{0 \leq t \leq T} E_{j+1/2}(t) \leq \frac{1}{2} \mathfrak{Z}_{0,j+1}^2, \quad (336)$$

$$\int_0^T E_{j+1/2}(t) dt \leq \frac{1}{2} \mathfrak{X}_{0,j+1}^2. \quad (337)$$

If  $m > j + \frac{1}{2}$  is an integer, then

$$\sup_{0 \leq t \leq T} E_{j+1/2, \geq K}(t) \leq \frac{1}{2} K^{2j+1-2m} \mathfrak{Z}_{0,m}^2, \quad (338)$$

$$\int_0^T E_{j+1/2, \geq K}(t) dt \leq \frac{1}{2} K^{2j+1-2m} \mathfrak{X}_{0,m}^2. \quad (339)$$

For  $k \in \mathbb{N}$ ,

$$\|H^{(k)}\|_{L^1(0,T)} \leq \frac{1}{2} \sum_{a=0}^k \binom{k}{a} \mathfrak{X}_{a,1} \mathfrak{X}_{k-a,1}. \quad (340)$$

For  $j, \ell \in \mathbb{N}_0$ , define

$$\begin{aligned} \mathfrak{E}_{j,\ell}^{[1]} &:= \frac{1}{2} \sum_{a=0}^{\ell} \binom{\ell}{a} \mathfrak{X}_{a,j+1} \mathfrak{X}_{\ell-a,j+1}, \\ \mathfrak{E}_{j,\ell}^{[\infty]} &:= \frac{1}{2} \sum_{a=0}^{\ell} \binom{\ell}{a} \mathfrak{Z}_{a,j+1} \mathfrak{Z}_{\ell-a,j+1}, \end{aligned} \quad (341)$$

$$\begin{aligned} \mathfrak{E}_{j,\ell}^{[1]} &:= \mathfrak{E}_{j,\ell+1}^{[1]} + 2\nu \mathfrak{E}_{j+1,\ell}^{[1]}, \\ \mathfrak{E}_{j,\ell}^{[\infty]} &:= \mathfrak{E}_{j,\ell+1}^{[\infty]} + 2\nu \mathfrak{E}_{j+1,\ell}^{[\infty]}. \end{aligned} \quad (342)$$

Then the complete differentiated energy and production rows obey

$$\left\| \partial_t^\ell E_{j+1/2} \right\|_{L^1(0,T)} \leq \mathfrak{E}_{j,\ell}^{[1]}, \quad \sup_{0 \leq t \leq T} \left| \partial_t^\ell E_{j+1/2}(t) \right| \leq \mathfrak{E}_{j,\ell}^{[\infty]}, \quad (343)$$

$$\left\| \partial_t^\ell \mathcal{P}_{j+1/2} \right\|_{L^1(0,T)} \leq \mathfrak{E}_{j,\ell}^{[1]}, \quad \sup_{0 \leq t \leq T} \left| \partial_t^\ell \mathcal{P}_{j+1/2}(t) \right| \leq \mathfrak{E}_{j,\ell}^{[\infty]}. \quad (344)$$

Every current in (64) now has a finite-order majorant retaining temporal depth, spatial height,

viscosity, and horizon. In particular,

$$\|H'\|_{L^1(0,T)} \leq \mathfrak{X}_{1,1}\mathfrak{X}_{0,1}, \quad (345)$$

$$\int_0^T \|\Lambda^{3/2}u(t)\|_2^2 dt \leq \mathfrak{X}_{0,2}^2, \quad (346)$$

$$\|\mathcal{P}_{1/2}\|_{L^1(0,T)} \leq \mathfrak{X}_{1,1}\mathfrak{X}_{0,1} + \nu\mathfrak{X}_{0,2}^2, \quad (347)$$

$$|H(T) - H(t)| \leq \min \left\{ \mathfrak{Z}_{0,1}\mathfrak{X}_{1,1}(T-t)^{1/2}, \mathfrak{Z}_{0,1}\mathfrak{Z}_{1,1}(T-t) \right\}, \quad 0 \leq t \leq T. \quad (348)$$

Finally,

$$\|u\|_{L^2(I_T; L^\infty)} \leq S_{2,0,\infty}\mathfrak{X}_{0,2}, \quad (349)$$

$$\int_0^T \|\nabla u(t)\|_\infty dt \leq S_{3,1,\infty}T^{1/2}\mathfrak{X}_{0,3}, \quad (350)$$

$$\int_0^T \|\nabla \times u(t)\|_\infty dt \leq 2S_{3,1,\infty}T^{1/2}\mathfrak{X}_{0,3}, \quad (351)$$

$$\sup_{0 \leq t \leq T} \|u(t)\|_{H^m}^2 \leq \|u_0\|_{H^m}^2 \exp\left(2K_m S_{3,1,\infty}T^{1/2}\mathfrak{X}_{0,3}\right), \quad m \geq 3. \quad (352)$$

Every right-hand side is an explicit algebraic expression in the displayed analytic constants, the initial-jet recursion (176)–(177), and the finite collection of canonical rectangle coordinates  $\mathfrak{B}_{a,b}[u_0, \nu; T]$  selected by the stated indices. Equations (175) and (178)–(215) specify every intermediate operation. Equation (204) rewrites each such rectangle coordinate through the finite height-coordinate recursion (194)–(202). Since (196) first bounds those height coordinates by the rectangle coordinates, this is a re-expression of the same finite same-history information rather than a new estimate from the critical-height rows. This is an exact coordinate-level catalogue for general data; it does not claim a second, independent bound of those rectangle coordinates by a finite list of datum norms.

Proposition 7.6 displays the exact powers of  $\nu$  and the rescaled horizon  $\nu T$ . For every fixed choice of the displayed indices, the definitions select only finitely many rectangle coordinates. Under the explicit critical smallness condition (266), Proposition 8.4 eliminates the rectangle coordinates and gives finite formulas solely in the displayed datum norms,  $\nu$ ,  $T$ ,  $N$ , and  $M$ .

*Proof.* The Fourier multiplier inequality

$$\|\partial_x^\alpha f\|_{H^\ell} \leq \|f\|_{H^{\ell+|\alpha|}},$$

followed by Lemma 7.8, proves (317)–(320). The Banach identity

$$J_{n,\alpha}(t) - J_{n,\alpha}(r_0) = \int_{r_0}^t J_{n+1,\alpha}(\tau) d\tau$$

in the two indicated Sobolev spaces and Cauchy–Schwarz prove the square-root branches of (321) and (322). The same identity, followed by the pointwise bound (209) for  $V_{n+1}$ , proves their linear branches. Interpolation between the low modulus and

$$\|J_{n,\alpha}(t) - J_{n,\alpha}(r_0)\|_{H^s} \leq 2\mathfrak{Z}_{n,r}$$

proves (323). The fixed Sobolev embedding constant in (168) proves (324) and (325).

Apply the recurrence

$$Q_n = R_i R_j \sum_{a=0}^n \binom{n}{a} V_{a,i} V_{n-a,j}$$

term by term at order  $r$ , using (166). Cauchy–Schwarz in time gives (326); placing either factor in  $L_t^\infty H_x^{r+1}$  gives (327); placing both factors there gives (328) and (329). After subtracting the recurrence at times  $t$  and  $r_0$ , split each product difference into two terms and apply (322); this gives the square-root branch of (330). Since  $\partial_t \Pi_{n,\alpha} = \Pi_{n+1,\alpha}$ , the Banach identity and (328) at row  $n+1$  give its linear branch. Sobolev embedding proves (331).

For every  $f \in H^{\ell+r_1}$ ,

$$\|P_{\geq K} f\|_{H^\ell} \leq K^{-r_1} \|f\|_{H^{\ell+r_1}}.$$

Apply this inequality at spatial order  $s+r_1$  to (208), (209), (326), and (328). The four resulting estimates are (332)–(335).

The inequalities

$$\|f\|_{\dot{H}^{j+1/2}} \leq \|f\|_{H^{j+1}}, \quad |\xi|^{2j+1} \leq K^{2j+1-2m} |\xi|^{2m} \quad (|\xi| \geq K),$$

give (336)–(339). The exact identity

$$H^{(k)}(t) = \frac{1}{2} \sum_{a=0}^k \binom{k}{a} \operatorname{Re} \left\langle \Lambda^{1/2} V_a(t), \Lambda^{1/2} V_{k-a}(t) \right\rangle_{L^2}$$

and Cauchy–Schwarz in time give (340) and (345). More generally,

$$\partial_t^\ell E_{j+1/2} = \frac{1}{2} \sum_{a=0}^\ell \binom{\ell}{a} \operatorname{Re} \left\langle \Lambda^{j+1/2} V_a, \Lambda^{j+1/2} V_{\ell-a} \right\rangle_{L^2}.$$

Cauchy–Schwarz in time and the pointwise estimates (208)–(209) prove (343). Differentiating  $\mathcal{P}_{j+1/2} = \partial_t E_{j+1/2} + 2\nu E_{j+3/2}$  then proves (344). The next two estimates follow from

$$\mathcal{P}_{1/2} = H' + \nu \left\| \Lambda^{3/2} u \right\|_2^2.$$

For the terminal critical-height modulus,

$$|H(T) - H(t)| \leq \frac{1}{2} (\|u(T)\|_{\dot{H}^{1/2}} + \|u(t)\|_{\dot{H}^{1/2}}) \|u(T) - u(t)\|_{\dot{H}^{1/2}},$$

and the Banach identity in  $H^1$  prove the square-root branch of (348).

The identity

$$H' = \operatorname{Re} \left\langle \Lambda^{1/2} V_1, \Lambda^{1/2} V_0 \right\rangle_{L^2},$$

together with (209), proves its linear branch.

Sobolev embedding and (208) prove (349)–(351). Lemma 2.8 gives

$$\frac{d}{dt} \|u\|_{H^m}^2 \leq 2K_m \|\nabla u\|_\infty \|u\|_{H^m}^2.$$

Gronwall’s inequality and (350) prove (352). □

## References

- [1] J.-P. Aubin, *Un théorème de compacité*, C. R. Acad. Sci. Paris **256** (1963), 5042–5044.
- [2] L. Caffarelli, R. Kohn, and L. Nirenberg, *Partial regularity of suitable weak solutions of the Navier–Stokes equations*, Communications on Pure and Applied Mathematics **35** (1982), 771–831, doi:10.1002/cpa.3160350604.
- [3] L. Escauriaza, G. A. Seregin, and V. Šverák,  *$L_{3,\infty}$ -solutions of Navier–Stokes equations and backward uniqueness*, Russian Mathematical Surveys **58** (2003), no. 2, 211–250, doi:10.1070/RM2003v058n02ABEH000609.
- [4] C. Fefferman, *Existence and smoothness of the Navier–Stokes equation*, in *The Millennium Prize Problems*, Clay Mathematics Institute, American Mathematical Society, 2006, pp. 57–67, Clay problem statement.
- [5] H. Fujita and T. Kato, *On the Navier–Stokes initial value problem. I*, Archive for Rational Mechanics and Analysis **16** (1964), 269–315, doi:10.1007/BF00276188.
- [6] E. Hopf, *Über die Anfangswertaufgabe für die hydrodynamischen Grundgleichungen*, Mathematische Nachrichten **4** (1950/51), 213–231, doi:10.1002/mana.3210040121.
- [7] J. Leray, *Sur le mouvement d’un liquide visqueux emplissant l’espace*, Acta Mathematica **63** (1934), 193–248, doi:10.1007/BF02547354.
- [8] E. H. Lieb, *Sharp constants in the Hardy–Littlewood–Sobolev and related inequalities*, Annals of Mathematics **118** (1983), 349–374, doi:10.2307/2007032.
- [9] T. Kato, *Strong  $L^p$ -solutions of the Navier–Stokes equation in  $\mathbb{R}^m$ , with applications to weak solutions*, Mathematische Zeitschrift **187** (1984), 471–480, EuDML record.
- [10] H. Koch and D. Tataru, *Well-posedness for the Navier–Stokes equations*, Advances in Mathematics **157** (2001), no. 1, 22–35, doi:10.1006/aima.2000.1937.
- [11] J.-L. Lions and E. Magenes, *Non-Homogeneous Boundary Value Problems and Applications*, vol. I, Springer, 1972, doi:10.1007/978-3-642-65161-8.
- [12] J. Serrin, *On the interior regularity of weak solutions of the Navier–Stokes equations*, Archive for Rational Mechanics and Analysis **9** (1962), 187–195, doi:10.1007/BF00253344.
- [13] J. Simon, *Compact sets in the space  $L^p(0, T; B)$* , Annali di Matematica Pura ed Applicata **146** (1987), 65–96, doi:10.1007/BF01762360.
- [14] T. Tao, *Localisation and compactness properties of the Navier–Stokes global regularity problem*, Analysis & PDE **6** (2013), 25–107, doi:10.2140/apde.2013.6.25.