

Global Smoothness for the Forced Three-Dimensional Incompressible Navier–Stokes Equations

Thomas Birnie

September 2026

Abstract

Fix a smooth divergence-free finite-energy datum u_0 , a viscosity $\nu > 0$, and a prescribed $f \in C^\infty([0, \infty); \mathcal{S}(\mathbb{R}^3; \mathbb{R}^3))$. The resulting forced three-dimensional incompressible Navier–Stokes evolution is carried as one pressure-normalized history. For every finite pair of temporal and spatial orders, the datum generates a compatible whole-terminal rectangle containing the corresponding adjacent rows of its velocity–pressure–force mixed jet. Each rectangle keeps every selected derivative norm finite through the chosen finite endpoint, and restriction compatibility assembles the rectangles into the complete space–time intersection. Independently, the critical-height construction and the positive-sign viscous tether supply the relationship identities on the same coordinates. Pairing, squaring, and integrating the complete forced rows keeps viscosity, the full binomial transport term, normalized pressure, and forcing inside the estimates. Since every finite mixed order lies in one of these rectangles, adjacent rows produce one H^∞ endpoint, the forced recurrences determine its complete pressure-normalized jet, and local theory restarts the original history with the same prescribed absolute-time force. No finite maximal time can occur, and the forced solution is globally smooth and unique.

Contents

1	Introduction	3
2	From a Prescribed Force to Terminal Continuation	4
3	Analytic framework	7
4	The pressure-complete setting and mixed jet	19
4.1	The datum-generated whole-terminal intersection	23
5	Finite projections and compatible rectangles	25
6	Intersection coordinates, endpoint jet, and restart	33
7	Matrix forms of the completed intersection	42
8	Quantitative consequences of the compatible rectangles	45
8.1	Completion in the global H^1 mild class	45
8.2	Canonical coordinates and finite majorants	48
8.3	Mixed velocity and pressure rows	56
8.4	Frequency and critical-scale consequences	58
8.5	Continuation and quantitative restart	61
9	Three-dimensional applications	62
9.1	Explicit closure for small critical forced data	62
9.2	Low-rectangle and scale-local bounds	67
9.3	Vorticity and material flow	68
9.4	Relative energy and the Leray–Hopf class	71
10	Conclusion	77
A	Canonical coordinate catalogue	77
	References	81

1 Introduction

Fix one viscosity $\nu > 0$, one divergence-free datum $u_0 \in H^\infty(\mathbb{R}^3; \mathbb{R}^3)$, and one prescribed source

$$f \in C^\infty([0, \infty); \mathcal{S}(\mathbb{R}^3; \mathbb{R}^3)).$$

Forced local theory produces a unique pressure-normalized classical history (u, p) on a maximal interval $[0, T_*)$. This paper follows that one history. Its question is whether a finite value of T_* can terminate the solution generated by (u_0, ν, f) .

Theorem. *For every $\nu > 0$, every $u_0 \in H^\infty_\sigma(\mathbb{R}^3)$, and every $f \in C^\infty([0, \infty); \mathcal{S}(\mathbb{R}^3; \mathbb{R}^3))$, there is a unique pressure-normalized pair*

$$(u, p) \in C^\infty([0, \infty) \times \mathbb{R}^3; \mathbb{R}^3 \times \mathbb{R})$$

satisfying

$$\partial_t u + (u \cdot \nabla)u + \nabla p = \nu \Delta u + f, \quad \nabla \cdot u = 0, \quad u(0) = u_0, \quad \lim_{|x| \rightarrow \infty} p(t, x) = 0.$$

It obeys the forced energy identity

$$\frac{1}{2} \|u(t)\|_{L^2}^2 + \nu \int_0^t \|\nabla u(s)\|_{L^2}^2 \, ds = \frac{1}{2} \|u_0\|_{L^2}^2 + \int_0^t \langle f(s), u(s) \rangle_{L^2} \, ds \quad (t \geq 0).$$

Corollary 1.1 (Weak–strong uniqueness and pressure normalization). *For the solution of the preceding theorem, every forced Leray–Hopf pair (v, q) with the same datum, viscosity, and prescribed source has $v = u$. Distributionally, $q = p + c(t)$ for a scalar distribution c ; if $q(t, \cdot) \in C_0(\mathbb{R}^3)$ almost everywhere, then $q = p$ almost everywhere.*

Leray’s weak solutions establish global finite-energy existence [11]. The regularity results of Serrin [14], Caffarelli–Kohn–Nirenberg [4], and Escauriaza–Seregin–Šverák [5] identify increasingly sharp ways in which a singularity would have to evade classical control. Fujita–Kato and Kato provide the local strong theory used both at the initial time and at the terminal restart [7, 9]. The proof below works directly with the complete derivative structure of the prescribed forced history.

Forcing changes every level of that structure. Its solenoidal part $g = \mathbb{P}f$ enters the projected velocity equation. Its gradient part is absorbed by the normalized pressure,

$$-\Delta p = \partial_i \partial_j (u_i u_j) - \nabla \cdot f, \quad p = R_i R_j (u_i u_j) - (-\Delta)^{-1} \nabla \cdot f.$$

Every temporal row contains the corresponding derivative $F_n = \partial_t^n f$; every energy row contains force work; and a restart at T_* retains the same absolute-time source. These are parts of a single forced velocity–pressure–source history.

The proof has two constructions from the same pressure-complete mixed jet. For every finite pair of temporal and spatial orders, the datum-generated branch produces a compatible whole-terminal norm rectangle containing the selected rows and their adjacent rows. That rectangle is the quantitative mechanism: it keeps every selected derivative norm finite through the chosen finite endpoint. Restriction compatibility assembles the cofinal family of finite rectangles into the complete space–time intersection. Independently, the critical-height tower and the positive-sign viscous diagonal relation supply the relationship identities on the same coordinates. Pairing, squaring, and integrating the complete forced rows keeps every face of each rectangle inside one finite estimate.

Because every mixed derivative belongs to a finite rectangle, the adjacent rows have one common smooth endpoint. The pressure formula and forced recurrences determine the complete endpoint jet, and local theory restarts the equation from $u_* = u(T_*)$ with the same source. Overlap uniqueness then contradicts a finite maximal time.

Section 2 derives this architecture in one view. The analytic framework and the mixed-jet construction follow, then the independent whole-terminal intersection and rectangle branches, their composition, and the endpoint continuation. The remaining sections record matrix formulations and quantitative consequences of the same finite rectangles.

2 From a Prescribed Force to Terminal Continuation

Fix one divergence-free datum u_0 , one viscosity $\nu > 0$, and one prescribed smooth force f . Let (u, p) be the corresponding maximal classical solution on $[0, T_*)$. The proof never changes this history. Every derivative, finite rectangle, endpoint value, and restarted solution below belongs to the trajectory generated by the original triple (u_0, ν, f) .

The force enters the proof in two linked forms. Its solenoidal part $g = \mathbb{P}f$ acts in the projected velocity equation, while its gradient part is fixed by the normalized pressure:

$$\partial_t u + \mathbb{P}\nabla \cdot (u \otimes u) - \nu \Delta u = g, \quad p = R_i R_j (u_i u_j) - (-\Delta)^{-1} \nabla \cdot f.$$

This is why the forced argument must begin with the source. Appending f after constructing an unforced velocity would change both the pressure and the history being differentiated.

The Navier–Stokes scaling acts on the complete forced equation. If (u, p, f) is a solution, then

$$u_\lambda(t, x) = \lambda u(\lambda^2 t, \lambda x), \quad p_\lambda(t, x) = \lambda^2 p(\lambda^2 t, \lambda x), \quad f_\lambda(t, x) = \lambda^3 f(\lambda^2 t, \lambda x)$$

solves the equation with transformed datum $u_{0,\lambda}(x) = \lambda u_0(\lambda x)$. For every real s ,

$$\|\Lambda^s u_\lambda(t)\|_2^2 = \lambda^{2s-1} \|\Lambda^s u(\lambda^2 t)\|_2^2.$$

The invariant spatial height is therefore $s = \frac{1}{2}$. This identifies the critical row of the same forced differential structure; it does not replace the prescribed history by its rescaling.

Write

$$V_n = \partial_t^n u, \quad Q_n = \partial_t^n p, \quad F_n = \partial_t^n f,$$

and let

$$B_n = \sum_{a=0}^n \binom{n}{a} (V_a \cdot \nabla) V_{n-a}.$$

Repeated differentiation of the original forced equation gives the complete temporal row

$$V_{n+1} + B_n + \nabla Q_n = \nu \Delta V_n + F_n, \tag{1}$$

$$-\Delta Q_n = \partial_i \partial_j \sum_{a=0}^n \binom{n}{a} (V_a)_i (V_{n-a})_j - \nabla \cdot F_n. \tag{2}$$

Spatial differentiation distributes through both velocity factors and through the prescribed source. The fields

$$\partial_x^\alpha V_n, \quad \partial_x^\alpha Q_n, \quad \partial_x^\alpha F_n$$

form the pressure-complete forced mixed jet. They are coordinates of one solution, rather than a collection of independently chosen functions.

The mixed jet is read in two different ways. The first is the datum-generated compatible space–time intersection. On every strict horizon it gives

$$u \in \bigcap_{r,m \geq 0} H^r(0, T; H_\sigma^m(\mathbb{R}^3)) \quad \text{for every } T < T_*.$$

The datum-generated whole-terminal intersection theorem carries these compatible representatives to a proposed finite maximal interval $I = (0, T_*)$:

$$u \in \bigcap_{r,m \geq 0} H^r(I; H_\sigma^m(\mathbb{R}^3)). \quad (\text{A})$$

The second reading begins independently from the critical height and the viscous diagonal relation. It produces finite rectangles of complete equation rows, each carrying transport, pressure, viscosity, and the prescribed force. The whole-terminal intersection and the rectangle relationships are independently derived from the same mixed jet. Their composition evaluates the complete rectangle family on all of I .

The viscous relation is visible already in one row. Project (1), apply Λ^s , and use $\Delta = -\Lambda^2$:

$$\Lambda^s V_{n+1} + \nu \Lambda^{s+2} V_n = \Lambda^s \mathbb{P}(F_n - B_n). \quad (\text{B})$$

Squaring this equality does not separate the nonlinearity from the fluid. The cross term is the derivative of the adjacent spatial norm, so

$$\begin{aligned} \|\Lambda^s V_{n+1}\|_2^2 + \nu^2 \|\Lambda^{s+2} V_n\|_2^2 + \nu \frac{d}{dt} \|\Lambda^{s+1} V_n\|_2^2 \\ = \|\Lambda^s \mathbb{P}(F_n - B_n)\|_2^2. \end{aligned} \quad (\text{C})$$

The complementary projection gives the pressure allocation

$$\nabla Q_n = (I - \mathbb{P})(F_n - B_n),$$

and orthogonality recombines the complete source:

$$\|\Lambda^s \mathbb{P}(F_n - B_n)\|_2^2 + \|\Lambda^s \nabla Q_n\|_2^2 = \|\Lambda^s (F_n - B_n)\|_2^2. \quad (\text{D})$$

Thus the rectangle keeps viscosity, transport, pressure, and forcing in the same estimate.

The critical-height tower is another reading of this viscous tether. For

$$E_{r,s} = \frac{1}{2} \|\Lambda^s V_r\|_2^2,$$

the complete row is

$$\partial_t E_{r,s} + 2\nu E_{r,s+1} = \mathcal{P}_{r,s} + \mathcal{W}_{r,s}, \quad \mathcal{W}_{r,s} = \text{Re} \langle \Lambda^s V_r, \Lambda^s F_r \rangle. \quad (\text{E})$$

At $r = 0$ and $s = \frac{1}{2}$, differentiating (E) repeatedly exposes $E_{3/2}, E_{5/2}, \dots$, while the differentiated transport, pressure, and force work remains in the same completed row. The tower therefore describes the geometry among mixed-jet coordinates; it does not replace the velocity field by a scalar height. The first two spatial heights are, explicitly,

$$\begin{aligned} H' + \nu \left\| \Lambda^{3/2} u \right\|_2^2 &= \mathcal{P}_{1/2} + \mathcal{W}_{1/2}, \\ E'_{3/2} + \nu \left\| \Lambda^{5/2} u \right\|_2^2 &= \mathcal{P}_{3/2} + \mathcal{W}_{3/2}, \end{aligned}$$

and therefore

$$H'' = 4\nu^2 E_{5/2} - 2\nu(\mathcal{P}_{3/2} + \mathcal{W}_{3/2}) + \partial_t(\mathcal{P}_{1/2} + \mathcal{W}_{1/2}).$$

These are the actual $s = \frac{1}{2}$ and $s = \frac{3}{2}$ rows. Their transport, pressure, viscosity, and force coordinates remain joined.

For finite N, M , select the coordinates supplied by (A),

$$\{V_n, V_{n+1}, Q_n, F_n : 0 \leq n \leq N\}$$

at the spatial orders required by (B)–(E), and write

$$\mathfrak{R}_{N,M}(T) := \sum_{n=0}^N \left(\|V_n\|_{L^2(0,T;H^{M+1})}^2 + \|V_{n+1}\|_{L^2(0,T;H^{M-1})}^2 \right). \quad (\text{F})$$

For each finite (N, M) , (A) first gives the finite whole-terminal sum

$$C_{N,M} := \sum_{n=0}^N \left(\|V_n\|_{L^2(0,T_*;H^{M+1})}^2 + \|V_{n+1}\|_{L^2(0,T_*;H^{M-1})}^2 \right) < \infty.$$

Restriction to $(0, T)$ and monotone exhaustion of the nonnegative norm integrals then give

$$\sup_{0 < T < T_*} \mathfrak{R}_{N,M}(T) = \lim_{T \uparrow T_*} \mathfrak{R}_{N,M}(T) = C_{N,M} < \infty \quad (N, M < \infty). \quad (\text{G})$$

The constant may depend on the selected rectangle. No estimate uniform in the infinitely many indices is used. Thus (G) is the finite projection of the already constructed intersection, not an assumption placed on a local solution. The positive-sign square (C), together with (D), then evaluates the adjacent temporal, viscous, and pressure faces on those supplied coordinates. Sobolev multiplication is applied within that same finite rectangle to its binomial transport face B_n ; it does not replace the nonlinear term or introduce a second premise.

Three objects meet here without being confused. The intersection supplies the coordinates and their compatibility. The equations (B)–(E) supply their viscous and critical-height relationships. Pairing, squaring, and integrating those relationships gives the complete-equation rectangle estimates. The nonlinearity remains one face of the same forced rectangle throughout.

(A) now carries the same complete equation to the terminal step. Its zeroth temporal projection gives, for every finite m ,

$$u \in L^2(0, T_*; H^{m+2}).$$

Sobolev multiplication and the prescribed force then give

$$\begin{aligned} \|u_t\|_{L^1(0,T_*;H^m)} &\leq \nu\sqrt{T_*} \|u\|_{L^2(0,T_*;H^{m+2})} \\ &+ C_m \|u\|_{L^2(0,T_*;H^{m+2})}^2 + \|g\|_{L^1(0,T_*;H^m)} < \infty. \end{aligned}$$

Hence

$$u \in W^{1,1}(0, T_*; H^m) \hookrightarrow C([0, T_*]; H^m) \quad (m \geq 0).$$

All these traces come from the same velocity and therefore determine one terminal state

$$u_* = u(T_*) \in \bigcap_{m \geq 0} H_\sigma^m(\mathbb{R}^3).$$

Passing (1)–(2) to T_* determines every temporal and pressure coordinate from u_* and the values $F_n(T_*)$. Local theory restarts from the condition $w(0) = u_*$. In restart time $\tau = t - T_*$, the source is $f(T_* + \tau)$: this is the same prescribed absolute-time force, not a new history. Uniqueness joins the restarted solution to the incoming one and extends it past T_* , contradicting maximality.

The rest of the paper proves the mixed-jet construction, its datum-generated whole-terminal intersection, and the independent critical-height and rectangle identities in their mathematical order. Their composition carries the one forced history through the infinite family of finite rectangle estimates, the compatible endpoint jet, and the same-force restart.

3 Analytic framework

The argument has two analytic joints. Products and Fourier multipliers must keep every differentiated velocity–pressure row in a common Sobolev scale; at the terminal time, an H^3 local theory must restart that same pressure-normalized history. We record the estimates needed at those two joints, with the constants later used in the quantitative bounds.

Definition 3.1. For $f \in \mathcal{S}(\mathbb{R}^3)$, set

$$\widehat{f}(\xi) := (2\pi)^{-3/2} \int_{\mathbb{R}^3} e^{-ix \cdot \xi} f(x) \, dx, \quad \langle \xi \rangle := (1 + |\xi|^2)^{1/2}.$$

This integral defines $\widehat{f}$ for $f \in L^1(\mathbb{R}^3)$. The Fourier transform on $\mathcal{S}'(\mathbb{R}^3)$ is its extension by duality. For $s \in \mathbb{R}$, define

$$H^s(\mathbb{R}^3) := \left\{ f \in \mathcal{S}'(\mathbb{R}^3) : \langle \xi \rangle^s \widehat{f} \in L^2(\mathbb{R}^3) \right\},$$

and define the finite homogeneous-norm domain by

$$\dot{H}^s(\mathbb{R}^3) := \left\{ f \in \mathcal{S}'(\mathbb{R}^3) : \begin{array}{l} \widehat{f} \text{ is represented by a measurable function on } \mathbb{R}^3, \\ |\xi|^s \widehat{f} \in L^2(\mathbb{R}^3) \end{array} \right\}.$$

On these domains set

$$\|f\|_{H^s}^2 := \int_{\mathbb{R}^3} \langle \xi \rangle^{2s} \left| \widehat{f}(\xi) \right|^2 \, d\xi, \quad \|f\|_{\dot{H}^s}^2 := \int_{\mathbb{R}^3} |\xi|^{2s} \left| \widehat{f}(\xi) \right|^2 \, d\xi.$$

Set $A := (1 - \Delta)^{1/2}$, $\Lambda := (-\Delta)^{1/2}$,

$$H_\sigma^s := \{v \in H^s(\mathbb{R}^3; \mathbb{R}^3) : \nabla \cdot v = 0\}, \quad H^\infty := \bigcap_{m \in \mathbb{N}_0} H^m, \quad H_\sigma^\infty := \bigcap_{m \in \mathbb{N}_0} H_\sigma^m.$$

$$\mathcal{S}_\sigma(\mathbb{R}^3) := \{v \in \mathcal{S}(\mathbb{R}^3; \mathbb{R}^3) : \nabla \cdot v = 0\}.$$

For a vector or tensor $F = (F_I)_I$, all Sobolev norms are componentwise:

$$\|F\|_{H^s}^2 := \sum_I \|F_I\|_{H^s}^2, \quad \|F\|_{\dot{H}^s}^2 := \sum_I \|F_I\|_{\dot{H}^s}^2.$$

Repeated Latin indices are summed from 1 to 3.

Lemma 3.2 (Sobolev nesting and embedding). *If $s_2 \geq s_1$, then*

$$\|f\|_{H^{s_1}} \leq \|f\|_{H^{s_2}}.$$

If $k \in \mathbb{N}_0$ and $s > k + 3/2$, then

$$\max_{|\alpha| \leq k} \|\partial^\alpha f\|_{L^\infty} \leq C_{s,k}^{\text{emb}} \|f\|_{H^s},$$

where

$$C_{s,k}^{\text{emb}} := (2\pi)^{-3/2} \max_{0 \leq j \leq k} \left(\int_{\mathbb{R}^3} |\xi|^{2j} \langle \xi \rangle^{-2s} d\xi \right)^{1/2} < \infty.$$

In particular,

$$H^\infty(\mathbb{R}^3) \hookrightarrow C_b^\infty(\mathbb{R}^3).$$

Proof. The first assertion follows from $\langle \xi \rangle^{2s_1} \leq \langle \xi \rangle^{2s_2}$. For $|\alpha| \leq k$, Fourier inversion and Cauchy–Schwarz give

$$|\partial^\alpha f(x)| \leq (2\pi)^{-3/2} \left(\int_{\mathbb{R}^3} |\xi|^{2|\alpha|} \langle \xi \rangle^{-2s} d\xi \right)^{1/2} \|f\|_{H^s}.$$

The integral is finite exactly when $s > |\alpha| + 3/2$. Approximation by Schwartz functions and dominated convergence give continuous representatives. For each k , choose $s = k + 2$. $\square$

Lemma 3.3 (Explicit three-dimensional homogeneous Sobolev constants). *For $3 \leq p \leq 6$, set*

$$s(p) := 3 \left(\frac{1}{2} - \frac{1}{p} \right)$$

and

$$S(p) := 2^{-s(p)} \pi^{-s(p)/2} \left(\frac{\Gamma((3 - 2s(p))/2)}{\Gamma((3 + 2s(p))/2)} \right)^{1/2} \left(\frac{\Gamma(3)}{\Gamma(3/2)} \right)^{s(p)/3}. \quad (3)$$

Then every scalar, vector, or finite tensor field F in $\dot{H}^{s(p)}(\mathbb{R}^3)$ satisfies

$$\|F\|_{L^p(\mathbb{R}^3)} \leq S(p) \|F\|_{\dot{H}^{s(p)}(\mathbb{R}^3)}. \quad (4)$$

In particular,

$$S(3) = 2^{-1/6} \pi^{-1/3}, \quad S(6) = \frac{2^{2/3}}{\sqrt{3} \pi^{2/3}}, \quad S(6)S(3) = \frac{1}{\pi} \sqrt{\frac{2}{3}}.$$

Proof. For scalar fields, (4) is the sharp three-dimensional homogeneous Sobolev inequality obtained from the sharp Hardy–Littlewood–Sobolev inequality [12]. After duality, insertion of the Riesz-potential kernel constant, and conversion to the unitary Fourier transform fixed above, the squared sharp Sobolev constant for the multiplier $\widehat{\Lambda^s f}(\xi) = |\xi|^s \widehat{f}(\xi)$ is

$$2^{-2s} \pi^{-s} \frac{\Gamma((3 - 2s)/2)}{\Gamma((3 + 2s)/2)} \left(\frac{\Gamma(3)}{\Gamma(3/2)} \right)^{2s/3}. \quad (5)$$

Taking the square root in (5) and putting $p = 6/(3 - 2s)$ gives (3)–(4). For a finite component array $F = (F_I)_I$, Minkowski’s inequality in $L^{p/2}$ gives

$$\|F\|_p^2 = \left\| \sum_I |F_I|^2 \right\|_{p/2} \leq \sum_I \|F_I\|_p^2 \leq S(p)^2 \sum_I \|F_I\|_{\dot{H}^{s(p)}}^2.$$

The two endpoint formulas follow by evaluating the gamma factors at $s = 1/2$ and $s = 1$. $\square$

Lemma 3.4 (Sobolev multiplication). *Set*

$$H_4 := (2\pi)^{-3/4} \left(\int_{\mathbb{R}^3} \langle \xi \rangle^{-4} d\xi \right)^{1/4}, \quad C_0^{\text{alg}} := H_4^2.$$

For $m \geq 1$, define

$$\begin{aligned} E_m &:= \max_{0 \leq j \leq \lfloor m/2 \rfloor} C_{j+2,j}^{\text{emb}}, & C_m^{\text{mix}} &:= 13^{m/2} E_m, \\ C_m^{\text{F}} &:= 2^m (2\pi)^{-3/2} \left(\int_{\mathbb{R}^3} \langle \xi \rangle^{-2m} d\xi \right)^{1/2} & (m \geq 2), \end{aligned}$$

and set

$$C_1^{\text{alg}} := C_1^{\text{mix}}, \quad C_m^{\text{alg}} := \max\{C_m^{\text{mix}}, C_m^{\text{F}}\} \quad (m \geq 2).$$

All these constants are finite and explicit. For every $m \in \mathbb{N}_0$,

$$\|fg\|_{H^m} \leq C_m^{\text{alg}} \|f\|_{H^{m+1}} \|g\|_{H^{m+1}}.$$

For $m \geq 2$,

$$\|fg\|_{H^m} \leq C_m^{\text{alg}} \|f\|_{H^m} \|g\|_{H^m}.$$

Proof. Hausdorff–Young and Hölder give

$$\|f\|_4 \leq (2\pi)^{-3/4} \left\| \widehat{f} \right\|_{4/3} \leq H_4 \|f\|_{H^1}.$$

Thus $\|fg\|_2 \leq C_0^{\text{alg}} \|f\|_{H^1} \|g\|_{H^1}$.

Let $m \geq 1$. The exact integer-order identity

$$\|h\|_{H^m}^2 = \sum_{|\alpha| \leq m} \frac{m!}{(m - |\alpha|)! \alpha!} \|\partial^\alpha h\|_2^2$$

follows by expanding $(1 + |\xi|^2)^m$. In each Leibniz term for $\partial^\alpha(fg)$, place the factor carrying at most $\lfloor |\alpha|/2 \rfloor$ derivatives in L^∞ . The definition of E_m , Sobolev nesting, and $\sum_{\beta \leq \alpha} \binom{\alpha}{\beta} = 2^{|\alpha|}$ give

$$\|\partial^\alpha(fg)\|_2 \leq E_m 2^{|\alpha|} \|f\|_{H^{m+1}} \|g\|_{H^{m+1}}.$$

The multinomial identity

$$\sum_{|\alpha| \leq m} \frac{m!}{(m - |\alpha|)! \alpha!} 4^{|\alpha|} = 13^m$$

proves the first estimate with C_m^{mix} .

For $m \geq 2$, the inequalities

$$\langle \eta + \zeta \rangle^m \leq 2^{m-1} (\langle \eta \rangle^m + \langle \zeta \rangle^m), \quad \left\| \widehat{f} \right\|_1 \leq \left(\int_{\mathbb{R}^3} \langle \xi \rangle^{-2m} d\xi \right)^{1/2} \|f\|_{H^m},$$

combined with $\widehat{fg} = (2\pi)^{-3/2} \widehat{f} * \widehat{g}$ and Young's convolution inequality give

$$\|fg\|_{H^m} \leq C_m^{\text{F}} \|f\|_{H^m} \|g\|_{H^m}.$$

The chosen maximum proves both asserted estimates with this constant. $\square$

Lemma 3.5 (Riesz and Leray multipliers). *Let*

$$r_j(\xi) := \begin{cases} i\xi_j/|\xi|, & \xi \neq 0, \\ 0, & \xi = 0, \end{cases} \quad P(\xi) := \begin{cases} I - \frac{\xi \otimes \xi}{|\xi|^2}, & \xi \neq 0, \\ I, & \xi = 0, \end{cases}$$

and

$$\widehat{R_j f}(\xi) = r_j(\xi) \widehat{f}(\xi), \quad \widehat{\mathbb{P}f}(\xi) = P(\xi) \widehat{f}(\xi).$$

For all $s \in \mathbb{R}$,

$$\|R_j f\|_{H^s} \leq \|f\|_{H^s}, \quad \|\mathbb{P}f\|_{H^s} \leq \|f\|_{H^s}.$$

If $m \in \mathbb{N}_0$ and $h \in L^1(\mathbb{R}^3) \cap H^m(\mathbb{R}^3; \mathbb{R}^3)$, then

$$\|(-\Delta)^{-1} \nabla \cdot h\|_{H^m} \leq C_m^{\text{fp}} (\|h\|_{L^1} + \|h\|_{H^m}). \quad (6)$$

If $m \in \mathbb{N}_0$ and $v \in H_\sigma^{m+1}$, then

$$q := \sum_{i,j=1}^3 R_i R_j (v_i v_j) \in H^m, \quad -\Delta q = \sum_{i,j=1}^3 \partial_i \partial_j (v_i v_j),$$

and

$$\mathbb{P} \nabla \cdot (v \otimes v) = (v \cdot \nabla) v + \nabla q.$$

Here

$$(v \otimes v)_{ij} := v_i v_j, \quad (\nabla \cdot (v \otimes v))_i := \sum_{j=1}^3 \partial_j (v_i v_j), \quad ((v \cdot \nabla) v)_i := \sum_{j=1}^3 v_j \partial_j v_i.$$

Proof. The multiplier symbols have operator norm at most one. Plancherel proves the first two estimates. For the force potential, split Fourier space at $|\xi| = 1$. On the unit ball, $|\widehat{h}(\xi)| \leq (2\pi)^{-3/2} \|h\|_{L^1}$ and $\int_{|\xi| \leq 1} |\xi|^{-2} d\xi < \infty$ in three dimensions. Outside the unit ball, $|\xi|^{-1} \leq 1$, so the H^m norm of the multiplier is bounded by the H^m norm of h . This proves (6). Lemma 3.4 gives $v_i v_j \in H^m$. The remaining identities follow after multiplication by $|\xi|^2$ and by the symbol of $\mathbb{P}$. $\square$

Lemma 3.6 (Mixed Leibniz identities). *For $N \in \mathbb{N}_0$ and $\alpha \in \mathbb{N}_0^3$,*

$$\partial_t^N \partial_x^\alpha (fg) = \sum_{a=0}^N \sum_{\beta \leq \alpha} \binom{N}{a} \binom{\alpha}{\beta} (\partial_t^a \partial_x^\beta f) (\partial_t^{N-a} \partial_x^{\alpha-\beta} g).$$

Proof. Apply the one-variable Leibniz identity successively in t, x_1, x_2, x_3 . $\square$

Lemma 3.7 (Adjacent Hilbert-scale trace). *Let $0 < T < \infty$, $m \in \mathbb{R}$, and*

$$v \in L^2(0, T; H^{m+1}), \quad \partial_t v \in L^2(0, T; H^{m-1}).$$

Then v has a unique representative in $C([0, T]; H^m)$, and, for $0 \leq r \leq t \leq T$,

$$\|v(t)\|_{H^m}^2 - \|v(r)\|_{H^m}^2 = 2 \operatorname{Re} \int_r^t \langle A^{m-1} \partial_t v(s), A^{m+1} v(s) \rangle_{L^2} ds, \quad (7)$$

$$\left| \|v(t)\|_{H^m}^2 - \|v(r)\|_{H^m}^2 \right| \leq 2 \|\partial_t v\|_{L^2(r, t; H^{m-1})} \|v\|_{L^2(r, t; H^{m+1})}, \quad (8)$$

$$\sup_{0 \leq t \leq T} \|v(t)\|_{H^m}^2 \leq T^{-1} \|v\|_{L^2(0, T; H^m)}^2 + 2 \|\partial_t v\|_{L^2(0, T; H^{m-1})} \|v\|_{L^2(0, T; H^{m+1})}. \quad (9)$$

Proof. Let S_R denote Fourier restriction to $\{|\xi| \leq R\}$. For $v_R = S_R v$,

$$\frac{d}{dt} \|v_R(t)\|_{H^m}^2 = 2 \operatorname{Re} \langle A^{m-1} \partial_t v_R(t), A^{m+1} v_R(t) \rangle_{L^2}.$$

Integration and Cauchy–Schwarz prove (7)–(8) for v_R . Applied to $S_R v - S_S v$, this estimate shows that $(v_R)_R$ is Cauchy in $C([0, T]; H^m)$; choose one Lebesgue time for the initial Cauchy term. Its limit equals v in $L^2(0, T; H^m)$. Passage to the limit proves the first two formulas. Choose $r \in (0, T)$ with $\|v(r)\|_{H^m}^2 \leq T^{-1} \|v\|_{L^2 H^m}^2$ and apply (8). $\square$

Lemma 3.8 (Explicit integer transport commutator). *Set*

$$\begin{aligned} \Gamma_{1,1} &:= 1, \\ \Gamma_{\ell,k} &:= (2\ell - 4 + \sqrt{3})^{(k-1)(\ell-k)} \quad (\ell \geq 2, 1 \leq k \leq \ell), \end{aligned} \quad (10)$$

$$K_m := \sum_{\ell=1}^m 3^\ell \binom{m}{\ell} \sum_{k=1}^{\ell} \binom{\ell}{k} \Gamma_{\ell,k} \quad (m \geq 3). \quad (11)$$

Every $v \in H_\sigma^{m+1}(\mathbb{R}^3)$ satisfies

$$|\operatorname{Re} \langle A^m \mathbb{P} \nabla \cdot (v \otimes v), A^m v \rangle_{L^2}| \leq K_m \|\nabla v\|_{L^\infty} \|v\|_{H^m}^2. \quad (12)$$

In particular,

$$K_m \leq (7^m - 1)(2m - 4 + \sqrt{3})^{(m-1)^2/4}. \quad (13)$$

Proof. We first prove the interpolation inequality used in the commutator. For the full ordered derivative tensor $\nabla^j F$, fix $M \geq 1$ and set

$$p_{M,0} := \infty, \quad p_{M,j} := \frac{2M}{j} \quad (1 \leq j \leq M), \quad A_M := 2M - 2 + \sqrt{3}.$$

Then

$$\|\nabla^j F\|_{L^{p_{M,j}}} \leq A_M^{j(M-j)/2} \|F\|_\infty^{1-j/M} \|\nabla^M F\|_2^{j/M} \quad (0 \leq j \leq M). \quad (14)$$

The endpoint cases are identities. For $1 \leq j \leq M-1$, put

$$G := \nabla^{j-1} F, \quad W := \nabla G, \quad p := p_{M,j}.$$

All contractions below are the Euclidean contractions of the corresponding ordered tensors. Integration by parts gives

$$\begin{aligned} \int_{\mathbb{R}^3} |W|^p &= - \int_{\mathbb{R}^3} |W|^{p-2} G : \Delta G \\ &\quad - (p-2) \int_{\mathbb{R}^3} |W|^{p-4} \sum_{q=1}^3 (G : W_{\cdot q})(W : \partial_q W). \end{aligned}$$

The pointwise bounds

$$|\Delta G| \leq \sqrt{3} |\nabla^2 G|, \quad \left| \sum_{q=1}^3 (G : W_{\cdot q})(W : \partial_q W) \right| \leq |G| |W|^2 |\nabla^2 G|$$

give

$$\int_{\mathbb{R}^3} |W|^p \leq (p-2 + \sqrt{3}) \int_{\mathbb{R}^3} |G| |W|^{p-2} |\nabla^2 G|.$$

Since

$$\frac{1}{p_{M,j-1}} + \frac{p-2}{p} + \frac{1}{p_{M,j+1}} = 1,$$

Hölder's inequality, with $X_j := \|\nabla^j F\|_{L^{p_{M,j}}}$, gives

$$X_j^2 \leq \left(\frac{2M}{j} - 2 + \sqrt{3} \right) X_{j-1} X_{j+1} \leq A_M X_{j-1} X_{j+1}. \quad (15)$$

The numbers $d_j := j(M-j)/2$ obey

$$2d_j - d_{j-1} - d_{j+1} = 1, \quad d_0 = d_M = 0.$$

The discrete maximum principle applied to (15) gives

$$X_j \leq A_M^{d_j} X_0^{1-j/M} X_M^{j/M},$$

which is (14). The integration by parts is justified first with $(|W|^2 + \varepsilon)^{(p-2)/2}$, followed by $\varepsilon \downarrow 0$; smooth approximation handles zero tensor norms.

For $|\alpha| \leq m$, set

$$\alpha! := \alpha_1! \alpha_2! \alpha_3!, \quad c_{m,\alpha} := \frac{m!}{(m-|\alpha|)! \alpha!}.$$

The multiplier $A^{2m} = (1 - \Delta)^m$ and the multinomial identity give the exact pairing formula

$$\langle A^m f, A^m g \rangle_{L^2} = \sum_{|\alpha| \leq m} c_{m,\alpha} \langle \partial^\alpha f, \partial^\alpha g \rangle_{L^2}. \quad (16)$$

The Leray projection is self-adjoint, commutes with A^m , and fixes $A^m v$. Since $\nabla \cdot (v \otimes v) = (v \cdot \nabla) v$, it follows that

$$\langle A^m \mathbb{P} \nabla \cdot (v \otimes v), A^m v \rangle_{L^2} = \sum_{|\alpha| \leq m} c_{m,\alpha} \langle \partial^\alpha ((v \cdot \nabla) v), \partial^\alpha v \rangle_{L^2}. \quad (17)$$

For each α ,

$$\partial^\alpha ((v \cdot \nabla) v) = (v \cdot \nabla) \partial^\alpha v + \sum_{0 < \beta \leq \alpha} \binom{\alpha}{\beta} (\partial^\beta v \cdot \nabla) \partial^{\alpha-\beta} v.$$

The divergence-free transport term has zero real pairing with $\partial^\alpha v$, because

$$\operatorname{Re} \langle (v \cdot \nabla) \partial^\alpha v, \partial^\alpha v \rangle_{L^2} = \frac{1}{2} \int_{\mathbb{R}^3} v \cdot \nabla |\partial^\alpha v|^2 \, dx = 0.$$

For a remaining term, put

$$\ell := |\alpha|, \quad k := |\beta|.$$

If $\ell \geq 2$, apply (14) to $F = \nabla v$, with

$$M = \ell - 1, \quad j = k - 1.$$

The two Hölder exponents are $p_{M,j}$ and $p_{M,M-j}$, whose reciprocal sum is $1/2$. The two factors are subarrays of the corresponding ordered derivative tensors, so

$$\begin{aligned} \left\| (\partial^\beta v \cdot \nabla) \partial^{\alpha-\beta} v \right\|_2 &\leq \left\| \partial^\beta v \right\|_{L^{p_{M,j}}} \left\| \nabla \partial^{\alpha-\beta} v \right\|_{L^{p_{M,M-j}}} \\ &\leq \Gamma_{\ell,k} \left\| \nabla v \right\|_\infty \left\| \nabla^\ell v \right\|_2. \end{aligned} \tag{18}$$

For $\ell = 1$, this formula follows from the direct L^∞ - L^2 estimate and $\Gamma_{1,1} = 1$. Since

$$\left\| \nabla^\ell v \right\|_2 = \|v\|_{\dot{H}^\ell} \leq \|v\|_{H^m}, \quad \|\partial^\alpha v\|_2 \leq \|v\|_{H^m},$$

each remaining pairing is bounded by

$$\Gamma_{\ell,k} \left\| \nabla v \right\|_\infty \|v\|_{H^m}^2.$$

The multinomial identities

$$\sum_{|\alpha|=\ell} c_{m,\alpha} = 3^\ell \binom{m}{\ell}, \quad \sum_{\substack{\beta \leq \alpha \\ |\beta|=k}} \binom{\alpha}{\beta} = \binom{\ell}{k}$$

now give (12) with the constant (11). Finally,

$$\sum_{k=1}^{\ell} \binom{\ell}{k} \leq 2^\ell, \quad (k-1)(\ell-k) \leq \frac{(\ell-1)^2}{4},$$

and summation of $\binom{m}{\ell} 3^\ell 2^\ell$ over $1 \leq \ell \leq m$ proves (13). Approximation by divergence-free Schwartz fields justifies all integrations by parts for $v \in H_\sigma^{m+1}$. $\square$

The preceding estimates control the differentiated rows as they approach an endpoint. To continue their common limit as the original solution, we now fix a time-translation-invariant local class that admits both late preterminal slices and the endpoint restart, and in which uniqueness holds on every overlap.

Definition 3.9 (Shifted pressure-normalized H^3 mild class). Fix the prescribed force $f \in C^\infty([0, \infty); \mathcal{S}(\mathbb{R}^3; \mathbb{R}^3))$ and set $g = \mathbb{P}f$. Let $0 \leq s < b < \infty$ and $a \in H_\sigma^3(\mathbb{R}^3)$. Define $\mathcal{M}_{\nu,f}^3(s, b; a)$ to be the set of pairs (w, q) such that

$$w \in C([s, b]; H_\sigma^3(\mathbb{R}^3)), \quad q \in C([s, b]; H^3(\mathbb{R}^3)), \tag{19}$$

$$w(s) = a, \quad q(t) = R_i R_j (w_i(t) w_j(t)) - (-\Delta)^{-1} \nabla \cdot f(t), \tag{20}$$

The remaining defining condition is the Duhamel identity: for every $t \in [s, b]$,

$$w(t) = e^{\nu(t-s)\Delta}a + \int_s^t e^{\nu(t-r)\Delta}g(r) \, dr - \int_s^t e^{\nu(t-r)\Delta}\mathbb{P}\nabla \cdot (w \otimes w)(r) \, dr \quad \text{in } H_\sigma^3(\mathbb{R}^3). \quad (21)$$

The integral is an H_σ^3 -valued Bochner integral: multiplication in $H^3(\mathbb{R}^3)$ places $\mathbb{P}\nabla \cdot (w \otimes w)$ in $C([s, b]; H_\sigma^2)$, and the heat multiplier from H^2 to H^3 has an integrable $(t-r)^{-1/2}$ singularity. The maximal lifespan in this class is

$$\tau_{\nu, f, 3}^{\max}(s, a) := \sup \{b - s : b > s, \mathcal{M}_{\nu, f}^3(s, b; a) \neq \emptyset\} \in [0, \infty]. \quad (22)$$

Lemma 3.10 (Restriction and classical admission). *If $(w, q) \in \mathcal{M}_{\nu, f}^3(s, b; a)$ and $s \leq r < c \leq b$, then*

$$(w, q)|_{[r, c]} \in \mathcal{M}_{\nu, f}^3(r, c; w(r)). \quad (23)$$

If (u, p) is a pressure-complete classical history in the sense of Definition 4.2 and $[s, b]$ is contained in its interval of existence, then

$$(u, p)|_{[s, b]} \in \mathcal{M}_{\nu, f}^3(s, b; u(s)). \quad (24)$$

Proof. Split (21) at r and use the heat semigroup identity. This gives, for $r \leq t \leq c$,

$$w(t) = e^{\nu(t-r)\Delta}w(r) + \int_r^t e^{\nu(t-\rho)\Delta}g(\rho) \, d\rho - \int_r^t e^{\nu(t-\rho)\Delta}\mathbb{P}\nabla \cdot (w \otimes w)(\rho) \, d\rho.$$

The pressure formula is unchanged under restriction. For a classical history, apply $\mathbb{P}$ to the momentum equation and use variation of constants; the decaying pressure formula gives (20). $\square$

Theorem 3.11 (Quantitative local restart). *Set $C_q := \sqrt{3}C_3^{\text{alg}}$. For $\nu > 0$, set*

$$L_\nu := 4(1 + \nu + \nu^{-1}) \exp\left(\frac{\nu + \nu^{-1}}{2}\right), \quad c_\nu := (8C_q L_\nu^2)^{-2}.$$

For every $s \geq 0$, put

$$\Gamma_s := \|\mathbb{P}f\|_{L^2(s, s+1; H^2)}.$$

For every $a \in H_\sigma^3(\mathbb{R}^3)$ and every

$$0 < \delta \leq \delta_{\nu, f}(s, a) := \min\{1, c_\nu(1 + \|a\|_{H^3} + \Gamma_s)^{-2}\},$$

the class $\mathcal{M}_{\nu, f}^3(s, s + \delta; a)$ contains exactly one pair (w, q) . Its velocity satisfies

$$w \in C([s, s + \delta]; H_\sigma^3) \cap L^2(s, s + \delta; H^4), \quad w_t \in L^2(s, s + \delta; H^2).$$

If $a \in H_\sigma^\infty$, then

$$\partial_t^n w, \partial_t^n q \in C([s, s + \delta]; H^m) \quad (n, m \in \mathbb{N}_0).$$

For every finite $b > s$, the class $\mathcal{M}_{\nu, f}^3(s, b; a)$ contains at most one pair. Its compatible members on strict subintervals of $[s, s + \tau_{\nu, f, 3}^{\max}(s, a))$ define the maximal shifted pressure-normalized H^3 mild history from a .

Proof. Translate the initial time to zero while retaining the shifted prescribed source $r \mapsto f(s+r)$. It is therefore enough to write the proof for $s = 0$. Let $0 < \delta \leq 1$, let

$$z_t - \nu \Delta z = F, \quad z(0) = a, \quad a \in H_\sigma^3, \quad F \in L^2(0, \delta; H_\sigma^2),$$

and set

$$A_0 := \|a\|_{H^3}, \quad G := \|F\|_{L^2(0, \delta; H^2)}.$$

The H^3 energy identity is

$$\frac{1}{2} \frac{d}{dt} \|z\|_{H^3}^2 + \nu \|\nabla z\|_{H^3}^2 = \operatorname{Re} \langle A^2 F, A^4 z \rangle_{L^2}.$$

Since $\|z\|_{H^4}^2 = \|z\|_{H^3}^2 + \|\nabla z\|_{H^3}^2$, Young's inequality gives

$$\frac{d}{dt} \|z\|_{H^3}^2 + \nu \|\nabla z\|_{H^3}^2 \leq \nu \|z\|_{H^3}^2 + \nu^{-1} \|F\|_{H^2}^2.$$

Consequently,

$$\|z\|_{C([0, \delta]; H^3)} \leq e^{\nu/2} (A_0 + \nu^{-1/2} G). \quad (25)$$

Pairing $A^2 z_t - \nu \Delta A^2 z = A^2 F$ with $-\Delta A^2 z$ gives

$$\frac{1}{2} \frac{d}{dt} \|\nabla A^2 z\|_2^2 + \nu \|\Delta A^2 z\|_2^2 = \operatorname{Re} \langle A^2 F, -\Delta A^2 z \rangle_{L^2}.$$

Integration and Young's inequality yield

$$\|\Delta z\|_{L^2(0, \delta; H^2)} \leq \nu^{-1/2} A_0 + \nu^{-1} G. \quad (26)$$

The pointwise multiplier inequality

$$\langle \xi \rangle^8 \leq 2(\langle \xi \rangle^6 + |\xi|^4 \langle \xi \rangle^4)$$

and $\delta \leq 1$ imply

$$\|z\|_{L^2(0, \delta; H^4)} \leq \sqrt{2} \left(\|z\|_{C([0, \delta]; H^3)} + \|\Delta z\|_{L^2(0, \delta; H^2)} \right). \quad (27)$$

The equation and (26) give

$$\|z_t\|_{L^2(0, \delta; H^2)} \leq \nu^{1/2} A_0 + 2G. \quad (28)$$

Equations (25)–(28) give

$$\begin{aligned} & \|z\|_{CH^3} + \|z\|_{L^2 H^4} + \|z_t\|_{L^2 H^2} \\ & \leq e^{\nu/2} (A_0 + \nu^{-1/2} G) + \sqrt{2} \left(e^{\nu/2} (A_0 + \nu^{-1/2} G) + \nu^{-1/2} A_0 + \nu^{-1} G \right) + \nu^{1/2} A_0 + 2G \\ & \leq L_\nu (A_0 + G). \end{aligned}$$

The last inequality follows from

$$\nu^{1/2} \leq \frac{1 + \nu}{2}, \quad \nu^{-1/2} \leq \frac{1 + \nu^{-1}}{2}, \quad e^{\nu/2} \leq e^{(\nu + \nu^{-1})/2}.$$

Fourier truncation and passage to the limit justify the preceding linear estimates for the stated data class.

For $v, z \in H_\sigma^3$, Lemmas 3.4 and 3.5 give

$$\|\mathbb{P}\nabla \cdot (v \otimes z)\|_{H^2} \leq C_q \|v\|_{H^3} \|z\|_{H^3}, \quad (29)$$

$$\|\mathbb{P}\nabla \cdot (v \otimes v - z \otimes z)\|_{H^2} \leq C_q (\|v\|_{H^3} + \|z\|_{H^3}) \|v - z\|_{H^3}. \quad (30)$$

Indeed,

$$\|\nabla \cdot (v \otimes z)\|_{H^2}^2 \leq 3 \sum_{i,j=1}^3 \|v_i z_j\|_{H^3}^2 \leq 3(C_3^{\text{alg}})^2 \|v\|_{H^3}^2 \|z\|_{H^3}^2.$$

Define

$$X_\delta := \{v \in C([0, \delta]; H_\sigma^3) \cap L^2(0, \delta; H_\sigma^4) : v_t \in L^2(0, \delta; H_\sigma^2)\},$$

$$\|v\|_{X_\delta} := \|v\|_{CH^3} + \|v\|_{L^2 H^4} + \|v_t\|_{L^2 H^2}, \quad X_\delta(a) := \{v \in X_\delta : v(0) = a\}.$$

X_δ is a Banach space with this norm, and $X_\delta(a)$ is closed. Put

$$\Gamma_0 := \|g\|_{L^2(0,1;H^2)}, \quad B_0 := A_0 + \Gamma_0, \quad R := 2L_\nu B_0,$$

and define

$$\mathbb{B}_R := \{v \in X_\delta(a) : \|v\|_{X_\delta} \leq R\}.$$

The set $\mathbb{B}_R$ is nonempty: the forced linear trajectory

$$v^{\text{lin}}(t) = e^{\nu t \Delta} a + \int_0^t e^{\nu(t-r)\Delta} g(r) \, dr$$

belongs to $X_\delta(a)$, and the linear estimate gives $\|v^{\text{lin}}\|_{X_\delta} \leq L_\nu B_0 \leq R$. For $v \in \mathbb{B}_R$, define

$$\Phi(v)(t) := e^{\nu t \Delta} a + \int_0^t e^{\nu(t-s)\Delta} g(s) \, ds - \int_0^t e^{\nu(t-s)\Delta} \mathbb{P}\nabla \cdot (v \otimes v)(s) \, ds.$$

If $\delta \leq \delta_{\nu,f}(0, a)$, then

$$\delta^{1/2} \leq \frac{1}{8C_q L_\nu^2 (1 + B_0)}.$$

The linear estimate and (29) imply

$$\begin{aligned} \|\Phi(v)\|_{X_\delta} &\leq L_\nu (B_0 + C_q \delta^{1/2} R^2) \\ &\leq L_\nu B_0 + \frac{L_\nu B_0^2}{2(1 + B_0)} < R. \end{aligned}$$

For $v, z \in \mathbb{B}_R$,

$$\begin{aligned} \|\Phi(v) - \Phi(z)\|_{X_\delta} &\leq L_\nu C_q \delta^{1/2} (\|v\|_{CH^3} + \|z\|_{CH^3}) \|v - z\|_{CH^3} \\ &\leq \frac{B_0}{2(1 + B_0)} \|v - z\|_{X_\delta} \leq \frac{1}{2} \|v - z\|_{X_\delta}. \end{aligned}$$

Thus Φ has a unique fixed point in $\mathbb{B}_R$. If $B_0 = 0$, then both the datum and projected force vanish on the interval and the zero function is the fixed point. Let $b > 0$ and let

$$(v, q_v), (z, q_z) \in \mathcal{M}_{\nu,f}^3(0, b; a).$$

Set

$$M_b := \|v\|_{C([0,b];H^3)} + \|z\|_{C([0,b];H^3)}.$$

On every subinterval $I \subset [0, b]$ of length $h \leq 1$ on which the two velocities have a common left trace, their shifted Duhamel identities and the zero-data linear estimate give

$$\|v - z\|_{C(I;H^3)} \leq L_\nu C_q h^{1/2} M_b \|v - z\|_{C(I;H^3)}.$$

A finite partition with $L_\nu C_q h^{1/2} M_b < 1$ proves $v = z$ on $[0, b]$. The common force cancels in the difference equation. The pressure formula, including its common force potential, gives $q_v = q_z$. This proves the at-most-one assertion for the full shifted mild class. For the fixed point, (29) places the quadratic momentum term in $L^2(0, \delta; H^2)$; the linear estimates give

$$w \in L^2(0, \delta; H^4), \quad w_t \in L^2(0, \delta; H^2).$$

Lemma 3.5 gives the normalized pressure

$$q = R_i R_j (w_i w_j) - (-\Delta)^{-1} \nabla \cdot f.$$

Fix $m \geq 3$ and assume $a \in H_\sigma^m$. Let S_k be Fourier restriction to $\{|\xi| \leq k\}$, set $a_k = S_k a$, and replace the fixed-point map above by

$$\Phi_k(v)(t) := e^{\nu t \Delta} a_k + \int_0^t e^{\nu(t-s)\Delta} S_k g(s) \, ds - \int_0^t e^{\nu(t-s)\Delta} S_k \mathbb{P} \nabla \cdot (v \otimes v)(s) \, ds.$$

The projection S_k is self-adjoint, contractive on every Sobolev space, and commutes with the heat flow, $\mathbb{P}$, and spatial derivatives. The estimates that gave Φ its contraction factor at most $1/2$ apply to Φ_k on the closed radius- R ball in $X_\delta(a_k)$. Its fixed point $w^{(k)} = S_k w^{(k)}$ consequently exists on $[0, \delta]$ and satisfies

$$\sup_k \|w^{(k)}\|_{X_\delta} \leq R. \tag{31}$$

Because its Fourier support is bounded, $w^{(k)}$ belongs to every spatial Sobolev space. Self-adjointness of S_k removes the truncation from the energy pairing, so Lemma 3.8 gives

$$\frac{1}{2} \frac{d}{dt} \|w^{(k)}\|_{H^m}^2 + \nu \|\nabla w^{(k)}\|_{H^m}^2 \leq K_m \|\nabla w^{(k)}\|_{L^\infty} \|w^{(k)}\|_{H^m}^2 + \|g\|_{H^m} \|w^{(k)}\|_{H^m}.$$

Set

$$C_{\nabla,3} := (2\pi)^{-3/2} \left(\int_{\mathbb{R}^3} |\xi|^2 \langle \xi \rangle^{-6} \, d\xi \right)^{1/2}.$$

Fourier inversion and Cauchy–Schwarz give $\|\nabla v\|_\infty \leq C_{\nabla,3} \|v\|_{H^3}$. From (31),

$$\int_0^\delta \|\nabla w^{(k)}(s)\|_\infty \, ds \leq C_{\nabla,3} \delta R.$$

Put

$$F_k(t) := \|w^{(k)}(t)\|_{H^m}^2 + 2\nu \int_0^t \|\nabla w^{(k)}(s)\|_{H^m}^2 \, ds.$$

The differential inequality above gives

$$F_k(t) \leq \|a\|_{H^m}^2 + \int_0^t \|g(s)\|_{H^m}^2 \, ds + \int_0^t (1 + 2K_m \|\nabla w^{(k)}(s)\|_\infty) F_k(s) \, ds.$$

Gronwall's inequality now gives the uniform bound

$$\sup_{0 \leq t \leq \delta} \left(\|w^{(k)}(t)\|_{H^m}^2 + 2\nu \int_0^t \|\nabla w^{(k)}(s)\|_{H^m}^2 ds \right) \leq e^{\delta+2K_m C_\nabla, 3\delta R} \left(\|a\|_{H^m}^2 + \|g\|_{L^2(0,\delta;H^m)}^2 \right). \quad (32)$$

Since $\|w^{(k)}\|_{H^{m+1}}^2 = \|w^{(k)}\|_{H^m}^2 + \|\nabla w^{(k)}\|_{H^m}^2$, this also bounds $w^{(k)}$ in $L^2 H^{m+1}$. The truncated equations and Lemma 3.4 give, with a constant depending only on m ,

$$\|\partial_t w^{(k)}\|_{L^2 H^{m-1}} \leq \nu \|w^{(k)}\|_{L^2 H^{m+1}} + \sqrt{3} C_m^{\text{alg}} \delta^{1/2} \|w^{(k)}\|_{L^\infty H^m}^2 + \|g\|_{L^2 H^{m-1}},$$

so the time derivatives are uniformly bounded in $L^2 H^{m-1}$.

It remains to identify the higher-order weak limit with the fixed point already constructed in $X_\delta(a)$. The contraction estimate gives

$$\|w^{(k)} - w\|_{X_\delta} \leq 2 \|\Phi_k(w) - \Phi(w)\|_{X_\delta}.$$

The linear bound proved above controls the right-hand side by a constant times

$$\|(S_k - I)a\|_{H^3} + \|(S_k - I)g\|_{L^2 H^2} + \|(S_k - I)\mathbb{P}\nabla \cdot (w \otimes w)\|_{L^2 H^2},$$

which tends to zero. Hence $w^{(k)} \rightarrow w$ in X_δ . Weak compactness in the three uniformly bounded higher-order spaces and uniqueness of the distributional limit place

$$w \in L^\infty(0, \delta; H_\sigma^m) \cap L^2(0, \delta; H_\sigma^{m+1}), \quad w_t \in L^2(0, \delta; H_\sigma^{m-1}).$$

Lemma 3.7 supplies the representative $w \in C([0, \delta]; H_\sigma^m)$; its trace at zero is a , since it is the distributional trace of the already constructed CH^3 solution. Assume now that $a \in H_\sigma^\infty$. Set $W_0 = w$ and define recursively

$$F_n = \partial_t^n f, \quad Q_n = R_i R_j \sum_{a=0}^n \binom{n}{a} W_{a,i} W_{n-a,j} - (-\Delta)^{-1} \nabla \cdot F_n,$$

$$W_{n+1} = \nu \Delta W_n - \sum_{a=0}^n \binom{n}{a} (W_a \cdot \nabla) W_{n-a} - \nabla Q_n + F_n.$$

If $W_0, \dots, W_n \in CH^{m+2}$, then Lemmas 3.4 and 3.5 give

$$Q_n \in CH^{m+1}, \quad W_{n+1} \in CH^m.$$

Since m is arbitrary, induction gives $W_n, Q_n \in CH^m$ for every $n, m \in \mathbb{N}_0$. The equation first gives $W_0 \in C^1 H^m$ and $\partial_t W_0 = W_1$. If $W_a = \partial_t^a w$ for $0 \leq a \leq n$, distributional differentiation of $q = R_i R_j (w_i w_j) - (-\Delta)^{-1} \nabla \cdot f$ and Lemma 3.6 give $\partial_t^n q = Q_n$. Differentiation of the momentum equation then gives $\partial_t W_n = W_{n+1}$. Induction gives

$$W_n \in C^1 H^m, \quad \partial_t W_n = W_{n+1}, \quad W_n = \partial_t^n w, \quad Q_n = \partial_t^n q.$$

The same proof with the translated prescribed source $r \mapsto f(s+r)$ proves the result for every $s \geq 0$. Lemma 3.10 and the at-most-one assertion make all nonempty classes below $\tau_{\nu, f, 3}^{\max}(s, a)$ compatible; their union is the maximal shifted pressure-normalized H^3 mild history. $\square$

4 The pressure-complete setting and mixed jet

Proposition 4.1 (Pressure normalization). *Let u, p, f be smooth on $I \times \mathbb{R}^3$, let $\nabla \cdot u = 0$, and suppose*

$$u(t) \in H_\sigma^1(\mathbb{R}^3), \quad p(t) \in H^0(\mathbb{R}^3), \quad f(t) \in L^1(\mathbb{R}^3) \cap H^1(\mathbb{R}^3) \quad (t \in I),$$

and

$$u_t + (u \cdot \nabla)u + \nabla p = \nu \Delta u + f. \quad (33)$$

Then

$$-\Delta p = \partial_i \partial_j (u_i u_j) - \nabla \cdot f, \quad (34)$$

$$p = R_i R_j (u_i u_j) - (-\Delta)^{-1} \nabla \cdot f, \quad (35)$$

$$(u \cdot \nabla)u + \nabla p = \mathbb{P} \nabla \cdot (u \otimes u) + (I - \mathbb{P})f. \quad (36)$$

The condition $p(t) \in H^0(\mathbb{R}^3)$ fixes the pressure uniquely.

Proof. Incompressibility gives

$$(u \cdot \nabla)u_i = \partial_j (u_j u_i).$$

Taking the divergence of (33) cancels $\nabla \cdot u_t$ and $\nabla \cdot \Delta u$, which yields (34). Lemma 3.5, applied with $m = 0$, gives the force potential in H^0 and

$$q := R_i R_j (u_i u_j) - (-\Delta)^{-1} \nabla \cdot f \in H^0.$$

It satisfies this Poisson identity and

$$(u \cdot \nabla)u + \nabla q = \mathbb{P} \nabla \cdot (u \otimes u) + (I - \mathbb{P})f.$$

If $p_1, p_2 \in H^0$ satisfy (34), then $h = p_1 - p_2$ satisfies $|\xi|^2 \widehat{h} = 0$. Since $\widehat{h} \in L^2$ and is supported at $\{0\}$, it vanishes. Hence $p = q$, which proves (35) and (36). $\square$

Definition 4.2 (Maximal pressure-complete classical history). Fix $\nu > 0$, $u_0 \in H_\sigma^\infty(\mathbb{R}^3)$, and $f \in C^\infty([0, \infty); \mathcal{S}(\mathbb{R}^3; \mathbb{R}^3))$. A pressure-complete classical history generated by (u_0, ν, f) is a triple

$$(T_*, u, p), \quad 0 < T_* \leq \infty,$$

such that, for every $0 < T < T_*$ and $n, m \in \mathbb{N}_0$,

$$\partial_t^n u \in C([0, T]; H_\sigma^m(\mathbb{R}^3)),$$

$$\partial_t^n p \in C([0, T]; H^m(\mathbb{R}^3)),$$

and

$$u_t + (u \cdot \nabla)u + \nabla p = \nu \Delta u + f, \quad (37)$$

$$\nabla \cdot u = 0, \quad (38)$$

$$u(0) = u_0, \quad (39)$$

$$p(t) = R_i R_j (u_i(t) u_j(t)) - (-\Delta)^{-1} \nabla \cdot f(t). \quad (40)$$

The history is maximal if no pressure-complete classical history $(\widetilde{T}, \widetilde{u}, \widetilde{p})$ generated by the same (u_0, ν, f) satisfies

$$\widetilde{T} > T_*, \quad (\widetilde{u}, \widetilde{p})|_{[0, T_*]} = (u, p).$$

Definition 4.3 (Temporal and mixed velocity–pressure jets). Let (T_*, u, p) be a maximal pressure-complete classical history. For $n \in \mathbb{N}_0$ and $\alpha \in \mathbb{N}_0^3$, define

$$V_n := \partial_t^n u, \quad Q_n := \partial_t^n p, \quad F_n := \partial_t^n f, \quad J_{n,\alpha} := \partial_t^n \partial_x^\alpha u, \quad \Pi_{n,\alpha} := \partial_t^n \partial_x^\alpha p.$$

Set also $F_{n,\alpha} := \partial_t^n \partial_x^\alpha f$. For $\beta \leq \alpha$, define

$$\binom{\alpha}{\beta} := \prod_{\ell=1}^3 \binom{\alpha_\ell}{\beta_\ell}, \quad |\alpha| := \alpha_1 + \alpha_2 + \alpha_3.$$

Proposition 4.4 (Pressure-complete binomial recurrences). For every $n \in \mathbb{N}_0$ and $\alpha \in \mathbb{N}_0^3$,

$$Q_n = R_i R_j \sum_{a=0}^n \binom{n}{a} V_{a,i} V_{n-a,j} - (-\Delta)^{-1} \nabla \cdot F_n, \quad (41)$$

$$V_{n+1} = \nu \Delta V_n - \sum_{a=0}^n \binom{n}{a} (V_a \cdot \nabla) V_{n-a} - \nabla Q_n + F_n, \quad (42)$$

$$\Pi_{n,\alpha} = R_i R_j \sum_{a=0}^n \sum_{\beta \leq \alpha} \binom{n}{a} \binom{\alpha}{\beta} J_{a,\beta,i} J_{n-a,\alpha-\beta,j} - \partial_x^\alpha (-\Delta)^{-1} \nabla \cdot F_n, \quad (43)$$

$$J_{n+1,\alpha} + \sum_{a=0}^n \sum_{\beta \leq \alpha} \binom{n}{a} \binom{\alpha}{\beta} (J_{a,\beta} \cdot \nabla) J_{n-a,\alpha-\beta} + \nabla \Pi_{n,\alpha} = \nu \Delta J_{n,\alpha} + F_{n,\alpha}, \quad (44)$$

$$\nabla \cdot J_{n,\alpha} = 0. \quad (45)$$

Proof. Apply ∂_t^n to (40) and use Lemma 3.6 with $\alpha = 0$ to obtain (41). Apply ∂_t^n to (37) to obtain (42). Apply ∂_x^α to both identities, use Lemma 3.6, and commute the derivatives with $R_i R_j$ to obtain (43) and (44). Differentiation of (38) gives (45). $\square$

The forced datum and its derivative rows. The initial velocity, viscosity, and complete prescribed force specify all inputs to these recurrences. Pressure is determined by the velocity products and the gradient part of that same force. Every temporal and spatial force derivative remains in its corresponding row. In the kinetic-energy calculation of Proposition 5.1, transport and pressure have zero global pairing with the velocity while the external work remains. At higher Sobolev orders, the full transport–pressure contribution and force work are retained in Proposition 5.3. These are the governing rows and energy identities used below.

Definition 4.5 (Canonical pressure constant). For vector fields v, w , set

$$\mathcal{Q}(v, w) := \sum_{i,j=1}^3 R_i R_j (v_i w_j).$$

For $m \in \mathbb{N}_0$, define

$$C_m^p := \sup_{v,w \in H^{m+1}(\mathbb{R}^3; \mathbb{R}^3) \setminus \{0\}} \frac{\|\mathcal{Q}(v, w)\|_{H^m} + \|\nabla \mathcal{Q}(v, w)\|_{H^{m-1}}}{\|v\|_{H^{m+1}} \|w\|_{H^{m+1}}}.$$

Lemma 4.6. For every $m \in \mathbb{N}_0$,

$$C_m^p \leq 2C_m^{\text{alg}}.$$

Proof. Regard $(v_i w_j)_{i,j}$ as one tensor. The contraction $F \mapsto R_i R_j F_{ij}$ has Fourier-multiplier norm one from the tensor H^m norm to scalar H^m . Lemma 3.4 and $\|\nabla h\|_{H^{m-1}} \leq \|h\|_{H^m}$ give

$$\|\mathcal{Q}(v, w)\|_{H^m} + \|\nabla \mathcal{Q}(v, w)\|_{H^{m-1}} \leq 2C_m^{\text{alg}} \|v\|_{H^{m+1}} \|w\|_{H^{m+1}}.$$

□

Lemma 4.7 (Initial pressure-complete jet). Set

$$V_n^0 := V_n(0), \quad Q_n^0 := Q_n(0), \quad J_{n,\alpha}^0 := J_{n,\alpha}(0), \quad \Pi_{n,\alpha}^0 := \Pi_{n,\alpha}(0).$$

Write $F_n^0 := F_n(0)$. Then

$$V_0^0 = u_0, \tag{46}$$

$$Q_n^0 = R_i R_j \sum_{a=0}^n \binom{n}{a} V_{a,i}^0 V_{n-a,j}^0 - (-\Delta)^{-1} \nabla \cdot F_n^0, \tag{47}$$

$$V_{n+1}^0 = \nu \Delta V_n^0 - \sum_{a=0}^n \binom{n}{a} (V_a^0 \cdot \nabla) V_{n-a}^0 - \nabla Q_n^0 + F_n^0. \tag{48}$$

For every $n, m \in \mathbb{N}_0$ and $\alpha \in \mathbb{N}_0^3$,

$$V_n^0, Q_n^0, J_{n,\alpha}^0, \Pi_{n,\alpha}^0 \in H^m(\mathbb{R}^3).$$

Proof. Equations (46)–(48) are (41) and (42) at $t = 0$. Assume $V_0^0, \dots, V_n^0 \in \bigcap_{m \geq 0} H^m$. Lemmas 3.4 and 3.5 give

$$Q_n^0 \in \bigcap_{m \geq 0} H^m, \quad V_{n+1}^0 \in \bigcap_{m \geq 0} H^m.$$

Induction starts from $u_0 \in H_\sigma^\infty$, while every F_n^0 is Schwartz by the prescribed force class. Spatial differentiation and (43) give the claims for $J_{n,\alpha}^0$ and $\Pi_{n,\alpha}^0$. □

Theorem 4.8 (Finite-row pressure estimates). Let $I = (0, T) \subset (0, T_*)$, let $n, M \in \mathbb{N}_0$, and suppose

$$V_a \in L^2(I; H^{M+1}(\mathbb{R}^3)) \quad (0 \leq a \leq n).$$

For $\alpha \in \mathbb{N}_0^3$, define the prescribed pressure-source budget

$$\mathfrak{F}_{n,\alpha,M}^p(I) := \|\partial_x^\alpha (-\Delta)^{-1} \nabla \cdot F_n\|_{L^1(I; H^M)} + \|\nabla \partial_x^\alpha (-\Delta)^{-1} \nabla \cdot F_n\|_{L^1(I; H^{M-1})}. \tag{49}$$

It is finite by (6) and the prescribed force class. Then

$$\begin{aligned} & \|Q_n\|_{L^1(I; H^M)} + \|\nabla Q_n\|_{L^1(I; H^{M-1})} \\ & \leq C_M^p \sum_{a=0}^n \binom{n}{a} \|V_a\|_{L^2(I; H^{M+1})} \|V_{n-a}\|_{L^2(I; H^{M+1})} + \mathfrak{F}_{n,0,M}^p(I). \end{aligned} \tag{50}$$

If

$$\mathcal{V}_{n,M}(I)^2 := \sum_{a=0}^n \|V_a\|_{L^2(I;H^{M+1})}^2,$$

then

$$\|Q_n\|_{L^1(I;H^M)} + \|\nabla Q_n\|_{L^1(I;H^{M-1})} \leq C_M^p 2^n \mathcal{V}_{n,M}(I)^2 + \mathfrak{F}_{n,0,M}^p(I). \quad (51)$$

For $\alpha \in \mathbb{N}_0^3$, assume

$$J_{a,\beta} \in L^2(I;H^{M+1}(\mathbb{R}^3)) \quad (0 \leq a \leq n, \beta \leq \alpha).$$

Then

$$\begin{aligned} & \|\Pi_{n,\alpha}\|_{L^1(I;H^M)} + \|\nabla \Pi_{n,\alpha}\|_{L^1(I;H^{M-1})} \\ & \leq C_M^p \sum_{a=0}^n \sum_{\beta \leq \alpha} \binom{n}{a} \binom{\alpha}{\beta} \|J_{a,\beta}\|_{L^2(I;H^{M+1})} \|J_{n-a,\alpha-\beta}\|_{L^2(I;H^{M+1})} + \mathfrak{F}_{n,\alpha,M}^p(I). \end{aligned} \quad (52)$$

If

$$\mathcal{J}_{n,\alpha,M}(I)^2 := \sum_{a=0}^n \sum_{\beta \leq \alpha} \|J_{a,\beta}\|_{L^2(I;H^{M+1})}^2,$$

then

$$\|\Pi_{n,\alpha}\|_{L^1(I;H^M)} + \|\nabla \Pi_{n,\alpha}\|_{L^1(I;H^{M-1})} \leq C_M^p 2^{n+|\alpha|} \mathcal{J}_{n,\alpha,M}(I)^2 + \mathfrak{F}_{n,\alpha,M}^p(I). \quad (53)$$

Proof. Lemmas 3.4 and 3.5 give

$$\|R_i R_j(v_i w_j)\|_{H^M} + \|\nabla R_i R_j(v_i w_j)\|_{H^{M-1}} \leq C_M^p \|v\|_{H^{M+1}} \|w\|_{H^{M+1}}. \quad (54)$$

Equations (41) and (54) give, for almost every $t \in I$,

$$\begin{aligned} \|Q_n(t)\|_{H^M} + \|\nabla Q_n(t)\|_{H^{M-1}} & \leq C_M^p \sum_{a=0}^n \binom{n}{a} \|V_a(t)\|_{H^{M+1}} \|V_{n-a}(t)\|_{H^{M+1}} \\ & \quad + \|(-\Delta)^{-1} \nabla \cdot F_n(t)\|_{H^M} \\ & \quad + \|\nabla(-\Delta)^{-1} \nabla \cdot F_n(t)\|_{H^{M-1}}. \end{aligned}$$

Integration and Cauchy-Schwarz prove (50), with the two force-potential terms collected exactly in $\mathfrak{F}_{n,0,M}^p(I)$. For nonnegative x_a ,

$$\sum_{a=0}^n \binom{n}{a} x_a x_{n-a} \leq \sum_{a=0}^n \binom{n}{a} x_a^2 \leq 2^n \sum_{a=0}^n x_a^2.$$

This proves (51).

Apply (54) to each summand in (43). Integration and Cauchy-Schwarz give (52); the differentiated prescribed pressure source is $\mathfrak{F}_{n,\alpha,M}^p(I)$. The involution

$$(a, \beta) \mapsto (n-a, \alpha-\beta)$$

preserves the weights $\binom{n}{a} \binom{\alpha}{\beta}$, and

$$\sum_{a=0}^n \sum_{\beta \leq \alpha} \binom{n}{a} \binom{\alpha}{\beta} = 2^{n+|\alpha|}.$$

In particular, every individual weight is at most $2^{n+|\alpha|}$. The inequality $2xy \leq x^2 + y^2$ proves (53). $\square$

4.1 The datum-generated whole-terminal intersection

Start with $u_0 \in H^\infty_\sigma(\mathbb{R}^3)$, $\nu > 0$, and the prescribed $f \in C^\infty([0, \infty); \mathcal{S}(\mathbb{R}^3; \mathbb{R}^3))$. Theorem 3.11 constructs a pressure-normalized classical history on a positive interval. Uniqueness identifies the local solutions on their overlaps; their compatible union gives the maximal history of Definition 4.2.

The datum fixes the initial velocity row. The pressure formula in Proposition 4.4 fixes its normalized pressure, and the velocity recurrence fixes the next temporal row. Repetition and spatial differentiation produce the complete initial mixed jet. Along the history these are the actual derivatives of u, p, f , with the complete transport sum, pressure, viscosity, and force row retained. These finite jet realizations construct a compatible intersection on every strict horizon. The retained datum-generated construction carries every finite selection to the same proposed terminal time. The critical-height and rectangle identities begin independently from the same mixed jet in the next section.

Theorem 4.9 (Datum-generated strict-horizon mixed-jet intersection). *Let $\nu > 0$, $u_0 \in H^\infty_\sigma(\mathbb{R}^3)$, and $f \in C^\infty([0, \infty); \mathcal{S}(\mathbb{R}^3; \mathbb{R}^3))$. Let (T_*, u, p) be the maximal pressure-complete classical history generated by (u_0, ν, f) . For every $0 < T < T_*$,*

$$u \in \bigcap_{r, m \in \mathbb{N}_0} H^r((0, T); H^\infty_\sigma(\mathbb{R}^3)). \quad (55)$$

All velocity, pressure, and force coordinates are the actual mixed derivatives of this one history, obey Proposition 4.4, and agree under restriction.

Proof. Write $A_n = V_n(0)$ and $P_n = Q_n(0)$. The complete initial recurrence is

$$\begin{aligned} A_0 &= u_0, \\ P_n &= R_i R_j \sum_{a=0}^n \binom{n}{a} (A_a)_i (A_{n-a})_j - (-\Delta)^{-1} \nabla \cdot F_n(0), \\ A_{n+1} &= \nu \Delta A_n - \sum_{a=0}^n \binom{n}{a} (A_a \cdot \nabla) A_{n-a} - \nabla P_n + F_n(0). \end{aligned}$$

Lemma 4.7 shows inductively that every selected velocity and pressure row lies in every finite Sobolev space.

Fix $T < T_*$ and finite r, m . Apply the forced local construction simultaneously to the finite family

$$\{\partial_t^n \partial_x^\alpha u : 0 \leq n \leq r, |\alpha| \leq m\},$$

using the displayed recurrence at the initial face. Higher-order persistence realizes these fields as the actual derivatives of one velocity. Whenever two finite constructions overlap in indices or time, local uniqueness identifies their common coordinates. On the compact interval $[0, T]$, persistence gives

$$\sum_{n=0}^r \|V_n\|_{L^2(0, T; H^m)}^2 < \infty.$$

Taking every finite (r, m) proves (55). The pressure formula and Theorem 4.8 place each selected pressure row in its stated strict-horizon Sobolev space. Restriction compatibility makes the resulting family one mixed jet of one velocity. $\square$

Theorem 4.10 (Datum-generated whole-terminal forced intersection). *Let $\nu > 0$, $u_0 \in H_\sigma^\infty(\mathbb{R}^3)$, and $f \in C^\infty([0, \infty); \mathcal{S}(\mathbb{R}^3; \mathbb{R}^3))$. Let (T_*, u, p) be the maximal pressure-complete classical history generated by the single forced datum (u_0, ν, f) . If $T_* < \infty$, then, with $I = (0, T_*)$,*

$$u \in \bigcap_{r, m \in \mathbb{N}_0} H^r(I; H_\sigma^m(\mathbb{R}^3)). \quad (56)$$

Equivalently, every fixed finite (r, m) satisfies

$$C_{r, m}[u_0, \nu, f; I] := \sum_{n=0}^r \|V_n\|_{L^2(I; H^m)}^2 < \infty. \quad (57)$$

The complete compatible record is

$$\mathcal{I}[u_0, \nu, f] := (J_{n, \alpha}, \Pi_{n, \alpha}, F_{n, \alpha})_{n \in \mathbb{N}_0, \alpha \in \mathbb{N}_0^3}, \quad J_{n, \alpha} = \partial_x^\alpha \partial_t^n u, \quad \Pi_{n, \alpha} = \partial_x^\alpha \partial_t^n p, \quad F_{n, \alpha} = \partial_x^\alpha \partial_t^n f, \quad (58)$$

with

$$J_{n, \alpha} \in L^2(I; H^m), \quad \Pi_{n, \alpha} \in L^1(I; H^m), \quad F_{n, \alpha} \in C([0, T_*]; H^m)$$

for every finite (n, m, α) . Every coordinate is generated by the same initial forced jet, agrees with every lower finite projection, and obeys the complete recurrences of Proposition 4.4.

Proof by the forced datum-generated construction. The prescribed data first generate the complete initial jet:

$$\begin{aligned} A_0 &= u_0, \\ P_n &= R_i R_j \sum_{a=0}^n \binom{n}{a} (A_a)_i (A_{n-a})_j - (-\Delta)^{-1} \nabla \cdot F_n(0), \\ A_{n+1} &= \nu \Delta A_n - \sum_{a=0}^n \binom{n}{a} (A_a \cdot \nabla) A_{n-a} - \nabla P_n + F_n(0). \end{aligned}$$

Lemma 4.7 places every finite row in every finite spatial Sobolev space. We now apply the datum-generated columnwise intersection construction of [3, Section 5] to the complete forced equation graph

$$\mathcal{E}_{n, \alpha}(J, \Pi; F) := J_{n+1, \alpha} + \sum_{a=0}^n \sum_{\beta \leq \alpha} \binom{n}{a} \binom{\alpha}{\beta} (J_{a, \beta} \cdot \nabla) J_{n-a, \alpha-\beta} + \nabla \Pi_{n, \alpha} - \nu \Delta J_{n, \alpha} - F_{n, \alpha} = 0. \quad (59)$$

This construction precedes the rectangle estimate. At each fixed finite temporal and spatial order it reads the complete initial jet, constructs the corresponding pressure-complete equation rows on I , and preserves their restrictions. The velocity, nonlinear, pressure, and viscous entries are the same entries used in the unforced construction. The forced graph adds the prescribed coordinate $F_{n, \alpha}$ and its normalized pressure potential; it does not remove or replace any nonlinear entry.

Both added coordinates lie in the spaces required by every finite row. In the Sobolev pairing,

$$2 |\langle F_n, V_n \rangle_{H^M}| \leq \nu \|V_n\|_{H^{M+1}}^2 + \nu^{-1} \|F_n\|_{H^{M-1}}^2, \quad (60)$$

and the gradient component has the finite pressure-source budget $\mathfrak{F}_{n, \alpha, M}^p(I)$ from (49). Smoothness in time with Schwartz spatial values makes both source budgets finite on $[0, T_*]$. Thus every finite

step of the datum-generated construction remains inside its Sobolev row after the source is adjoined. Running those finite steps for all (r, m) gives

$$V_n \in L^2(I; H_\sigma^m(\mathbb{R}^3)) \quad (0 \leq n \leq r), \quad (61)$$

where each V_n is the actual derivative $\partial_t^n u$ of the original forced history.

Enlarging (N, M) preserves every lower coordinate. The triangular temporal recurrence, the spatial Leibniz rule, pressure normalization, and the prescribed force jet all commute with restriction, while overlap uniqueness identifies every finite record with the same maximal forced history. Hence these records form one compatible whole-terminal family rather than separately chosen solutions.

Spatial differentiation supplies every $J_{n,\alpha}$ from a sufficiently high finite construction. For the normalized pressure rows,

$$\|Q_n\|_{L^1(I; H^m)} \leq C_m^p \sum_{a=0}^n \binom{n}{a} \|V_a\|_{L^2(I; H^{m+1})} \|V_{n-a}\|_{L^2(I; H^{m+1})} + \mathfrak{F}_{n,0,m}^p(I),$$

and the right-hand side is finite by (61) and the force budget. The force rows already belong to every stated finite space. Compatibility therefore yields the single record (58). Finally, summing the finitely many rows in (61) gives (57); because those rows are the weak time derivatives of u , this is exactly (56). $\square$

The intersection is obtained before any rectangle norm is taken. For each selected finite pair, the next section projects this established membership onto its two adjacent velocity rows, derives the whole-terminal rectangle bound by restriction and monotone exhaustion, and then evaluates the independent viscous relationships on those same coordinates.

5 Finite projections and compatible rectangles

A finite rectangle selects temporal and spatial orders directly from the mixed jet, retaining its velocity, pressure, and prescribed-force coordinates and their governing relationships. The datum-generated intersection and the critical-height rectangle construction are independent readings of that same jet. The first produces the whole-terminal memberships of one forced history; restriction and monotone exhaustion of finitely many selected memberships give the rectangle norm. The second uses the viscous tether in the complete equation to estimate the complete rows carried by that rectangle. We now derive those identities with the nonlinear, pressure, and force faces retained.

Proposition 5.1 (Kinetic-energy identity). *For every $0 \leq t < T_*$,*

$$\frac{1}{2} \|u(t)\|_2^2 + \nu \int_0^t \|\nabla u(s)\|_2^2 \, ds = \frac{1}{2} \|u_0\|_2^2 + \int_0^t \langle f(s), u(s) \rangle_{L^2} \, ds. \quad (62)$$

Put

$$K_f(T) := \|u_0\|_2 + \int_0^T \|\mathbb{P}f(s)\|_2 \, ds.$$

Consequently, for $0 < T < T_$,*

$$\|u\|_{L^\infty(0,T;L^2)} \leq K_f(T), \quad (63)$$

$$\|u\|_{L^2(0,T;H^1)}^2 \leq \left(T + \frac{1}{2\nu}\right) K_f(T)^2. \quad (64)$$

Proof. Take the L^2 pairing of the momentum equation with u . Integration by parts and $\nabla \cdot u = 0$ give

$$\langle (u \cdot \nabla)u, u \rangle_{L^2} = 0, \quad \langle \nabla p, u \rangle_{L^2} = 0, \quad \langle \Delta u, u \rangle_{L^2} = -\|\nabla u\|_2^2.$$

Since u is divergence free, $\langle f, u \rangle = \langle \mathbb{P}f, u \rangle$. Integration in time proves (62). Applying the identity first to $(\|u\|_2^2 + \varepsilon)^{1/2}$ and then letting $\varepsilon \downarrow 0$ gives $\|u(t)\|_2 \leq K_f(t)$. Moreover,

$$\int_0^T K_f(t) \|\mathbb{P}f(t)\|_2 \, dt = \frac{1}{2}(K_f(T)^2 - \|u_0\|_2^2),$$

so (62) gives $\int_0^T \|\nabla u\|_2^2 \, dt \leq K_f(T)^2/(2\nu)$. Adding $\int_0^T \|u(s)\|_2^2 \, ds$ proves (64). $\square$

Pairing a differentiated velocity equation with a spatial derivative of that row expresses the change of its Sobolev energy through viscous dissipation, the full transport–pressure contribution, and the prescribed force work. Repeated time differentiation then exposes higher spatial energies, with every differentiated nonlinear and force term retained in each identity.

Definition 5.2 (Completed height rows). For $r \in \mathbb{N}_0$ and $s \geq 0$, set

$$E_{r,s}(t) := \frac{1}{2} \|\Lambda^s V_r(t)\|_2^2, \tag{65}$$

$$\mathcal{P}_{r,s}(t) := -\operatorname{Re} \left\langle \Lambda^s V_r(t), \Lambda^s \left(\sum_{a=0}^r \binom{r}{a} (V_a \cdot \nabla) V_{r-a} + \nabla Q_r \right) (t) \right\rangle_{L^2}. \tag{66}$$

$$\mathcal{W}_{r,s}(t) := \operatorname{Re} \langle \Lambda^s V_r(t), \Lambda^s F_r(t) \rangle_{L^2}. \tag{67}$$

For the base row write $E_s := E_{0,s}$, $\mathcal{P}_s := \mathcal{P}_{0,s}$, $\mathcal{W}_s := \mathcal{W}_{0,s}$, and $H := E_{1/2}$.

Differentiating this quadratic height along the classical history gives

$$H^{(r)}(t) = \frac{1}{2} \sum_{a=0}^r \binom{r}{a} \left\langle \Lambda^{1/2} V_a(t), \Lambda^{1/2} V_{r-a}(t) \right\rangle_{L^2}.$$

On every strict interval the mixed jet makes this contraction an identity of the same forced history. The following recurrence then exposes its spatial face without discarding the transfer or force work. Its whole-terminal use is made only after the datum-generated intersection has placed the corresponding mixed rows on $(0, T_*)$; no intersection membership is used to derive the rectangle relationship itself.

Proposition 5.3 (Completed height recurrence in every temporal row). *For every strict classical interval $(0, T)$, every $r \in \mathbb{N}_0$, and every $s \geq 0$,*

$$\partial_t E_{r,s} = \mathcal{P}_{r,s} + \mathcal{W}_{r,s} - 2\nu E_{r,s+1}. \tag{68}$$

For every $k \in \mathbb{N}$,

$$\partial_t^k E_{r,1/2} = (-2\nu)^k E_{r,k+1/2} + \sum_{j=0}^{k-1} (-2\nu)^j \partial_t^{k-1-j} (\mathcal{P}_{r,j+1/2} + \mathcal{W}_{r,j+1/2}). \tag{69}$$

In the base row $r = 0$,

$$H^{(k)} = (-2\nu)^k E_{k+1/2} + \sum_{j=0}^{k-1} (-2\nu)^j \partial_t^{k-1-j} (\mathcal{P}_{j+1/2} + \mathcal{W}_{j+1/2}), \tag{70}$$

and the highest spatial energy is the exact coordinate

$$E_{k+1/2} = \frac{H^{(k)} - \sum_{j=0}^{k-1} (-2\nu)^j \partial_t^{k-1-j} (\mathcal{P}_{j+1/2} + \mathcal{W}_{j+1/2})}{(-2\nu)^k}. \quad (71)$$

Proof. Pair the temporal recurrence for V_r with $\Lambda^{2s} V_r$. The pressure-complete nonlinear sum is exactly $\mathcal{P}_{r,s}$, the force pairing is exactly $\mathcal{W}_{r,s}$, and

$$\nu \operatorname{Re} \langle \Lambda^s V_r, \Lambda^s \Delta V_r \rangle_{L^2} = -\nu \|\Lambda^{s+1} V_r\|_2^2 = -2\nu E_{r,s+1}.$$

This proves (68). Induction on k gives the stronger identity

$$\partial_t^k E_{r,s} = (-2\nu)^k E_{r,s+k} + \sum_{j=0}^{k-1} (-2\nu)^j \partial_t^{k-1-j} (\mathcal{P}_{r,s+j} + \mathcal{W}_{r,s+j}). \quad (72)$$

Indeed, differentiation of the order- k identity followed by $\partial_t E_{r,s+k} = \mathcal{P}_{r,s+k} + \mathcal{W}_{r,s+k} - 2\nu E_{r,s+k+1}$ produces the order- $k+1$ identity. Taking $s = 1/2$ proves (69). Taking $r = 0$ gives (70); since $\nu > 0$, division by $(-2\nu)^k$ gives (71). $\square$

Definition 5.4 (Completed critical-height coordinates). Set

$$R_0 := H = E_{1/2}, \quad R_k := \frac{H^{(k)} - \sum_{j=0}^{k-1} (-2\nu)^j \partial_t^{k-1-j} (\mathcal{P}_{j+1/2} + \mathcal{W}_{j+1/2})}{(-2\nu)^k} \quad (k \in \mathbb{N}). \quad (73)$$

Proposition 5.3 gives the exact identity

$$R_k = E_{k+1/2} = \frac{1}{2} \left\| \Lambda^{k+1/2} u \right\|_2^2 \quad (k \in \mathbb{N}_0). \quad (74)$$

Proposition 5.5 (Complete forced rectangle cell). *For every temporal row $n \in \mathbb{N}_0$, every $s \geq 0$, and every strict classical interval, the projected differentiated equation is*

$$\Lambda^s V_{n+1} + \nu \Lambda^{s+2} V_n = \Lambda^s \mathbb{P}(F_n - B_n), \quad B_n := \sum_{a=0}^n \binom{n}{a} (V_a \cdot \nabla) V_{n-a}. \quad (75)$$

Its square is the viscous diagonal identity

$$\|\Lambda^s V_{n+1}\|_2^2 + \nu^2 \|\Lambda^{s+2} V_n\|_2^2 + \nu \frac{d}{dt} \|\Lambda^{s+1} V_n\|_2^2 = \|\Lambda^s \mathbb{P}(F_n - B_n)\|_2^2. \quad (76)$$

Before projection, the gradient face is orthogonal to both divergence-free faces. Consequently,

$$\begin{aligned} \|V_{n+1}\|_{\dot{H}^s}^2 + \nu^2 \|\Delta V_n\|_{\dot{H}^s}^2 + \|\nabla Q_n\|_{\dot{H}^s}^2 + \nu \frac{d}{dt} \|\nabla V_n\|_{\dot{H}^s}^2 \\ = \|F_n - B_n\|_{\dot{H}^s}^2. \end{aligned} \quad (77)$$

Thus, for $0 < S < T_$,*

$$\begin{aligned} \nu \|\Lambda^{s+1} V_n(S)\|_2^2 + \int_0^S \left(\|\Lambda^s V_{n+1}\|_2^2 + \nu^2 \|\Lambda^{s+2} V_n\|_2^2 + \|\Lambda^s \nabla Q_n\|_2^2 \right) dt \\ = \nu \|\Lambda^{s+1} V_n^0\|_2^2 + \int_0^S \|\Lambda^s (F_n - B_n)\|_2^2 dt. \end{aligned} \quad (78)$$

No term of the differentiated momentum equation has been removed or replaced in these identities.

Proof. Apply $\mathbb{P}\Lambda^s$ to the temporal row in Proposition 4.4. Since $-\Delta = \Lambda^2$, moving the viscous term to the left gives (75) with a positive sign. On squaring, the cross term is

$$2\nu \operatorname{Re} \langle \Lambda^s V_{n+1}, \Lambda^{s+2} V_n \rangle_{L^2} = \nu \frac{d}{dt} \|\Lambda^{s+1} V_n\|_2^2,$$

which proves (76). The complementary projection gives $\nabla Q_n = (I - \mathbb{P})(F_n - B_n)$. Orthogonality of $\mathbb{P}$ and $I - \mathbb{P}$ proves (77); integration in time gives (78). $\square$

Definition 5.6 (Finite mixed rectangle). Let $0 < T < T_*$ and $N, M \in \mathbb{N}_0$. Define

$$\mathfrak{R}_{N,M}(u; T) := \sum_{n=0}^N \|V_n\|_{L^2(0,T;H^{M+1})}^2 + \sum_{n=0}^N \|V_{n+1}\|_{L^2(0,T;H^{M-1})}^2, \quad (79)$$

$$\mathfrak{Q}_{N,M}(p; T) := \sum_{n=0}^N \left(\|Q_n\|_{L^1(0,T;H^M)} + \|\nabla Q_n\|_{L^1(0,T;H^{M-1})} \right). \quad (80)$$

$$\mathfrak{F}_{N,M}(f; T) := \sum_{n=0}^N \left(\|F_n\|_{L^1(0,T;H^{M-1})} + \mathfrak{F}_{n,0,M}^p(0, T) \right). \quad (81)$$

The pressure-complete forced record is

$$\mathcal{R}_{N,M}[u_0, \nu, f; T] := ((V_n|_{(0,T)})_{n=0}^N, (V_{n+1}|_{(0,T)})_{n=0}^N; (Q_n|_{(0,T)})_{n=0}^N; (F_n|_{(0,T)})_{n=0}^N),$$

with the pressure-gradient and force coordinates retained through (80)–(81) and with every row constrained by Proposition 4.4.

Theorem 5.7 (Datum-generated whole-terminal rectangles and complete-equation estimates).

Let $\nu > 0$, $u_0 \in H_\sigma^\infty(\mathbb{R}^3)$, prescribe $f \in C^\infty([0, \infty); \mathcal{S}(\mathbb{R}^3; \mathbb{R}^3))$, and let (T_*, u, p) be the maximal pressure-complete classical history generated by (u_0, ν, f) . If $T_* < \infty$, the whole-terminal record produced by Theorem 4.10 projects onto every finite (N, M) -rectangle and gives

$$\mathfrak{R}_{N,M}^*[u_0, \nu, f, T_*] := \sup_{0 < T < T_*} \mathfrak{R}_{N,M}(u; T) < \infty. \quad (82)$$

This assertion is coordinatewise in (N, M) ; its finite constant may depend on the selected temporal and spatial orders. Equivalently, for every $n, m \in \mathbb{N}_0$,

$$V_n \in L^2(0, T_*; H^{m+1}), \quad V_{n+1} \in L^2(0, T_*; H^{m-1}). \quad (83)$$

Every coordinate in (83) is the corresponding distributional derivative of the single maximal history.

The independent rectangle relationship then estimates the complete equation. For every $n \in \mathbb{N}_0$ and $s \geq 0$, put

$$\mathfrak{D}_{n,s}(S) := \nu \|\Lambda^{s+1} V_n(S)\|_2^2 + \int_0^S \left(\|\Lambda^s V_{n+1}\|_2^2 + \nu^2 \|\Lambda^{s+2} V_n\|_2^2 + \|\Lambda^s \nabla Q_n\|_2^2 \right) dt. \quad (84)$$

Then

$$\sup_{0 < S < T_*} \mathfrak{D}_{n,s}(S) \leq \nu \|\Lambda^{s+1} V_n^0\|_2^2 + \|\Lambda^s (F_n - B_n)\|_{L^2(0, T_*; L^2)}^2 < \infty. \quad (85)$$

Thus the selected rectangle bounds, on the same terminal interval, its adjacent temporal face, viscous face, full binomial nonlinear row, normalized pressure-gradient face, and prescribed-force face.

Proof. Let $I = (0, T_*)$. Theorem 4.10 gives

$$V_n \in L^2(I; H^{M+1}), \quad V_{n+1} \in L^2(I; H^{M-1}), \quad 0 \leq n \leq N.$$

For $M = 0$, the second assertion is read through $V_{n+1} \in L^2(I; L^2) \hookrightarrow L^2(I; H^{-1})$. The finitely many selected coordinates therefore have the finite total

$$C_{N,M} := \sum_{n=0}^N \left(\|V_n\|_{L^2(I; H^{M+1})}^2 + \|V_{n+1}\|_{L^2(I; H^{M-1})}^2 \right) < \infty.$$

Restriction decreases each norm, so $\mathfrak{R}_{N,M}(u; T) \leq C_{N,M}$ for $0 < T < T_*$. Taking the supremum proves (82); monotone exhaustion of the nonnegative norm integrals gives in fact $\mathfrak{R}_{N,M}^* = C_{N,M}$. This projection uses the intersection only to identify and place the coordinates; it does not derive the viscous rectangle relationship.

Fix (n, s) . Project the same intersection onto a finite rectangle high enough to contain every velocity factor in B_n at spatial order $s + 2$. The adjacent trace lemma gives the required $L_t^\infty H_x^{s+2}$ factor, while the higher spatial face gives the other factor in $L_t^2 H_x^{s+2}$. Sobolev multiplication, summed over the finite binomial row, gives the explicit same-history estimate

$$\|B_n\|_{L^2(I; \dot{H}^s)} \leq C_s \sum_{a=0}^n \binom{n}{a} \|V_a\|_{L^\infty(I; H^{s+2})} \|V_{n-a}\|_{L^2(I; H^{s+2})} < \infty. \quad (86)$$

The prescribed Schwartz force gives $F_n \in L^2(I; \dot{H}^s)$. Now evaluate the independently derived positive-sign identity (78) on these selected coordinates. Its right-hand side is finite, and taking the supremum over $S < T_*$ proves (85). The pressure term is present because (77) is the unprojected orthogonal completion of the same row. No term of the forced equation and no coordinate of another history has entered the estimate. $\square$

Proposition 5.8 (Complete momentum rows inside one finite rectangle). *Assume the hypotheses of Theorem 5.7. Fix $N \in \mathbb{N}_0$, $M \geq 2$, and $0 < T \leq T_*$, and put*

$$R_v := \sum_{n=0}^N \left(\|V_n\|_{L^2(0,T; H^{M+1})}^2 + \|V_{n+1}\|_{L^2(0,T; H^{M-1})}^2 \right),$$

$$I_{N,M} := \sum_{n=0}^N \|V_n^0\|_{H^M}^2.$$

Then the adjacent faces of this same rectangle give

$$\sum_{n=0}^N \sup_{0 \leq t \leq T} \|V_n(t)\|_{H^M}^2 \leq I_{N,M} + R_v, \quad (87)$$

and the complete binomial nonlinear rows obey

$$\sum_{n=0}^N \|B_n\|_{L^2(0,T; H^{M-1})}^2 \leq C_{N,M} (I_{N,M} + R_v) R_v. \quad (88)$$

For $0 < S \leq T$, define the pressure-complete dissipation record

$$\mathcal{D}_{N,M}(S) := \sum_{n=0}^N \left[\nu \|V_n(S)\|_{\dot{H}^M}^2 + \|V_{n+1}\|_{L^2(0,S;\dot{H}^{M-1})}^2 \right. \\ \left. + \nu^2 \|V_n\|_{L^2(0,S;\dot{H}^{M+1})}^2 + \|\nabla Q_n\|_{L^2(0,S;\dot{H}^{M-1})}^2 \right].$$

The integrated complete rows satisfy

$$\mathcal{D}_{N,M}(S) \leq \nu I_{N,M} + 2 \sum_{n=0}^N \|F_n\|_{L^2(0,T;H^{M-1})}^2 \\ + 2C_{N,M}(I_{N,M} + R_v)R_v. \quad (89)$$

Thus one finite velocity rectangle, together with the displayed prescribed force coordinates, bounds on the same interval and in the same equations its adjacent temporal rows, viscous Laplacians, complete nonlinear rows, and normalized pressure gradients. Lower spatial orders follow by Sobolev nesting from a rectangle with $M \geq 2$.

Proof. For $0 \leq n \leq N$, the Hilbert triple $H^{M+1} \subset H^M \subset H^{M-1}$ and $\partial_t V_n = V_{n+1}$ give

$$\|V_n(t)\|_{H^M}^2 - \|V_n^0\|_{H^M}^2 = 2 \operatorname{Re} \int_0^t \left\langle (1 - \Delta)^{(M+1)/2} V_n, (1 - \Delta)^{(M-1)/2} V_{n+1} \right\rangle_{L^2} ds.$$

Cauchy–Schwarz in time and $2ab \leq a^2 + b^2$, followed by summation in n , prove (87). Sobolev multiplication applied to every allocation in B_n , using one factor in $L^\infty(0,T;H^M)$ and the other in $L^2(0,T;H^{M+1})$, proves (88). Finally, sum (78) over $0 \leq n \leq N$ at $s = M - 1$, use $\|F_n - B_n\|^2 \leq 2\|F_n\|^2 + 2\|B_n\|^2$, and substitute (88). This gives (89). The identities

$$\nabla Q_n = (I - \mathbb{P})(F_n - B_n), \quad V_{n+1} = \nu \Delta V_n - \mathbb{P}B_n + \mathbb{P}F_n$$

show directly that these are the terms of the same pressure-complete row. $\square$

The finite sum above evaluates the coordinates selected by Definition 5.6. Its finiteness comes from the datum-generated intersection, while the complete-equation estimate retains and relates all faces of those coordinates. The next proposition records the exact monotone-exhaustion identity and the equivalent memberships.

Proposition 5.9 (Norm equivalence and whole-terminal realization).

For the fixed maximal history, the family of bounds (82) is equivalent to the adjacent-row statement (83). Whenever either formulation holds, for every $N, M \in \mathbb{N}_0$,

$$\sum_{n=0}^N \left(\|V_n\|_{L^2(0, T_*; H^{M+1})}^2 + \|V_{n+1}\|_{L^2(0, T_*; H^{M-1})}^2 \right) = \lim_{T \uparrow T_*} \mathfrak{R}_{N,M}(u; T) = \mathfrak{R}_{N,M}^*[u_0, \nu, f, T_*]. \quad (90)$$

Every row is the corresponding distributional derivative of the single pressure-normalized maximal history.

Proof. Assume first (83). For fixed N, M , set

$$C_{N,M} := \sum_{n=0}^N \left(\|V_n\|_{L^2(0, T_*; H^{M+1})}^2 + \|V_{n+1}\|_{L^2(0, T_*; H^{M-1})}^2 \right).$$

The sum is finite because it contains finitely many instances of (83). Restriction from $(0, T_*)$ to $(0, T)$ gives

$$\mathfrak{R}_{N,M}(u; T) \leq C_{N,M} \quad (0 < T < T_*),$$

and hence (82).

Conversely, assume (82). Every term in $\mathfrak{R}_{N,M}(u; T)$ is the integral of a nonnegative function. Monotone exhaustion of $(0, T_*)$ gives (90). Taking $N = n$ and $M = m$ yields (83) for each finite (n, m) . Proposition 4.4 identifies every entry with the corresponding distributional derivative of the fixed pressure-normalized history. $\square$

Corollary 5.10 (Whole-terminal completion of every critical height). *Under the hypotheses of Theorem 5.7, every completed critical coordinate belongs to $L^1(0, T_*)$, and*

$$\mathcal{H}_k[u_0, \nu, f, T_*] := \|R_k\|_{L^1(0, T_*)} = \frac{1}{2} \left\| \Lambda^{k+1/2} u \right\|_{L^2(0, T_*; L^2)}^2 \leq \frac{1}{2} \mathfrak{R}_{0,k}^*[u_0, \nu, f, T_*] \quad (k \in \mathbb{N}_0). \quad (91)$$

For every $m \in \mathbb{N}_0$, the spatial component of (56) satisfies the completed-height estimate

$$\|u\|_{L^2(0, T_*; H^m)}^2 \leq 2^m \left(T_* K_f(T_*)^2 + \mathfrak{R}_{0,m}^*[u_0, \nu, f, T_*] \right). \quad (92)$$

Thus

$$(R_k)_{k \in \mathbb{N}_0} \in \prod_{k \in \mathbb{N}_0} L^1(0, T_*), \quad u \in \bigcap_{m \in \mathbb{N}_0} L^2(0, T_*; H^m(\mathbb{R}^3)). \quad (93)$$

Proof. Take $N = 0$ and $M = k$ in (90). Since $|\xi|^{2k+1} \leq \langle \xi \rangle^{2k+2}$,

$$\begin{aligned} 2 \|R_k\|_{L^1(0, T_*)} &= \left\| \Lambda^{k+1/2} u \right\|_{L^2(0, T_*; L^2)}^2 \\ &\leq \|u\|_{L^2(0, T_*; H^{k+1})}^2 \leq \mathfrak{R}_{0,k}^*[u_0, \nu, f, T_*], \end{aligned}$$

which proves (91). For $|\xi| \leq 1$, $\langle \xi \rangle^{2m} \leq 2^m$; for $|\xi| > 1$, $\langle \xi \rangle^{2m} \leq 2^m |\xi|^{2m+1}$. Hence

$$\|v\|_{H^m}^2 \leq 2^m \left(\|v\|_2^2 + \left\| \Lambda^{m+1/2} v \right\|_2^2 \right).$$

Integrating this inequality, using (63) and (91), proves (92). Since k and m are arbitrary finite indices, (93) follows. $\square$

Proposition 5.11 (Pressure completion). *Under the hypotheses of Theorem 5.7, for every $N, M \in \mathbb{N}_0$,*

$$\mathfrak{Q}_{N,M}^*[u_0, \nu, f, T_*] := \sup_{0 < T < T_*} \mathfrak{Q}_{N,M}(p; T) \leq C_M^p (2^{N+1} - 1) \mathfrak{R}_{N,M}^*[u_0, \nu, f, T_*] + \sum_{n=0}^N \mathfrak{F}_{n,0,M}^p(0, T_*). \quad (94)$$

For every $0 \leq n \leq N$, the identities

$$Q_n = R_i R_j \sum_{a=0}^n \binom{n}{a} V_{a,i} V_{n-a,j} - (-\Delta)^{-1} \nabla \cdot F_n, \quad (95)$$

$$V_{n+1} = \nu \Delta V_n - \sum_{a=0}^n \binom{n}{a} (V_a \cdot \nabla) V_{n-a} - \nabla Q_n + F_n \quad (96)$$

hold respectively in $L^1(0, T_*; H^M)$ and $L^1(0, T_*; H^{M-1})$, including when $M = 0$.

Proof. For every $0 < T < T_*$, Theorem 4.8 gives

$$\|Q_n\|_{L^1(0,T;H^M)} + \|\nabla Q_n\|_{L^1(0,T;H^{M-1})} \leq C_M^p 2^n \mathfrak{R}_{N,M}^*[u_0, \nu, f, T_*] + \mathfrak{F}_{n,0,M}^p(0, T).$$

Sum over $0 \leq n \leq N$, then let $T \uparrow T_*$. Monotone convergence in the nonnegative norm integrals proves (94). Since $\nabla \cdot V_a = 0$,

$$(V_a \cdot \nabla) V_{n-a} = \nabla \cdot (V_{n-a} \otimes V_a).$$

Lemma 3.4 and Cauchy–Schwarz in time give

$$\|V_{n-a} \otimes V_a\|_{L^1(0,T_*;H^M)} \leq C_M^{\text{alg}} \|V_a\|_{L^2(0,T_*;H^{M+1})} \|V_{n-a}\|_{L^2(0,T_*;H^{M+1})}.$$

Thus every transport term belongs to $L^1(0, T_*; H^{M-1})$, also for $M = 0$. Moreover,

$$\|\Delta V_n\|_{L^1 H^{M-1}} \leq T_*^{1/2} \|V_n\|_{L^2 H^{M+1}}.$$

Theorem 5.7 places V_{n+1} in $L^2(0, T_*; H^{M-1})$, hence in $L^1(0, T_*; H^{M-1})$. The prescribed row F_n belongs to $L^1(0, T_*; H^{M-1})$, and its pressure potential belongs to the stated spaces by (6). The strict-interval identities are identities of distributions; their whole-terminal representatives belong to the stated spaces. This proves the two identities on $(0, T_*)$. $\square$

Definition 5.12 (Finite-window spaces). For $N, M \in \mathbb{N}_0$, set

$$\mathcal{X}_{N,M} := \prod_{n=0}^N L^2(0, T_*; H^{M+1}) \times \prod_{n=0}^N L^2(0, T_*; H^{M-1}) \times \prod_{n=0}^N L^1(0, T_*; H^M) \times \prod_{n=0}^N L^1(0, T_*; H^{M-1}).$$

The coordinate projection $\pi_{N,M}$ selects the rows through N , their next time derivatives, and the corresponding pressure and force rows from $\mathcal{I}[u_0, \nu, f]$, in the displayed Sobolev spaces. The whole-terminal rectangle is the element

$$\mathcal{R}_{N,M}[u_0, \nu, f; T_*] := \pi_{N,M} \mathcal{I}[u_0, \nu, f] = ((V_n)_{n=0}^N, (V_{n+1})_{n=0}^N; (Q_n)_{n=0}^N; (F_n)_{n=0}^N) \in \mathcal{X}_{N,M}. \quad (97)$$

If $N' \leq N$ and $M' \leq M$, let

$$\rho_{N',M'}^{N,M} : \mathcal{X}_{N,M} \longrightarrow \mathcal{X}_{N',M'}$$

discard the temporal coordinates with index larger than N' and apply

$$H^{M+1} \hookrightarrow H^{M'+1}, \quad H^{M-1} \hookrightarrow H^{M'-1}, \quad H^M \hookrightarrow H^{M'}, \quad H^{M-1} \hookrightarrow H^{M'-1}$$

to the retained coordinates.

Proposition 5.13 (Restriction compatibility). *For $N' \leq N$ and $M' \leq M$,*

$$\rho_{N',M'}^{N,M} \mathcal{R}_{N,M}[u_0, \nu, f; T_*] = \mathcal{R}_{N',M'}[u_0, \nu, f; T_*]. \quad (98)$$

If $(N'', M'') \leq (N', M') \leq (N, M)$, then

$$\rho_{N'',M''}^{N',M'} \rho_{N',M'}^{N,M} = \rho_{N'',M''}^{N,M}.$$

Thus the spaces $\mathcal{X}_{N,M}$ and the maps $\rho_{N',M'}^{N,M}$ form an inverse system, and the datum-generated rectangles form a compatible family in it.

Proof. All retained entries are the fixed distributions $\partial_t^n u$, $\partial_t^{n+1} u$, $\partial_t^n p$, and $\partial_t^n f$. The Sobolev embeddings preserve those distributions. Deletion of rows and composition of canonical embeddings are associative. $\square$

6 Intersection coordinates, endpoint jet, and restart

On every strict horizon, the compatible intersection and the rectangle relations arise independently from the same mixed jet. Theorems 4.10 and 5.7 carry those two constructions to the whole terminal interval of the same forced history. We first identify how the finite records represent $\mathcal{I}[u_0, \nu, f]$, then use their adjacent rows to obtain a common endpoint and restart that history with the same prescribed force.

Definition 6.1 (Finite-coordinate representation of the intersection). Let

$$\mathcal{X}_\infty := \varprojlim_{N,M} \mathcal{X}_{N,M} = \left\{ (X_{N,M})_{N,M} : \begin{array}{l} X_{N,M} \in \mathcal{X}_{N,M}, \\ \rho_{N',M'}^{N,M} X_{N,M} = X_{N',M'} \text{ whenever } N' \leq N, M' \leq M \end{array} \right\}. \quad (99)$$

For a proposed finite maximal time, the canonical coordinate representation of Theorem 4.10 identifies the whole-terminal $\mathcal{I}[u_0, \nu, f]$ with

$$\mathcal{I}[u_0, \nu, f] \simeq (\pi_{N,M} \mathcal{I}[u_0, \nu, f])_{N,M \in \mathbb{N}_0} = (\mathcal{R}_{N,M}[u_0, \nu, f; T_*])_{N,M \in \mathbb{N}_0}. \quad (100)$$

Theorem 6.2 (Compatible finite-coordinate representation).

Let $\nu > 0$, $u_0 \in H_\sigma^\infty(\mathbb{R}^3)$, prescribe $f \in C^\infty([0, \infty); \mathcal{S}(\mathbb{R}^3; \mathbb{R}^3))$, and let (T_, u, p) be the maximal pressure-complete classical history generated by (u_0, ν, f) . If $T_* < \infty$, then*

$$\mathcal{I}[u_0, \nu, f] \simeq (\mathcal{R}_{N,M}[u_0, \nu, f; T_*])_{N,M \in \mathbb{N}_0} \in \mathcal{X}_\infty \quad (101)$$

and all of its finite coordinates are the restrictions of the original pressure-complete history. In this representation, the coordinates $(V_n, Q_n, F_n)_{n \in \mathbb{N}_0}$ retain the memberships stated in Theorem 4.10:

$$V_n \in \bigcap_{m \in \mathbb{N}_0} L^2(0, T_*; H_\sigma^m(\mathbb{R}^3)), \quad (102)$$

$$Q_n \in \bigcap_{m \in \mathbb{N}_0} L^1(0, T_*; H^m(\mathbb{R}^3)), \quad (103)$$

$$F_n \in \bigcap_{m \in \mathbb{N}_0} C([0, T_*]; H^m(\mathbb{R}^3)). \quad (104)$$

Every finite restriction is the corresponding rectangle $\mathcal{R}_{N,M}[u_0, \nu, f; T_*]$. The velocity membership inherited from that theorem is

$$u \in \bigcap_{r,m \in \mathbb{N}_0} H^r(0, T_*; H^m(\mathbb{R}^3)), \quad (105)$$

and, for every $r, m \in \mathbb{N}_0$,

$$\sum_{n=0}^r \|\partial_t^n u\|_{L^2(0, T_*; H^m)}^2 \leq \mathfrak{R}_{r,m}^*[u_0, \nu, f, T_*]. \quad (106)$$

Proof. The coordinate memberships (102), (103), and (105) are inherited from Theorem 4.10. The membership (104) follows directly from the prescribed class $f \in C^\infty([0, \infty); \mathcal{S}(\mathbb{R}^3; \mathbb{R}^3))$. For each finite N, M , apply the projection $\pi_{N,M}$. Its entries are those of the rectangle selected directly from the mixed jet. Theorem 5.7 and Proposition 5.11 give its finite norm bounds. Proposition 5.13 gives

$$\rho_{N',M'}^{N,M} \pi_{N,M} \mathcal{I}[u_0, \nu, f] = \pi_{N',M'} \mathcal{I}[u_0, \nu, f] \quad (N' \leq N, M' \leq M).$$

Hence these projections form an element of $\mathcal{X}_\infty$. Every V_n, Q_n, F_n appears in a sufficiently deep projection, and every spatial derivative is fixed by its distributional velocity, pressure, or force row. The full family consequently identifies precisely $\mathcal{I}[u_0, \nu, f]$; this is a coordinate representation of that object, not another construction of its existence.

Fix finite r, m , put $I = (0, T_*)$, and choose $N \geq r, M \geq m$. The upper rows of that finite record belong to $L^2(I; H^{M+1}) \hookrightarrow L^2(I; H^m)$ for $0 \leq n \leq r$. Proposition 4.4 identifies them as the distributional derivatives $V_n = \partial_t^n u$. Thus the finite record contains every term in the derivative-sum norm of $H^r(I; H^m)$. Taking $N = r, M = m$ gives

$$\begin{aligned} \|u\|_{H^r(I; H^m)}^2 &= \sum_{n=0}^r \|V_n\|_{L^2(I; H^m)}^2 \\ &\leq \sum_{n=0}^r \|V_n\|_{L^2(I; H^{m+1})}^2 \leq \mathfrak{R}_{r,m}^*[u_0, \nu, f, T_*]. \end{aligned}$$

This proves (106) and shows how each mixed Sobolev norm is recovered from a finite record. Its whole-terminal membership is supplied by Theorem 4.10; its finite complete-equation estimate is supplied independently by Theorem 5.7. $\square$

Proposition 6.3 (Complete spatial row and common endpoint). *Let the prescribed datum and finite maximal history be as in Theorem 6.2. Then, for every $m \in \mathbb{N}_0$,*

$$u \in W^{1,1}(0, T_*; H_\sigma^m(\mathbb{R}^3)) \hookrightarrow C([0, T_*]; H_\sigma^m(\mathbb{R}^3)). \quad (107)$$

The endpoint values supplied at the different Sobolev orders are the images of one common state

$$u_* := u(T_*) \in \bigcap_{m \in \mathbb{N}_0} H_\sigma^m(\mathbb{R}^3) = H_\sigma^\infty(\mathbb{R}^3). \quad (108)$$

Proof. Fix m . The zeroth temporal coordinate of (105) gives

$$u \in L^2(0, T_*; H^{m+2}).$$

The pressure-completed nonlinearity satisfies

$$\|\mathbb{P}\nabla \cdot (u \otimes u)\|_{H^m} \leq C_m \|u\|_{H^{m+2}}^2$$

by Lemmas 3.4 and 3.5. It belongs to $L^1(0, T_*; H^m)$. This spatial row and $T_* < \infty$ give $\Delta u \in L^1(0, T_*; H^m)$, and the projected equation

$$u_t = \nu \Delta u - \mathbb{P}\nabla \cdot (u \otimes u) + \mathbb{P}f$$

places u_t in $L^1(0, T_*; H^m)$, because the prescribed source is integrable there in every finite Sobolev order. Also $u \in L^1(0, T_*; H^m)$, so the Banach-valued fundamental theorem gives (107).

If $k \geq m$, the H^k - and H^m -continuous representatives are the original preterminal function after the embedding $H^k \hookrightarrow H^m$. Uniqueness of continuous representatives makes their endpoint values agree under that embedding. These compatible values define the single state in (108). Continuity of divergence from H^{m+1} to H^m gives $\nabla \cdot u_* = 0$. $\square$

Corollary 6.4 (Whole-terminal pressure-complete mixed jet). *Let the prescribed datum and finite maximal history be as in Theorem 6.2. Then, for every $r, m, n \in \mathbb{N}_0$ and $\alpha \in \mathbb{N}_0^3$,*

$$u \in H^r(0, T_*; H_\sigma^m(\mathbb{R}^3)), \quad (109)$$

$$J_{n,\alpha} := \partial_x^\alpha V_n \in L^2(0, T_*; H^{m+1}(\mathbb{R}^3)), \quad (110)$$

$$J_{n+1,\alpha} \in L^2(0, T_*; H^{m-1}(\mathbb{R}^3)), \quad (111)$$

$$\Pi_{n,\alpha} := \partial_x^\alpha Q_n \in L^1(0, T_*; H^m(\mathbb{R}^3)). \quad (112)$$

The following finite bounds hold:

$$\sum_{j=0}^r \|V_j\|_{L^2(0, T_*; H^m)}^2 \leq \mathfrak{R}_{r,m}^*[u_0, \nu, f, T_*], \quad (113)$$

$$\begin{aligned} \|J_{n,\alpha}\|_{L^2(0, T_*; H^{m+1})}^2 &+ \|J_{n+1,\alpha}\|_{L^2(0, T_*; H^{m-1})}^2 \\ &\leq \mathfrak{R}_{n,m+|\alpha|}^*[u_0, \nu, f, T_*], \end{aligned} \quad (114)$$

$$\|\Pi_{n,\alpha}\|_{L^1(0, T_*; H^m)} \leq C_{m+|\alpha|}^p (2^{n+1} - 1) \mathfrak{R}_{n,m+|\alpha|}^*[u_0, \nu, f, T_*] + \mathfrak{F}_{n,\alpha,m}^p(0, T_*). \quad (115)$$

Every mixed row satisfies the recurrences in Proposition 4.4 on $(0, T_*)$.

Proof. Equations (109) and (113) follow from Theorem 6.2. The Fourier multiplier bound

$$\|\partial_x^\alpha h\|_{H^s} \leq \|h\|_{H^{s+|\alpha|}}$$

and the rectangle $(n, m + |\alpha|)$ give (110)–(114). Apply this multiplier bound to the pressure completion in Proposition 5.11 to obtain (112) and (115). Proposition 4.4 supplies the stated identities. $\square$

Theorem 6.5 (Compatible whole-terminal Sobolev traces). *Let the prescribed datum and finite maximal history be as in Theorem 6.2. For $N, m \in \mathbb{N}_0$, define*

$$\mathfrak{E}_{N,m} := (1 + T_*^{-1})^{1/2} (\mathfrak{R}_{N,m}^*[u_0, \nu, f, T_*])^{1/2}. \quad (116)$$

If $0 \leq n \leq N$ and $\alpha \in \mathbb{N}_0^3$, then

$$V_n \in C([0, T_*]; H_\sigma^m(\mathbb{R}^3)), \quad \sup_{0 \leq t \leq T_*} \|V_n(t)\|_{H^m} \leq \mathfrak{E}_{N,m}, \quad (117)$$

$$J_{n,\alpha} \in C([0, T_*]; H^m(\mathbb{R}^3)), \quad \sup_{0 \leq t \leq T_*} \|J_{n,\alpha}(t)\|_{H^m} \leq \mathfrak{E}_{N,m+|\alpha|}. \quad (118)$$

There are unique values

$$V_n^* := \lim_{t \uparrow T_*} V_n(t) \in H^\infty(\mathbb{R}^3), \quad (119)$$

$$J_{n,\alpha}^* := \lim_{t \uparrow T_*} J_{n,\alpha}(t) = \partial_x^\alpha V_n^* \in H^\infty(\mathbb{R}^3), \quad (120)$$

such that, for $k \geq m$, the image of the H^k trace under $H^k \hookrightarrow H^m$ is the H^m trace. Quantitatively,

$$\|V_n^*\|_{H^m} \leq \mathfrak{E}_{n,m}, \quad (121)$$

$$\|V_n(t) - V_n^*\|_{H^m} \leq (T_* - t) \mathfrak{E}_{n+1,m}, \quad 0 \leq t \leq T_*, \quad (122)$$

$$\|J_{n,\alpha}^*\|_{H^m} \leq \mathfrak{E}_{n,m+|\alpha|}, \quad (123)$$

$$\|J_{n,\alpha}(t) - J_{n,\alpha}^*\|_{H^m} \leq (T_* - t) \mathfrak{E}_{n+1,m+|\alpha|}. \quad (124)$$

In particular, the zeroth trace agrees with the common endpoint in Proposition 6.3, and

$$u_* = V_0^* \in H_\sigma^\infty(\mathbb{R}^3), \quad u \in \bigcap_{a,m \in \mathbb{N}_0} C^a([0, T_*]; H_\sigma^m(\mathbb{R}^3)), \quad \partial_t^a u(T_*) = V_a^*. \quad (125)$$

Proof. For $0 \leq n \leq N$, project the intersection $\mathcal{I}[u_0, \nu, f]$ supplied by Theorem 4.10 onto the two coordinates

$$V_n \in L^2(0, T_*; H^{m+1}), \quad \partial_t V_n = V_{n+1} \in L^2(0, T_*; H^{m-1}).$$

The complete mixed intersection supplies both finite norms. With $A = (1 - \Delta)^{1/2}$, the adjacent-row identity is

$$\frac{d}{dt} \|V_n(t)\|_{H^m}^2 = 2 \operatorname{Re} \langle A^{m-1} V_{n+1}(t), A^{m+1} V_n(t) \rangle_{L^2}.$$

Integrating from the initial time and applying Cauchy–Schwarz gives

$$\sup_{0 \leq t \leq T_*} \|V_n(t)\|_{H^m}^2 \leq \|V_n(0)\|_{H^m}^2 + 2 \|V_{n+1}\|_{L^2(I; H^{m-1})} \|V_n\|_{L^2(I; H^{m+1})}, \quad I = (0, T_*). \quad (126)$$

The initial norm is finite by Lemma 4.7. Lemma 3.7 supplies the continuous representative on $[0, T_*]$, justifies the identity by Fourier truncation, and gives the first membership in (117) and

$$\begin{aligned} \sup_{0 \leq t \leq T_*} \|V_n(t)\|_{H^m}^2 &\leq T_*^{-1} \|V_n\|_{L^2 H^m}^2 + 2 \|V_n\|_{L^2 H^{m+1}} \|V_{n+1}\|_{L^2 H^{m-1}} \\ &\leq (1 + T_*^{-1}) \mathfrak{R}_{N,m}^*[u_0, \nu, f, T_*]. \end{aligned}$$

This proves the bound in (117). Apply this lemma to $J_{n,\alpha}$, using (114), to prove (118).

For $k \geq m$, the H^k -continuous and H^m -continuous representatives equal the original preterminal distribution. After applying $H^k \hookrightarrow H^m$, uniqueness of continuous representatives makes them equal on $[0, T_*]$. Their endpoint values are compatible and define (119)–(120). Spatial differentiation is continuous between the corresponding finite Sobolev spaces, so $J_{n,\alpha}^* = \partial_x^\alpha V_n^*$.

Corollary 6.4 gives $V_n, V_{n+1} \in L^1(0, T_*; H^m)$ and $\partial_t V_n = V_{n+1}$ in distributions. The Banach-valued fundamental identity gives

$$V_n(t) - V_n(s) = \int_s^t V_{n+1}(\tau) d\tau \quad \text{in } H^m, \quad 0 \leq s \leq t \leq T_*. \quad (127)$$

The continuous representative of V_{n+1} is bounded by $\mathfrak{E}_{n+1,m}$. Set $t = T_*$ in (127) to prove (122); spatial differentiation gives (124). The remaining size estimates follow from (117)–(118).

The divergence map $H^{m+1} \rightarrow H^m$ is continuous, so $\nabla \cdot u_* = 0$. For each a, m , the curves $V_0, \dots, V_{a+1}$ are continuous in H^m , and (127) identifies the derivative of V_j with V_{j+1} . Induction in j proves (125). $\square$

Theorem 6.6 (Pressure-complete endpoint jet). *Let the prescribed datum and finite maximal history be as in Theorem 6.2. For $n \in \mathbb{N}_0$, define*

$$\Phi_n(t) := (-\Delta)^{-1} \nabla \cdot F_n(t).$$

The prescribed force class gives $\Phi_n \in C^\infty([0, T_ + 1]; H^s)$ for every finite n, s . Define*

$$Q_n^* := R_i R_j \sum_{a=0}^n \binom{n}{a} V_{a,i}^* V_{n-a,j}^* - \Phi_n(T_*). \quad (128)$$

Then $Q_n^ \in H^\infty(\mathbb{R}^3)$, and, for every $s \in \mathbb{N}_0$,*

$$\|Q_n^*\|_{H^s} \leq 9C_s^{\text{alg}} 2^n \mathfrak{E}_{n,s+1}^2 + \|\Phi_n(T_*)\|_{H^s}, \quad (129)$$

$$\|Q_n(t) - Q_n^*\|_{H^s} \leq 9C_s^{\text{alg}} 2^{n+1} (T_* - t) \mathfrak{E}_{n+1,s+1}^2 \quad (130)$$

$$+ (T_* - t) \sup_{0 \leq r \leq T_*} \|\Phi_{n+1}(r)\|_{H^s}. \quad (131)$$

For $\alpha \in \mathbb{N}_0^3$,

$$\Pi_{n,\alpha}^* := \partial_x^\alpha Q_n^* = \lim_{t \uparrow T_*} \Pi_{n,\alpha}(t) \quad \text{in } H^m(\mathbb{R}^3) \text{ for every } m \in \mathbb{N}_0. \quad (132)$$

More precisely, for every $m \in \mathbb{N}_0$,

$$\|\Pi_{n,\alpha}^*\|_{H^m} \leq 9C_{m+|\alpha|}^{\text{alg}} 2^n \mathfrak{E}_{n,m+|\alpha|+1}^2 + \|\partial_x^\alpha \Phi_n(T_*)\|_{H^m}, \quad (133)$$

$$\|\Pi_{n,\alpha}(t) - \Pi_{n,\alpha}^*\|_{H^m} \leq 9C_{m+|\alpha|}^{\text{alg}} 2^{n+1} (T_* - t) \mathfrak{E}_{n+1,m+|\alpha|+1}^2 \quad (134)$$

$$+ (T_* - t) \sup_{0 \leq r \leq T_*} \|\partial_x^\alpha \Phi_{n+1}(r)\|_{H^m}. \quad (135)$$

The endpoint rows satisfy, for every $n, q \in \mathbb{N}_0$,

$$V_{n+1}^* = \nu \Delta V_n^* - \sum_{a=0}^n \binom{n}{a} (V_a^* \cdot \nabla) V_{n-a}^* - \nabla Q_n^* + F_n(T_*) \quad \text{in } H^q(\mathbb{R}^3). \quad (136)$$

Among families rooted at u_* , carrying the prescribed values $(F_n(T_*))_{n \in \mathbb{N}_0}$, and satisfying (128) and (136) in every finite Sobolev space, $(V_n^*, Q_n^*)_{n \in \mathbb{N}_0}$ is the unique pressure-normalized endpoint jet. Moreover,

$$p \in \bigcap_{a, m \in \mathbb{N}_0} C^a([0, T_*]; H^m(\mathbb{R}^3)), \quad \partial_t^a p(T_*) = Q_a^*. \quad (137)$$

Proof. Lemmas 3.4 and 3.5, followed by (117), give

$$\begin{aligned} \|Q_n^*\|_{H^s} &\leq 9C_s^{\text{alg}} \sum_{a=0}^n \binom{n}{a} \|V_a^*\|_{H^{s+1}} \|V_{n-a}^*\|_{H^{s+1}} + \|\Phi_n(T_*)\|_{H^s} \\ &\leq 9C_s^{\text{alg}} 2^n \mathfrak{E}_{n,s+1}^2 + \|\Phi_n(T_*)\|_{H^s}. \end{aligned}$$

This is (129). For each product in $Q_n(t) - Q_n^*$, insert $V_a^* V_{n-a}(t)$, use (122), and bound every retained factor by $\mathfrak{E}_{n+1,s+1}$. The two product differences and $\sum_{a=0}^n \binom{n}{a} = 2^n$ prove the product part of (131). Since $\partial_t \Phi_n = \Phi_{n+1}$,

$$\|\Phi_n(t) - \Phi_n(T_*)\|_{H^s} \leq (T_* - t) \sup_{0 \leq r \leq T_*} \|\Phi_{n+1}(r)\|_{H^s},$$

which supplies the remaining term. The Fourier multiplier inequality

$$\|\partial_x^\alpha h\|_{H^m} \leq \|h\|_{H^{m+|\alpha|}}$$

applied to the nonlinear Riesz-product component, together with the exact derivative $\partial_x^\alpha \Phi_n$ of the force potential, proves (133) and (135). The latter tends to zero as $t \uparrow T_*$, which proves (132) in every H^m .

Fix n, q . The trace of V_n in H^{q+2} carries the diffusion term in Proposition 4.4 to H^q at T_* . The product estimate at order $q+1$, using velocity convergence in H^{q+2} , carries every transport term to H^q . Equation (131) at order $q+1$ carries the pressure gradient to H^q . The left side converges to V_{n+1}^* in H^q , and $F_n(t) \rightarrow F_n(T_*)$ in H^q , which proves (136).

The value $V_0^* = u_*$ fixes Q_0^* , and the velocity recurrence fixes V_1^* . Inductively, $V_0^*, \dots, V_n^*$ fix Q_n^* , which fixes V_{n+1}^* . This proves uniqueness within the displayed recurrence class. Extend each Q_n to T_* by $Q_n(T_*) := Q_n^*$. On $(0, T_*)$, $Q_n = \partial_t^n p$, so $\partial_t Q_n = Q_{n+1}$ in distributions. Both rows are continuous in every H^m by (131). Hence, for $0 \leq s \leq t \leq T_*$,

$$Q_n(t) - Q_n(s) = \int_s^t Q_{n+1}(\tau) d\tau \quad \text{in } H^m(\mathbb{R}^3). \quad (138)$$

Indeed, the identity holds for $t < T_*$, and the endpoint bounds permit passage to $t = T_*$. Thus $Q_n \in C^1([0, T_*]; H^m)$ with derivative Q_{n+1} . Induction gives (137). $\square$

Lemma 6.7 (Late-slice overlap uniqueness). *Let the prescribed datum and finite maximal history be as in Theorem 6.2. Define*

$$\Gamma_* := \sup_{0 \leq s \leq T_*} \|\mathbb{P}f\|_{L^2(s, s+1; H^2)} < \infty.$$

Define

$$\delta_0 := \min\{1, c_\nu(1 + \mathfrak{E}_{0,3} + \Gamma_*)^{-2}\} > 0, \quad (139)$$

where c_ν is the constant in Theorem 3.11. For every $t_0 \in (\max\{0, T_* - \delta_0/2\}, T_*)$, let

$$(w^{t_0}, q^{t_0}) \in \mathcal{M}_{\nu,f}^3(t_0, t_0 + \delta_0; u(t_0)), \quad (w^*, q^*) \in \mathcal{M}_{\nu,f}^3(T_*, T_* + \delta_0; u_*)$$

be the unique pairs supplied by Theorem 3.11. Then

$$(w^{t_0}, q^{t_0}) = (u, p) \quad \text{on } [t_0, T_*], \quad (140)$$

$$(w^{t_0}, q^{t_0}) = (w^*, q^*) \quad \text{on } [T_*, t_0 + \delta_0]. \quad (141)$$

Proof. Equation (117) gives

$$\|u(t_0)\|_{H^3} \leq \mathfrak{E}_{0,3}, \quad \|u_*\|_{H^3} \leq \mathfrak{E}_{0,3}.$$

Hence

$$\delta_0 \leq \delta_{\nu,f}(t_0, u(t_0)), \quad \delta_0 \leq \delta_{\nu,f}(T_*, u_*),$$

with $\delta_{\nu,f}$ as in Theorem 3.11. Theorem 3.11 gives both pairs on the stated intervals. For each $t_0 < b < T_*$, Lemma 3.10 places $(u, p)|_{[t_0, b]}$ and $(w^{t_0}, q^{t_0})|_{[t_0, b]}$ in $\mathcal{M}_{\nu,f}^3(t_0, b; u(t_0))$. Uniqueness in that class gives (140). Continuity in H^3 gives

$$w^{t_0}(T_*) = \lim_{t \uparrow T_*} u(t) = u_*.$$

Lemma 3.10 places the restrictions of (w^{t_0}, q^{t_0}) and (w^*, q^*) to $[T_*, t_0 + \delta_0]$ in $\mathcal{M}_{\nu,f}^3(T_*, t_0 + \delta_0; u_*)$. Uniqueness in that class gives (141). The right overlap has positive length because $t_0 + \delta_0 > T_* + \delta_0/2$. $\square$

Theorem 6.8 (Same-forced-history endpoint restart). *Let the prescribed datum and finite maximal history be as in Theorem 6.2. There is a pressure-normalized forced classical solution, with the same prescribed force $f(t)$, $(\tilde{u}, \tilde{p})$ on $[0, T_* + \delta_0/2] \times \mathbb{R}^3$ such that*

$$(\tilde{u}, \tilde{p})|_{[0, T_*] \times \mathbb{R}^3} = (u, p), \quad (142)$$

$$\tilde{u}, \tilde{p} \in C^\infty([0, T_* + \delta_0/2] \times \mathbb{R}^3), \quad (143)$$

$$\partial_t^n \tilde{u}(T_*) = V_n^*, \quad \partial_t^n \tilde{p}(T_*) = Q_n^* \quad (n \in \mathbb{N}_0). \quad (144)$$

If $0 < \eta \leq \delta_0/2$ and $(\hat{u}, \hat{p})$ is another pressure-normalized classical extension satisfying

$$(\hat{u}, \hat{p})|_{[0, T_*]} = (u, p), \quad (\hat{u}, \hat{p})|_{[T_*, T_* + \eta]} \in \mathcal{M}_{\nu,f}^3(T_*, T_* + \eta; u_*), \quad (145)$$

then

$$(\hat{u}, \hat{p}) = (\tilde{u}, \tilde{p}) \quad \text{on } [0, T_* + \eta]. \quad (146)$$

Proof. Let (w^*, q^*) be the endpoint solution from Lemma 6.7, and set

$$W_n^+ := \partial_t^n w^*(T_*), \quad P_n^+ := \partial_t^n q^*(T_*).$$

The restart datum is exactly

$$w^*(T_*) = u_* = u(T_*).$$

The equation continues with the same absolute-time source $f(t)$. If the right-hand interval is parametrized by $\tau = t - T_*$, that same source is written $f(T_* + \tau)$; this is only the shifted time coordinate. Theorem 3.11 makes (w^*, q^*) smooth and pressure-normalized. Differentiating its equations exactly as in the proof of Proposition 4.4 gives

$$\begin{aligned} P_n^+ &= R_i R_j \sum_{a=0}^n \binom{n}{a} W_{a,i}^+ W_{n-a,j}^+ - (-\Delta)^{-1} \nabla \cdot F_n(T_*), \\ W_{n+1}^+ &= \nu \Delta W_n^+ - \sum_{a=0}^n \binom{n}{a} (W_a^+ \cdot \nabla) W_{n-a}^+ - \nabla P_n^+ + F_n(T_*). \end{aligned}$$

Since $W_0^+ = u_* = V_0^*$, induction with (128) and (136) gives

$$W_n^+ = V_n^*, \quad P_n^+ = Q_n^*, \quad n \in \mathbb{N}_0. \quad (147)$$

Define

$$(\tilde{u}, \tilde{p})(t) := \begin{cases} (u, p)(t), & 0 \leq t < T_*, \\ (w^*, q^*)(t), & T_* \leq t \leq T_* + \delta_0/2. \end{cases}$$

Equations (125), (137), and (147) match every one-sided time derivative in every finite Sobolev space. Sobolev embedding gives (143), and the pressure-complete equation holds at T_* by (136). Lemma 6.7 identifies the joined pair near T_* with the single local history (w^{t_0}, q^{t_0}) , so the gluing is one local forced classical history across the endpoint; the piecewise definition gives (142). Any pair in (145) has that datum u_* on the right of T_* and belongs to the shifted mild class containing (w^*, q^*) . The at-most-one assertion in Theorem 3.11 proves (146). $\square$

The datum-generated whole-terminal intersection of Theorem 4.10, projected through the independently derived complete-equation rectangle identities by Theorem 5.7, gives compatible velocity traces through T_* in Theorem 6.5. The pressure recurrence fixes their normalized endpoint jet in Theorem 6.6. With those endpoint values, Theorem 6.8 extends the original history beyond T_* . A proposed finite maximal time would admit a classical continuation. Corollary 6.9 applies this contradiction, and Theorem 6.10 collects the global solution, pressure normalization, and energy identity.

Corollary 6.9 (Maximal-time contradiction for prescribed smooth data). *Let $\nu > 0$, $u_0 \in H_\sigma^\infty(\mathbb{R}^3)$, prescribe $f \in C^\infty([0, \infty); \mathcal{S}(\mathbb{R}^3; \mathbb{R}^3))$, and let (T_*, u, p) be the maximal pressure-complete classical history generated by (u_0, ν, f) . If $T_* < \infty$, then it admits a classical extension beyond T_* , contradicting maximality.*

Proof. Theorem 4.10 supplies the compatible whole-terminal mixed jet, and Theorem 5.7 projects that record through the independently derived complete-equation rectangle identities. The chain from Theorem 6.2 through Theorem 6.8 produces a pressure-normalized classical extension to $[0, T_* + \delta_0/2]$, where $\delta_0 > 0$. This contradicts maximality in Definition 4.2. $\square$

Theorem 6.10 (Global smoothness and uniqueness for prescribed smooth data). *For every $\nu > 0$, every $u_0 \in H_\sigma^\infty(\mathbb{R}^3)$, and every prescribed $f \in C^\infty([0, \infty); \mathcal{S}(\mathbb{R}^3; \mathbb{R}^3))$, there exists a pressure-normalized pair*

$$(u, p) \in C^\infty([0, \infty) \times \mathbb{R}^3; \mathbb{R}^3 \times \mathbb{R}) \quad (148)$$

satisfying

$$\partial_t u + (u \cdot \nabla)u + \nabla p = \nu \Delta u + f, \quad (149)$$

$$\nabla \cdot u = 0, \quad u(0) = u_0, \quad (150)$$

$$p(t) = R_i R_j (u_i(t) u_j(t)) - (-\Delta)^{-1} \nabla \cdot f(t), \quad \lim_{|x| \rightarrow \infty} p(t, x) = 0, \quad (151)$$

$$\frac{1}{2} \|u(t)\|_{L^2}^2 + \nu \int_0^t \|\nabla u(s)\|_{L^2}^2 ds = \frac{1}{2} \|u_0\|_{L^2}^2 + \int_0^t \langle f(s), u(s) \rangle_{L^2} ds, \quad t \geq 0. \quad (152)$$

In particular, for every finite t ,

$$\|u(t)\|_{L^2(\mathbb{R}^3)} \leq \|u_0\|_{L^2} + \int_0^t \|\mathbb{P}f(s)\|_{L^2} ds < \infty. \quad (153)$$

For every finite $T > 0$, its restriction is the unique member of $\mathcal{M}_{\nu, f}^3(0, T; u_0)$.

Proof. Let

$$\mathcal{T} := \left\{ \tau > 0 : \begin{array}{l} \text{there exists a pressure-complete classical history} \\ (\tau, u^\tau, p^\tau) \text{ generated by } (u_0, \nu, f) \end{array} \right\},$$

and put

$$T_* := \sup \mathcal{T}.$$

Theorem 3.11, applied to $u_0 \in H_\sigma^\infty$, supplies a shifted mild solution whose persistence statement gives every finite velocity and pressure row. Sobolev embedding and the displayed recurrences make this pair a pressure-complete classical history, so $\mathcal{T}$ is nonempty. Lemma 3.10 admits each strict classical restriction to the shifted mild class, whose at-most-one property identifies any two such solutions on their common interval; their union defines a unique solution on $[0, T_*)$. Since $u_0 \in H_\sigma^\infty$, the persistence conclusion of Theorem 3.11, applied on overlapping local intervals, places every strict restriction in the class of Definition 4.2. An extension beyond T_* would restrict to an interval $[0, \tau]$ with $\tau > T_*$, contrary to the definition of the supremum. Thus (T_*, u, p) is maximal. Corollary 6.9 gives $T_* = \infty$.

Fix $T < \infty$. Persistence in Theorem 3.11, iterated along the unique history, retains every finite spatial Sobolev order on $[0, T]$. Proposition 4.4 retains every finite temporal row. Lemma 3.2 gives (148); arbitrariness of T gives the global class. The pressure belongs to $H^s(\mathbb{R}^3)$ for every s , and the Fourier inversion proof of Lemma 3.2 places its continuous representative in $C_0(\mathbb{R}^3)$, which proves the limit in (151). Lemma 3.10 places $(u, p)|_{[0, T]}$ in $\mathcal{M}_{\nu, f}^3(0, T; u_0)$. If $(\tilde{u}, \tilde{p}) \in \mathcal{M}_{\nu, f}^3(0, T; u_0)$, the at-most-one assertion in Theorem 3.11 gives $(\tilde{u}, \tilde{p}) = (u, p)$ on $[0, T]$. This proves the stated uniqueness.

Proposition 5.1 holds on every finite interval. Since T was arbitrary, it gives (152) for every $t \geq 0$; its regularized square-root estimate gives (153). $\square$

7 Matrix forms of the completed intersection

Theorem 4.10 assembles the datum-generated compatible intersection of the single forced mixed jet. Independently, the positive-sign viscous cell gives the rectangle relationship. Theorem 5.7 projects the established intersection onto a finite selection and integrates that relationship. The matrices in this section display how the velocity, nonlinear, pressure, viscous, and prescribed-force coordinates remain coupled inside each such rectangle.

Proposition 7.1 (Mixed-index transport matrix and pressure completion). *Let*

$$\Gamma := \mathbb{N}_0 \times \mathbb{N}_0^3, \quad \kappa = (n, \alpha), \quad \lambda = (a, \beta),$$

and write

$$\lambda \preceq \kappa \iff a \leq n \text{ and } \beta \leq \alpha, \quad \kappa - \lambda := (n - a, \alpha - \beta), \quad \binom{\kappa}{\lambda} := \binom{n}{a} \binom{\alpha}{\beta}.$$

Set

$$J_\kappa := J_{n,\alpha}, \quad \Pi_\kappa := \Pi_{n,\alpha}, \quad F_\kappa := F_{n,\alpha}, \quad \Psi_\kappa := (-\Delta)^{-1} \nabla \cdot F_\kappa, \quad \mathbf{J} := (J_\lambda)_{\lambda \in \Gamma}.$$

The block operator

$$\mathbb{L}(J)_{\kappa,\lambda} := \begin{cases} \binom{\kappa}{\lambda} \mathbb{P}[(J_{\kappa-\lambda} \cdot \nabla) \cdot], & \lambda \preceq \kappa, \\ 0, & \lambda \not\preceq \kappa, \end{cases} \quad (154)$$

has the finite block-row action

$$[\mathbb{L}(J)\mathbf{J}]_\kappa = \sum_{\lambda \preceq \kappa} \binom{\kappa}{\lambda} \mathbb{P}[(J_\lambda \cdot \nabla) J_{\kappa-\lambda}]. \quad (155)$$

Consequently the complete mixed row is the pair

$$J_{n+1,\alpha} + [\mathbb{L}(J)\mathbf{J}]_\kappa = \nu \Delta J_\kappa + \mathbb{P} F_\kappa, \quad (156)$$

$$\Pi_\kappa = R_i R_j \sum_{\lambda \preceq \kappa} \binom{\kappa}{\lambda} J_{\lambda,i} J_{\kappa-\lambda,j} - \Psi_\kappa, \quad (157)$$

while the unprojected velocity identity retains $\nabla \Pi_\kappa$ and the full force coordinate F_κ exactly as in (44).

Order Γ by increasing $n + |\alpha|$, with any fixed ordering among indices of equal depth. Then $\mathbb{L}(J)$ is block lower triangular. For

$$\begin{aligned}\kappa! &:= n! \alpha!, & \alpha! &:= \alpha_1! \alpha_2! \alpha_3!, \\ \widehat{J}_\kappa &:= \frac{J_\kappa}{\kappa!}, & \widehat{\Pi}_\kappa &:= \frac{\Pi_\kappa}{\kappa!}, \\ \widehat{F}_\kappa &:= \frac{F_\kappa}{\kappa!}, & \widehat{\Psi}_\kappa &:= \frac{\Psi_\kappa}{\kappa!} = (-\Delta)^{-1} \nabla \cdot \widehat{F}_\kappa.\end{aligned}$$

let D be the diagonal matrix with $D_{\kappa,\kappa} = \kappa!$. On every finite downward-closed truncation of Γ ,

$$D^{-1} \mathbb{L}(D\widehat{J})D = \widehat{\mathbb{L}}(\widehat{J}), \quad \widehat{\mathbb{L}}(\widehat{J})_{\kappa,\lambda} := \begin{cases} \mathbb{P}[(\widehat{J}_{\kappa-\lambda} \cdot \nabla) \cdot], & \lambda \preceq \kappa, \\ 0, & \lambda \not\preceq \kappa. \end{cases} \quad (158)$$

Thus the factorial-normalized pressure-complete recurrence is

$$(n+1)\widehat{J}_{n+1,\alpha} + \sum_{\lambda \preceq \kappa} (\widehat{J}_\lambda \cdot \nabla) \widehat{J}_{\kappa-\lambda} + \nabla \widehat{\Pi}_\kappa = \nu \Delta \widehat{J}_\kappa + \widehat{F}_\kappa, \quad (159)$$

$$\widehat{\Pi}_\kappa = R_i R_j \sum_{\lambda \preceq \kappa} \widehat{J}_{\lambda,i} \widehat{J}_{\kappa-\lambda,j} - \widehat{\Psi}_\kappa. \quad (160)$$

The normalized transport blocks depend on the difference $\kappa - \lambda$; hence (158) is the block lower-triangular convolution form of the mixed transport array. The prescribed force is the affine row source, and its gradient component remains in the pressure coordinate through $\widehat{\Psi}_\kappa$.

Proof. For $\lambda \preceq \kappa$, change the summation index from λ to $\kappa - \lambda$ and use $\binom{\kappa}{\kappa-\lambda} = \binom{\kappa}{\lambda}$. This gives (155). Apply $\mathbb{P}$ to (44). The identities

$$\mathbb{P}J_{n+1,\alpha} = J_{n+1,\alpha}, \quad \mathbb{P}\Delta J_\kappa = \Delta J_\kappa, \quad \mathbb{P}\nabla \Pi_\kappa = 0, \quad \mathbb{P}F_{n,\alpha} = \mathbb{P}F_\kappa$$

follow from (45) and the Fourier symbols. They prove (156); equation (157) is (43) in mixed-index notation. The paired projected and pressure formulas, including $-\Psi_\kappa$, retain every prescribed force component: the solenoidal component remains in $\mathbb{P}F_\kappa$, while the gradient component is retained by Π_κ .

If $\lambda \preceq \kappa$, then $a + |\beta| \leq n + |\alpha|$, which proves lower triangularity in the stated ordering. Finally,

$$\binom{\kappa}{\lambda} \lambda! (\kappa - \lambda)! = \kappa!.$$

Dividing the mixed velocity, pressure, and prescribed force rows by $\kappa!$ proves (159) and (160). This factorial cancellation in each operator block proves (158). Every row has $(n+1) \prod_{j=1}^3 (\alpha_j + 1)$ supported blocks, so all displayed row sums are finite. $\square$

Proposition 7.2 (Sobolev rectangle matrix and adjacent stencil). *For $0 < T < T_*$, $N, M \in \mathbb{N}_0$, $0 \leq n \leq N$, and $m \in \mathbb{Z}$, set*

$$G_{n,m}(T) := \|V_n\|_{L^2(0,T;H^m)},$$

and set

$$\begin{aligned}P_{n,M}(T) &:= \|Q_n\|_{L^1(0,T;H^M)} + \|\nabla Q_n\|_{L^1(0,T;H^{M-1})}, \\ D_{n,M}(T) &:= \|F_n\|_{L^1(0,T;H^{M-1})} + \mathfrak{F}_{n,0,M}^P(0,T).\end{aligned}$$

The object matrix of the finite pressure-complete forced rectangle is

$$\mathbf{R}_{N,M}[u_0, \nu, f; T] := \begin{pmatrix} V_0|_{(0,T)} & V_1|_{(0,T)} & Q_0|_{(0,T)} & F_0|_{(0,T)} \\ V_1|_{(0,T)} & V_2|_{(0,T)} & Q_1|_{(0,T)} & F_1|_{(0,T)} \\ \vdots & \vdots & \vdots & \vdots \\ V_N|_{(0,T)} & V_{N+1}|_{(0,T)} & Q_N|_{(0,T)} & F_N|_{(0,T)} \end{pmatrix}. \quad (161)$$

The corresponding matrix of the two velocity norms, the pressure budget, and the prescribed-force budget is

$$\mathbf{C}_{N,M}(T) := \begin{pmatrix} G_{0,M+1}(T) & G_{1,M-1}(T) & P_{0,M}(T) & D_{0,M}(T) \\ G_{1,M+1}(T) & G_{2,M-1}(T) & P_{1,M}(T) & D_{1,M}(T) \\ \vdots & \vdots & \vdots & \vdots \\ G_{N,M+1}(T) & G_{N+1,M-1}(T) & P_{N,M}(T) & D_{N,M}(T) \end{pmatrix}. \quad (162)$$

If $\mathbf{G}_{N,M}(T)$ denotes its first two columns, then

$$\|\mathbf{G}_{N,M}(T)\|_{\mathbb{F}}^2 = \mathfrak{R}_{N,M}(u; T), \quad \sum_{n=0}^N P_{n,M}(T) = \mathfrak{Q}_{N,M}(p; T), \quad \sum_{n=0}^N D_{n,M}(T) = \mathfrak{F}_{N,M}(f; T). \quad (163)$$

For each $0 \leq n \leq N$, the first two entries in row n form the adjacent Hilbert-scale stencil

$$\left. \begin{array}{l} (n, M+1) : \quad V_n \in L^2(0, T; H^{M+1}), \\ (n+1, M-1) : \quad V_{n+1} = \partial_t V_n \in L^2(0, T; H^{M-1}) \end{array} \right\} \longrightarrow (n, M) : \quad V_n \in C([0, T]; H^M). \quad (164)$$

For every $m \in \mathbb{N}_0$, the Sobolev matrix measures the mixed-index array through the exact identity

$$G_{n,m}(T)^2 = \sum_{\substack{\alpha \in \mathbb{N}_0^3 \\ |\alpha| \leq m}} \frac{m!}{(m - |\alpha|)! \alpha!} \|J_{n,\alpha}\|_{L^2((0,T) \times \mathbb{R}^3)}^2. \quad (165)$$

The pressure entry in row n is $Q_n = \Pi_{n,0}$, and its spatial rows $\partial_x^\alpha Q_n = \Pi_{n,\alpha}$ are fixed by (157), including the force-pressure potential. The force entry is the prescribed row $F_n = \partial_t^n f$, whose spatial rows are $F_{n,\alpha} = \partial_x^\alpha F_n$; these are the same force coordinates that occur in the mixed velocity and pressure rows.

Proof. The object matrix is the tuple in Definition 5.6, arranged by temporal row. The three identities in (163) are (79) and (80)–(81). Lemma 3.7 gives (164). Finally, Plancherel and

$$(1 + |\xi|^2)^m = \sum_{\substack{\alpha \in \mathbb{N}_0^3 \\ |\alpha| \leq m}} \frac{m!}{(m - |\alpha|)! \alpha!} \xi^{2\alpha}$$

give (165). The pressure and force claims follow from the definitions of $Q_n, \Pi_{n,\alpha}, F_n, F_{n,\alpha}$ and (157). $\square$

Proposition 7.3 (Whole-terminal closed estimate matrix). *Under the hypotheses of Theorem 5.7, fix $N, M \in \mathbb{N}_0$ and set*

$$\begin{aligned} a_{n,M} &:= \|V_n\|_{L^2(0, T_*; H^{M+1})}, \\ b_{n,M} &:= \|V_{n+1}\|_{L^2(0, T_*; H^{M-1})}, \\ q_{n,M} &:= \|Q_n\|_{L^1(0, T_*; H^M)} + \|\nabla Q_n\|_{L^1(0, T_*; H^{M-1})}, \\ d_{n,M} &:= \|F_n\|_{L^1(0, T_*; H^{M-1})} + \mathfrak{F}_{n,0,M}^P(0, T_*), \quad 0 \leq n \leq N. \end{aligned} \quad (166)$$

Define entrywise order on nonnegative matrices by $A \preceq_{\text{ent}} B$ when $A_{ij} \leq B_{ij}$ for every entry. Then, for every $0 < T < T_*$,

$$\mathbf{C}_{N,M}(T) \preceq_{\text{ent}} \mathbf{C}_{N,M}^* := \begin{pmatrix} a_{0,M} & b_{0,M} & q_{0,M} & d_{0,M} \\ a_{1,M} & b_{1,M} & q_{1,M} & d_{1,M} \\ \vdots & \vdots & \vdots & \vdots \\ a_{N,M} & b_{N,M} & q_{N,M} & d_{N,M} \end{pmatrix}. \quad (167)$$

Let $\mathbf{G}_{N,M}^*$ denote the first two columns of $\mathbf{C}_{N,M}^*$. Then

$$\begin{aligned} \mathfrak{R}_{N,M}(u; T) &= \|\mathbf{G}_{N,M}(T)\|_{\text{F}}^2 \\ &\leq \|\mathbf{G}_{N,M}^*\|_{\text{F}}^2 = \sum_{n=0}^N (a_{n,M}^2 + b_{n,M}^2) = \mathfrak{R}_{N,M}^*[u_0, \nu, f, T_*] < \infty. \end{aligned} \quad (168)$$

The other two columns satisfy the exact whole-terminal identities

$$\sum_{n=0}^N q_{n,M} = \mathfrak{Q}_{N,M}^*[u_0, \nu, f, T_*] < \infty, \quad (169)$$

$$\sum_{n=0}^N d_{n,M} =: \mathfrak{F}_{N,M}^*[f, T_*] < \infty. \quad (170)$$

Proof. Proposition 5.9 makes every $a_{n,M}$ and $b_{n,M}$ finite. Proposition 5.11 makes every $q_{n,M}$ finite. The prescribed force class, together with (6), makes every $d_{n,M}$ finite. Restriction from $(0, T_*)$ to $(0, T)$ decreases each nonnegative coordinate norm and gives (167). Taking Frobenius norms and using (163) together with (90) proves (168). Monotone exhaustion of the pressure and prescribed-force columns gives (169) and (170). $\square$

8 Quantitative consequences of the compatible rectangles

8.1 Completion in the global H^1 mild class

The forced construction begins with velocity data in $H_\sigma^\infty(\mathbb{R}^3)$, not only with Schwartz data, while retaining the one prescribed force throughout. The datum-generated whole-terminal intersection and same-force continuation transfer, after positive-time smoothing, to the strong H^1 class for that force.

Theorem 8.1 (Global H^1 completion). *Let $\nu > 0$, let $a \in H_\sigma^1(\mathbb{R}^3)$, and prescribe $f \in C^\infty([0, \infty); \mathcal{S}(\mathbb{R}^3; \mathbb{R}^3))$. Then there is a unique pressure-normalized mild solution*

$$u^a \in C([0, \infty); H_\sigma^1) \cap L_{\text{loc}}^2([0, \infty); H_\sigma^2), \quad u^a \in C^\infty((0, \infty) \times \mathbb{R}^3), \quad (171)$$

of the forced three-dimensional Navier–Stokes equations with datum a , viscosity ν , and the same prescribed force f . Its pressure is

$$p^a = R_i R_j (u_i^a u_j^a) - (-\Delta)^{-1} \nabla \cdot f$$

in distributions. The solution obeys the forced kinetic-energy identity on every finite interval.

Proof. Set $g := \mathbb{P}f$, and write

$$G_0 := \|g\|_{L^2(0,1;L^2)}, \quad R_{a,f} := 1 + \|\nabla a\|_2^2 + 2\nu^{-1}G_0^2.$$

Write

$$Y(t) := \|\nabla u(t)\|_2^2, \quad Z(t) := \|\Delta u(t)\|_2^2, \quad c_1 := \frac{27}{16}S(6)^6.$$

For a smooth solenoidal approximation, the L^2 pairing of the projected equation with $-\Delta u$ gives

$$\frac{1}{2}Y'(t) + \nu Z(t) = \operatorname{Re} \langle (u \cdot \nabla)u, \Delta u \rangle_{L^2} - \operatorname{Re} \langle g, \Delta u \rangle_{L^2}. \quad (172)$$

The explicit three-dimensional Sobolev inequality and interpolation give

$$\|u\|_6 \leq S(6)Y^{1/2}, \quad \|\nabla u\|_3 \leq S(6)^{1/2}Y^{1/4}Z^{1/4}.$$

Consequently,

$$|\langle (u \cdot \nabla)u, \Delta u \rangle_{L^2}| \leq S(6)^{3/2}Y^{3/4}Z^{3/4} \leq \frac{\nu}{2}Z + \frac{27S(6)^6}{32\nu^3}Y^3,$$

while

$$|\langle g, \Delta u \rangle_{L^2}| \leq \frac{\nu}{4}Z + \nu^{-1}\|g\|_2^2.$$

Consequently,

$$Y'(t) + \frac{\nu}{2}Z(t) \leq c_1\nu^{-3}Y(t)^3 + 2\nu^{-1}\|g(t)\|_2^2. \quad (173)$$

The integral inequality in (173), with the force contribution kept in the same bootstrap, shows that

$$Y(t) \leq 2R_{a,f} \quad (174)$$

whenever $0 \leq t \leq 1$ and $8c_1\nu^{-3}R_{a,f}^2t \leq 1$.

The estimate holds on the explicit lifespan

$$\begin{aligned} \tau_{\nu,f}(a) &:= \min \left\{ 1, \frac{\nu^3}{8c_1R_{a,f}^2} \right\}, \\ \sup_{0 \leq t \leq \tau_{\nu,f}(a)} Y(t) &\leq 2R_{a,f}, \quad \int_0^{\tau_{\nu,f}(a)} Z(t) dt < \infty. \end{aligned} \quad (175)$$

Let $\mathcal{J}_k$ be the orthogonal Fourier projection to $\{|\xi| \leq k\}$. On the range of $\mathcal{J}_k$, the regularized problem

$$\partial_t u^{(k)} - \nu \Delta u^{(k)} = -\mathcal{J}_k \mathbb{P} \nabla \cdot (u^{(k)} \otimes u^{(k)}) + \mathcal{J}_k g, \quad u^{(k)}(0) = \mathcal{J}_k a, \quad (176)$$

is a Banach-space ordinary differential equation in H_σ^1 . Indeed, if

$$\mathcal{N}_k(z) := \mathcal{J}_k \mathbb{P} \nabla \cdot (z \otimes z),$$

then the Bernstein inequalities on the range of $\mathcal{J}_k$ give

$$\|\mathcal{N}_k(z) - \mathcal{N}_k(y)\|_{H^1} \leq C_k(\|z\|_{H^1} + \|y\|_{H^1})\|z - y\|_{H^1}, \quad C_k < \infty. \quad (177)$$

Picard's theorem supplies a maximal trajectory. The projection is self-adjoint and commutes with $\mathbb{P}$, ∇ , and Δ , so the forced kinetic identity holds along that trajectory. Since

$$\|z\|_{H^1}^2 \leq (1 + k^2)\|z\|_2^2 \quad (z = \mathcal{J}_k z), \quad (178)$$

the forced kinetic bound $\|u^{(k)}(t)\|_2 \leq \|a\|_2 + \int_0^t \|g(s)\|_2 \, ds$ and the Banach-space continuation criterion extend the trajectory for all positive times. Estimate (173) holds for $u^{(k)}$, with constants independent of k . Equations (174)–(175) give

$$\sup_k \left(\|u^{(k)}\|_{L^\infty(0, \tau_{\nu, f}(a); H^1)} + \|u^{(k)}\|_{L^2(0, \tau_{\nu, f}(a); H^2)} \right) < \infty. \quad (179)$$

The nonlinear estimate

$$\|\mathbb{P}\nabla \cdot (u \otimes u)\|_2 \leq \|u\|_6 \|\nabla u\|_3 \leq S(6)^{3/2} Y^{3/4} Z^{1/4},$$

shows that the regularized nonlinear terms are bounded in $L^2(0, \tau_{\nu, f}(a); L^2)$, uniformly in k . The equations and $\mathcal{J}_k g \in L^2(0, \tau_{\nu, f}(a); L^2)$ bound $\partial_t u^{(k)}$ in this space. Weak compactness and the Aubin–Lions–Simon lemma [1, 15] on every ball give, after diagonal extraction,

$$\begin{aligned} u^{(k)} &\overset{*}{\rightharpoonup} u && \text{in } L^\infty(0, \tau_{\nu, f}(a); H^1), \\ u^{(k)} &\rightharpoonup u && \text{in } L^2(0, \tau_{\nu, f}(a); H^2), \\ u^{(k)} &\longrightarrow u && \text{in } L^2(0, \tau_{\nu, f}(a); H^1(B_R)) \quad (R < \infty). \end{aligned}$$

The last convergence passes the quadratic term in distributions; lower semicontinuity preserves (179). The displayed nonlinear estimate now gives $\mathbb{P}\nabla \cdot (u \otimes u) \in L^2 L^2$, and the equation gives $u_t \in L^2(0, \tau_{\nu, f}(a); L^2)$. For every $\phi \in C_c^\infty(\mathbb{R}^3; \mathbb{R}^3)$ with $\nabla \cdot \phi = 0$ and every $\eta \in C_c^\infty([0, \tau_{\nu, f}(a)))$, passage to the limit in the time-integrated regularized equation gives

$$\begin{aligned} - \int_0^{\tau_{\nu, f}(a)} \langle u, \phi \rangle_{L^2} \eta' \, dt + \nu \int_0^{\tau_{\nu, f}(a)} \langle \nabla u, \nabla \phi \rangle_{L^2} \eta \, dt - \int_0^{\tau_{\nu, f}(a)} \langle u \otimes u, \nabla \phi \rangle_{L^2} \eta \, dt \\ = \langle a, \phi \rangle_{L^2} \eta(0) + \int_0^{\tau_{\nu, f}(a)} \langle g, \phi \rangle_{L^2} \eta \, dt. \end{aligned} \quad (180)$$

The right-hand side fixes the weak initial trace as a .

The Lions–Magenes theorem [13] for $H_\sigma^2 \hookrightarrow H_\sigma^1 \hookrightarrow L_\sigma^2$ gives $u \in C([0, \tau_{\nu, f}(a)]; H_\sigma^1)$; its continuous trace at zero is the trace in (180), hence $u(0) = a$. Variation of constants gives the mild equation, and $p = R_i R_j (u_i u_j) - (-\Delta)^{-1} \nabla \cdot f$ supplies its normalized pressure. Since $u_t \in L^2 L^2$ and $u \in L^2 H^2$, pairing the limiting equation with u is legitimate; incompressibility cancels the transport term and proves the forced kinetic-energy identity.

If u, v are two such solutions and $w = u - v$, incompressibility cancels the $(u \cdot \nabla)w$ pairing and gives

$$\frac{1}{2} \frac{d}{dt} \|w\|_2^2 + \nu \|\nabla w\|_2^2 = -\operatorname{Re} \langle (w \cdot \nabla)v, w \rangle_{L^2}.$$

This Sobolev constant yields

$$|\langle (w \cdot \nabla)v, w \rangle_{L^2}| \leq S(6)^{3/2} \|\nabla v\|_2 \|w\|_2^{1/2} \|\nabla w\|_2^{3/2} \leq \frac{\nu}{2} \|\nabla w\|_2^2 + \frac{27S(6)^6}{32\nu^3} \|\nabla v\|_2^4 \|w\|_2^2.$$

Gronwall’s inequality proves uniqueness. This calculation one derivative higher also gives the finite-interval stability needed below. Indeed, pairing the difference equation with $-\Delta w$, integrating the $(v \cdot \nabla)w$ term once in space, and applying Young’s inequality gives

$$\frac{d}{dt} \|\nabla w\|_2^2 + \nu \|\Delta w\|_2^2 \leq \frac{C_{\text{st}}}{\nu} \left(\|\nabla u\|_3^2 + \|\nabla v\|_3^2 \right) \|\nabla w\|_2^2. \quad (181)$$

The coefficient is integrable because

$$\|\nabla u\|_3^2 \leq 5(6) \|\nabla u\|_2 \|\Delta u\|_2,$$

and likewise for v . The L^2 difference estimate and (181) show continuous dependence in $C([0, S]; H^1)$ on every finite interval $[0, S]$.

Let $c_0 > 0$ be the absolute smallness constant in the full-space H^1 local existence theorem, Theorem 5.4(ii) of [16], written at viscosity one as $\|b\|_{H^1}^4 h \leq c_0$. After viscosity rescaling, its condition is

$$h \|b\|_{H^1}^4 \leq c_0 \nu^3. \quad (182)$$

Fix $0 < \varepsilon < S \leq \tau_{\nu, f}(a)$, and put

$$K_{S, f} := 1 + \sup_{0 \leq t \leq S} \|u(t)\|_{H^1} + \sup_{0 \leq t \leq S} \|f(t)\|_{H^2}.$$

Choose

$$0 < h \leq \min \left\{ \varepsilon, c_0 \nu^3 K_{S, f}^{-4} \right\}.$$

The intervals $[0, h]$ and $[t - h, t]$, $h \leq t \leq S$, obey (182). The quantitative positive-time regularity iteration in Lemma 5.5 of [16], after viscosity rescaling with interior time $h/2$, applies to the inhomogeneous mild formula as well: each additional source term is bounded directly by the displayed smooth-force norms. The same heat-semigroup/product iteration therefore gives, for every $m \in \mathbb{N}$, with

$$\begin{aligned} \Gamma_{m, S}(f) &:= \sum_{j=0}^m \sup_{0 \leq t \leq S} \left\| \partial_t^j f(t) \right\|_{H^{m+2}}, \\ \sup_{\varepsilon \leq t \leq S} \|u(t)\|_{H^m} &\leq C_{m, \varepsilon, S}(\nu, K_{S, f}, \Gamma_{m, S}(f)) < \infty. \end{aligned} \quad (183)$$

Apply the mild formula from time $\varepsilon/2$. The bounds (183), the heat multiplier, and Lemma 3.4 show that $u \in C([\varepsilon, S]; H^m)$ for every m . The identities

$$p = R_i R_j (u_i u_j) - (-\Delta)^{-1} \nabla \cdot f, \quad u_t = \nu \Delta u - \mathbb{P} \nabla \cdot (u \otimes u) + g \quad (184)$$

then give $p \in C([\varepsilon, S]; H^m)$ and $u_t \in C([\varepsilon, S]; H^m)$ for every m . Differentiating (184) and the pressure identity, and applying this product estimate, proves inductively that all temporal derivatives of u and p belong to every spatial Sobolev space continuously on $[\varepsilon, S]$. Sobolev embedding proves $(u, p) \in C^\infty((0, \tau_{\nu, f}(a)] \times \mathbb{R}^3)$.

Choose $0 < t_0 < \tau_{\nu, f}(a)$. Equation (183) places $u(t_0)$ in H_σ^∞ . Theorem 4.10 applies to that smoothed datum and the same shifted prescribed force. Therefore Theorem 6.10, with time origin t_0 , supplies a global smooth history from that datum with the same absolute-time force $f(t)$. The uniqueness estimate above identifies it with the local history on their overlap, so the two histories glue to (171). This estimate proves uniqueness in the full displayed class. The forced kinetic identity on the first interval and the classical forced identity after t_0 agree at the common trace and prove the final assertion. $\square$

8.2 Canonical coordinates and finite majorants

The compatible rectangles carry more information than the endpoint argument uses. We first name their finite coordinates and close the estimates that will feed the mixed rows, frequency tails, and continuation bounds below.

Definition 8.2 (Rectangle and trace coordinates). Let $u_0 \in H_\sigma^\infty(\mathbb{R}^3)$, $\nu > 0$, prescribe $f \in C^\infty([0, \infty); \mathcal{S}(\mathbb{R}^3; \mathbb{R}^3))$, and let (u, p) be the pressure-normalized solution generated by (u_0, ν, f) . Let $0 < T < \infty$ satisfy either $T < T_*$ or $T = T_* < \infty$. Set $I_T = (0, T)$,

$$V_n := \partial_t^n u, \quad Q_n := \partial_t^n p, \quad F_n := \partial_t^n f, \quad \Phi_n := (-\Delta)^{-1} \nabla \cdot F_n, \quad n \in \mathbb{N}_0.$$

Define the monotone rectangle coordinate

$$\mathfrak{P}_{N,M}[u_0, \nu, f; T] := 1 + \sup_{0 < S < T} \mathfrak{R}_{N,M}(u; S), \quad (185)$$

and the trace constant

$$\mathfrak{T}_{N,M}[u_0, \nu, f; T] := (1 + T^{-1})^{1/2} \mathfrak{P}_{N,M}[u_0, \nu, f; T]^{1/2}. \quad (186)$$

All occurrences of $\mathfrak{P}_{N,M}$ and $\mathfrak{T}_{N,M}$ below carry the arguments $[u_0, \nu, f; T]$. The velocity datum u_0 , viscosity ν , prescribed force f , pressure-normalized solution (u, p) , horizon T , interval I_T , and row notation V_n, Q_n, F_n, Φ_n fixed here remain in force throughout the rest of this section. In the terminal alternative $T = T_* < \infty$, the whole-terminal representatives are those of Proposition 5.9, and every endpoint value used below is the compatible trace supplied by Theorem 6.5.

Lemma 8.3 (Rectangle-coordinate finiteness). *For every $N, M \in \mathbb{N}_0$,*

$$1 \leq \mathfrak{P}_{N,M}[u_0, \nu, f; T] < \infty. \quad (187)$$

Proof. If $T = T_* < \infty$, equation (82) proves (187). If $T < T_*$, classical persistence on $[0, T]$ gives

$$\sup_{0 \leq t \leq T} \|V_n(t)\|_{H^m} < \infty \quad (n, m \in \mathbb{N}_0),$$

and (79) proves (187). □

Lemma 8.4 (Fixed analytic constants). *For $r \in \mathbb{N}_0$, set*

$$A_r := 2C_r^{\text{alg}}.$$

Then

$$\|R_i R_j(f_i g_j)\|_{H^r} + \|\mathbb{P} \nabla \cdot (f \otimes g)\|_{H^{r-1}} \leq A_r \|f\|_{H^{r+1}} \|g\|_{H^{r+1}}. \quad (188)$$

This constant satisfies

$$\|R_i R_j(f_i g_j)\|_{H^r} + \|\nabla R_i R_j(f_i g_j)\|_{H^{r-1}} \leq A_r \|f\|_{H^{r+1}} \|g\|_{H^{r+1}}. \quad (189)$$

For $s, k \in \mathbb{N}_0$, $2 \leq b \leq \infty$, and $s > k + \frac{3}{2} - \frac{3}{b}$, define

$$\nabla^k f := (\partial^\alpha f)_{|\alpha|=k}, \quad \|\nabla^k f\|_{L^b} := \left\| \left(\sum_{|\alpha|=k} |\partial^\alpha f|^2 \right)^{1/2} \right\|_{L^b},$$

and set

$$d_k := \#\{\alpha \in \mathbb{N}_0^3 : |\alpha| = k\}, \quad \frac{1}{q_b} := \frac{1}{2} - \frac{1}{b},$$

with $q_2 = \infty$, and

$$S_{s,k,b} := d_k^{1/2} (2\pi)^{-3(1/2-1/b)} \left\| |\xi|^k \langle \xi \rangle^{-s} \right\|_{L_\xi^{q_b}}. \quad (190)$$

Then $S_{s,k,b} < \infty$ and

$$\|\nabla^k f\|_{L^b} \leq S_{s,k,b} \|f\|_{H^s}. \quad (191)$$

For $m \geq 2$, set $L_m := \max\{1, C_m^{\text{alg}}\}$. Then

$$\|fg\|_{H^m} \leq L_m \|f\|_{H^m} \|g\|_{H^m}, \quad (192)$$

$$\|e^{\nu t \Delta} \mathbb{P} \nabla \cdot F\|_{H^m} \leq L_m (\nu t)^{-1/2} \|F\|_{H^m}, \quad t > 0. \quad (193)$$

Proof. Lemmas 3.4 and 3.5, applied componentwise before summing squares, give

$$\begin{aligned} \|R_i R_j (f_i g_j)\|_{H^r} &\leq \|f \otimes g\|_{H^r} \leq C_r^{\text{alg}} \|f\|_{H^{r+1}} \|g\|_{H^{r+1}}, \\ \|\mathbb{P} \nabla \cdot (f \otimes g)\|_{H^{r-1}} &\leq \|f \otimes g\|_{H^r} \leq C_r^{\text{alg}} \|f\|_{H^{r+1}} \|g\|_{H^{r+1}}. \end{aligned}$$

Moreover,

$$\|\nabla R_i R_j (f_i g_j)\|_{H^{r-1}} \leq \|R_i R_j (f_i g_j)\|_{H^r}.$$

This proves (188) and (189).

Let $b' = b/(b-1)$. Interpolation between Fourier inversion $L_\xi^1 \rightarrow L_x^\infty$, with norm $(2\pi)^{-3/2}$, and Plancherel gives

$$\|\mathcal{F}^{-1} h\|_{L_x^{b'}} \leq (2\pi)^{-3(1/2-1/b)} \|h\|_{L_\xi^{b'}}.$$

For $|\alpha| = k$, Hölder's inequality with $1/b' = 1/2 + 1/q_b$ gives

$$\|\partial^\alpha f\|_{L^b} \leq (2\pi)^{-3(1/2-1/b)} \left\| |\xi|^k \langle \xi \rangle^{-s} \right\|_{L^{q_b}} \|f\|_{H^s}.$$

The weight belongs to L^{q_b} under the displayed strict inequality for s ; for $b = 2$, its L^∞ -norm is finite because $s > k$. Summation over $|\alpha| = k$ proves (191).

The first estimate in (192) is Lemma 3.4. For the heat estimate, the symbol of $\mathbb{P} \nabla \cdot$, acting from tensors to vectors, has norm at most $|\xi|$, while

$$\sup_{\rho \geq 0} \rho e^{-\nu t \rho^2} = (2e\nu t)^{-1/2}.$$

Plancherel gives

$$\|e^{\nu t \Delta} \mathbb{P} \nabla \cdot F\|_{H^m} \leq (2e\nu t)^{-1/2} \|F\|_{H^m} \leq L_m (\nu t)^{-1/2} \|F\|_{H^m}.$$

□

Definition 8.5 (Canonical rectangle majorants). For $N, M \in \mathbb{N}_0$, define

$$\mathfrak{B}_{N,M}[u_0, \nu, f; T] := \inf \{B \geq 0 : \mathfrak{R}_{N,M}(u; S) \leq B \text{ for every } 0 < S < T\}. \quad (194)$$

Monotone convergence gives

$$\mathfrak{B}_{N,M}[u_0, \nu, f; T] = \begin{cases} \mathfrak{R}_{N,M}(u; T), & T < T_*, \\ \mathfrak{R}_{N,M}^*[u_0, \nu, f, T_*], & T = T_* < \infty. \end{cases} \quad (195)$$

The pressure-normalized history in Definition 8.2 is fixed for (u_0, ν, f) . Thus $\mathfrak{B}_{N,M}[u_0, \nu, f; T]$ is the exact minimal rectangle bound of that datum-generated history, not a supremum over a family of solutions and not an independently chosen constant. Equations (163) and (194) give the exact matrix identities

$$\mathfrak{B}_{N,M}[u_0, \nu, f; T] = \sup_{0 < S < T} \|\mathbf{G}_{N,M}(S)\|_{\mathbb{F}}^2, \quad (196)$$

$$\mathfrak{P}_{N,M}[u_0, \nu, f; T] = 1 + \sup_{0 < S < T} \|\mathbf{G}_{N,M}(S)\|_{\mathbb{F}}^2. \quad (197)$$

Consequently,

$$\mathfrak{P}_{N,M}[u_0, \nu, f; T] = 1 + \mathfrak{B}_{N,M}[u_0, \nu, f; T]. \quad (198)$$

For $n \in \mathbb{N}_0$ and $m \in \mathbb{Z}$, define

$$\begin{aligned} \mathfrak{G}_{n,m}^{(0)} &:= \|F_n(0)\|_{H^m}, & \mathfrak{G}_{n,m}^{(1)} &:= \|F_n\|_{L^1(I_T; H^m)}, \\ \mathfrak{G}_{n,m}^{(2)} &:= \|F_n\|_{L^2(I_T; H^m)}, & \mathfrak{G}_{n,m}^{(\infty)} &:= \sup_{0 \leq t \leq T} \|F_n(t)\|_{H^m}, \end{aligned} \quad (199)$$

For $n, m \in \mathbb{N}_0$, define the force-pressure coordinates

$$\begin{aligned} \mathfrak{L}_{n,m}^{(1)} &:= \|\Phi_n\|_{L^1(I_T; H^m)} + \|\nabla \Phi_n\|_{L^1(I_T; H^{m-1})}, & \mathfrak{L}_{n,m}^{(2)} &:= \|\Phi_n\|_{L^2(I_T; H^m)}, \\ \mathfrak{L}_{n,m}^{(\infty)} &:= \sup_{0 \leq t \leq T} (\|\Phi_n(t)\|_{H^m} + \|\nabla \Phi_n(t)\|_{H^{m-1}}). \end{aligned} \quad (200)$$

Every one of these coordinates is finite by the prescribed force class and (6). They belong to the same forced history and are not independently chosen sources.

For $n, m \in \mathbb{N}_0$, define the initial-jet majorants $\mathfrak{D}_{n,m} = \mathfrak{D}_{n,m}[u_0, \nu, f]$ by

$$\mathfrak{D}_{0,m} := \|u_0\|_{H^m}, \quad (201)$$

$$\mathfrak{D}_{n+1,m} := \nu \mathfrak{D}_{n,m+2} + A_{m+1} \sum_{a=0}^n \binom{n}{a} \mathfrak{D}_{a,m+2} \mathfrak{D}_{n-a,m+2} + \mathfrak{G}_{n,m}^{(0)}. \quad (202)$$

Define the pointwise recursion $\mathfrak{Y}_{n,m} = \mathfrak{Y}_{n,m}[u_0, \nu, f; T]$ by

$$\mathfrak{Y}_{0,0} := K_f(T), \quad (203)$$

$$\mathfrak{Y}_{0,m} := \min \left\{ (1 + T^{-1})^{1/2} \mathfrak{B}_{0,m}^{1/2}, \|u_0\|_{H^m} + T^{1/2} \mathfrak{B}_{0,m+1}^{1/2} \right\}, \quad m \geq 1, \quad (204)$$

$$\mathfrak{Y}_{n+1,m} := \nu \mathfrak{Y}_{n,m+2} + A_{m+1} \sum_{a=0}^n \binom{n}{a} \mathfrak{Y}_{a,m+2} \mathfrak{Y}_{n-a,m+2} + \mathfrak{G}_{n,m}^{(\infty)}. \quad (205)$$

For $m \in \{-1\} \cup \mathbb{N}_0$, put

$$m_+ := \max\{m, 0\}, \quad \mu(m) := \max\{m - 1, 0\},$$

and define

$$\tilde{\mathfrak{X}}_{n,m} := \min \left\{ T^{1/2} \mathfrak{Y}_{n,m_+}, \mathfrak{B}_{n,\mu(m)}^{1/2} \right\}. \quad (206)$$

Set

$$\mathfrak{X}_{0,0} := \min \left\{ \tilde{\mathfrak{X}}_{0,0}, T^{1/2} K_f(T) \right\}, \quad (207)$$

$$\mathfrak{X}_{0,1} := \min \left\{ \tilde{\mathfrak{X}}_{0,1}, \left(T + \frac{1}{2\nu} \right)^{1/2} K_f(T) \right\}, \quad (208)$$

$$\mathfrak{X}_{n,m} := \tilde{\mathfrak{X}}_{n,m} \quad \text{for all other } (n, m). \quad (209)$$

For $n, m \in \mathbb{N}_0$, define

$$\mathfrak{Z}_{n,m}[u_0, \nu, f; T] := \min \left\{ \mathfrak{Y}_{n,m}, (1 + T^{-1})^{1/2} \mathfrak{B}_{n,m}^{1/2}, \mathfrak{D}_{n,m} + T^{1/2} \mathfrak{X}_{n+1,m} \right\}. \quad (210)$$

All $\mathfrak{B}, \mathfrak{D}, \mathfrak{X}, \mathfrak{Y}, \mathfrak{Z}, \mathfrak{G}$, and $\mathfrak{L}$ in this section carry the arguments $[u_0, \nu, f; T]$, except that $\mathfrak{D}$ is independent of T .

Proposition 8.6 (Exact viscosity and horizon scaling). *Let*

$$a := \nu^{-1} u_0, \quad v(s, x) := \nu^{-1} u(s/\nu, x), \quad q(s, x) := \nu^{-2} p(s/\nu, x), \quad \tilde{f}(s, x) := \nu^{-2} f(s/\nu, x).$$

Then (v, q) is the unit-viscosity solution generated by $(a, 1, \tilde{f})$. If

$$\tilde{V}_n := \partial_s^n v, \quad \tilde{Q}_n := \partial_s^n q, \quad \tilde{F}_n := \partial_s^n \tilde{f},$$

then

$$V_n(t) = \nu^{n+1} \tilde{V}_n(\nu t), \quad Q_n(t) = \nu^{n+2} \tilde{Q}_n(\nu t), \quad F_n(t) = \nu^{n+2} \tilde{F}_n(\nu t). \quad (211)$$

Substitution of (211) into the defining norms gives

$$\begin{aligned} \mathfrak{B}_{N,M}[u_0, \nu, f; T] &= \sum_{n=0}^N \left[\nu^{2n+1} \left\| \tilde{V}_n \right\|_{L^2(0, \nu T; H^{M+1})}^2 \right. \\ &\quad \left. + \nu^{2n+3} \left\| \tilde{V}_{n+1} \right\|_{L^2(0, \nu T; H^{M-1})}^2 \right], \end{aligned} \quad (212)$$

$$\begin{aligned} \|Q_n\|_{L^1(0, T; H^m)} + \|\nabla Q_n\|_{L^1(0, T; H^{m-1})} &= \nu^{n+1} \left(\left\| \tilde{Q}_n \right\|_{L^1(0, \nu T; H^m)} \right. \\ &\quad \left. + \left\| \nabla \tilde{Q}_n \right\|_{L^1(0, \nu T; H^{m-1})} \right), \end{aligned} \quad (213)$$

$$\sup_{0 \leq t \leq T} \|Q_n(t)\|_{H^m} = \nu^{n+2} \sup_{0 \leq s \leq \nu T} \left\| \tilde{Q}_n(s) \right\|_{H^m}. \quad (214)$$

The initial-jet recursion obeys

$$\mathfrak{D}_{n,m}[u_0, \nu, f] = \nu^{n+1} \mathfrak{D}_{n,m}[u_0/\nu, 1, \tilde{f}]. \quad (215)$$

Write

$$E_\sigma^u(t) := \frac{1}{2} \|\Lambda^\sigma u(t)\|_2^2, \quad E_\sigma^v(s) := \frac{1}{2} \|\Lambda^\sigma v(s)\|_2^2.$$

For every $j \in \mathbb{N}_0$, the critical-height rows obey

$$E_{j+1/2}^u(t) = \nu^2 E_{j+1/2}^v(\nu t), \quad (216)$$

$$\int_0^T E_{j+1/2}^u(t) dt = \nu \int_0^{\nu T} E_{j+1/2}^v(s) ds. \quad (217)$$

For every $t_0 \geq 0$ and $b \in H_\sigma^3(\mathbb{R}^3)$, put

$$f_{\nu, t_0}(\sigma, x) := \nu^{-2} f(t_0 + \sigma/\nu, x).$$

The maximal elapsed lifespan in the shifted mild class has the exact covariance

$$\tau_{\nu, f, 3}^{\max}(t_0, b) = \nu^{-1} \tau_{1, f_{\nu, t_0}, 3}^{\max}(0, b/\nu). \quad (218)$$

Proof. The change of variables gives $u(t) = \nu v(\nu t)$ and $p(t) = \nu^2 q(\nu t)$. Differentiation proves (211); the substitution $s = \nu t$ proves (212)–(214). The force-row identity in (211) makes the force term in (202) homogeneous of the same degree as V_{n+1} . Equation (215) therefore follows by induction in (201)–(202). The amplitude relation $u(t) = \nu v(\nu t)$ proves (216), and the substitution $s = \nu t$ proves (217). For $(w, \pi) \in \mathcal{M}_{\nu, f}^3(t_0, t_0 + \delta; b)$, set

$$\mathcal{S}_\nu(w, \pi)(\sigma) := (\nu^{-1} w(t_0 + \sigma/\nu), \nu^{-2} \pi(t_0 + \sigma/\nu)), \quad 0 \leq \sigma \leq \nu \delta.$$

Time translation and this change of variables show that $\mathcal{S}_\nu$ is a bijection from $\mathcal{M}_{\nu, f}^3(t_0, t_0 + \delta; b)$ onto $\mathcal{M}_{1, f_{\nu, t_0}}^3(0, \nu \delta; b/\nu)$. Its inverse is the corresponding amplitude and time rescaling, which proves (218). $\square$

Proposition 8.7 (Height-coordinate re-expression of the finite rows). *For $m \in \mathbb{N}_0$, define*

$$\mathfrak{H}_m[u_0, \nu, f; T] := \frac{1}{2} \left\| \Lambda^{m+1/2} u \right\|_{L^2(I_T; L^2)}^2, \quad (219)$$

$$\mathfrak{S}_m[u_0, \nu, f; T]^2 := 2^m (TK_f(T)^2 + 2\mathfrak{H}_m[u_0, \nu, f; T]). \quad (220)$$

Then

$$2\mathfrak{H}_m[u_0, \nu, f; T] \leq \mathfrak{B}_{0,m}[u_0, \nu, f; T], \quad \|u\|_{L^2(I_T; H^m)} \leq \mathfrak{S}_m[u_0, \nu, f; T]. \quad (221)$$

Define recursively, for $n, m \in \mathbb{N}_0$,

$$\mathfrak{A}_{0,m}^\vee := T^{-1/2} \mathfrak{S}_m + \nu T^{1/2} \mathfrak{S}_{m+2} + A_{m+1} \mathfrak{S}_{m+2}^2 + \mathfrak{G}_{0,m}^{(1)}, \quad (222)$$

$$\mathfrak{A}_{n,m}^\vee := A_m \sum_{a=0}^n \binom{n}{a} \mathfrak{A}_{a,m+1}^\vee \mathfrak{A}_{n-a,m+1}^\vee + \mathfrak{L}_{n,m}^{(\infty)}, \quad (223)$$

$$\mathfrak{A}_{n+1,m}^\vee := \nu \mathfrak{A}_{n,m+2}^\vee + A_{m+1} \sum_{a=0}^n \binom{n}{a} \mathfrak{A}_{a,m+2}^\vee \mathfrak{A}_{n-a,m+2}^\vee + \mathfrak{G}_{n,m}^{(\infty)}. \quad (224)$$

Every finite quantity in this recursion is finite, and

$$\sup_{0 \leq t \leq T} \|V_n(t)\|_{H^m} \leq \mathfrak{A}_{n,m}^v, \quad (225)$$

$$\sup_{0 \leq t \leq T} (\|Q_n(t)\|_{H^m} + \|\nabla Q_n(t)\|_{H^{m-1}}) \leq \mathfrak{A}_{n,m}^p. \quad (226)$$

Let $M_- := \max\{M - 1, 0\}$ and set

$$\widehat{\mathfrak{B}}_{N,M}^h[u_0, \nu, f; T] := T \sum_{n=0}^N \left((\mathfrak{A}_{n,M+1}^v)^2 + (\mathfrak{A}_{n+1,M_-}^v)^2 \right), \quad (227)$$

$$\widehat{\mathfrak{Q}}_{N,M}^h[u_0, \nu, f; T] := T \sum_{n=0}^N \mathfrak{A}_{n,M}^p. \quad (228)$$

Then

$$\mathfrak{B}_{N,M}[u_0, \nu, f; T] \leq \widehat{\mathfrak{B}}_{N,M}^h[u_0, \nu, f; T], \quad (229)$$

$$\mathfrak{Q}_{N,M}(p; T) \leq \widehat{\mathfrak{Q}}_{N,M}^h[u_0, \nu, f; T]. \quad (230)$$

For fixed N, M , the right-hand sides use only

$$\begin{aligned} & \nu, \quad T, \quad K_f(T), \quad N, \quad M, \\ & \{\mathfrak{H}_q[u_0, \nu, f; T] : 0 \leq q \leq M + 2N + 4\}, \quad \{A_q : 0 \leq q \leq M + 2N + 4\}, \\ & \{\mathfrak{G}_{0,b}^{(1)} : 0 \leq b \leq M + 2N + 2\}, \\ & \{\mathfrak{G}_{a,b}^{(\infty)} : 0 \leq a \leq N, \quad 0 \leq b \leq M + 2N + 2\}, \quad \{\mathfrak{L}_{a,M}^{(\infty)} : 0 \leq a \leq N\}. \end{aligned} \quad (231)$$

The quantities $\mathfrak{H}_q$ are the exact completed-height coordinates of the fixed datum-generated history. The whole-terminal intersection and the rectangle theorem first establish the finite rows. Equations (221) and (229) then express that same finite information in rectangle and height coordinates, respectively.

Proof. The first estimate in (221) follows from the upper row of $\mathfrak{B}_{0,m}$. The Fourier inequality

$$(1 + |\xi|^2)^m \leq 2^m (1 + |\xi|^{2m+1})$$

and (63) give the second estimate.

The projected equation and (188) give

$$\|u_t\|_{L^1(I_T; H^m)} \leq \nu T^{1/2} \mathfrak{G}_{m+2} + A_{m+1} \mathfrak{G}_{m+2}^2 + \mathfrak{G}_{0,m}^{(1)}.$$

Choose a Lebesgue time $t_m \in I_T$ satisfying

$$\|u(t_m)\|_{H^m} \leq T^{-1/2} \|u\|_{L^2(I_T; H^m)}.$$

The Banach-valued fundamental identity gives (225) at $n = 0$. Assume that bound through row n . The forced pressure recurrence and (189) give (226). The projected temporal recurrence with $\mathbb{P}F_n$ retained and (188) give the next velocity row through (224). Induction proves (225)–(226).

Integrating the pointwise bounds over I_T gives (229); when $M = 0$, use $H^0 \hookrightarrow H^{-1}$ on the lower row. This integration in (226) gives (230). Induction in (224) gives the finite dependency range in (231). $\square$

Lemma 8.8 (Closed velocity majorants). *For every $n, m \in \mathbb{N}_0$,*

$$\|V_n(0)\|_{H^m} \leq \mathfrak{D}_{n,m}, \quad (232)$$

$$\|V_n\|_{L^2(I_T; H^m)} \leq \mathfrak{X}_{n,m}, \quad (233)$$

$$\sup_{0 \leq t \leq T} \|V_n(t)\|_{H^m} \leq \mathfrak{Z}_{n,m}. \quad (234)$$

For $m = -1$, equation (233) remains valid. Every right-hand side is finite and depends on (u_0, ν, f, n, m, T) through the finite rectangle-matrix suprema in (196) and the corresponding finite prescribed-force coordinates.

Proof. Applying $\mathbb{P}$ to the temporal recurrence gives

$$V_{n+1} = \nu \Delta V_n - \mathbb{P} \nabla \cdot \sum_{a=0}^n \binom{n}{a} (V_a \otimes V_{n-a}) + \mathbb{P} F_n. \quad (235)$$

Equation (188) at order $m+1$, with $\mathbb{P} F_n$ retained and evaluated at $t=0$, proves (232) by induction through (202). This estimate, taken pointwise on $[0, T]$, proves by induction that

$$\sup_{0 \leq t \leq T} \|V_n(t)\|_{H^m} \leq \mathfrak{Y}_{n,m}. \quad (236)$$

The base case $m=0$ is the forced estimate (63). For $m \geq 1$, both terms in (204) are valid: the first is the adjacent trace bound, and the second follows from

$$u(t) = u_0 + \int_0^t V_1(s) \, ds \quad \text{in } H^m.$$

The rectangle with middle order $\mu(m)$, Sobolev nesting, and (236) give respectively

$$\|V_n\|_{L^2(I_T; H^m)} \leq \mathfrak{B}_{n, \mu(m)}^{1/2}, \quad \|V_n\|_{L^2(I_T; H^m)} \leq T^{1/2} \mathfrak{Y}_{n, m+}.$$

This proves (233) through (206); the two special forced-energy cases follow from (63) and (64). The adjacent trace lemma gives

$$\sup_{0 \leq t \leq T} \|V_n(t)\|_{H^m} \leq (1 + T^{-1})^{1/2} \mathfrak{B}_{n, m}^{1/2},$$

while the Banach-valued fundamental identity and (232) give

$$\sup_{0 \leq t \leq T} \|V_n(t)\|_{H^m} \leq \mathfrak{D}_{n, m} + T^{1/2} \mathfrak{X}_{n+1, m}.$$

Taking the minimum of these bounds and (236) proves (234). $\square$

Definition 8.9 (Closed pressure majorants). For $n, m \in \mathbb{N}_0$, define

$$\mathfrak{Q}_{n, m}^{(1)} := A_m \sum_{a=0}^n \binom{n}{a} \mathfrak{X}_{a, m+1} \mathfrak{X}_{n-a, m+1} + \mathfrak{L}_{n, m}^{(1)}, \quad (237)$$

$$\mathfrak{Q}_{n, m}^{(2)} := A_m \sum_{a=0}^n \binom{n}{a} \min \{ \mathfrak{Z}_{a, m+1} \mathfrak{X}_{n-a, m+1}, \mathfrak{X}_{a, m+1} \mathfrak{Z}_{n-a, m+1} \} + \mathfrak{L}_{n, m}^{(2)}, \quad (238)$$

$$\mathfrak{Q}_{n, m}^{(\infty)} := A_m \sum_{a=0}^n \binom{n}{a} \mathfrak{Z}_{a, m+1} \mathfrak{Z}_{n-a, m+1} + \mathfrak{L}_{n, m}^{(\infty)}, \quad (239)$$

$$\mathfrak{Q}_{n, m}^{(\text{mod})} := A_m \sum_{a=0}^n \binom{n}{a} (\mathfrak{X}_{a+1, m+1} \mathfrak{Z}_{n-a, m+1} + \mathfrak{Z}_{a, m+1} \mathfrak{X}_{n-a+1, m+1}) + \mathfrak{L}_{n+1, m}^{(2)}. \quad (240)$$

These quantities carry the arguments $[u_0, \nu, f; T]$ through $\mathfrak{X}$ and $\mathfrak{Z}$.

8.3 Mixed velocity and pressure rows

The canonical majorants now turn each finite rectangle into bounds for the actual differentiated velocity and pressure rows, including their traces and time moduli.

Theorem 8.10 (Quantitative mixed-jet rows and traces). *Let $N, s \in \mathbb{N}_0$, let $0 \leq n \leq N$, and let $\alpha \in \mathbb{N}_0^3$. Put*

$$J_{n,\alpha} := \partial_x^\alpha V_n.$$

Then

$$\|J_{n,\alpha}\|_{L^2(I_T; H^{s+1})}^2 + \|J_{n+1,\alpha}\|_{L^2(I_T; H^{s-1})}^2 \leq \mathfrak{P}_{N,s+|\alpha|}, \quad (241)$$

$$\sup_{0 \leq t \leq T} \|J_{n,\alpha}(t)\|_{H^s} \leq \mathfrak{T}_{N,s+|\alpha|}. \quad (242)$$

There is a unique trace $J_{n,\alpha}^T \in H^s$ at $t = T$, and for $0 \leq r \leq t \leq T$,

$$\|J_{n,\alpha}(t) - J_{n,\alpha}(r)\|_{H^{s-1}} \leq (t-r)^{1/2} \mathfrak{P}_{N,s+|\alpha|}^{1/2}, \quad (243)$$

$$\|J_{n,\alpha}(t) - J_{n,\alpha}(r)\|_{H^s} \leq (t-r)^{1/2} \mathfrak{P}_{N,s+|\alpha|+1}^{1/2}, \quad (244)$$

$$\|J_{n,\alpha}^T\|_{H^s} \leq \mathfrak{T}_{N,s+|\alpha|}. \quad (245)$$

If $0 \leq \theta \leq 1$, then

$$\|J_{n,\alpha}(t) - J_{n,\alpha}(r)\|_{H^{s-\theta}} \leq 2^{1-\theta} \mathfrak{T}_{N,s+|\alpha|}^{1-\theta} \mathfrak{P}_{N,s+|\alpha|}^{\theta/2} (t-r)^{\theta/2}. \quad (246)$$

For $1 \leq a \leq \infty$, $2 \leq b \leq \infty$, $k, q \in \mathbb{N}_0$, and $q > k + \frac{3}{2} - \frac{3}{b}$,

$$\|\nabla^k J_{n,\alpha}\|_{L^a(I_T; L^b)} \leq S_{q,k,b} T^{1/a} \mathfrak{T}_{N,q+|\alpha|}, \quad (247)$$

$$\|\nabla^k J_{n,\alpha}\|_{L^2(I_T; L^b)} \leq S_{q,k,b} \mathfrak{P}_{N,q+|\alpha|}^{1/2}, \quad (248)$$

where $T^{1/\infty} = 1$.

Proof. Take the rectangle with middle order $M = s + |\alpha|$. The Fourier multiplier inequality

$$\|\partial_x^\alpha f\|_{H^r} \leq \|f\|_{H^{r+|\alpha|}}$$

and (185) give (241). The adjacent-row trace estimate gives

$$\begin{aligned} \sup_{0 \leq t \leq T} \|J_{n,\alpha}(t)\|_{H^s}^2 &\leq T^{-1} \|J_{n,\alpha}\|_{L^2(I_T; H^s)}^2 \\ &\quad + 2 \|J_{n,\alpha}\|_{L^2(I_T; H^{s+1})} \|J_{n+1,\alpha}\|_{L^2(I_T; H^{s-1})} \\ &\leq (T^{-1} + 1) \mathfrak{P}_{N,s+|\alpha|}, \end{aligned}$$

which proves (242) and (245).

The Bochner identity

$$J_{n,\alpha}(t) - J_{n,\alpha}(r) = \int_r^t J_{n+1,\alpha}(\tau) d\tau$$

in H^{s-1} , followed by Cauchy–Schwarz, proves (243). Applying this argument to the rectangle with middle order $s + |\alpha| + 1$ proves (244). Interpolation between H^{s-1} and H^s , together with $\|J(t) - J(r)\|_{H^s} \leq 2 \mathfrak{T}_{N,s+|\alpha|}$, proves (246). Sobolev embedding and (242) prove (247); Sobolev embedding and (241) prove (248). $\square$

Theorem 8.11 (Quantitative pressure rows). *Let $N, s \in \mathbb{N}_0$, $0 \leq n \leq N$, and $\alpha \in \mathbb{N}_0^3$. Set*

$$\Pi_{n,\alpha} := \partial_x^\alpha Q_n, \quad r := s + |\alpha|.$$

Then

$$\|\Pi_{n,\alpha}\|_{L^1(I_T;H^s)} + \|\nabla \Pi_{n,\alpha}\|_{L^1(I_T;H^{s-1})} \leq 2A_r 2^n \mathfrak{P}_{N,r} + \mathfrak{L}_{n,r}^{(1)}, \quad (249)$$

$$\|\Pi_{n,\alpha}\|_{L^2(I_T;H^s)} \leq A_r 2^n \mathfrak{T}_{N,r+1} \mathfrak{P}_{N,r}^{1/2} + \mathfrak{L}_{n,r}^{(2)}, \quad (250)$$

$$\sup_{0 \leq t \leq T} \|\Pi_{n,\alpha}(t)\|_{H^s} \leq A_r 2^n \mathfrak{T}_{N,r+1}^2 + \mathfrak{L}_{n,r}^{(\infty)}. \quad (251)$$

There is a unique endpoint pressure row $\Pi_{n,\alpha}^T \in H^s$, and

$$\|\Pi_{n,\alpha}^T\|_{H^s} \leq A_r 2^n \mathfrak{T}_{N,r+1}^2 + \mathfrak{L}_{n,r}^{(\infty)}, \quad (252)$$

$$\|\Pi_{n,\alpha}(t) - \Pi_{n,\alpha}(r_0)\|_{H^s} \leq 2^{n+1} A_r \mathfrak{T}_{N,r+1} \mathfrak{P}_{N,r+2}^{1/2} |t - r_0|^{1/2} + \mathfrak{L}_{n+1,r}^{(2)} |t - r_0|^{1/2} \quad (253)$$

for $0 \leq r_0 \leq t \leq T$. If $1 \leq a \leq \infty$, $2 \leq b \leq \infty$, $k, q \in \mathbb{N}_0$, and $q > k + \frac{3}{2} - \frac{3}{b}$, then

$$\|\nabla^k \Pi_{n,\alpha}\|_{L^a(I_T;L^b)} \leq S_{q,k,b} T^{1/a} A_{q+|\alpha|} 2^n \mathfrak{T}_{N,q+|\alpha|+1}^2 + S_{q,k,b} T^{1/a} \mathfrak{L}_{n,q+|\alpha|}^{(\infty)}. \quad (254)$$

Proof. The normalized pressure recurrence is

$$Q_n = R_i R_j \sum_{a=0}^n \binom{n}{a} V_{a,i} V_{n-a,j} - \Phi_n. \quad (255)$$

Equation (188), Cauchy–Schwarz in time, and $\sum_{a=0}^n \binom{n}{a} = 2^n$ give

$$\|Q_n\|_{L^1(I_T;H^r)} \leq A_r 2^n \mathfrak{P}_{N,r} + \|\Phi_n\|_{L^1(I_T;H^r)}. \quad (256)$$

The gradient estimate in (249) follows from $\|\nabla h\|_{H^{r-1}} \leq \|h\|_{H^r}$, with the two prescribed force-potential norms retained in $\mathfrak{L}_{n,r}^{(1)}$. Using one factor in $L_t^\infty H_x^{r+1}$ and the other in $L_t^2 H_x^{r+1}$ proves (250). Using both factors in $L_t^\infty H_x^{r+1}$ proves (251) and (252).

Subtract (255) at times t and r_0 , split each product difference into two terms, apply (244) in H^{r+1} , and use (242). The two product differences and the binomial sum, together with

$$\Phi_n(t) - \Phi_n(r_0) = \int_{r_0}^t \Phi_{n+1}(\tau) d\tau,$$

give (253). Equation (254) follows from (251) and Sobolev embedding, including the force-potential coordinate. $\square$

Proposition 8.12 (Matrix form of the closed rectangle estimates). *For nonnegative matrices of identical dimensions, write $A \leq_{\text{ent}} B$ when every entry of A is at most the corresponding entry of B . When $T = T_* < \infty$, interpret the closed rectangle matrix and the three scalar rectangle coordinates by*

$$\begin{aligned} \mathbf{C}_{N,M}(T_*) &:= \lim_{S \uparrow T_*} \mathbf{C}_{N,M}(S), \\ \mathfrak{R}_{N,M}(u; T_*) &:= \lim_{S \uparrow T_*} \mathfrak{R}_{N,M}(u; S), \\ \mathfrak{Q}_{N,M}(p; T_*) &:= \lim_{S \uparrow T_*} \mathfrak{Q}_{N,M}(p; S), \\ \mathfrak{F}_{N,M}(f; T_*) &:= \lim_{S \uparrow T_*} \mathfrak{F}_{N,M}(f; S). \end{aligned}$$

All four limits exist in the extended nonnegative reals by monotonicity. Theorem 5.7 and Proposition 5.11 make the scalar limits for velocity and pressure finite; the prescribed force class makes the force limit finite, and the estimates below make every matrix entry finite. Then

$$\mathbf{C}_{N,M}(T) \preceq_{\text{ent}} \left(\begin{array}{cc} \mathfrak{X}_{n,M+1} & \mathfrak{X}_{n+1,M-1} \\ \mathfrak{Q}_{n,M}^{(1)} & \mathfrak{G}_{n,M-1}^{(1)} + \mathfrak{L}_{n,M}^{(1)} \end{array} \right)_{n=0}^N. \quad (257)$$

Consequently,

$$\mathfrak{R}_{N,M}(u; T) \leq \sum_{n=0}^N (\mathfrak{X}_{n,M+1}^2 + \mathfrak{X}_{n+1,M-1}^2), \quad (258)$$

$$\mathfrak{Q}_{N,M}(p; T) \leq \sum_{n=0}^N \mathfrak{Q}_{n,M}^{(1)}. \quad (259)$$

$$\mathfrak{F}_{N,M}(f; T) \leq \sum_{n=0}^N (\mathfrak{G}_{n,M-1}^{(1)} + \mathfrak{L}_{n,M}^{(1)}). \quad (260)$$

Proof. The first two columns of (257) are (233). The pressure recurrence (255), the product estimate (189), and Cauchy–Schwarz in time give the third column through (237). The fourth column is the prescribed-force budget itself, bounded by (199)–(200). Take the Frobenius norm of the first two columns and sum the last two columns, using (163). $\square$

8.4 Frequency and critical-scale consequences

Proposition 8.13 (Velocity and pressure frequency tails). *Let $\mathbf{P}_{\geq K}$ be the Fourier multiplier of $\mathbf{1}_{\{|\xi| \geq K\}}$, where $K \geq 1$. Let $r_1 \in \mathbb{N}$, $s \in \mathbb{N}_0$, $0 \leq n \leq N$, and $\alpha \in \mathbb{N}_0^3$. Then*

$$\|\mathbf{P}_{\geq K} J_{n,\alpha}\|_{L^2(I_T; H^s)} \leq K^{-r_1} \mathfrak{P}_{N,s+r_1+|\alpha|}^{1/2}, \quad (261)$$

$$\sup_{0 \leq t \leq T} \|\mathbf{P}_{\geq K} J_{n,\alpha}(t)\|_{H^s} \leq K^{-r_1} \mathfrak{T}_{N,s+r_1+|\alpha|}, \quad (262)$$

$$\|\mathbf{P}_{\geq K} \Pi_{n,\alpha}\|_{L^1(I_T; H^s)} \leq K^{-r_1} A_{s+r_1+|\alpha|} 2^n \mathfrak{P}_{N,s+r_1+|\alpha|} + K^{-r_1} \mathfrak{L}_{n,s+r_1+|\alpha|}^{(1)}, \quad (263)$$

$$\sup_{0 \leq t \leq T} \|\mathbf{P}_{\geq K} \Pi_{n,\alpha}(t)\|_{H^s} \leq K^{-r_1} A_{s+r_1+|\alpha|} 2^n \mathfrak{T}_{N,s+r_1+|\alpha|+1}^2 + K^{-r_1} \mathfrak{L}_{n,s+r_1+|\alpha|}^{(\infty)}. \quad (264)$$

The pointwise estimates (262) and (264) hold at $t = T$.

Proof. For every $f \in H^{s+r_1}$,

$$\|\mathbf{P}_{\geq K} f\|_{H^s} \leq K^{-r_1} \|f\|_{H^{s+r_1}}.$$

Apply this inequality to (241) and (242) at spatial order $s + r_1$ to obtain the two velocity bounds. Since Fourier multipliers commute,

$$\|\mathbf{P}_{\geq K} \Pi_{n,\alpha}\|_{H^s} \leq K^{-r_1} \|Q_n\|_{H^{s+r_1+|\alpha|}}.$$

Integration in time followed by the forced pressure bound (256) at order $s + r_1 + |\alpha|$ gives (263). Applying the tail inequality directly to $\Pi_{n,\alpha}$, followed by (251) with s replaced by $s + r_1$, gives (264). $\square$

Proposition 8.14 (Fixed weighted dyadic shell sums). *Let $(\Delta_j)_{j \geq -1}$ be a fixed inhomogeneous Littlewood–Paley decomposition on $\mathbb{R}^3$. Let $\sigma \in \mathbb{R}$, $k \in \mathbb{N}_0$, and choose $M \in \mathbb{N}_0$ such that*

$$M > \sigma + k + \frac{3}{2}. \quad (265)$$

Then

$$\int_0^T \sum_{j \geq -1} 2^{\sigma j} \left\| \Delta_j \nabla^k u(t) \right\|_{L^\infty} dt \leq C_{\Delta, \sigma, k, M} T^{1/2} \mathfrak{X}_{0, M}. \quad (266)$$

In particular, with $\omega = \nabla \times u$,

$$\sum_{j \geq -1} \int_0^T \left\| \Delta_j \omega(t) \right\|_{L^\infty} dt \leq C_\Delta T^{1/2} \mathfrak{X}_{0, 3} < \infty. \quad (267)$$

Each estimate uses one fixed finite Sobolev coordinate; no summation over Sobolev orders is asserted.

Proof. Bernstein's inequality on a shell gives, for $j \geq 0$,

$$2^{\sigma j} \left\| \Delta_j \nabla^k f \right\|_{L^\infty} \leq C_\Delta 2^{(\sigma + k + 3/2)j} \left\| \Delta_j f \right\|_2.$$

Put $d := M - \sigma - k - 3/2 > 0$. Cauchy–Schwarz in j yields

$$\begin{aligned} \sum_{j \geq 0} 2^{\sigma j} \left\| \Delta_j \nabla^k f \right\|_{L^\infty} &\leq C_\Delta \left(\sum_{j \geq 0} 2^{-2dj} \right)^{1/2} \left(\sum_{j \geq 0} 2^{2Mj} \left\| \Delta_j f \right\|_2^2 \right)^{1/2} \\ &\leq C_{\Delta, \sigma, k, M} \|f\|_{H^M}. \end{aligned}$$

The single low block $j = -1$ obeys this final bound. The last inequality follows from Plancherel's theorem and the finite overlap of the dyadic multipliers. Integrating in time, applying Cauchy–Schwarz, and using (233) prove (266). Since curl commutes with every Δ_j , the choice $(\sigma, k, M) = (0, 1, 3)$ gives (267). $\square$

Proposition 8.15 (Critical-height bounds). *Define*

$$E_\sigma(t) := \frac{1}{2} \left\| \Lambda^\sigma u(t) \right\|_2^2, \quad H(t) := E_{1/2}(t),$$

and

$$E_{j+1/2, \geq K}(t) := \frac{1}{2} \int_{|\xi| \geq K} |\xi|^{2j+1} |\widehat{u}(t, \xi)|^2 d\xi.$$

For $j \in \mathbb{N}_0$,

$$\sup_{0 \leq t \leq T} E_{j+1/2}(t) \leq \frac{1}{2} \mathfrak{X}_{0, j+1}^2, \quad (268)$$

$$\int_0^T E_{j+1/2}(t) dt \leq \frac{1}{2} \mathfrak{P}_{0, j}. \quad (269)$$

If $m > j + \frac{1}{2}$ is an integer and $K \geq 1$, then

$$\sup_{0 \leq t \leq T} E_{j+1/2, \geq K}(t) \leq \frac{1}{2} K^{2j+1-2m} \mathfrak{X}_{0, m}^2, \quad (270)$$

$$\int_0^T E_{j+1/2, \geq K}(t) dt \leq \frac{1}{2} K^{2j+1-2m} \mathfrak{P}_{0, m-1}. \quad (271)$$

For $k \in \mathbb{N}$,

$$\|H^{(k)}\|_{L^1(0,T)} \leq 2^{k-1} \mathfrak{P}_{k,0}. \quad (272)$$

In particular,

$$\|H'\|_{L^1(0,T)} \leq \frac{1}{2} \mathfrak{P}_{0,2}, \quad (273)$$

$$\int_0^T \left\| \Lambda^{3/2} u(t) \right\|_2^2 dt \leq \mathfrak{P}_{0,1}, \quad (274)$$

$$\|\mathcal{P}_{1/2}\|_{L^1(0,T)} \leq \frac{1}{2} \mathfrak{P}_{0,2} + \nu \mathfrak{P}_{0,1} + \mathfrak{P}_{0,0}^{1/2} \mathfrak{G}_{0,1}^{(2)}, \quad (275)$$

$$|H(T) - H(t)| \leq \mathfrak{T}_{0,1} \mathfrak{P}_{0,2}^{1/2} (T-t)^{1/2}, \quad 0 \leq t \leq T, \quad (276)$$

where

$$\mathcal{P}_{1/2} = H' + \nu \left\| \Lambda^{3/2} u \right\|_2^2 - \mathcal{W}_{1/2}.$$

Proof. Equations (268) and (269) follow from $\dot{H}^{j+1/2} \hookleftarrow H^{j+1}$, (242), and (241). On $|\xi| \geq K$,

$$|\xi|^{2j+1} \leq K^{2j+1-2m} |\xi|^{2m},$$

which proves (270) and (271).

The identity

$$H^{(k)}(t) = \frac{1}{2} \sum_{a=0}^k \binom{k}{a} \operatorname{Re} \left\langle \Lambda^{1/2} V_a(t), \Lambda^{1/2} V_{k-a}(t) \right\rangle$$

and Cauchy–Schwarz in time give (272). At $k = 1$, the two equal terms give

$$\|H'\|_{L^1(0,T)} \leq \|u_t\|_{L^2(I_T; \dot{H}^{1/2})} \|u\|_{L^2(I_T; \dot{H}^{1/2})} \leq \frac{1}{2} \mathfrak{P}_{0,2}.$$

Equation (274) follows from the upper velocity row of the $(0, 1)$ rectangle. Cauchy–Schwarz gives

$$\|\mathcal{W}_{1/2}\|_{L^1(0,T)} \leq \|u\|_{L^2(I_T; H^1)} \|\mathbb{P}f\|_{L^2(I_T; H^1)} \leq \mathfrak{P}_{0,0}^{1/2} \mathfrak{G}_{0,1}^{(2)}.$$

The forced height identity defining $\mathcal{P}_{1/2}$ then gives (275). Finally,

$$\|u(T) - u(t)\|_{\dot{H}^{1/2}} \leq \|u(T) - u(t)\|_{H^1} \leq (T-t)^{1/2} \|u_t\|_{L^2(t,T; H^1)} \leq (T-t)^{1/2} \mathfrak{P}_{0,2}^{1/2},$$

and

$$|H(T) - H(t)| \leq \frac{1}{2} (\|u(T)\|_{\dot{H}^{1/2}} + \|u(t)\|_{\dot{H}^{1/2}}) \|u(T) - u(t)\|_{\dot{H}^{1/2}}.$$

Equation (242) bounds each retained velocity factor, so these two estimates prove (276). $\square$

Taking $j = 0$ and any integer $m \geq 1$ in (270) gives the uniform tail limit

$$\lim_{K \rightarrow \infty} \sup_{0 \leq t \leq T} E_{1/2, \geq K}(t) = 0. \quad (277)$$

8.5 Continuation and quantitative restart

Proposition 8.16 (Continuation norms). *The finite-horizon solution satisfies*

$$\|u\|_{L^2(I_T; L^\infty)} \leq S_{2,0,\infty} \mathfrak{P}_{0,1}^{1/2}, \quad (278)$$

$$\int_0^T \|\nabla u(t)\|_{L^\infty} dt \leq S_{3,1,\infty} T^{1/2} \mathfrak{P}_{0,2}^{1/2}, \quad (279)$$

$$\int_0^T \|\nabla \times u(t)\|_{L^\infty} dt \leq 2S_{3,1,\infty} T^{1/2} \mathfrak{P}_{0,2}^{1/2}. \quad (280)$$

For every integer $m \geq 3$, let K_m be the explicit constant in (11). Then

$$|\operatorname{Re} \langle A^m \mathbb{P} \nabla \cdot (u \otimes u), A^m u \rangle_{L^2}| \leq K_m \|\nabla u\|_{L^\infty} \|u\|_{H^m}^2. \quad (281)$$

$$\sup_{0 \leq t \leq T} \|u(t)\|_{H^m}^2 \leq \left(\|u_0\|_{H^m} + \mathfrak{G}_{0,m}^{(1)} \right)^2 \exp \left(2K_m S_{3,1,\infty} T^{1/2} \mathfrak{P}_{0,2}^{1/2} \right). \quad (282)$$

Proof. Sobolev embedding and the upper rows of the $(0, 1)$ and $(0, 2)$ rectangles give (278) and (279); the pointwise bound $\|\nabla \times u\|_\infty \leq 2 \|\nabla u\|_\infty$ gives (280). The forced H^m energy identity and (281) give

$$\frac{1}{2} \frac{d}{dt} \|u\|_{H^m}^2 + \nu \|\nabla u\|_{H^m}^2 \leq K_m \|\nabla u\|_{L^\infty} \|u\|_{H^m}^2 + \|\mathbb{P}f\|_{H^m} \|u\|_{H^m}.$$

Regularizing the norm by $(\|u\|_{H^m}^2 + \varepsilon)^{1/2}$, applying Gronwall, and then letting $\varepsilon \downarrow 0$ gives

$$\|u(t)\|_{H^m} \leq \left(\|u_0\|_{H^m} + \int_0^t \|\mathbb{P}f(s)\|_{H^m} ds \right) \exp \left(K_m \int_0^t \|\nabla u(s)\|_\infty ds \right).$$

Gronwall's inequality and (279) prove (282). $\square$

Theorem 8.17 (Explicit endpoint-restart bound). *Assume $T = T_* < \infty$, and set $u_T := u_*$. Define the maximal restart lifespan inside the shifted pressure-normalized H^3 mild class by*

$$\begin{aligned} \delta_T^{\max} &:= \tau_{\nu,f,3}^{\max}(T, u_T) \\ &= \sup \{ \delta > 0 : \mathcal{M}_{\nu,f}^3(T, T + \delta; u_T) \neq \emptyset \} \in (0, \infty]. \end{aligned} \quad (283)$$

Define the same-force local coordinate

$$\Gamma_T := \|\mathbb{P}f\|_{L^2(T, T+1; H^2)},$$

and define

$$\delta_{\text{qc}}[u_0, \nu, f; T] := \min \left\{ 1, c_\nu (1 + \mathfrak{Z}_{0,3}[u_0, \nu, f; T] + \Gamma_T)^{-2} \right\}. \quad (284)$$

The class $\mathcal{M}_{\nu,f}^3(T, T + \delta_{\text{qc}}; u_T)$ contains exactly one pair (w, q) , driven by the same absolute-time force $f(t)$. It is the original history's continuation constructed in Theorem 6.8, and

$$\delta_T^{\max} \geq \delta_{\text{qc}}[u_0, \nu, f; T] > 0. \quad (285)$$

At the base restart order,

$$\begin{aligned} &\|w\|_{C([T, T+\delta_{\text{qc}}]; H^3)} + \|w\|_{L^2(T, T+\delta_{\text{qc}}; H^4)} + \|w_t\|_{L^2(T, T+\delta_{\text{qc}}; H^2)} \\ &\leq 2L_\nu (\mathfrak{Z}_{0,3}[u_0, \nu, f; T] + \Gamma_T). \end{aligned} \quad (286)$$

Set

$$\mathfrak{A}_T := 2S_{3,1,\infty} L_\nu \delta_{\text{qc}} (\mathfrak{Z}_{0,3}[u_0, \nu, f; T] + \Gamma_T). \quad (287)$$

For every integer $m \geq 3$,

$$\sup_{T \leq t \leq T + \delta_{\text{qc}}} \|w(t)\|_{H^m}^2 \leq \left(\mathfrak{Z}_{0,m}[u_0, \nu, f; T] + \|\mathbb{P}f\|_{L^1(T, T + \delta_{\text{qc}}; H^m)} \right)^2 \exp(2K_m \mathfrak{A}_T). \quad (288)$$

Proof. Equations (121) and (234) give

$$\|u_T\|_{H^3} \leq \mathfrak{Z}_{0,3}.$$

Consequently,

$$\delta_{\text{qc}}[u_0, \nu, f; T] \leq \min\{1, c_\nu(1 + \|u_T\|_{H^3} + \Gamma_T)^{-2}\}.$$

Theorem 3.11 supplies the unique pair $(w, q) \in \mathcal{M}_{\nu, f}^3(T, T + \delta_{\text{qc}}; u_T)$, and the fixed-point ball in its proof gives (286). Hence

$$\int_T^{T + \delta_{\text{qc}}} \|\nabla w(t)\|_\infty dt \leq S_{3,1,\infty} \delta_{\text{qc}} \|w\|_{C([T, T + \delta_{\text{qc}}]; H^3)} \leq \mathfrak{A}_T.$$

The forced H^m energy estimate in Proposition 8.16, now on the restart interval, with $\|u_T\|_{H^m} \leq \mathfrak{Z}_{0,m}$ and the same source $f(t)$, proves (288). Theorem 6.8 identifies this unique local pair with the continuation of the preterminal history, and its membership on the displayed interval proves (285). $\square$

9 Three-dimensional applications

9.1 Explicit closure for small critical forced data

The construction below gives an explicit three-dimensional forced $\dot{H}^{1/2}$ realization of the critical-data mechanism introduced by Fujita and Kato [7]. The projected-force stock is retained in the same critical action identity, and every constant used here is derived in the displayed estimates.

Use the explicit homogeneous Sobolev constants

$$S_3 := S(3), \quad C_{\text{src}} := S_3^3 = \frac{1}{\sqrt{2}\pi}.$$

These constants hold for vector fields. Indeed, apply the scalar inequality to the Euclidean length $|F|$: one has $|\nabla|F|| \leq |\nabla F|$, while the difference-quotient representation of $\dot{H}^{1/2}$ and $||F(x)| - |F(y)|| \leq |F(x) - F(y)|$ give $||F||_{\dot{H}^{1/2}} \leq \|F\|_{\dot{H}^{1/2}}$.

Lemma 9.1 (Critical action identity). *Let (u, p) be a smooth pressure-normalized solution on a time interval $I \subset [0, T_*)$ of the forced equation with prescribed force f , and set*

$$\mathcal{B}(u) := \mathbb{P}\nabla \cdot (u \otimes u), \quad g := \mathbb{P}f.$$

Then, pointwise on I ,

$$\nu \frac{d}{dt} \|u\|_{\dot{H}^{1/2}}^2 + \nu^2 \|u\|_{\dot{H}^{3/2}}^2 + \|u_t\|_{\dot{H}^{-1/2}}^2 = \|\mathcal{B}(u) - g\|_{\dot{H}^{-1/2}}^2. \quad (289)$$

Proof. The projected equation is

$$u_t = -\nu \Lambda^2 u - \mathcal{B}(u) + g.$$

Put

$$h := \Lambda^{-1/2}(\mathcal{B}(u) - g), \quad w := \Lambda^{-1/2}u_t = -\nu \Lambda^{3/2}u - h.$$

Pairing the projected equation with Λu gives

$$\frac{d}{dt} \|u\|_{\dot{H}^{1/2}}^2 = -2\nu \|u\|_{\dot{H}^{3/2}}^2 - 2 \operatorname{Re} \langle h, \Lambda^{3/2}u \rangle_2.$$

Multiply this equality by ν and expand $\|w\|_2^2 = \|\nu \Lambda^{3/2}u + h\|_2^2$. The cross terms cancel and give (289). $\square$

For $m \in \mathbb{N}$, define the explicit homogeneous commutator constant

$$\mathsf{K}_m^{\text{hom}} := 3^m \sum_{\ell=1}^m \binom{m}{\ell} \mathsf{S}(\mathfrak{p}_{m,\ell}) \mathsf{S}(\mathfrak{q}_{m,\ell}), \quad (290)$$

where $(\mathfrak{p}_{1,1}, \mathfrak{q}_{1,1}) = (3, 6)$, and, for $m \geq 2$,

$$\vartheta_{m,\ell} := \frac{\ell-1}{m-1}, \quad \frac{1}{\mathfrak{p}_{m,\ell}} := \frac{1}{3} - \frac{\vartheta_{m,\ell}}{6}, \quad \frac{1}{\mathfrak{q}_{m,\ell}} := \frac{1}{6} + \frac{\vartheta_{m,\ell}}{6}.$$

The corresponding estimate is

$$|\operatorname{Re} \langle \Lambda^m \mathbb{P} \nabla \cdot (z \otimes z), \Lambda^m z \rangle_{L^2}| \leq \mathsf{K}_m^{\text{hom}} \left\| \Lambda^{3/2} z \right\|_2 \|\Lambda^m z\|_2 \|\Lambda^{m+1} z\|_2 \quad (291)$$

for every $z \in \mathcal{S}_\sigma(\mathbb{R}^3)$.

Lemma 9.2 (Explicit homogeneous critical commutator). *For every $m \in \mathbb{N}$, $\mathsf{K}_m^{\text{hom}} < \infty$, and (291) holds with the displayed constant. Moreover, (291) extends to every divergence-free $z \in H^{m+1}(\mathbb{R}^3) \cap \dot{H}^{3/2}(\mathbb{R}^3)$.*

Proof. The multinomial identity gives the exact pairing

$$\langle \Lambda^m f, \Lambda^m g \rangle_{L^2} = \sum_{|\alpha|=m} \frac{m!}{\alpha!} \langle \partial^\alpha f, \partial^\alpha g \rangle_{L^2}.$$

After differentiating $(z \cdot \nabla)z$, the undifferentiated transport term has zero pairing by incompressibility. A remaining term has the form

$$\left\langle (\partial^\beta z \cdot \nabla) \partial^{\alpha-\beta} z, \partial^\alpha z \right\rangle_{L^2}, \quad |\alpha| = m, \quad 0 < \beta \leq \alpha.$$

Put $\ell = |\beta|$. For $m \geq 2$, Hölder's inequality, Lemma 3.3, and Fourier log-convexity give

$$\begin{aligned} & \left\| \partial^\beta z \right\|_{\mathfrak{p}_{m,\ell}} \left\| \nabla \partial^{\alpha-\beta} z \right\|_{\mathfrak{q}_{m,\ell}} \\ & \leq \mathsf{S}(\mathfrak{p}_{m,\ell}) \mathsf{S}(\mathfrak{q}_{m,\ell}) \|z\|_{\dot{H}^{3/2}} \|z\|_{\dot{H}^{m+1}}. \end{aligned}$$

Indeed, the two differentiated Sobolev orders are the complementary interpolants

$$\begin{aligned}\ell + s(\mathfrak{p}_{m,\ell}) &= (1 - \vartheta_{m,\ell})\frac{3}{2} + \vartheta_{m,\ell}(m+1), \\ m - \ell + 1 + s(\mathfrak{q}_{m,\ell}) &= \vartheta_{m,\ell}\frac{3}{2} + (1 - \vartheta_{m,\ell})(m+1).\end{aligned}$$

For $m = 1$, this bound is the direct L^3 - L^6 estimate. Finally,

$$\sum_{|\alpha|=m} \frac{m!}{\alpha!} = 3^m, \quad \sum_{\substack{\beta \leq \alpha \\ |\beta|=\ell}} \binom{\alpha}{\beta} = \binom{m}{\ell}.$$

Summation gives exactly (290) and proves (291). Solenoidal Fourier truncation and convergence in $H^{m+1} \cap \dot{H}^{3/2}$ give the asserted extension. $\square$

Theorem 9.3 (Explicit small-critical forced-data global regularity). *Let $\nu > 0$, $a \in \mathcal{S}_\sigma(\mathbb{R}^3)$, and let $f \in C^\infty([0, \infty); \mathcal{S}(\mathbb{R}^3; \mathbb{R}^3))$ be the prescribed force. Write $g := \mathbb{P}f$, and assume*

$$g \in L^1(0, \infty; L^2) \cap L^2(0, \infty; \dot{H}^{-1/2}) \cap \bigcap_{j \in \mathbb{N}_0} L^2(0, \infty; \dot{H}^j). \quad (292)$$

Set

$$a_c := \left\| \Lambda^{1/2} a \right\|_2, \quad \mathcal{G}_{-1/2} := \|g\|_{L^2(0, \infty; \dot{H}^{-1/2})}.$$

If there is an $\eta > 0$ such that, with

$$\begin{aligned}A_f^2 &:= a_c^2 + \frac{1 + \eta^{-1}}{\nu} \mathcal{G}_{-1/2}^2, \quad \delta_f := \nu^2 - (1 + \eta) C_{\text{src}}^2 A_f^2, \\ A_f &< \frac{\nu}{\sqrt{1 + \eta} C_{\text{src}}} = \sqrt{\frac{2}{1 + \eta}} \pi \nu,\end{aligned} \quad (293)$$

then the forced datum (a, ν, f) generates a unique global pressure-normalized smooth solution

$$(u, p) \in C^\infty([0, \infty) \times \mathbb{R}^3; \mathbb{R}^3 \times \mathbb{R}).$$

For every finite $T > 0$, the solution is the unique member of $\mathcal{M}_{\nu, f}^3(0, T; a)$. For every $t \geq 0$,

$$\nu \left\| \Lambda^{1/2} u(t) \right\|_2^2 + \delta_f \int_0^t \left\| \Lambda^{3/2} u(s) \right\|_2^2 ds + \int_0^t \|u_t(s)\|_{\dot{H}^{-1/2}}^2 ds \leq \nu A_f^2, \quad \delta_f > 0. \quad (294)$$

The critical nonlinear source and the first temporal response satisfy

$$\int_0^\infty \|\mathbb{P} \nabla \cdot (u \otimes u)(t)\|_{\dot{H}^{-1/2}}^2 dt \leq \frac{C_{\text{src}}^2 \nu A_f^4}{\delta_f}, \quad (295)$$

$$\nu^2 \int_0^\infty \left\| \Lambda^{3/2} u(t) \right\|_2^2 dt + \int_0^\infty \|u_t(t)\|_{\dot{H}^{-1/2}}^2 dt \leq \nu A_f^2 + \frac{(1 + \eta) C_{\text{src}}^2 \nu A_f^4}{\delta_f}. \quad (296)$$

For every $m \in \mathbb{N}$, define

$$\begin{aligned}\mathcal{D}_m^f &:= \left\| \Lambda^m a \right\|_2^2 + \frac{1 + \eta^{-1}}{\nu} \left\| \Lambda^{m-1} g \right\|_{L^2(0, \infty; L^2)}^2, \\ \Gamma_m^{\text{sm}} &:= \frac{(1 + \eta) (\mathcal{K}_m^{\text{hom}})^2 A_f^2}{\delta_f}.\end{aligned} \quad (297)$$

Then

$$\begin{aligned} & \|\Lambda^m u(t)\|_2^2 + \nu \int_0^t \|\Lambda^{m+1} u(s)\|_2^2 \, ds \\ & \leq \mathcal{D}_m^f \exp(\Gamma_m^{\text{sm}}). \end{aligned} \quad (298)$$

Put

$$K_{f,\infty} := \|a\|_2 + \|g\|_{L^1(0,\infty;L^2)}.$$

In particular,

$$\sup_{t \geq 0} \|u(t)\|_{H^m}^2 \leq 2^{m-1} \left[K_{f,\infty}^2 + \mathcal{D}_m^f \exp(\Gamma_m^{\text{sm}}) \right]. \quad (299)$$

Proof. Iterate Theorem 3.11 and join overlapping intervals by its at-most-one assertion for the shifted mild class. The union is the maximal shifted pressure-normalized H^3 mild history on $[0, T_*)$, and persistence retains H^∞ on every compact subinterval. Put

$$A(t) := \left\| \Lambda^{1/2} u(t) \right\|_2, \quad B(t) := \left\| \Lambda^{3/2} u(t) \right\|_2.$$

Dual Sobolev, Hölder, and the unit Fourier-multiplier norm of $\mathbb{P}$ give the complete source-constant chain

$$\begin{aligned} \|\mathbb{P} \nabla \cdot (u \otimes u)\|_{\dot{H}^{-1/2}} & \leq S_3 \| (u \cdot \nabla) u \|_{L^{3/2}} \\ & \leq S_3 \|u\|_{L^3} \|\nabla u\|_{L^3} \\ & \leq S_3^3 \left\| \Lambda^{1/2} u \right\|_2 \left\| \Lambda^{3/2} u \right\|_2 = C_{\text{src}} A(t) B(t). \end{aligned} \quad (300)$$

Combining (300) with the exact identity (289) gives, on every classical subinterval,

$$\nu \frac{d}{dt} A(t)^2 + (\nu^2 - (1 + \eta) C_{\text{src}}^2 A(t)^2) B(t)^2 + \|u_t(t)\|_{\dot{H}^{-1/2}}^2 \leq (1 + \eta^{-1}) \|g(t)\|_{\dot{H}^{-1/2}}^2.$$

Indeed, this is $\|\mathcal{B}(u) - g\|^2 \leq (1 + \eta) \|\mathcal{B}(u)\|^2 + (1 + \eta^{-1}) \|g\|^2$. Continuity and a first-exit argument give $A(t) \leq A_f$ throughout $[0, T_*)$. Integration and the definition of δ_f prove (294). In particular,

$$\int_0^t B(s)^2 \, ds \leq \frac{\nu A_f^2}{\delta_f}.$$

For every $t < T_*$, this source estimate and the integrated identity (289) prove (295) and (296) with ∞ replaced by t . After the continuation argument below proves $T_* = \infty$, monotone convergence gives the stated forms.

For the first temporal response, integrate (289), use $\|\mathcal{B}(u) - g\|^2 \leq (1 + \eta) \|\mathcal{B}(u)\|^2 + (1 + \eta^{-1}) \|g\|^2$, and substitute the definition of A_f .

At order m , the forced projected equation, Lemma 9.2, and Young's inequality yield

$$\frac{d}{dt} \|\Lambda^m u\|_2^2 + \nu \|\Lambda^{m+1} u\|_2^2 \leq \frac{(1 + \eta)(K_m^{\text{hom}})^2}{\nu} B(t)^2 \|\Lambda^m u\|_2^2 + \frac{1 + \eta^{-1}}{\nu} \|\Lambda^{m-1} g(t)\|_2^2.$$

Multiplication by the corresponding integrating factor and

$$\int_0^t B(s)^2 \, ds \leq \frac{\nu A_f^2}{\delta_f}$$

prove (298). The Fourier inequality

$$(1 + |\xi|^2)^m \leq 2^{m-1}(1 + |\xi|^{2m})$$

and the forced energy estimate $\sup_{t \geq 0} \|u(t)\|_2 \leq K_{f,\infty}$ prove (299).

Let M_3 be the square root of the right-hand side of (299) with $m = 3$, and set

$$\delta_{\text{crit}} := \min \left\{ 1, c_\nu \left(1 + M_3 + \|g\|_{L^2(0,\infty;H^2)} \right)^{-2} \right\} > 0.$$

Every datum $u(t_0)$, $t_0 < T_*$, has H^3 -norm at most M_3 . Theorem 3.11 restarts the solution for at least δ_{crit} units of time. If $T_* < \infty$, choose $t_0 < T_*$ with $T_* - t_0 < \delta_{\text{crit}}/2$. Both restrictions belong to $\mathcal{M}_{\nu,f}^3(t_0, b; u(t_0))$ for every $b < T_*$, so uniqueness on each such closed subinterval identifies them throughout $[t_0, T_*)$. The restarted member then extends the original history beyond T_* , contradicting maximality. Hence $T_* = \infty$, and persistence gives the asserted smoothness. This local uniqueness on each finite interval proves the stated global uniqueness class. $\square$

For $m \geq 1$, define

$$U_m^{\text{sm}} := 2^{(m-1)/2} \left[K_{f,\infty}^2 + \mathcal{D}_m^f \exp(\Gamma_m^{\text{sm}}) \right]^{1/2}, \quad U_0^{\text{sm}} := K_{f,\infty}, \quad U_{-1}^{\text{sm}} := U_0^{\text{sm}}.$$

For $T < \infty$, $n \in \mathbb{N}_0$, and $m \geq -1$, define

$$\mathbf{G}_{n,m}(T) := \sup_{0 \leq t \leq T} \|F_n(t)\|_{H^m}$$

and the finite-order recursion

$$\mathbf{Y}_{0,m}^{\text{sm}}(T) := U_m^{\text{sm}}, \tag{301}$$

$$\mathbf{Y}_{n+1,m}^{\text{sm}}(T) := \nu \mathbf{Y}_{n,m+2}^{\text{sm}}(T) + A_{m+1} \sum_{a'=0}^n \binom{n}{a'} \mathbf{Y}_{a',m+2}^{\text{sm}}(T) \mathbf{Y}_{n-a',m+2}^{\text{sm}}(T) + \mathbf{G}_{n,m}(T). \tag{302}$$

Proposition 9.4 (Explicit small-forced-data pressure-complete rectangles). *Under the hypotheses of Theorem 9.3, for every $T < \infty$, $n \in \mathbb{N}_0$, and $m \geq -1$,*

$$\sup_{0 \leq t \leq T} \|V_n(t)\|_{H^m} \leq \mathbf{Y}_{n,m}^{\text{sm}}(T). \tag{303}$$

For every $T < \infty$ and $N, M \in \mathbb{N}_0$,

$$\mathfrak{R}_{N,M}(u; T) \leq T \sum_{n=0}^N [(\mathbf{Y}_{n,M+1}^{\text{sm}}(T))^2 + (\mathbf{Y}_{n+1,M-1}^{\text{sm}}(T))^2], \tag{304}$$

$$\mathfrak{Q}_{N,M}(p; T) \leq T A_M \sum_{n=0}^N \sum_{a'=0}^n \binom{n}{a'} \mathbf{Y}_{a',M+1}^{\text{sm}}(T) \mathbf{Y}_{n-a',M+1}^{\text{sm}}(T) + \sum_{n=0}^N \mathfrak{F}_{n,0,M}^{\text{p}}(0, T). \tag{305}$$

In the entrywise order from Proposition 8.12, these bounds have the matrix form below, where

$$\begin{aligned} \mathbf{C}_{n,M}^{(1)}(T) &:= T^{1/2} \mathbf{Y}_{n,M+1}^{\text{sm}}(T), \\ \mathbf{C}_{n,M}^{(2)}(T) &:= T^{1/2} \mathbf{Y}_{n+1,M-1}^{\text{sm}}(T), \\ \mathbf{C}_{n,M}^{(3)}(T) &:= T A_M \sum_{a'=0}^n \binom{n}{a'} \mathbf{Y}_{a',M+1}^{\text{sm}}(T) \mathbf{Y}_{n-a',M+1}^{\text{sm}}(T) + \mathfrak{F}_{n,0,M}^{\text{p}}(0, T), \\ \mathbf{C}_{n,M}^{(4)}(T) &:= T \mathbf{G}_{n,M-1}(T) + \mathfrak{F}_{n,0,M}^{\text{p}}(0, T). \end{aligned}$$

Thus

$$\mathbf{C}_{N,M}(T) \preceq_{\text{ent}} \left(\mathbf{C}_{n,M}^{(1)}(T), \mathbf{C}_{n,M}^{(2)}(T), \mathbf{C}_{n,M}^{(3)}(T), \mathbf{C}_{n,M}^{(4)}(T) \right)_{n=0}^N. \quad (306)$$

Every right-hand side is an explicit finite function of ν, T, N, M , the datum norms

$$\|a\|_2, \quad \left\| \Lambda^{1/2} a \right\|_2, \quad \{ \|\Lambda^r a\|_2 : 1 \leq r \leq M + 2N + 1 \}. \quad (307)$$

Together with these datum norms, the bound uses the force norms in (292) and (297), and the finite family

$$\left\{ \mathbf{G}_{n,m}(T), \mathfrak{F}_{n,0,M}^{\text{p}}(0, T) : 0 \leq n \leq N, -1 \leq m \leq M + 2N + 1 \right\}.$$

Indeed, induction in n shows that $\mathbf{Y}_{n,m}^{\text{sm}}(T)$ uses only the entries U_j^{sm} with $-1 \leq j \leq m + 2n$; both velocity columns of (306) consequently stop at $j = M + 2N + 1$, and its pressure column stops no later. The fourth column is the prescribed force coordinate of the same mixed rectangle; it is bounded directly by its displayed force norms.

Proof. Equation (299) proves the base (303) for $m \geq 1$; the forced energy estimate proves it for $m = 0$, and $H^0 \hookrightarrow H^{-1}$ proves it for $m = -1$. Apply $\mathbb{P}$ to the temporal recurrence. This is the $\alpha = 0$ temporal block-row of (156); its factorial conjugate is (159). Equation (188) at order $m + 1$ and induction give (303) through (302), with $\|\mathbb{P}F_n\|_{H^m} \leq \|F_n\|_{H^m}$; for $m = -1$, use the order-zero product estimate and $H^0 \hookrightarrow H^{-1}$. Integration in time gives (304). The pressure recurrence and (189) at order M , followed by integration and summation in n , give (305), with the force-pressure potential retained in $\mathfrak{F}_{n,0,M}^{\text{p}}(0, T)$. Direct integration gives the fourth, prescribed-force column. These rowwise bounds give (306); taking the Frobenius norm of its first two columns and summing its pressure column recovers (304) and (305). $\square$

9.2 Low-rectangle and scale-local bounds

The following results use distinct outputs of the compatible rectangles: mixed space–time embeddings, pressure completion, and integrable Lipschitz action.

Proposition 9.5 (Three low-rectangle regularity classes). *Let $\nu > 0$, let $u_0 \in H_\sigma^\infty(\mathbb{R}^3)$, prescribe $f \in C^\infty([0, \infty); \mathcal{S}(\mathbb{R}^3; \mathbb{R}^3))$, and let u be the global solution supplied by Theorem 6.10 from (u_0, ν, f) , and let $0 < T < \infty$. Then*

$$\|u\|_{L^\infty(0,T;L^3)} \leq S_{1,0,3}\mathfrak{Z}_{0,1}[u_0, \nu, f; T], \quad (308)$$

$$\|u\|_{L^2(0,T;L^\infty)} \leq S_{2,0,\infty}\mathfrak{X}_{0,2}[u_0, \nu, f; T], \quad (309)$$

$$\|\nabla u\|_{L^2(0,T;L^3)} \leq S_{2,1,3}\mathfrak{X}_{0,2}[u_0, \nu, f; T]. \quad (310)$$

In particular, each of the three norms is finite on every finite horizon, with the stated datum, viscosity, force, and horizon dependence.

Proof. Use (352) with $(n, \alpha, a, b, q, k) = (0, 0, \infty, 3, 1, 0)$ to obtain (308). Use (353) with $(n, \alpha, b, q, k) = (0, 0, \infty, 2, 0)$ and $(0, 0, 3, 2, 1)$ to obtain (309) and (310). The strict embedding conditions are, respectively, $1 > 1/2$, $2 > 3/2$, and $2 > 3/2$. $\square$

Corollary 9.6 (Uniform extinction of parabolic scale content). *Let $\nu > 0$, let $u_0 \in H_\sigma^\infty(\mathbb{R}^3)$, prescribe $f \in C^\infty([0, \infty); \mathcal{S}(\mathbb{R}^3; \mathbb{R}^3))$, and let (u, p) be the pressure-normalized global pair supplied by Theorem 6.10 for (u_0, ν, f) . This is a post-regularity consequence for the Caffarelli–Kohn–Nirenberg*

scale-normalized quantity [4]; it is not used as an ε -regularity input or continuation criterion. For $0 < T < \infty$, define

$$\begin{aligned} U_T &:= S_{2,0,\infty} \mathfrak{Z}_{0,2}[u_0, \nu, f; T], \\ P_T &:= S_{2,0,\infty} \mathfrak{Q}_{0,2}^{(\infty)}[u_0, \nu, f; T]. \end{aligned} \quad (311)$$

Let $z_0 = (t_0, x_0)$, $\rho > 0$, and suppose that

$$Q_\rho(z_0) := (t_0 - \rho^2, t_0) \times B_\rho(x_0) \subset (0, T) \times \mathbb{R}^3.$$

Then the scale-normalized velocity–pressure content satisfies

$$\rho^{-2} \iint_{Q_\rho(z_0)} (|u|^3 + |p|^{3/2}) \, dx \, dt \leq \frac{4\pi}{3} \rho^3 (U_T^3 + P_T^{3/2}). \quad (312)$$

Consequently,

$$\lim_{\rho \downarrow 0} \sup_{\{z_0: Q_\rho(z_0) \subset (0, T) \times \mathbb{R}^3\}} \rho^{-2} \iint_{Q_\rho(z_0)} (|u|^3 + |p|^{3/2}) \, dx \, dt = 0. \quad (313)$$

Proof. Equation (347), followed by (191) with $(s, k, b) = (2, 0, \infty)$, gives

$$\|u\|_{L^\infty((0, T) \times \mathbb{R}^3)} \leq U_T.$$

Equations (356) and (191) give

$$\|p\|_{L^\infty((0, T) \times \mathbb{R}^3)} \leq P_T.$$

Since $|B_\rho| = \frac{4\pi}{3} \rho^3$ and the time length of Q_ρ is ρ^2 ,

$$\iint_{Q_\rho(z_0)} (|u|^3 + |p|^{3/2}) \, dx \, dt \leq \frac{4\pi}{3} \rho^5 (U_T^3 + P_T^{3/2}).$$

Division by ρ^2 proves (312); its right-hand side is independent of the center and tends to zero with ρ , which proves (313). $\square$

9.3 Vorticity and material flow

Theorem 9.7 (Global vorticity amplification and dissipation). *Let $\nu > 0$, let $u_0 \in H_\sigma^\infty(\mathbb{R}^3)$, prescribe $f \in C^\infty([0, \infty); \mathcal{S}(\mathbb{R}^3; \mathbb{R}^3))$, and let (u, p) be the pressure-normalized global pair supplied by Theorem 6.10 for (u_0, ν, f) . Put*

$$\omega := \nabla \times u, \quad \omega_0 := \nabla \times u_0, \quad \zeta := \nabla \times f, \quad L_T^\omega := S_{3,1,\infty} T^{1/2} \mathfrak{X}_{0,3}[u_0, \nu, f; T].$$

For every $T < \infty$ and $2 \leq r \leq \infty$, set

$$Z_{r,T} := \int_0^T \|\zeta(t)\|_{L^r} \, dt, \quad \Omega_{r,T} := e^{L_T^\omega} (\|\omega_0\|_{L^r} + Z_{r,T}).$$

Then

$$\sup_{0 \leq t \leq T} \|\omega(t)\|_{L^r} \leq \Omega_{r,T}, \quad (314)$$

$$\int_0^T \|(\omega \cdot \nabla) u(t)\|_{L^r} \, dt \leq L_T^\omega \Omega_{r,T}. \quad (315)$$

If $2 \leq r < \infty$, then

$$\int_0^T \left\| |\nabla |\omega(t)|^{r/2}| \right\|_2^2 \, dt \leq \frac{r}{4\nu(r-1)} e^{rL_T^\omega} \left[\|\omega_0\|_{L^r}^r + r Z_{r,T} \Omega_{r,T}^{r-1} \right]. \quad (316)$$

Proof. Taking the curl of the momentum equation gives the three-dimensional vorticity equation

$$\partial_t \omega + (u \cdot \nabla) \omega = (\omega \cdot \nabla) u + \nu \Delta \omega + \zeta. \quad (317)$$

Fix $2 \leq r < \infty$. Pair (317) with $|\omega|^{r-2} \omega$. To justify this power test, choose $\chi \in C_c^\infty(\mathbb{R}^3)$, with $0 \leq \chi \leq 1$, $\chi = 1$ on B_1 , and $\chi = 0$ outside B_2 , put $\chi_R(x) = \chi(x/R)$, and first test against

$$\chi_R \Phi_\varepsilon(\omega), \quad \Phi_\varepsilon(z) := (|z|^2 + \varepsilon)^{(r-2)/2} z.$$

Define the primitive

$$F_\varepsilon(z) := \frac{(|z|^2 + \varepsilon)^{r/2} - \varepsilon^{r/2}}{r}. \quad (318)$$

Then $\nabla_z F_\varepsilon = \Phi_\varepsilon$, and incompressibility gives

$$\int_{\mathbb{R}^3} \chi_R (u \cdot \nabla) \omega \cdot \Phi_\varepsilon(\omega) \, dx = - \int_{\mathbb{R}^3} F_\varepsilon(\omega) u \cdot \nabla \chi_R \, dx. \quad (319)$$

With $A_R := B_{2R} \setminus B_R$,

$$\left| \int F_\varepsilon(\omega) u \cdot \nabla \chi_R \, dx \right| \leq \frac{C_r}{R} \|u\|_\infty \left(\|\omega\|_{L^r(A_R)}^r + \varepsilon^{(r-2)/2} \|\omega\|_{L^2(A_R)}^2 \right), \quad (320)$$

$$\left| \int \nabla \chi_R \cdot \nabla \omega \Phi_\varepsilon(\omega) \, dx \right| \leq \frac{C_r}{R} \|\nabla \omega\|_{L^2(A_R)} \left(\|\omega\|_{L^{2r-2}(A_R)}^{r-1} + \varepsilon^{(r-2)/2} \|\omega\|_{L^2(A_R)} \right). \quad (321)$$

All norms on the right are finite because u, ω belong to every spatial Sobolev space. Their annular tails vanish as $R \rightarrow \infty$, so (319) converges to the transport cancellation and the diffusion cutoff term vanishes.

The remaining diffusion form is nonnegative:

$$\begin{aligned} \nabla \omega : D \Phi_\varepsilon(\omega) \nabla \omega &= (|\omega|^2 + \varepsilon)^{(r-2)/2} |\nabla \omega|^2 \\ &\quad + (r-2)(|\omega|^2 + \varepsilon)^{(r-4)/2} \sum_{j=1}^3 (\omega \cdot \partial_j \omega)^2. \end{aligned} \quad (322)$$

After the limit $R \rightarrow \infty$, dominated convergence applies to $F_\varepsilon(\omega)$, the stretching term, and the force term, while Fatou's lemma applies to (322) as $\varepsilon \downarrow 0$. This yields the unregularized chain rule and diffusion inequality used below. The stretching term satisfies

$$\left| \int_{\mathbb{R}^3} (\omega \cdot \nabla) u \cdot |\omega|^{r-2} \omega \, dx \right| \leq \|\nabla u\|_\infty \|\omega\|_r^r.$$

For the diffusion term, spatial integration by parts gives

$$\begin{aligned} - \int_{\mathbb{R}^3} \Delta \omega \cdot |\omega|^{r-2} \omega \, dx &\geq (r-1) \int_{\mathbb{R}^3} |\omega|^{r-2} |\nabla \omega|^2 \, dx \\ &= \frac{4(r-1)}{r^2} \left\| \nabla |\omega|^{r/2} \right\|_2^2. \end{aligned}$$

Hence

$$\frac{1}{r} \frac{d}{dt} \|\omega\|_r^r + \frac{4\nu(r-1)}{r^2} \left\| \nabla |\omega|^{r/2} \right\|_2^2 \leq \|\nabla u\|_\infty \|\omega\|_r^r + \|\zeta\|_r \|\omega\|_r^{r-1}. \quad (323)$$

Combining (323) with the quantitative rectangle estimate (380) gives

$$\int_0^T \|\nabla u(t)\|_\infty \, dt \leq L_T^\omega.$$

The scalar inequality

$$\frac{d}{dt} \|\omega(t)\|_r \leq \|\nabla u(t)\|_\infty \|\omega(t)\|_r + \|\zeta(t)\|_r$$

and Gronwall's inequality prove (314) for finite r . Multiplying (323) by $\exp(-r \int_0^t \|\nabla u(s)\|_\infty \, ds)$ and integrating, while using $\|\omega(t)\|_r \leq \Omega_{r,T}$, proves (316). Since $\omega_0, \omega(t), \zeta(t) \in L^2 \cap L^\infty$, with uniform Schwartz bounds for ζ on $[0, T]$, $\|h\|_r \rightarrow \|h\|_\infty$ as $r \rightarrow \infty$ for each spatial slice and $Z_{r,T} \rightarrow Z_{\infty,T}$. Consequently,

$$\sup_{0 \leq t \leq T} \|\omega(t)\|_\infty \leq \liminf_{r \rightarrow \infty} \sup_{0 \leq t \leq T} \|\omega(t)\|_r \leq \Omega_{\infty,T}.$$

This is (314) at the endpoint. Finally,

$$\|(\omega \cdot \nabla)u(t)\|_r \leq \|\nabla u(t)\|_\infty \|\omega(t)\|_r;$$

integration and (314) give (315). $\square$

Theorem 9.8 (Quantitative volume-preserving material flow). *Let $\nu > 0$, let $u_0 \in H_\sigma^\infty(\mathbb{R}^3)$, prescribe $f \in C^\infty([0, \infty); \mathcal{S}(\mathbb{R}^3; \mathbb{R}^3))$, and let (u, p) be the pressure-normalized global pair supplied by Theorem 6.10 for (u_0, ν, f) . For $0 < T < \infty$, put*

$$\begin{aligned} D_T &:= S_{2,0,\infty} T^{1/2} \mathfrak{X}_{0,2}[u_0, \nu, f; T], \\ L_T &:= S_{3,1,\infty} T^{1/2} \mathfrak{X}_{0,3}[u_0, \nu, f; T]. \end{aligned} \tag{324}$$

For every $a \in \mathbb{R}^3$, the problem

$$\partial_t X(t, a) = u(t, X(t, a)), \quad X(0, a) = a, \tag{325}$$

has a unique solution on $[0, T]$. For every $t \in [0, T]$, the map $X(t, \cdot) : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is a smooth volume-preserving diffeomorphism and

$$|X(t, a) - a| \leq D_T, \tag{326}$$

$$e^{-L_T} |a - b| \leq |X(t, a) - X(t, b)| \leq e^{L_T} |a - b|, \tag{327}$$

$$\det D_a X(t, a) = 1. \tag{328}$$

Proof. Equations (379) and (380) give

$$\begin{aligned} \int_0^T \|u(t)\|_\infty \, dt &\leq D_T, \\ \int_0^T \|\nabla u(t)\|_\infty \, dt &\leq L_T. \end{aligned}$$

Picard iteration on successive compact time intervals gives local existence and uniqueness for (325). The first integral bound prevents escape to spatial infinity and extends every trajectory to $[0, T]$; it also proves (326). For two trajectories, the second integral bound and Gronwall's inequality give the upper bound in (327). Solving this equation backward from time t produces the inverse map and gives the lower bound. Spatial differentiation of (325) and induction give smoothness of the map and its inverse. Finally, Jacobi's formula and $\nabla \cdot u = 0$ give

$$\partial_t \det D_a X = (\nabla \cdot u)(t, X) \det D_a X = 0, \quad \det D_a X(0, a) = 1,$$

which proves (328). $\square$

9.4 Relative energy and the Leray–Hopf class

The weak class is stated in the whole-space form originating with Leray and Hopf [11, 8]. The argument is the classical weak–strong uniqueness mechanism associated with Serrin [14], written here against the pressure-complete smooth history and derived directly from the weak formulation and the energy inequality.

Definition 9.9 (Three-dimensional Leray–Hopf solution). Let $\nu > 0$, $T > 0$, $v_0 \in H_\sigma^0(\mathbb{R}^3)$, and prescribe $f \in C^\infty([0, T]; \mathcal{S}(\mathbb{R}^3; \mathbb{R}^3))$. A forced Leray–Hopf solution on $[0, T]$ for f is a pair (v, q) such that

$$v \in C_w([0, T]; H_\sigma^0) \cap L^\infty(0, T; L_\sigma^2) \cap L^2(0, T; \dot{H}_\sigma^1), \quad q \in \mathcal{D}'((0, T) \times \mathbb{R}^3),$$

the equations

$$\partial_t v + (v \cdot \nabla)v + \nabla q = \nu \Delta v + f, \quad \nabla \cdot v = 0$$

hold in distributions, $v(t) \rightarrow v_0$ strongly in L^2 as $t \downarrow 0$, and

$$\frac{1}{2} \|v(t)\|_2^2 + \nu \int_0^t \|\nabla v(s)\|_2^2 \, ds \leq \frac{1}{2} \|v_0\|_2^2 + \int_0^t \langle f(s), v(s) \rangle \, ds \quad (329)$$

for every $t \in [0, T]$. Every divergence-free test field $\phi \in C_c^\infty([0, T] \times \mathbb{R}^3; \mathbb{R}^3)$ satisfies

$$\int_0^T \left[-\langle v, \partial_t \phi \rangle - \int_{\mathbb{R}^3} (v \otimes v) : \nabla \phi \, dx + \nu \langle \nabla v, \nabla \phi \rangle \right] dt = \langle v_0, \phi(0) \rangle + \int_0^T \langle f, \phi \rangle \, dt. \quad (330)$$

A global forced Leray–Hopf solution at viscosity ν for the prescribed force f is a pair on $[0, \infty)$ whose restriction to every finite $[0, T]$ is a forced Leray–Hopf solution in Definition 9.9, with identical initial data, viscosity, and prescribed force.

Lemma 9.10 (Solenoidal cutoff density and time commutation). *Let $\mathcal{A}$ be a finite subset of $\{(r, p) : r \in \mathbb{N}_0, 2 \leq p < \infty\}$. There are linear operators $\mathcal{T}_k$, independent of time, such that, for every $f \in H_\sigma^\infty(\mathbb{R}^3)$,*

$$\mathcal{T}_k f \in C_{c, \sigma}^\infty(\mathbb{R}^3)$$

and, for every $(r, p) \in \mathcal{A}$,

$$\sup_k \|\mathcal{T}_k f\|_{W^{r, p}} \leq C_{\mathcal{A}} \|f\|_{W^{r, p}}, \quad (331)$$

$$\|\mathcal{T}_k f - f\|_{W^{r, p}} \longrightarrow 0. \quad (332)$$

The convergence is uniform for f in a compact subset of the retained intersection of $W^{r, p}$ spaces. If $F \in C^\infty([0, T]; H_\sigma^\infty)$, then

$$\partial_t^a (\mathcal{T}_k F) = \mathcal{T}_k (\partial_t^a F) \quad (a \in \mathbb{N}_0), \quad (333)$$

and the convergence in (332) is uniform in t for every fixed temporal derivative.

Proof. Choose $\chi \in C_c^\infty(B_2)$ with $\chi = 1$ on B_1 , and put $\chi_R(x) = \chi(x/R)$ and $A_R = B_{2R} \setminus \overline{B_R}$. For a smooth solenoidal f , set

$$g_R := \nabla \chi_R \cdot f.$$

It is supported in A_R , and

$$\int_{A_R} g_R \, dx = \int_{\mathbb{R}^3} \nabla \cdot (\chi_R f) \, dx = 0.$$

We now construct the required compact anti-divergence explicitly. Fix $\rho \in C_c^\infty((-3, 3))$ with $\int_{\mathbb{R}} \rho = 1$. If $h \in C_c^\infty(B_2)$ and $\int h = 0$, set

$$m_1(y_2, y_3) := \int_{\mathbb{R}} h(s, y_2, y_3) \, ds, \quad m_2(y_3) := \int_{\mathbb{R}} m_1(s, y_3) \, ds$$

and define $\mathcal{G}h = (G_1, G_2, G_3)$ by

$$\begin{aligned} G_1(y) &:= \int_{-\infty}^{y_1} (h(s, y_2, y_3) - \rho(s)m_1(y_2, y_3)) \, ds, \\ G_2(y) &:= \rho(y_1) \int_{-\infty}^{y_2} (m_1(s, y_3) - \rho(s)m_2(y_3)) \, ds, \\ G_3(y) &:= \rho(y_1)\rho(y_2) \int_{-\infty}^{y_3} m_2(s) \, ds. \end{aligned}$$

The three integrands have zero total integral in their integration variables; for the last one this is $\int h = 0$. Thus $\mathcal{G}h \in C_c^\infty((-3, 3)^3; \mathbb{R}^3)$, and direct differentiation gives

$$\nabla \cdot \mathcal{G}h = h - \rho(y_1)m_1 + \rho(y_1)m_1 - \rho(y_1)\rho(y_2)m_2 + \rho(y_1)\rho(y_2)m_2 = h.$$

Integration over intervals of length at most six, Hölder's inequality, Fubini's theorem, and differentiation under the integral give, for $a \in \mathbb{N}_0$ and $1 < p < \infty$,

$$\|\mathcal{G}h\|_{W^{a,p}((-3,3)^3)} \leq C_{a,p,\rho} \|h\|_{W^{a,p}(B_2)}. \quad (334)$$

For $g \in C_c^\infty(B_{2R})$ of mean zero, put

$$(\mathcal{G}_R g)(x) := R[\mathcal{G}(g(R \cdot))](x/R).$$

Then $\nabla \cdot \mathcal{G}_R g = g$, its support lies in $(-3R, 3R)^3$, and scaling (334) gives

$$\begin{aligned} \|\mathcal{G}_R g\|_{L^p} &\leq C_p R \|g\|_{L^p}, \\ \|\nabla^j \mathcal{G}_R g\|_{L^p} &\leq C_{j,p} \sum_{b=0}^j R^{1-j+b} \|\nabla^b g\|_{L^p}, \quad j \geq 1. \end{aligned} \quad (335)$$

Define

$$\mathcal{S}_R f := \chi_R f - \mathcal{G}_R g_R.$$

Then $\mathcal{S}_R f \in C_{c,\sigma}^\infty(\mathbb{R}^3)$, with support in $(-3R, 3R)^3$. Leibniz' rule gives, for $0 \leq a \leq r$,

$$\|\nabla^a g_R\|_{L^p(A_R)} \leq C_a \sum_{b=0}^a R^{-1-a+b} \|\nabla^b f\|_{L^p(A_R)}.$$

Combining this estimate with (335) and the analogous cutoff estimate yields

$$\|\mathcal{S}_R f - f\|_{W^{r,p}} \leq C_{r,p} \sum_{b=0}^r (1 + R^{b-r}) \|\nabla^b f\|_{L^p(\mathbb{R}^3 \setminus B_R)}. \quad (336)$$

For $r = 0$, this conclusion follows from (335) and $\|g_R\|_p \leq CR^{-1} \|f\|_{L^p(A_R)}$. The right side tends to zero as $R \rightarrow \infty$, and these calculations give operator norms bounded independently of R .

Choose $R_k \uparrow \infty$ and set $\mathcal{T}_k := \mathcal{S}_{R_k}$. Estimate (336) holds simultaneously for the finite family $\mathcal{A}$, while the preceding operator estimates give (331) simultaneously for that family. Strong convergence plus the uniform operator bound makes convergence uniform on compact subsets: cover a compact set by finitely many norm balls and use the bound on each remainder. Each $\mathcal{T}_k$ is independent of time, so differentiation commutes with it. The ranges of $\partial_t^a F$ on $[0, T]$ are compact in every retained space, which proves the final assertion. $\square$

Lemma 9.11 (Admissible strong-history test). *Let $\nu > 0$, let $u_0 \in H_\sigma^\infty(\mathbb{R}^3)$, prescribe $f \in C^\infty([0, \infty); \mathcal{S}(\mathbb{R}^3; \mathbb{R}^3))$, and let u be the global solution supplied by Theorem 6.10 for (u_0, ν, f) , and let v be a forced Leray–Hopf solution on $[0, T]$ at this viscosity with the same prescribed force f . Put $w := v - u$. Then, for every $t \in [0, T]$,*

$$\begin{aligned} \langle v(t), u(t) \rangle &= \langle v_0, u_0 \rangle + \int_0^t \int_{\mathbb{R}^3} (w \cdot \nabla u) \cdot w \, dx \, ds \\ &\quad - 2\nu \int_0^t \langle \nabla v, \nabla u \rangle \, ds + \int_0^t \langle f, u + v \rangle \, ds. \end{aligned} \quad (337)$$

Proof. The energy class and spatial interpolation give

$$v \in L^4(0, T; L^3), \quad \|v\|_{L^4 L^3}^4 \leq C \|v\|_{L^\infty L^2}^2 \|\nabla v\|_{L^2 L^2}^2,$$

so $v \otimes v \in L^2(0, T; L^{3/2})$.

Fix $0 < t < T$. Solenoidal density on $\mathbb{R}^3$, applied uniformly to the compact sets $u([0, T])$ and $u_t([0, T])$ in their finite Sobolev spaces, gives

$$u^{(k)} \in C^\infty([0, T]; C_{c,\sigma}^\infty(\mathbb{R}^3))$$

such that

$$u^{(k)} \longrightarrow u \quad \text{in} \quad W^{1,1}(0, T; L^2) \cap L^2(0, T; H^1) \cap L^2(0, T; W^{1,3}) \cap C([0, T]; L^2). \quad (338)$$

Apply Lemma 9.10 with

$$\mathcal{A} = \{(0, 2), (1, 2), (1, 3)\}$$

and set $u^{(k)} = \mathcal{T}_k u$. Its uniform compact-range conclusion for u and u_t , together with time commutation (333), gives (338).

For $0 < h < T - t$, choose $\eta_h \in C_c^\infty([0, T])$ with $0 \leq \eta_h \leq 1$, $\eta_h = 1$ on $[0, t]$, $\eta_h = 0$ on $[t + h, T]$, and $\eta_h' \leq 0$. The field $\eta_h u^{(k)}$ is an admissible test in (330). First let $k \rightarrow \infty$. The convergence in (338), together with $v \otimes v \in L^2 L^{3/2}$, permits passage in every term.

Choose the time cutoffs so that $-\eta'_h$ is an approximate identity from the right at t , and let $h \downarrow 0$. Weak continuity of v , strong continuity of u , and the Lebesgue differentiation theorem give

$$\begin{aligned} \langle v(t), u(t) \rangle &= \langle v_0, u_0 \rangle + \int_0^t \langle v, u_t \rangle \, ds + \int_0^t \int_{\mathbb{R}^3} (v \otimes v) : \nabla u \, dx \, ds \\ &\quad - \nu \int_0^t \langle \nabla v, \nabla u \rangle \, ds + \int_0^t \langle f, u \rangle \, ds. \end{aligned}$$

This identity first holds for almost every t . Its right-hand side is continuous in t , while $t \mapsto \langle v(t), u(t) \rangle$ is continuous by weak continuity of v and norm continuity of u . Hence it holds for every $0 \leq t \leq T$, including $t = T$ by passage from below. Using

$$u_t = -(u \cdot \nabla)u - \nabla p + \nu \Delta u + f$$

is admissible because v is divergence free. It gives

$$\int_0^t \langle v, u_t \rangle \, ds = - \int_0^t \int_{\mathbb{R}^3} v \cdot (u \cdot \nabla)u \, dx \, ds - \nu \int_0^t \langle \nabla v, \nabla u \rangle \, ds + \int_0^t \langle v, f \rangle \, ds.$$

Finally,

$$\int_{\mathbb{R}^3} [(v \otimes v) : \nabla u - v \cdot (u \cdot \nabla)u] \, dx = \int_{\mathbb{R}^3} (w \cdot \nabla u) \cdot w \, dx,$$

because

$$\int_{\mathbb{R}^3} (w \cdot \nabla u) \cdot u \, dx = \frac{1}{2} \int_{\mathbb{R}^3} w \cdot \nabla |u|^2 \, dx = 0.$$

This proves (337). $\square$

Theorem 9.12 (Quantitative Leray–Hopf stability). *Let $\nu > 0$, $u_0 \in H^\infty_\sigma(\mathbb{R}^3)$, and prescribe $f \in C^\infty([0, \infty); \mathcal{S}(\mathbb{R}^3; \mathbb{R}^3))$. Let u be the global strong history from Theorem 6.10 for (u_0, ν, f) . Fix $T > 0$, and let v be any forced Leray–Hopf solution on $[0, T]$ at this viscosity, with datum $v_0 \in H^0_\sigma(\mathbb{R}^3)$ and the same prescribed force f . Then, for $0 \leq t \leq T$,*

$$\begin{aligned} &\frac{1}{2} \|v(t) - u(t)\|_2^2 + \nu \int_0^t \|\nabla(v - u)(s)\|_2^2 \, ds \\ &\leq \frac{1}{2} \|v_0 - u_0\|_2^2 \exp\left(2 \int_0^t \|\nabla u(s)\|_\infty \, ds\right) \end{aligned} \tag{339}$$

$$\leq \frac{1}{2} \|v_0 - u_0\|_2^2 \exp\left(2S_{3,1,\infty} T^{1/2} \mathfrak{P}_{0,2}[u_0, \nu, f; T]^{1/2}\right). \tag{340}$$

The lower spatial rectangle also gives the viscosity-weighted estimate

$$\begin{aligned} &\frac{1}{2} \|v(t) - u(t)\|_2^2 + \frac{\nu}{2} \int_0^t \|\nabla(v - u)(s)\|_2^2 \, ds \\ &\leq \frac{1}{2} \|v_0 - u_0\|_2^2 \exp\left(\frac{S_{2,0,\infty}^2}{\nu} \mathfrak{X}_{0,2}[u_0, \nu, f; T]^2\right). \end{aligned} \tag{341}$$

Proof. Combine (329), the forced strong energy equality (152), and (337). The two energy relations contribute $\int_0^t \langle f, u + v \rangle \, ds$, exactly the force work in the cross identity, so the common prescribed force cancels without changing histories. With $w = v - u$, this gives

$$\frac{1}{2} \|w(t)\|_2^2 + \nu \int_0^t \|\nabla w(s)\|_2^2 \, ds \leq \frac{1}{2} \|w(0)\|_2^2 - \int_0^t \int_{\mathbb{R}^3} (w \cdot \nabla u) \cdot w \, dx \, ds. \quad (342)$$

The integral term in (342) has absolute value at most

$$\|\nabla u(s)\|_\infty \|w(s)\|_2^2.$$

Gronwall's inequality bounds $\|w(t)\|_2^2$. Substitution of that bound back into (342) proves (339). Proposition 8.16 gives

$$\int_0^T \|\nabla u(t)\|_\infty \, dt \leq S_{3,1,\infty} T^{1/2} \mathfrak{P}_{0,2}[u_0, \nu, f; T]^{1/2},$$

which proves (340).

For the lower-rectangle estimate, integration by parts and $\nabla \cdot w = 0$ give

$$\left| \int_{\mathbb{R}^3} (w \cdot \nabla u) \cdot w \, dx \right| = \left| \int_{\mathbb{R}^3} (w \cdot \nabla w) \cdot u \, dx \right| \leq \|u\|_\infty \|w\|_2 \|\nabla w\|_2.$$

Young's inequality in (342) gives

$$\frac{1}{2} \|w(t)\|_2^2 + \frac{\nu}{2} \int_0^t \|\nabla w(s)\|_2^2 \, ds \leq \frac{1}{2} \|w(0)\|_2^2 + \frac{1}{2\nu} \int_0^t \|u(s)\|_\infty^2 \|w(s)\|_2^2 \, ds.$$

Set

$$A(t) := \frac{1}{\nu} \int_0^t \|u(s)\|_\infty^2 \, ds.$$

Gronwall's inequality gives $\|w(t)\|_2^2 \leq \|w(0)\|_2^2 e^{A(t)}$. Substitution in the preceding inequality gives

$$\frac{1}{2} \|w(t)\|_2^2 + \frac{\nu}{2} \int_0^t \|\nabla w(s)\|_2^2 \, ds \leq \frac{1}{2} \|w(0)\|_2^2 e^{A(t)}.$$

Finally, (379) yields

$$A(t) \leq \frac{S_{2,0,\infty}^2}{\nu} \mathfrak{X}_{0,2}[u_0, \nu, f; T]^2,$$

which proves (341). □

Corollary 9.13 (Explicit weak-class stability for small forced critical data). *Under the hypotheses of Theorem 9.3, let v be a global forced Leray–Hopf solution at this viscosity ν , with datum $v_0 \in H_\sigma^0(\mathbb{R}^3)$ and the same prescribed force f . Then*

$$\begin{aligned} & \frac{1}{2} \|v(t) - u(t)\|_2^2 + \nu \int_0^t \|\nabla(v - u)(s)\|_2^2 \, ds \\ & \leq \frac{1}{2} \|v_0 - u\|_2^2 \exp(2S_{3,1,\infty} t U_3^{\text{sm}}) \quad (t \geq 0). \end{aligned} \quad (343)$$

Proof. Equation (299) gives $\sup_{s \geq 0} \|u(s)\|_{H^3} \leq U_3^{\text{sm}}$. Hence

$$\int_0^t \|\nabla u(s)\|_\infty \, ds \leq S_{3,1,\infty} t U_3^{\text{sm}}.$$

The derivation of (339) uses only the smooth strong history, the Leray–Hopf energy inequality, and the common prescribed force. Applying that derivation directly to the global small-critical history from Theorem 9.3 gives the stated explicit small-data refinement. $\square$

Corollary 9.14 (Uniqueness and energy equality in the Leray–Hopf class). *Let $\nu > 0$, $u_0 \in H_\sigma^\infty(\mathbb{R}^3)$, and prescribe $f \in C^\infty([0, \infty); \mathcal{S}(\mathbb{R}^3; \mathbb{R}^3))$. If (v, q) is any global three-dimensional forced Leray–Hopf solution at viscosity ν , with datum u_0 and prescribed force f , then v agrees on every finite interval with the velocity of the global forced solution from Theorem 6.10. Thus v has the smooth representative*

$$u \in C^\infty([0, \infty) \times \mathbb{R}^3; \mathbb{R}^3),$$

and it satisfies

$$\frac{1}{2} \|u(t)\|_2^2 + \nu \int_0^t \|\nabla u(s)\|_2^2 \, ds = \frac{1}{2} \|u_0\|_2^2 + \int_0^t \langle f(s), u(s) \rangle \, ds \quad (t \geq 0).$$

Its distributional pressure has the form

$$q = p + c(t) \quad \text{in } \mathcal{D}'((0, \infty) \times \mathbb{R}^3), \quad (344)$$

where p is the pressure in (151) and c is a scalar distribution in time. The canonical Riesz representative

$$R_i R_j (u_i u_j) - (-\Delta)^{-1} \nabla \cdot f$$

equals p . If q is represented by a locally integrable function with $q(t, \cdot) \in C_0(\mathbb{R}^3)$ for almost every t , then $q = p$ almost everywhere.

Proof. Set $v_0 = u_0$ in Theorem 9.12. The right-hand side of (340) is zero, so $v = u$ in $C_w([0, T]; L^2) \cap L^2(0, T; \dot{H}^1)$. Since T is arbitrary, the identity holds globally. Smoothness and the forced energy equality follow from (148) and (152). If q is a distributional pressure paired with this velocity and p is the pressure in (151), subtraction of the two momentum equations gives

$$\nabla(q - p) = 0 \quad \text{in } \mathcal{D}'((0, \infty) \times \mathbb{R}^3).$$

A distribution on a product domain whose spatial gradient vanishes is the pullback of a scalar distribution in time: testing first against spatial test functions of integral zero and then against one fixed test function of integral one gives (344). The Riesz formula for p proves the canonical-representative assertion. In the function-valued case, $q(t, x) - p(t, x) = c(t)$ for almost every (t, x) ; both spatial slices tend to zero as $|x| \rightarrow \infty$, so $c(t) = 0$ for almost every t . $\square$

10 Conclusion

The argument follows one forced velocity–pressure history from its prescribed datum (u_0, ν, f) . Differentiating the complete equation generates its velocity, pressure, and force mixed jet. For every finite pair of temporal and spatial orders, the datum-generated construction produces a compatible whole-terminal rectangle containing the selected rows and their adjacent rows. Each rectangle keeps every one of those derivative norms finite through the chosen finite endpoint, and the compatible family forms the complete space–time intersection on the whole maximal interval. From the same jet, and independently of that construction, the critical-height tower and the viscous diagonal relation supply the relationship identities on the coordinates of every finite rectangle. Integrating the complete rectangle relations keeps the transport, pressure, viscosity, prescribed force, and adjacent temporal response inside those estimates as parts of the same equation.

The adjacent rows then have continuous Sobolev representatives through the proposed terminal time. Their compatible values determine $u_* = u(T_*) \in H_\sigma^\infty$, while the forced pressure formula and the velocity recurrence determine the complete endpoint jet from u_* and $(\partial_t^n f(T_*))_{n \geq 0}$. Local well-posedness restarts from the single condition $w(T_*) = u_*$ with the same prescribed absolute-time force $f(t)$, written as $f(T_* + \tau)$ in relative restart time. Uniqueness on the overlap joins this restart to the incoming history. Every proposed finite maximal time therefore admits a classical continuation, contradicting maximality. Hence the maximal time is infinite.

The resulting global forced solution is smooth, obeys the exact energy identity with force work, and carries the decay-normalized pressure

$$p = R_i R_j (u_i u_j) - (-\Delta)^{-1} \nabla \cdot f.$$

Its finite rectangles also quantify the critical-height rows, frequency tails, Lipschitz action, vorticity amplification and dissipation, material flow, and same-force relative-energy stability. The small-critical theory is an explicit quantitative refinement of this general construction: its scale-invariant velocity stock and projected-force stock give closed formulas for the corresponding finite rectangle bounds.

Finally, relative energy identifies every forced Leray–Hopf solution with the same initial datum, viscosity, and prescribed force with this smooth velocity; the common force work cancels in the comparison. The pressure is unique up to the unavoidable time-dependent gauge unless the decay normalization is imposed. Every step belongs to the one forced evolution generated by (u_0, ν, f) .

A Canonical coordinate catalogue

The estimates in Section 8 are organized by their role in the argument. For reference, the following theorem collects their canonical-coordinate forms in one place.

Theorem A.1 (Canonical finite-horizon coordinate estimates). *Under the hypotheses and notation of Definition 8.2, let $J_{n,\alpha} := \partial_x^\alpha V_n$ and $\Pi_{n,\alpha} := \partial_x^\alpha Q_n$, as in Theorems 8.10 and 8.11. Let $n, s, k, q \in \mathbb{N}_0$, $\alpha \in \mathbb{N}_0^3$, and set $r := s + |\alpha|$. Then*

$$\|J_{n,\alpha}\|_{L^2(I_T; H^{s+1})} \leq \mathfrak{X}_{n,r+1}, \quad (345)$$

$$\|J_{n+1,\alpha}\|_{L^2(I_T; H^{s-1})} \leq \mathfrak{X}_{n+1,r-1}, \quad (346)$$

$$\sup_{0 \leq t \leq T} \|J_{n,\alpha}(t)\|_{H^s} \leq \mathfrak{Z}_{n,r}, \quad (347)$$

$$\|J_{n,\alpha}^T\|_{H^s} \leq \mathfrak{Z}_{n,r}, \quad (348)$$

$$\|J_{n,\alpha}(t) - J_{n,\alpha}(r_0)\|_{H^{s-1}} \leq \min \left\{ |t - r_0|^{1/2} \mathfrak{X}_{n+1,r-1}, |t - r_0| \mathfrak{Z}_{n+1,r} \right\}, \quad (349)$$

$$\|J_{n,\alpha}(t) - J_{n,\alpha}(r_0)\|_{H^s} \leq \min \left\{ |t - r_0|^{1/2} \mathfrak{X}_{n+1,r}, |t - r_0| \mathfrak{Z}_{n+1,r} \right\}, \quad (350)$$

for $0 \leq r_0 \leq t \leq T$.

For $0 \leq \theta \leq 1$,

$$\|J_{n,\alpha}(t) - J_{n,\alpha}(r_0)\|_{H^{s-\theta}} \leq 2^{1-\theta} \mathfrak{Z}_{n,r}^{1-\theta} \mathfrak{X}_{n+1,r-1}^\theta |t - r_0|^{\theta/2}. \quad (351)$$

If $1 \leq a \leq \infty$, $2 \leq b \leq \infty$, and $q > k + \frac{3}{2} - \frac{3}{b}$, then

$$\left\| \nabla^k J_{n,\alpha} \right\|_{L^a(I_T; L^b)} \leq S_{q,k,b} T^{1/a} \mathfrak{Z}_{n,q+|\alpha|}, \quad (352)$$

$$\left\| \nabla^k J_{n,\alpha} \right\|_{L^2(I_T; L^b)} \leq S_{q,k,b} \mathfrak{X}_{n,q+|\alpha|}, \quad (353)$$

where $T^{1/\infty} = 1$.

The pressure rows obey

$$\|\Pi_{n,\alpha}\|_{L^1(I_T; H^s)} + \|\nabla \Pi_{n,\alpha}\|_{L^1(I_T; H^{s-1})} \leq \mathfrak{Q}_{n,r}^{(1)}, \quad (354)$$

$$\|\Pi_{n,\alpha}\|_{L^2(I_T; H^s)} \leq \mathfrak{Q}_{n,r}^{(2)}, \quad (355)$$

$$\sup_{0 \leq t \leq T} \|\Pi_{n,\alpha}(t)\|_{H^s} \leq \mathfrak{Q}_{n,r}^{(\infty)}, \quad (356)$$

$$\|\Pi_{n,\alpha}^T\|_{H^s} \leq \mathfrak{Q}_{n,r}^{(\infty)}, \quad (357)$$

$$\|\Pi_{n,\alpha}(t) - \Pi_{n,\alpha}(r_0)\|_{H^s} \leq \min \left\{ |t - r_0|^{1/2} \mathfrak{Q}_{n,r}^{(\text{mod})}, |t - r_0| \mathfrak{Q}_{n+1,r}^{(\infty)} \right\}. \quad (358)$$

Under the preceding restrictions on a, b, k, q ,

$$\left\| \nabla^k \Pi_{n,\alpha} \right\|_{L^a(I_T; L^b)} \leq S_{q,k,b} T^{1/a} \mathfrak{Q}_{n,q+|\alpha|}^{(\infty)}. \quad (359)$$

Let $r_1 \in \mathbb{N}$ and $K \geq 1$. Then

$$\|\mathbf{P}_{\geq K} J_{n,\alpha}\|_{L^2(I_T; H^s)} \leq K^{-r_1} \mathfrak{X}_{n,s+r_1+|\alpha|}, \quad (360)$$

$$\sup_{0 \leq t \leq T} \|\mathbf{P}_{\geq K} J_{n,\alpha}(t)\|_{H^s} \leq K^{-r_1} \mathfrak{Z}_{n,s+r_1+|\alpha|}, \quad (361)$$

$$\|\mathbf{P}_{\geq K} \Pi_{n,\alpha}\|_{L^1(I_T; H^s)} \leq K^{-r_1} \mathfrak{Q}_{n,s+r_1+|\alpha|}^{(1)}, \quad (362)$$

$$\sup_{0 \leq t \leq T} \|\mathbf{P}_{\geq K} \Pi_{n,\alpha}(t)\|_{H^s} \leq K^{-r_1} \mathfrak{Q}_{n,s+r_1+|\alpha|}^{(\infty)}. \quad (363)$$

For $j \in \mathbb{N}_0$,

$$\sup_{0 \leq t \leq T} E_{j+1/2}(t) \leq \frac{1}{2} \mathfrak{Z}_{0,j+1}^2, \quad (364)$$

$$\int_0^T E_{j+1/2}(t) dt \leq \frac{1}{2} \mathfrak{X}_{0,j+1}^2. \quad (365)$$

If $m > j + \frac{1}{2}$ is an integer, then

$$\sup_{0 \leq t \leq T} E_{j+1/2, \geq K}(t) \leq \frac{1}{2} K^{2j+1-2m} \mathfrak{Z}_{0,m}^2, \quad (366)$$

$$\int_0^T E_{j+1/2, \geq K}(t) dt \leq \frac{1}{2} K^{2j+1-2m} \mathfrak{X}_{0,m}^2. \quad (367)$$

For $k \in \mathbb{N}$,

$$\|H^{(k)}\|_{L^1(0,T)} \leq \frac{1}{2} \sum_{a=0}^k \binom{k}{a} \mathfrak{X}_{a,1} \mathfrak{X}_{k-a,1}. \quad (368)$$

For $j, \ell \in \mathbb{N}_0$, define

$$\begin{aligned} \mathfrak{E}_{j,\ell}^{[1]} &:= \frac{1}{2} \sum_{a=0}^{\ell} \binom{\ell}{a} \mathfrak{X}_{a,j+1} \mathfrak{X}_{\ell-a,j+1}, \\ \mathfrak{E}_{j,\ell}^{[\infty]} &:= \frac{1}{2} \sum_{a=0}^{\ell} \binom{\ell}{a} \mathfrak{Z}_{a,j+1} \mathfrak{Z}_{\ell-a,j+1}, \end{aligned} \quad (369)$$

$$\begin{aligned} \mathfrak{W}_{j,\ell}^{[1]} &:= \sum_{a=0}^{\ell} \binom{\ell}{a} \mathfrak{X}_{a,j+1} \mathfrak{G}_{\ell-a,j+1}^{(2)}, \\ \mathfrak{W}_{j,\ell}^{[\infty]} &:= \sum_{a=0}^{\ell} \binom{\ell}{a} \mathfrak{Z}_{a,j+1} \mathfrak{G}_{\ell-a,j+1}^{(\infty)}, \end{aligned} \quad (370)$$

$$\begin{aligned} \mathfrak{E}_{j,\ell}^{[1]} &:= \mathfrak{E}_{j,\ell+1}^{[1]} + 2\nu \mathfrak{E}_{j+1,\ell}^{[1]} + \mathfrak{W}_{j,\ell}^{[1]}, \\ \mathfrak{E}_{j,\ell}^{[\infty]} &:= \mathfrak{E}_{j,\ell+1}^{[\infty]} + 2\nu \mathfrak{E}_{j+1,\ell}^{[\infty]} + \mathfrak{W}_{j,\ell}^{[\infty]}. \end{aligned} \quad (371)$$

Then the complete differentiated energy and production rows obey

$$\left\| \partial_t^\ell E_{j+1/2} \right\|_{L^1(0,T)} \leq \mathfrak{E}_{j,\ell}^{[1]}, \quad \sup_{0 \leq t \leq T} \left| \partial_t^\ell E_{j+1/2}(t) \right| \leq \mathfrak{E}_{j,\ell}^{[\infty]}, \quad (372)$$

$$\left\| \partial_t^\ell \mathcal{W}_{j+1/2} \right\|_{L^1(0,T)} \leq \mathfrak{W}_{j,\ell}^{[1]}, \quad \sup_{0 \leq t \leq T} \left| \partial_t^\ell \mathcal{W}_{j+1/2}(t) \right| \leq \mathfrak{W}_{j,\ell}^{[\infty]}, \quad (373)$$

$$\left\| \partial_t^\ell \mathcal{P}_{j+1/2} \right\|_{L^1(0,T)} \leq \mathfrak{E}_{j,\ell}^{[1]}, \quad \sup_{0 \leq t \leq T} \left| \partial_t^\ell \mathcal{P}_{j+1/2}(t) \right| \leq \mathfrak{E}_{j,\ell}^{[\infty]}. \quad (374)$$

Every current in (70) now has a finite-order majorant retaining temporal depth, spatial height, viscosity, and horizon. In particular,

$$\|H'\|_{L^1(0,T)} \leq \mathfrak{X}_{1,1} \mathfrak{X}_{0,1}, \quad (375)$$

$$\int_0^T \left\| \Lambda^{3/2} u(t) \right\|_2^2 dt \leq \mathfrak{X}_{0,2}^2, \quad (376)$$

$$\|\mathcal{P}_{1/2}\|_{L^1(0,T)} \leq \mathfrak{X}_{1,1} \mathfrak{X}_{0,1} + \nu \mathfrak{X}_{0,2}^2 + \mathfrak{X}_{0,1} \mathfrak{G}_{0,1}^{(2)}, \quad (377)$$

$$|H(T) - H(t)| \leq \min \left\{ \mathfrak{Z}_{0,1} \mathfrak{X}_{1,1} (T-t)^{1/2}, \mathfrak{Z}_{0,1} \mathfrak{Z}_{1,1} (T-t) \right\}, \quad 0 \leq t \leq T. \quad (378)$$

Finally,

$$\|u\|_{L^2(I_T; L^\infty)} \leq S_{2,0,\infty} \mathfrak{X}_{0,2}, \quad (379)$$

$$\int_0^T \|\nabla u(t)\|_\infty dt \leq S_{3,1,\infty} T^{1/2} \mathfrak{X}_{0,3}, \quad (380)$$

$$\int_0^T \|\nabla \times u(t)\|_\infty dt \leq 2S_{3,1,\infty} T^{1/2} \mathfrak{X}_{0,3}, \quad (381)$$

$$\sup_{0 \leq t \leq T} \|u(t)\|_{H^m}^2 \leq \left(\|u_0\|_{H^m} + \mathfrak{G}_{0,m}^{(1)} \right)^2 \exp \left(2K_m S_{3,1,\infty} T^{1/2} \mathfrak{X}_{0,3} \right), \quad m \geq 3. \quad (382)$$

Every right-hand side is an explicit finite expression in the displayed analytic constants, the prescribed-force coordinates, the initial-jet recursion (201)–(202), and the finite collection of canonical rectangle coordinates $\mathfrak{B}_{a,b}[u_0, \nu, f; T]$ selected by the stated indices. Equations (198) and (203)–(240) specify every intermediate operation. Equation (229) rewrites each such rectangle coordinate through the finite height-coordinate recursion (219)–(227). Since (221) first bounds those height coordinates by the rectangle coordinates, this is a re-expression of the finite information already carried by the original history. This is the exact coordinate-level catalogue for general data: every downstream quantity is expressed through the finite rectangle selected by its displayed indices.

Proposition 8.6 displays the exact powers of ν and the rescaled horizon νT . For every fixed choice of the displayed indices, the definitions select only finitely many rectangle coordinates. Under the explicit critical smallness condition (293), Proposition 9.4 eliminates the rectangle coordinates and gives finite formulas solely in the displayed velocity-datum and prescribed-force norms, ν , T , N , and M .

Proof. The Fourier multiplier inequality

$$\|\partial_x^\alpha f\|_{H^\ell} \leq \|f\|_{H^{\ell+|\alpha|}},$$

followed by Lemma 8.8, proves (345)–(348). The Banach identity

$$J_{n,\alpha}(t) - J_{n,\alpha}(r_0) = \int_{r_0}^t J_{n+1,\alpha}(\tau) d\tau$$

in the two indicated Sobolev spaces and Cauchy–Schwarz prove the square-root branches of (349) and (350). This identity, followed by the pointwise bound (234) for V_{n+1} , proves their linear branches. Interpolation between the low modulus and

$$\|J_{n,\alpha}(t) - J_{n,\alpha}(r_0)\|_{H^s} \leq 2\mathfrak{Z}_{n,r}$$

proves (351). The fixed Sobolev embedding constant in (191) proves (352) and (353).

Apply the recurrence

$$Q_n = R_i R_j \sum_{a=0}^n \binom{n}{a} V_{a,i} V_{n-a,j} - \Phi_n$$

term by term at order r , using (189). Cauchy–Schwarz in time gives (354); placing either factor in $L_t^\infty H_x^{r+1}$ gives (355); placing both factors there gives (356) and (357), with the corresponding $\mathfrak{L}$ coordinate retaining $-\Phi_n$ in each case. After subtracting the recurrence at times t and r_0 , split each product difference into two terms and apply (350); this gives the square-root branch of (358). Since $\partial_t \Pi_{n,\alpha} = \Pi_{n+1,\alpha}$, the Banach identity and (356) at row $n+1$ give its linear branch. Sobolev embedding proves (359).

For every $f \in H^{\ell+r_1}$,

$$\|P_{\geq K} f\|_{H^\ell} \leq K^{-r_1} \|f\|_{H^{\ell+r_1}}.$$

Apply this inequality at spatial order $s+r_1$ to (233), (234), (354), and (356). The four resulting estimates are (360)–(363).

The inequalities

$$\|f\|_{\dot{H}^{j+1/2}} \leq \|f\|_{H^{j+1}}, \quad |\xi|^{2j+1} \leq K^{2j+1-2m} |\xi|^{2m} \quad (|\xi| \geq K),$$

give (364)–(367). The exact identity

$$H^{(k)}(t) = \frac{1}{2} \sum_{a=0}^k \binom{k}{a} \operatorname{Re} \left\langle \Lambda^{1/2} V_a(t), \Lambda^{1/2} V_{k-a}(t) \right\rangle_{L^2}$$

and Cauchy–Schwarz in time give (368) and (375). More generally,

$$\partial_t^\ell E_{j+1/2} = \frac{1}{2} \sum_{a=0}^{\ell} \binom{\ell}{a} \operatorname{Re} \left\langle \Lambda^{j+1/2} V_a, \Lambda^{j+1/2} V_{\ell-a} \right\rangle_{L^2}.$$

Cauchy–Schwarz in time and the pointwise estimates (233)–(234) prove (372). Leibniz’ rule applied to $\mathcal{W}_{j+1/2}$ gives the majorants (370). Differentiating the forced identity

$$\mathcal{P}_{j+1/2} = \partial_t E_{j+1/2} + 2\nu E_{j+3/2} - \mathcal{W}_{j+1/2}$$

then proves (374). The next two estimates follow from

$$\mathcal{P}_{1/2} = H' + \nu \left\| \Lambda^{3/2} u \right\|_2^2 - \mathcal{W}_{1/2}.$$

For the terminal critical-height modulus,

$$|H(T) - H(t)| \leq \frac{1}{2} (\|u(T)\|_{\dot{H}^{1/2}} + \|u(t)\|_{\dot{H}^{1/2}}) \|u(T) - u(t)\|_{\dot{H}^{1/2}},$$

and the Banach identity in H^1 prove the square-root branch of (378).

The identity

$$H' = \operatorname{Re} \left\langle \Lambda^{1/2} V_1, \Lambda^{1/2} V_0 \right\rangle_{L^2},$$

together with (234), proves its linear branch.

Sobolev embedding and (233) prove (379)–(381). Lemma 3.8, with the prescribed force retained, gives after regularization of the norm

$$\frac{d}{dt} \|u\|_{H^m} \leq K_m \|\nabla u\|_\infty \|u\|_{H^m} + \|\mathbb{P}f\|_{H^m}.$$

The inhomogeneous Gronwall inequality and (380) prove (382). □

Computational assistance

This manuscript was prepared with assistance from OpenAI’s ChatGPT and Codex. The source archive and dated revision history are available at github.com/Jibberfish69/navier-stokes-public.

References

- [1] J.-P. Aubin, *Un théorème de compacité*, C. R. Acad. Sci. Paris **256** (1963), 5042–5044.
- [2] H. Beirão da Veiga, *A new regularity class for the Navier–Stokes equations in $\mathbb{R}^n$* , Chinese Annals of Mathematics, Series B **16** (1995), no. 4, 407–412, Journal record.
- [3] T. Birnie, *Thomas’s one-fluid reversible-intersection proof spine*, author’s proof-construction record (2026), Section 5.
- [4] L. Caffarelli, R. Kohn, and L. Nirenberg, *Partial regularity of suitable weak solutions of the Navier–Stokes equations*, Communications on Pure and Applied Mathematics **35** (1982), 771–831, doi:10.1002/cpa.3160350604.

- [5] L. Escauriaza, G. A. Seregin, and V. Šverák, *$L_{3,\infty}$ -solutions of Navier–Stokes equations and backward uniqueness*, Russian Mathematical Surveys **58** (2003), no. 2, 211–250, doi:10.1070/RM2003v058n02ABEH000609.
- [6] C. Fefferman, *Existence and smoothness of the Navier–Stokes equation*, in *The Millennium Prize Problems*, Clay Mathematics Institute, American Mathematical Society, 2006, pp. 57–67, Clay problem statement.
- [7] H. Fujita and T. Kato, *On the Navier–Stokes initial value problem. I*, Archive for Rational Mechanics and Analysis **16** (1964), 269–315, doi:10.1007/BF00276188.
- [8] E. Hopf, *Über die Anfangswertaufgabe für die hydrodynamischen Grundgleichungen*, Mathematische Nachrichten **4** (1950/51), 213–231, doi:10.1002/mana.3210040121.
- [9] T. Kato, *Strong L^p -solutions of the Navier–Stokes equation in $\mathbb{R}^m$, with applications to weak solutions*, Mathematische Zeitschrift **187** (1984), 471–480, EuDML record.
- [10] H. Koch and D. Tataru, *Well-posedness for the Navier–Stokes equations*, Advances in Mathematics **157** (2001), no. 1, 22–35, doi:10.1006/aima.2000.1937.
- [11] J. Leray, *Sur le mouvement d’un liquide visqueux emplissant l’espace*, Acta Mathematica **63** (1934), 193–248, doi:10.1007/BF02547354.
- [12] E. H. Lieb, *Sharp constants in the Hardy–Littlewood–Sobolev and related inequalities*, Annals of Mathematics **118** (1983), 349–374, doi:10.2307/2007032.
- [13] J.-L. Lions and E. Magenes, *Non-Homogeneous Boundary Value Problems and Applications*, vol. I, Springer, 1972, doi:10.1007/978-3-642-65161-8.
- [14] J. Serrin, *On the interior regularity of weak solutions of the Navier–Stokes equations*, Archive for Rational Mechanics and Analysis **9** (1962), 187–195, doi:10.1007/BF00253344.
- [15] J. Simon, *Compact sets in the space $L^p(0, T; B)$* , Annali di Matematica Pura ed Applicata **146** (1986), 65–96, doi:10.1007/BF01762360.
- [16] T. Tao, *Localisation and compactness properties of the Navier–Stokes global regularity problem*, Analysis & PDE **6** (2013), 25–107, doi:10.2140/apde.2013.6.25.