

# 1 Introduction

## 1.1 Clay Requirements and Definitions

Let  $\Omega$  be either  $\mathbb{R}^3$  or the periodic domain  $\mathbb{T}^3 = \mathbb{R}^3/\mathbb{Z}^3$ . Fix a viscosity  $\nu > 0$ , set the external force to zero, and choose a smooth divergence-free initial velocity  $u_0$ . On  $\mathbb{R}^3$ , require  $u_0$  and all of its spatial derivatives to decay rapidly at infinity; on  $\mathbb{T}^3$ , require periodicity instead. The velocity  $u(x, t) \in \mathbb{R}^3$  and pressure  $p(x, t) \in \mathbb{R}$  must satisfy

$$\partial_t u + (u \cdot \nabla)u + \nabla p = \nu \Delta u, \quad \nabla \cdot u = 0, \quad u(x, 0) = u_0(x).$$

The local theory produces a unique smooth solution on a short interval, and  $T_* \in (0, \infty]$  denotes the largest time to which that smooth solution can be continued. The positive existence-and-smoothness problem studied in this paper is exactly whether

$$T_* = \infty$$

for every admissible  $u_0$ .

Each term in the equation belongs to the same fluid motion. The derivative  $\partial_t u$  records the change of velocity at a fixed spatial point, while  $(u \cdot \nabla)u$  records the change caused by carrying velocity through a nonuniform field. The pressure gradient transmits the constraint imposed by incompressibility, and  $\nu \Delta u$  transfers momentum across spatial variation. The condition  $\nabla \cdot u = 0$  says that the flow preserves volume. The problem asks whether these coupled effects can evolve the initial fluid forever without producing a loss of smoothness.

## Smoothness

**Definition 1.1.1** (Smoothness). A velocity–pressure pair  $(u, p)$  is smooth through a finite time  $T > 0$  when

$$u \in C^\infty(\Omega \times [0, T]; \mathbb{R}^3), \quad p \in C^\infty(\Omega \times [0, T]).$$

Equivalently, every mixed spatial and temporal derivative of every finite order exists and is continuous through  $t = T$ . The pair is smooth for all time when it is smooth through every finite  $T > 0$ .

The word smooth has a precise all-orders meaning. It does not require one numerical constant to bound every derivative order simultaneously. It requires each finite derivative order to remain well defined and continuous whenever that order is examined.

## Regularity

**Definition 1.1.2** (Regularity). Regularity describes the type and degree of control available for a solution. In the classical scale,  $C^0$  means continuity,  $C^1$  adds continuous first derivatives, and  $C^k$  requires all mixed derivatives through total order  $k$ . Smoothness is the all-orders case  $C^\infty$ . Sobolev and mixed space–time bounds are other forms of regularity, expressed through norms that measure integral control of the field and its derivatives.

A regularity statement must name its level. Continuity of the velocity does not by itself control its gradient, and control of finitely many derivatives does not by itself give  $C^\infty$ . The historical continuation criteria below matter because they identify levels of regularity strong enough to recover all remaining finite orders from the equations.

## Continuation and Maximal Time

**Definition 1.1.3** (Continuation). Suppose  $(u, p)$  solves the Navier–Stokes equations on  $\Omega \times [0, T)$ . A continuation past  $T$  is a velocity–pressure pair  $(\tilde{u}, \tilde{p})$  solving the same equations with the same viscosity on  $\Omega \times [0, T + \varepsilon)$  for some  $\varepsilon > 0$ , agreeing with  $(u, p)$  before  $T$ , and belonging to the required regularity class on the larger interval.

For the Clay problem, the required class is  $C^\infty$ . The adjective maximal means that the smooth solution has no such continuation beyond  $T_*$ . Thus  $T_* < \infty$  can occur only if the solution loses enough regularity near  $T_*$  to defeat every applicable local restart.

## Finite Energy

On the whole space, the required solution also has bounded kinetic energy:

$$\sup_{t \geq 0} \int_{\mathbb{R}^3} |u(x, t)|^2 dx < \infty.$$

This condition keeps the total motion physically finite, yet it is much weaker than smoothness because it controls the velocity in  $L^2$  without directly controlling its pointwise values or higher derivatives. The distinction between finite energy and smoothness is where the historical estimates begin.

## What a Loss of Smoothness Means

At a finite endpoint  $T$ ,  $C^k$  regularity is lost when at least one required mixed derivative through order  $k$  fails to extend continuously to  $\Omega \times \{T\}$ . That derivative may become unbounded, or it may remain bounded while approaching incompatible limiting values. A failure at any finite order prevents continuation in  $C^\infty$ . The formal problem is whether the maximal smooth time can ever be finite.

## 1.2 Historical Background

A smooth local solution gives a genuine Navier–Stokes fluid for a short time, while the Clay problem asks whether that same solution can cross every finite time. The history of the problem can be read as a sequence of attempts to close that gap. Each major estimate controls one feature of the motion, reveals what remains free to concentrate, and leaves a sharper version of the continuation question behind it. The order below follows that tightening of the problem, with publication dates serving as secondary orientation.

## Leray's Global Energy Control

The first decisive calculation asks what the equations preserve before any derivative is placed under pointwise control. For a smooth solution, take the spatial inner product of the momentum equation with  $u$ . Incompressibility turns the transport term into a spatial divergence and makes the pressure term orthogonal to the velocity, while integration by parts turns viscosity into dissipation. The result is

$$\|u(t)\|_{L^2}^2 + 2\nu \int_0^t \|\nabla u(s)\|_{L^2}^2 ds = \|u_0\|_{L^2}^2.$$

The velocity stays bounded in  $L^2$  at each time, and viscosity pays for the total  $L^2$  gradient cost accumulated over the interval.

Leray used regularized equations, compactness, and weak lower semicontinuity to carry the corresponding inequality into a global weak solution (Leray 1934). On every finite interval, a Leray solution satisfies

$$u \in L^\infty(0, T; L^2(\Omega)) \cap L^2(0, T; H^1(\Omega)).$$

This is genuine global existence in the energy class. It does not settle the Clay problem because smoothness and uniqueness are not known for arbitrary three-dimensional Leray solutions.

The two terms in the estimate also carry different information in time. The  $L^2$  size of  $u$  is bounded at every instant, whereas the  $L^2$  size of  $\nabla u$  is controlled only after time integration. A function can have finite area under its graph and still form arbitrarily tall peaks on arbitrarily short intervals. The energy inequality leaves open the possibility that derivative activity concentrates near a finite time even though the total kinetic energy remains finite.

## Enstrophy and Vortex Stretching

For a divergence-free field on  $\mathbb{R}^3$  with sufficient decay, or on  $\mathbb{T}^3$ ,

$$\|\nabla u\|_{L^2}^2 = \|\omega\|_{L^2}^2, \quad \omega := \nabla \times u.$$

The quantity  $\|\omega\|_{L^2}^2$  is the enstrophy, so Leray's estimate controls the area under the enstrophy curve without controlling its height.

The vorticity equation shows why this distinction becomes serious in three dimensions:

$$\partial_t \omega + (u \cdot \nabla) \omega = (\omega \cdot \nabla) u + \nu \Delta \omega.$$

Transport moves vorticity with the fluid, and viscosity diffuses sharp spatial variation. The remaining term,  $(\omega \cdot \nabla) u$ , stretches and tilts vorticity through the surrounding velocity gradient. Its contribution to the enstrophy balance is

$$\frac{1}{2} \frac{d}{dt} \|\omega\|_{L^2}^2 + \nu \|\nabla \omega\|_{L^2}^2 = \int_{\Omega} (\omega \cdot \nabla) u \cdot \omega \, dx.$$

The stretching integral has no fixed sign. Viscosity continues to oppose spatial concentration, while the energy estimate alone gives no pointwise-in-time comparison proving that diffusion always defeats stretching.

This is the first clear shape of the three-dimensional obstruction. The velocity cannot escape the global energy bound, yet the derivatives that determine classical smoothness may rise through narrow peaks that the energy balance does not see individually. This leaves a precise question: how much derivative control would be enough to continue the smooth solution?

## Sobolev Regularity and Continuation

Sobolev spaces answer that question by measuring velocity and spatial derivatives in one norm. For an integer  $m \geq 0$ ,

$$H^m(\Omega) = \left\{ v : \partial^\alpha v \in L^2(\Omega) \text{ for every } |\alpha| \leq m \right\},$$

with the usual weak interpretation of the derivatives and the Fourier definition for fractional orders. An  $L^p$  norm records the spatial size and concentration of the function itself, while an  $H^m$  norm also records its variation through order  $m$ .

Sobolev embedding converts sufficiently high  $L^2$ -derivative control into ordinary continuous derivatives. In three dimensions (Cerdà 2010, Theorem 7.28),

$$H^s(\Omega) \hookrightarrow C^k(\Omega) \quad \text{whenever} \quad s > k + \frac{3}{2}.$$

One  $H^s$  bound buys only finitely many continuous derivatives. For example,  $s > 5/2$  gives

$$H^s(\Omega) \hookrightarrow W^{1,\infty}(\Omega),$$

so the velocity and its first spatial derivative are bounded and continuous.

The threshold  $s > 5/2$  is useful because the Lipschitz norm of  $u$  controls the way transport amplifies every higher derivative already present in the smooth datum. For each fixed finite  $m \geq s$ , the standard differentiated estimate has the form

$$\frac{1}{2} \frac{d}{dt} \|u\|_{H^m}^2 + \nu \|\nabla u\|_{H^m}^2 \leq C_m \|\nabla u\|_{L^\infty} \|u\|_{H^m}^2.$$

Once  $\|\nabla u\|_{L^\infty}$  is integrable in time, Gronwall's inequality propagates every fixed finite Sobolev order from the smooth initial datum. The constant may depend on  $m$ , which is enough because smoothness asks for every finite order separately and imposes no one numerical bound across the infinite family. Applying Sobolev embedding at successively higher orders then recovers every continuous spatial derivative.

The pressure follows from the velocity through the elliptic relation

$$-\Delta p = \sum_{i,j=1}^3 \partial_i \partial_j (u_i u_j),$$

with zero mean on the torus or the usual decay normalization on  $\mathbb{R}^3$ . Once all spatial derivatives of  $u$  and  $p$  are controlled, the momentum equation

$$\partial_t u = \nu \Delta u - (u \cdot \nabla) u - \nabla p$$

recovers the first time derivative. Repeatedly differentiating the pressure relation and the momentum equation then recovers every finite mixed spatial and temporal derivative. This is the continuation argument from a strong Sobolev bound to the  $C^\infty$  velocity–pressure pair required by the problem.

The local theories of Fujita–Kato and Kato supply the restart mechanism behind this argument (Fujita and Kato 1964; Kato 1984). Suppose a smooth solution exists on  $[0, T)$  and, for one  $s > 5/2$ ,

$$\sup_{0 \leq t < T} \|u(t)\|_{H^s} \leq M.$$

Choose an actual time  $t_0 < T$  close to  $T$ . Local theory restarted from  $u(t_0)$  has a lifespan bounded below in terms of  $s$ ,  $M$ , and  $\nu$ , so a sufficiently late choice of  $t_0$  produces a solution beyond  $T$ . Uniqueness identifies the restarted solution with the original one on their overlap. Thus a finite maximal smooth time  $T_*$  would force every continuation-grade  $H^s$  norm to become unbounded as  $t \uparrow T_*$ .

This criterion identifies the exact kind of estimate that would close the problem. The energy inequality supplies

$$u \in L_t^\infty L_x^2 \cap L_t^2 H_x^1,$$

while continuation asks for pointwise-in-time control at a substantially stronger derivative level. The distance between those two statements is the analytic room in which a finite-time singularity could still hide.

## Partial Regularity and the Shape of a Singular Set

Continuation theory describes what must diverge along a maximal classical solution, while the global weak theory asks where regularity can fail after that classical description is no longer available. Caffarelli, Kohn, and Nirenberg studied suitable weak solutions through the local energy inequality on shrinking parabolic cylinders (Caffarelli, Kohn, and Nirenberg 1982). For  $z_0 = (x_0, t_0)$ , write

$$Q_r(z_0) := B_r(x_0) \times (t_0 - r^2, t_0).$$

Their epsilon-regularity mechanism says that sufficiently small scale-invariant velocity and pressure quantities on  $Q_r(z_0)$  force regularity on a smaller cylinder.

At a singular point, the contrapositive applies at every sufficiently small scale. Every shrinking cylinder centred there must retain a definite amount of scale-invariant concentration, because one cylinder below the threshold would already make the point regular. A covering argument then confines the possible singular set to a set of zero one-dimensional parabolic Hausdorff measure. This is an exceptionally small set, yet even one singular point would defeat global smoothness.

Partial regularity changes the picture of failure. A counterexample cannot be a diffuse loss of control spread harmlessly across the fluid. It must concentrate velocity and pressure through

every smaller parabolic neighbourhood of at least one spacetime point. That conclusion creates the need for criteria that measure spatial intensity and temporal persistence together.

## Scale-Critical Continuation Criteria

The Ladyzhenskaya–Prodi–Serrin criteria provide that joint measurement (Prodi 1959; Serrin 1962; Serrin 1963; Ladyzhenskaya 1967). They give regularity through  $T$  whenever

$$u \in L^p(0, T; L^q(\Omega)), \quad \frac{2}{p} + \frac{3}{q} \leq 1, \quad q > 3.$$

The exponent  $q$  measures the spatial intensity of a concentration, and  $p$  measures how long that intensity can persist.

At one derivative, interpolation and Young’s inequality lead to

$$\frac{d}{dt} \|\nabla u\|_{L^2}^2 + \nu \|\Delta u\|_{L^2}^2 \leq C_{\nu, q} \|u\|_{L^q}^{\frac{2q}{q-3}} \|\nabla u\|_{L^2}^2.$$

The mixed-norm condition makes the coefficient on the right integrable in time, so Gronwall’s inequality keeps the enstrophy finite and continuation crosses  $T$ . At the equality

$$\frac{2}{p} + \frac{3}{q} = 1,$$

the criterion is invariant under the Navier–Stokes scaling

$$u_\lambda(x, t) := \lambda u(\lambda x, \lambda^2 t).$$

These critical exponents match exactly the balance between stronger spatial concentration and shorter parabolic duration.

The direct estimate above covers every  $q > 3$  and degenerates as  $q \downarrow 3$ . The endpoint  $q = 3$  is especially important because

$$\|u_\lambda(t)\|_{L^3} = \|u(\lambda^2 t)\|_{L^3}.$$

Magnifying a suspected singularity to unit scale does not weaken the norm used to measure it.

Escauriaza, Seregin, and Šverák proved that boundedness in  $L_t^\infty L_x^3$  still excludes a finite singular time (Escauriaza, G. A. Seregin, and Šverák 2003). Their proof assumes a first singular time, rescales around the concentration, extracts a nontrivial ancient limiting solution, and uses backward uniqueness and unique continuation to force that limit to vanish. The contradiction shows that

$$u \in L^\infty(0, T; L^3(\mathbb{R}^3))$$

is sufficient for regularity through  $T$ .

Seregin subsequently showed, for the smooth compactly supported data in his whole-space result, that a finite maximal time forces full divergence of the critical norm (G. Seregin 2012):

$$\lim_{t \uparrow T_*} \|u(t)\|_{L^3(\mathbb{R}^3)} = \infty.$$

The critical norm cannot spike along a thin sequence of times and repeatedly return below a fixed level. Every finite level must eventually be left behind as the alleged endpoint approaches.

Tao later quantified part of this conclusion (Tao 2021). Along infinitely many times approaching a finite maximal time, the  $L^3$  norm must exceed a positive power of a triple logarithm of the inverse time remaining. This rate is weak compared with a power law, yet it proves that a finite-time singularity would require a genuine cascade through successively smaller scales.

## Where the Problem Sits Today

The established theory entering this paper replaces an undefined breakdown with a sharply constrained alternative. If  $T_* < \infty$ , the kinetic energy remains bounded, while continuation-grade Sobolev norms and the critical  $L^3$  norm must diverge. Any singular point must retain scale-invariant concentration in every sufficiently small parabolic cylinder around it. The concentration may accelerate through shorter and shorter time intervals, because parabolic scale reduction also shortens the available time scale.

The basic scaling already explains why the global energy estimate cannot rule this out. At corresponding times,

$$\|u_\lambda(t)\|_{L^3} = \|u(\lambda^2 t)\|_{L^3}, \quad \|u_\lambda(t)\|_{L^2}^2 = \lambda^{-1} \|u(\lambda^2 t)\|_{L^2}^2.$$

Compression to a shorter length scale preserves the critical  $L^3$  size while reducing the  $L^2$  energy seen at that scale. A scale-independent ceiling on the critical norm cannot be read directly from the global energy bound.

A simple frequency-doubling picture makes the remaining timing problem visible. If an active frequency doubles at each stage, its spatial scale halves and its parabolic time scale quarters. The resulting durations form a convergent geometric series, so positive duration at every stage still leaves room for infinitely many stages before one finite time. This is the Zeno-cascade picture behind the endpoint threat. It is a heuristic picture of the allowed timing, not a construction of a Navier–Stokes singularity.

The public frontier has isolated the exact burden. Known theory says what must diverge, how concentrated the divergence must become, and which bounds would exclude it. It does not derive from arbitrary smooth data, fixed viscosity, and the equations a continuation-grade bound that remains finite all the way to every candidate endpoint. That is the direct estimate a Gold-standard proof would have to supply.

### 1.3 Gold Standard Proof

The historical results leave a clean direct route to global smoothness. Let  $[0, T_*)$  be the maximal interval of the smooth solution generated by  $u_0$ . A Gold-standard proof derives, for some fixed  $s_0 > 5/2$ ,

$$\sup_{0 \leq t < T_*} \|u(t)\|_{H^{s_0}(\Omega)} < \infty$$

from  $u_0$ ,  $\nu$ , and the Navier–Stokes equations themselves. This is the direct estimate absent from the classical Gold route surveyed above.

The payoff is larger than the single norm might suggest. Sobolev embedding gives  $H^{s_0} \hookrightarrow W^{1,\infty}$ , so the bound controls the velocity gradient that appears in every higher differentiated transport estimate. For each fixed finite  $m \geq s_0$ ,

$$\frac{1}{2} \frac{d}{dt} \|u\|_{H^m}^2 + \nu \|\nabla u\|_{H^m}^2 \leq C_m \|\nabla u\|_{L^\infty} \|u\|_{H^m}^2.$$

The initial datum already belongs to every  $H^m$ , and Gronwall's inequality propagates each finite order to  $T_*$ . The constants may depend on  $m$ , because  $C^\infty$  requires every finite derivative and imposes no common numerical bound on the entire infinite family. Higher Sobolev embeddings then recover all continuous spatial derivatives.

Pressure is recovered from the same velocity through

$$-\Delta p = \sum_{i,j=1}^3 \partial_i \partial_j (u_i u_j),$$

and the momentum equation recovers  $\partial_t u$ . Repeated differentiation recovers every higher mixed derivative of  $u$  and  $p$ . A late time slice can then be used as new initial data for local theory, and the uniform  $H^{s_0}$  bound gives enough lifespan to cross  $T_*$ . This contradicts maximality, so the direct estimate forces  $T_* = \infty$ .

The obstruction appears as soon as one tries to produce that estimate from the basic energy law. Energy gives

$$\|u(t)\|_{L^2}^2 + 2\nu \int_0^t \|\nabla u(s)\|_{L^2}^2 ds = \|u_0\|_{L^2}^2.$$

It controls the total time integral of the squared gradient. Raising the calculation by one spatial derivative gives

$$\frac{1}{2} \frac{d}{dt} \|\nabla u\|_{L^2}^2 + \nu \|\Delta u\|_{L^2}^2 = \int_{\Omega} (u \cdot \nabla) u \cdot \Delta u dx.$$

The transport term now survives, and the usual interpolation and Young inequalities give

$$\left| \int_{\Omega} (u \cdot \nabla) u \cdot \Delta u dx \right| \leq \frac{\nu}{2} \|\Delta u\|_{L^2}^2 + C\nu^{-3} \|\nabla u\|_{L^2}^6.$$

After viscosity absorbs the first term, the estimate asks for time control of the sixth power of the gradient. The energy identity supplies time control only of its second power. Tall, narrow peaks can respect the second-power integral while defeating the sixth-power integral, so this estimate does not close the first derivative level. That observation identifies an estimate gap; it does not assert that the fluid actually forms those peaks.

The Gold burden is easy to state and hard to pay. One must derive a continuation-grade bound from the coupled action of transport, pressure, incompressibility, and viscosity without assuming the strong quantity the estimate is meant to control. Once that payment is made, the Sobolev bootstrap, pressure recovery, and local restart complete the argument. Silver keeps the same smoothness claim and changes the direction of attack by asking what any alleged nonsmooth outcome must have ceased to satisfy.



## 1.4 Silver Standard Proof

Looking for estimates strong enough to satisfy the Gold-standard burden is genuinely difficult. The public theory surveyed above did not close that direct route, and once you dig your teeth into the problem, it becomes tempting to look for any other way of proving smoothness.

Assume, for a moment, that the problem has already been solved by the Gold-standard method.

That would mean smoothness follows from the Navier–Stokes equations and the properties built into them. It would have to come from the velocity, pressure, viscosity, incompressibility, and nonlinear transport all acting in unison to keep the fluid smooth. Any hypothetical blow-up scenario, or any other case that breaks smoothness, would then be unreachable while all of those requirements remain in force. The imagined nonsmooth object would have to stop describing the same fluid from the original problem statement. If every possible loss of smoothness stops satisfying that original problem, then none of those scenarios is a valid Navier–Stokes counterexample. If there are no valid counterexamples, the fluid must be smooth.

In short, the idea becomes this.

If

$$\text{Navier–Stokes} \implies \text{smooth},$$

then

$$\text{not smooth} \implies \text{not Navier–Stokes}.$$

The first implication was assumed only to reveal the logical shape of the second. A Silver proof has to establish the second implication directly by showing that every alleged nonsmooth outcome loses at least one requirement of the original fluid. The Silver Standard is a proof by contrapositive.

The beauty of the Silver method is that it no longer asks *how is the fluid smooth?* It asks only *is this nonsmooth object still Navier–Stokes?* Gold demands a mechanism strong enough to produce the continuation estimate, which is why it earns the higher podium title. Silver shifts the burden away from constructing that estimate and onto excluding every genuinely nonsmooth outcome from the original class. Its upside is the simpler logical target. Its downside is the apparent need to handle every possible way smoothness could fail. A single principle that reaches all of those cases would simplify that burden.

## References

- [1] Luis Caffarelli, Robert V. Kohn, and Louis Nirenberg. “Partial regularity of suitable weak solutions of the Navier–Stokes equations”. In: *Communications on Pure and Applied Mathematics* 35.6 (1982), pp. 771–831. DOI: 10.1002/cpa.3160350604.
- [2] Joan Cerdà. *Linear Functional Analysis*. Vol. 116. Graduate Studies in Mathematics. Providence, RI: American Mathematical Society, 2010. ISBN: 978-0-8218-5115-9.
- [3] L. Escauriaza, G. A. Seregin, and V. Šverák. “ $L_{3,\infty}$ -solutions of the Navier–Stokes equations and backward uniqueness”. In: *Russian Mathematical Surveys* 58.2 (2003), pp. 211–250. DOI: 10.1070/RM2003v058n02ABEH000609.

- [4] Hiroshi Fujita and Tosio Kato. “On the Navier–Stokes Initial Value Problem. I”. In: *Archive for Rational Mechanics and Analysis* 16.4 (1964), pp. 269–315. DOI: 10.1007/BF00276188.
- [5] Tosio Kato. “Strong  $L^p$ -solutions of the Navier–Stokes equation in  $\mathbb{R}^m$ , with applications to weak solutions”. In: *Mathematische Zeitschrift* 187.4 (1984), pp. 471–480. DOI: 10.1007/BF01174182.
- [6] O. A. Ladyzhenskaya. “On the uniqueness and on the smoothness of weak solutions of the Navier–Stokes equations”. In: *Zapiski Nauchnykh Seminarov LOMI* 5 (1967), pp. 169–185. URL: <https://www.mathnet.ru/eng/zns12228>.
- [7] Jean Leray. “Sur le mouvement d’un liquide visqueux emplissant l’espace”. In: *Acta Mathematica* 63 (1934), pp. 193–248. DOI: 10.1007/BF02547354.
- [8] Giovanni Prodi. “Un teorema di unicità per le equazioni di Navier–Stokes”. In: *Annali di Matematica Pura ed Applicata* 48.1 (1959), pp. 173–182. DOI: 10.1007/BF02410664.
- [9] Gregory Seregin. “A Certain Necessary Condition of Potential Blow Up for Navier–Stokes Equations”. In: *Communications in Mathematical Physics* 312.3 (2012), pp. 833–845. DOI: 10.1007/s00220-011-1391-x.
- [10] James Serrin. “On the interior regularity of weak solutions of the Navier–Stokes equations”. In: *Archive for Rational Mechanics and Analysis* 9 (1962), pp. 187–195. DOI: 10.1007/BF00253344.
- [11] James Serrin. “The Initial Value Problem for the Navier–Stokes Equations”. In: *Nonlinear Problems*. Madison, WI: University of Wisconsin Press, 1963, pp. 69–98.
- [12] Terence Tao. “Quantitative Bounds for Critically Bounded Solutions to the Navier–Stokes Equations”. In: *Nine Mathematical Challenges—An Elucidation*. Vol. 104. Proceedings of Symposia in Pure Mathematics. Providence, RI: American Mathematical Society, 2021, pp. 149–193. DOI: 10.1090/pspum/104/01874.