

STATISTISCHE METHODEN DER DATENANALYSE

Excercise Sheets

Physik

UNI WIEN

17. November 2023

1 Sheet

1.1 ML Estimator

1.1.1 Calculating the Estimator

The Joint probability density of an exponentially distributed sample is

$$g(x_1, \dots, x_m | \tau) = \prod_{i=1}^n \frac{1}{\tau} e^{-\frac{x_i}{\tau}} \quad (1.1)$$

The logarithm of this density interpreted as a likelihood function is

$$l(\tau) = \ln g(x_1, \dots, x_n | \tau) = \sum_{i=1}^n \ln \left(\frac{1}{\tau} e^{-\frac{x_i}{\tau}} \right) \quad (1.2)$$

$$= \sum_{i=1}^n -\ln(\tau) - \frac{1}{\tau} x_i \quad (1.3)$$

Maximizing the chance to draw this particular sample:

$$\frac{\partial l}{\partial \tau} = \sum_{i=1}^n -\frac{1}{\tau} + \frac{1}{\tau^2} x_i \stackrel{!}{=} 0 \quad (1.4)$$

$$\Leftrightarrow -n + \sum_{i=1}^n \frac{1}{\tau} x_i = 0 \quad (1.5)$$

$$\Leftrightarrow \hat{\tau} = \frac{1}{n} \sum_{i=1}^n x_i \quad (1.6)$$

Inserting $s = \sum x_i = 395.25$ and $n = 250$ yields

$$\hat{\tau} = 1.581 \quad (1.7)$$

1.1.2 Showing that the Estimator is efficient

Showing $\hat{\tau}$ is unbiased

$$E_{\tau} \left[\frac{1}{n} \sum_{i=1}^n x_i \right] = \frac{1}{n} \sum_{i=1}^n E[x_i] \quad (1.8)$$

$$= \frac{1}{n} \sum_{i=1}^n \tau \quad (1.9)$$

$$= \tau \quad (1.10)$$

Showing $\hat{\tau}$ is efficient Calculate the Fisher Information

$$I_{\tau} = E \left[-\frac{\partial^2 \ln g(x_1, \dots, x_n | \tau)}{\partial \tau^2} \right] \quad (1.11)$$

$$= E \left[-\sum_{i=1}^n \frac{1}{\tau^2} - \frac{2}{\tau^3} x_i \right] \quad (1.12)$$

$$= -\sum_{i=1}^n \frac{1}{\tau^2} - \frac{2}{\tau^3} E[x_i] \quad (1.13)$$

$$= -\sum_{i=1}^n \frac{1}{\tau^2} - \frac{2}{\tau^2} \quad (1.14)$$

$$= \sum_{i=1}^n \frac{1}{\tau^2} \quad (1.15)$$

$$= \frac{n}{\tau^2} \quad (1.16)$$

Comparing with estimator variance

$$V[\hat{\tau}] = V \left[\frac{1}{n} \sum_{i=1}^n x_i \right] \quad (1.17)$$

$$= \frac{\tau^2}{n} \quad (1.18)$$

Conclusion, estimator is as efficient as it gets.

1.2 Laplace distribution

1.2.1 Expectation Value and Variance

The Laplace distribution density is

$$f(x; m, s) = \frac{1}{2s} \exp \left[-\frac{|x - m|}{s} \right] \quad (1.19)$$

Expectation Value The expectation value of laplace distributed variable X is

$$E[X] = \int_{-\infty}^{\infty} \frac{x}{2s} \exp \left[-\frac{|x - m|}{s} \right] dx \quad (1.20)$$

$$= \frac{1}{2s} \int_{-\infty}^{\infty} (u + m) \exp \left[-\frac{|u|}{s} \right] du \quad (1.21)$$

$$= \frac{1}{2s} \int_{-\infty}^{\infty} u \exp \left[-\frac{|u|}{s} \right] du + m \quad (1.22)$$

$$= m \quad (1.23)$$

Variance

$$V[X] = E[(X - m)^2] \quad (1.24)$$

$$= \frac{1}{2s} \int_{-\infty}^{\infty} (x - m)^2 \exp\left\{-\frac{|x - m|}{s}\right\} dx \quad (1.25)$$

$$= \frac{1}{2s} \int_{-\infty}^{\infty} u^2 \exp\left\{-\frac{|u|}{s}\right\} du \quad (1.26)$$

$$= \frac{1}{s} \int_0^{\infty} u^2 \exp\left\{-\frac{u}{s}\right\} du \quad (1.27)$$

$$= s^3 \int_0^{\infty} v^2 e^{-v} dv \quad (1.28)$$

$$= 2s^3 \quad (1.29)$$

1.2.2 Estimators for m and s

For a given sample the joint density is

$$g(x_1, \dots, x_n | s, m) = \prod_{i=1}^n \frac{1}{2s} e^{-\frac{|x_i - m|}{s}} \quad (1.30)$$

The log likelihood function is

$$\ln g = \sum_{i=1}^n \ln \left(\frac{1}{2s} \exp\left\{-\frac{|x_i - m|}{s}\right\} \right) \quad (1.31)$$

$$= \sum_{i=1}^n \ln \left(\frac{1}{2s} \right) - \frac{|x_i - m|}{s} \quad (1.32)$$

$$= n \ln \left(\frac{1}{2s} \right) - \frac{1}{s} \sum_{i=1}^n |x_i - m| \quad (1.33)$$

The maximum likelihood estimator for m can now be found

$$\frac{\partial \ln g}{\partial s} = \frac{\partial}{\partial s} \left[n \ln \left(\frac{1}{2s} \right) - \frac{1}{s} \sum_{i=1}^n |x_i - m| \right] \quad (1.34)$$

$$= -\frac{n}{s} + \frac{1}{s^2} \sum_{i=1}^n |x_i - m| \stackrel{!}{=} 0 \quad (1.35)$$

$$\Rightarrow \hat{s} = \frac{1}{n} \sum_{i=1}^n |x_i - m| \quad (1.36)$$

The maximum likelihood estimator for s can also be found

$$\frac{\partial \ln g}{\partial m} = \frac{\partial}{\partial m} \left[n \ln \left(\frac{1}{2s} \right) - \frac{1}{s} \sum_{i=1}^n |x_i - m| \right] \quad (1.37)$$

$$= -\frac{1}{s} \sum_{i=1}^n \frac{\partial}{\partial m} |x_i - m| \quad (1.38)$$

$$= -\frac{1}{s} \sum_{i=1}^n \text{sgn}(x_i - m) \quad (1.39)$$

1.3 Survey

The multimodal distribution is defined as

$$f(n_A, n_B, n_C, n_D | p_A, p_B, p_C, p_D) = n! \prod_{i=A,B,C,D} \frac{1}{n_i!} p_i^{n_i} \quad (1.40)$$

with the log density

$$\ln f = \ln \left(n! \prod_{i=A,B,C,D} \frac{1}{n_i!} p_i^{n_i} \right) \quad (1.41)$$

$$= \ln(n!) + \sum_{i=A,B,C,D} \ln \left(\frac{1}{n_i!} \right) + \ln(p_i^{n_i}) \quad (1.42)$$

$$= \ln(n!) + \sum_{i=A,B,C,D} \ln \left(\frac{1}{n_i!} \right) + n_i \ln(p_i) \quad (1.43)$$

Now the estimators for the voter shares can be found with

$$\frac{\partial \ln f}{\partial p_i} = \frac{n_i}{p_i} \quad (1.44)$$