

QUANTUM MECHANICS I

Winter 2023, Jichao Yang

1 Vectors and Dirac Notation

Remark 1.1. *This course will use two books, Shanker is more advanced and will be assigned with reading, Griffiths has more problem and easy-going in terms of difficulty.*

Definition 1.2. A vector space V on a field \mathbb{F} is a set of vectors that are closed under addition and multiplication. That is, for any $v, w \in V$, $a, b \in \mathbb{F}$, $av + bw \in V$.

Remark 1.3. *From now on we will use V to denote a vector space, a, b, c, \dots to denote elements of \mathbb{F} , and u, v, w, \dots to denote elements of V .*

Definition 1.4. Vector addition and multiplication follow a set of axioms, which can be found in Shanker 1.1. Some of the most important ones are listed below:

1. $v + w = w + v$
2. $(u + v) + w = u + (v + w)$
3. $a \cdot (v + w) = a \cdot v + a \cdot w$

Definition 1.5. A real number space is a V over \mathbb{R} . A complex number space is a V over \mathbb{C} .

Remark 1.6. *From now on we will be limiting our discussion to complex number spaces.*

Example 1.7. Complex column vectors:

$$\begin{pmatrix} x_1 \\ \vdots \\ x_k \end{pmatrix} \tag{1}$$

Each $x_i \in \mathbb{C}$. Check that both operations are closed.

Example 1.8. Complex function vectors: $F = \{f : [0, L] \subset \mathbb{R} \rightarrow \mathbb{C}\}$. Define addition and scalar multiplication as function addition and multiplying function with scalar. Check function is closed under both operations.

Definition 1.9. v_1, \dots, v_n are linearly dependent if exists a_1, \dots, a_n , at least one of them non-zero, such that

$$\sum_{i=1}^n a_i \cdot v_i = 0$$

v_1, \dots, v_n are linearly independent otherwise.

Definition 1.10. The dimension of a vector field V is the maximum number of linearly independent vectors subset to V .

Remark 1.11. *Example 1.7 has dimension k , example 1.8 has infinite dimension.*

Definition 1.12. v_1, \dots, v_n span V if and only if exists a_1, \dots, a_n , at least one nonzero, such that

$$a_1 v_1 + \dots + a_n v_n = 0$$

Definition 1.13. v_1, \dots, v_n is a basis of V if and only if they span V and are linearly independent.

Corollary 1.14. If v_1, \dots, v_n are a basis for V , then any $v \in V$ can be uniquely expressed by a linear combination of v_1, \dots, v_n .

Definition 1.15. The complex conjugate of some complex number $a + bi$ is $a - bi$.

Definition 1.16. An inner product function is $(\cdot) : V \times V \rightarrow \mathbb{C}$, denoted as (w, v) such that

1. $(w, a_1 v_1 + a_2 v_2) = a_1 \cdot (w, v_1) + a_2 \cdot (w, v_2)$
2. $(v, w) \geq 0$, and only equal if v or w equals 0.
3. $(w, v) = (v, w)^*$

Corollary 1.17. $(a_1 w_1 + a_2 w_2, v) = a_1^* \cdot (w_1, v) + a_2^* \cdot (w_2, v)$

Proof. First note that $(c_1 \cdot c_2)^* = (c_1)^* \cdot (c_2)^*$.

$$\begin{aligned} ((a + bi)(c + di))^* &= (ac - bd + (ad + bc)i)^* \\ &= ac - bd - (ad + bc)i = (a - bi)(c - di) \\ &= (a + bi)^*(c + di)^* \end{aligned}$$

Hence the proof:

$$\begin{aligned} (a_1 w_1 + a_2 w_2, v) &= (v, a_1 w_1 + a_2 w_2)^* \\ &= (a_1(v, w_1) + a_2(v, w_2))^* = (a_1(v, w_1))^* + (a_2(v, w_2))^* \\ &= a_1^*(v, w_1)^* + a_2^*(v, w_2)^* = a_1^*(w_1, v) + a_2^*(w_2, v) \end{aligned}$$

□

Example 1.18. The inner product of example 1.7 is $(w, v) = w_1^* v_1 + \cdots + w_k^* v_k$.

Example 1.19. The inner product of example 1.8 is

$$(f, g) = \int_0^L f^*(x) \cdot g(x) dx$$

Definition 1.20. The norm (sometimes called length) of v denoted $\|v\|$ is

$$\|v\| = \sqrt{(v, v)}$$

Remark 1.21. *The inner product is a function that maps to \mathbb{C} , but $(v, v) \in \mathbb{R}$, so the square root for Definition 1.20 is the real number square root.*

Definition 1.22. v, w are orthogonal if and only if $(v, w) = 0$.

Definition 1.23. v_1, \dots, v_n are orthonormal if and only if for any v_i, v_j ,

$$(v_i, v_j) = \delta_{ij} = \text{bool}(i == j)$$

Where δ_{ij} is called the 'kronecker delta.'

Corollary 1.24. v_1, \dots, v_n are orthonormal if and only if they are orthogonal and have length 1.

Remark 1.25. *From now on we will limit the discussion to vector spaces with well-defined inner-products.*

Definition 1.26. A linear functional F is a linear function $F : V \rightarrow \mathbb{C}$.

Corollary 1.27. The set of linear functionals $\{F\}$ form a vector space.

Proof. Check closedness over addition and multiplication of linear functions. □

Definition 1.28. Given any vector space V , exists a corresponding vector space V' , named the dual space, which is the set of all linear functions on V .