

COMPSCI 589

Lecture 24: Principal Components Analysis

Benjamin M. Marlin

College of Information and Computer Sciences
University of Massachusetts Amherst

Slides by Benjamin M. Marlin (marlin@cs.umass.edu).
Created with support from National Science Foundation Award# IIS-1350522.

Outline


1 Review

2 Linear Algebra

3 PCA

4 Connection to SVD

Machine Learning Tasks



The Dimensionality Reduction Task


Definition: The Dimensionality Reduction Task

Given a collection of feature vectors $\mathbf{x}_i \in \mathbb{R}^p$, map the feature vectors into a lower dimensional space $\mathbf{z}_i \in \mathbb{R}^q$ where $q < p$ while preserving certain properties of the data.

The Dimensionality Reduction Task

Definition: The Dimensionality Reduction Task


Given a collection of feature vectors $\mathbf{x}_i \in \mathbb{R}^p$, map the feature vectors into a lower dimensional space $\mathbf{z}_i \in \mathbb{R}^q$ where $q < p$ while preserving certain properties of the data.



high-dim distribution




high-dim samples



estimated manifold

Linear Dimensionality Reduction

- The simplest dimensionality reduction methods assume that the observed high dimensional data vectors $\mathbf{x}_i \in \mathbb{R}^p$ lie on a q -dimensional linear manifold within \mathbb{R}^p .
- Mathematically, the linear sub-space assumption can be written as $\mathbf{X} = \mathbf{Z} \times \mathbf{B}$



Learning

- Minimize reconstruction error as

$$\min \|\mathbf{X} - \tilde{\mathbf{X}}\|_F^2$$

- Using encoder ($t(\mathbf{X})$) and decoder (S):

$$\min \sum_{i=1}^N \|X_i - S(t(X_i))\|_F^2$$

- Best encoder for fixed decoder $S(\lambda) = \mu + V_q \lambda$ is
 $\lambda = V_q^T (X - \mu)$

- Updated reconstruction error:

$$\min_{V_q} \sum_{i=1}^N \|X_i - V_q V_q^T (X_i)\|_F^2$$

Singular Value Decomposition

- SVD representation of

$$\mathbf{X} = \mathbf{U}\mathbf{D}\mathbf{V}^T$$

where \mathbf{D} is a $p \times p$ diagonal matrix with positive elements, \mathbf{U} is an $N \times p$ matrix such that $\mathbf{U}^T\mathbf{U} = \mathbf{I}$, and \mathbf{V} is a $p \times p$ matrix such that $\mathbf{V}^T\mathbf{V} = \mathbf{I}$.

- PCA using SVD

$$\|\mathbf{X} - \mathbf{U}\mathbf{D}_q^T\mathbf{V}_q^T\|_F^2$$

where $q < p$

Outline

1 Review

2 Linear Algebra

3 PCA

4 Connection to SVD

Eigenvectors

- Let $\mathbf{A} \in \mathbb{R}^{p \times p}$ be a matrix, $\mathbf{v} \in \mathbb{R}^p$ be a vector, and λ be scalar.

Eigenvectors

- Let $\mathbf{A} \in \mathbb{R}^{p \times p}$ be a matrix, $\mathbf{v} \in \mathbb{R}^p$ be a vector, and λ be scalar.
- If $\mathbf{A}\mathbf{v} = \lambda\mathbf{v}$ then \mathbf{v} is a right eigenvector of A with eigenvalue λ .

Eigenvectors

- Let $\mathbf{A} \in \mathbb{R}^{p \times p}$ be a matrix, $\mathbf{v} \in \mathbb{R}^p$ be a vector, and λ be scalar.
- If $\mathbf{A}\mathbf{v} = \lambda\mathbf{v}$ then \mathbf{v} is a right eigenvector of A with eigenvalue λ .
- If $\mathbf{A}^T\mathbf{v} = \lambda\mathbf{v}$ then \mathbf{v} is a left eigenvector of A with eigenvalue λ (equivalently $\mathbf{v}^T\mathbf{A} = \lambda\mathbf{v}^T$).

Eigenvectors

- Let $\mathbf{A} \in \mathbb{R}^{p \times p}$ be a matrix, $\mathbf{v} \in \mathbb{R}^p$ be a vector, and λ be scalar.
- If $\mathbf{A}\mathbf{v} = \lambda\mathbf{v}$ then \mathbf{v} is a right eigenvector of A with eigenvalue λ .
- If $\mathbf{A}^T\mathbf{v} = \lambda\mathbf{v}$ then \mathbf{v} is a left eigenvector of A with eigenvalue λ (equivalently $\mathbf{v}^T\mathbf{A} = \lambda\mathbf{v}^T$).
- If \mathbf{A} is symmetric so that $\mathbf{A} = \mathbf{A}^T$, then the left and right eigenvectors of \mathbf{A} are the same with the same eigenvalues.

Eigenvectors

- Let $\mathbf{A} \in \mathbb{R}^{p \times p}$ be a matrix, $\mathbf{v} \in \mathbb{R}^p$ be a vector, and λ be scalar.
- If $\mathbf{A}\mathbf{v} = \lambda\mathbf{v}$ then \mathbf{v} is a right eigenvector of A with eigenvalue λ .
- If $\mathbf{A}^T\mathbf{v} = \lambda\mathbf{v}$ then \mathbf{v} is a left eigenvector of A with eigenvalue λ (equivalently $\mathbf{v}^T\mathbf{A} = \lambda\mathbf{v}^T$).
- If \mathbf{A} is symmetric so that $\mathbf{A} = \mathbf{A}^T$, then the left and right eigenvectors of \mathbf{A} are the same with the same eigenvalues.

$$\begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix}$$

Eigenvectors

- Let $\mathbf{A} \in \mathbb{R}^{p \times p}$ be a matrix, $\mathbf{v} \in \mathbb{R}^p$ be a vector, and λ be scalar.
- If $\mathbf{A}\mathbf{v} = \lambda\mathbf{v}$ then \mathbf{v} is a right eigenvector of A with eigenvalue λ .
- If $\mathbf{A}^T\mathbf{v} = \lambda\mathbf{v}$ then \mathbf{v} is a left eigenvector of A with eigenvalue λ (equivalently $\mathbf{v}^T\mathbf{A} = \lambda\mathbf{v}^T$).
- If \mathbf{A} is symmetric so that $\mathbf{A} = \mathbf{A}^T$, then the left and right eigenvectors of \mathbf{A} are the same with the same eigenvalues.

$$\begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 3 \\ 3 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

Eigenvectors

- Let $\mathbf{A} \in \mathbb{R}^{p \times p}$ be a matrix, $\mathbf{v} \in \mathbb{R}^p$ be a vector, and λ be scalar.
- If $\mathbf{A}\mathbf{v} = \lambda\mathbf{v}$ then \mathbf{v} is a right eigenvector of A with eigenvalue λ .
- If $\mathbf{A}^T\mathbf{v} = \lambda\mathbf{v}$ then \mathbf{v} is a left eigenvector of A with eigenvalue λ (equivalently $\mathbf{v}^T\mathbf{A} = \lambda\mathbf{v}^T$).
- If \mathbf{A} is symmetric so that $\mathbf{A} = \mathbf{A}^T$, then the left and right eigenvectors of \mathbf{A} are the same with the same eigenvalues.

$$\begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 3 \\ 3 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 \end{bmatrix} \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix} = 3 \begin{bmatrix} 1 & 1 \end{bmatrix}$$

Eigenvectors

- Let $\mathbf{A} \in \mathbb{R}^{p \times p}$ be a matrix, $\mathbf{v} \in \mathbb{R}^p$ be a vector, and λ be scalar.
- If $\mathbf{A}\mathbf{v} = \lambda\mathbf{v}$ then \mathbf{v} is a right eigenvector of A with eigenvalue λ .
- If $\mathbf{A}^T\mathbf{v} = \lambda\mathbf{v}$ then \mathbf{v} is a left eigenvector of A with eigenvalue λ (equivalently $\mathbf{v}^T\mathbf{A} = \lambda\mathbf{v}^T$).
- If \mathbf{A} is symmetric so that $\mathbf{A} = \mathbf{A}^T$, then the left and right eigenvectors of \mathbf{A} are the same with the same eigenvalues.

$$\begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 3 \\ 3 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 \end{bmatrix} \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix} = 3 \begin{bmatrix} 1 & 1 \end{bmatrix}$$

- A full-rank (invertible) matrix $\mathbf{A} \in \mathbb{R}^{p \times p}$ will have p linearly independent eigenvectors.

Eigendecomposition

- Let $\mathbf{V} \in \mathbb{R}^{p \times p}$ be a matrix whose columns \mathbf{v}_d are p linearly independent eigenvectors of \mathbf{A} with Λ the corresponding diagonal matrix of eigenvalues such that $\Lambda_{dd} = \lambda_d$. Then:

$$\mathbf{A}\mathbf{V} = \mathbf{V}\Lambda$$

Eigendecomposition

- Let $\mathbf{V} \in \mathbb{R}^{p \times p}$ be a matrix whose columns \mathbf{v}_d are p linearly independent eigenvectors of \mathbf{A} with Λ the corresponding diagonal matrix of eigenvalues such that $\Lambda_{dd} = \lambda_d$. Then:

$$\mathbf{A}\mathbf{V} = \mathbf{V}\Lambda$$

$$\mathbf{A} = \mathbf{V}\Lambda\mathbf{V}^{-1}$$

Eigendecomposition

- Let $\mathbf{V} \in \mathbb{R}^{p \times p}$ be a matrix whose columns \mathbf{v}_d are p linearly independent eigenvectors of \mathbf{A} with Λ the corresponding diagonal matrix of eigenvalues such that $\Lambda_{dd} = \lambda_d$. Then:

$$\mathbf{A}\mathbf{V} = \mathbf{V}\Lambda$$

$$\mathbf{A} = \mathbf{V}\Lambda\mathbf{V}^{-1}$$

$$\mathbf{V}^{-1}\mathbf{A}\mathbf{V} = \Lambda$$

Eigendecomposition

- Let $\mathbf{V} \in \mathbb{R}^{p \times p}$ be a matrix whose columns \mathbf{v}_d are p linearly independent eigenvectors of \mathbf{A} with Λ the corresponding diagonal matrix of eigenvalues such that $\Lambda_{dd} = \lambda_d$. Then:

$$\mathbf{A}\mathbf{V} = \mathbf{V}\Lambda$$

$$\mathbf{A} = \mathbf{V}\Lambda\mathbf{V}^{-1}$$

$$\mathbf{V}^{-1}\mathbf{A}\mathbf{V} = \Lambda$$

- Without loss of generality, we can assume that

$$\lambda_1 > \lambda_2 > \dots > \lambda_n$$

Eigendecomposition of a Symmetric Matrix

- If \mathbf{A} is symmetric, we can choose p orthonormal eigenvectors so that $\|\mathbf{v}_d\|_2 = 1$, $\mathbf{v}_d^T \mathbf{v}_{d'} = 0$ and p real eigenvalues $\lambda_d \in \mathbb{R}$. This representation of \mathbf{A} is unique. As a result, we have:

$$\mathbf{A} = \mathbf{V} \mathbf{\Lambda} \mathbf{V}^T = \sum_{d=1}^p \lambda_d \mathbf{v}_d \mathbf{v}_d^T$$

Representation of a Vector in the Eigen Basis

- Similarly, if \mathbf{a} is an arbitrary vector, then we can also represent \mathbf{a} using the basis provided by the eigenvectors \mathbf{V} of a real symmetric matrix \mathbf{A} . We obtain:

$$\mathbf{a} = \sum_{d=1}^p \alpha_d \mathbf{v}_d \quad (1)$$

$$\alpha_d = \mathbf{a}^T \mathbf{v}_d \quad (2)$$

Outline

1 Review

2 Linear Algebra

3 PCA


4 Connection to SVD

Principal Component Analysis

- Given a data matrix $\mathbf{X} \in \mathbb{R}^{N \times p}$, the goal of Principal Component Analysis (PCA) is to identify the directions of maximum variance contained in the data.

Principal Component Analysis

- Given a data matrix $\mathbf{X} \in \mathbb{R}^{N \times p}$, the goal of Principal Component Analysis (PCA) is to identify the directions of maximum variance contained in the data.



Sample Variance in a Given Direction

- Let $\mathbf{w} \in \mathbb{R}^p$ such that $\|\mathbf{w}\|_2 = \sqrt{\mathbf{w}^T \mathbf{w}} = 1$.

Sample Variance in a Given Direction

- Let $\mathbf{w} \in \mathbb{R}^p$ such that $\|\mathbf{w}\|_2 = \sqrt{\mathbf{w}^T \mathbf{w}} = 1$.
- The sample estimate of the variance in the direction \mathbf{w} given the data set \mathbf{X} is given by the expression:

$$\frac{1}{N} \sum_{i=1}^N (\mathbf{X}_i \mathbf{w} - \mu)^2 \quad \text{where} \quad \mu = \frac{1}{N} \sum_{i=1}^N \mathbf{X}_i \mathbf{w}$$

Pre-Centering

- Under the assumption that the data are pre-centered so that $\frac{1}{N} \sum_{i=1}^N \mathbf{X}_i = 0$, this expression simplifies to:

$$\frac{1}{N} \sum_{i=1}^N (\mathbf{X}_i \mathbf{w})^2 = (\mathbf{X} \mathbf{w})^T (\mathbf{X} \mathbf{w}) = \mathbf{w}^T \mathbf{X}^T \mathbf{X} \mathbf{w}$$

The Direction of Maximum Variance

- Suppose we want to identify the direction \mathbf{w}_1 of maximum variance given the data matrix \mathbf{X} . We can formulate this optimization problem as follows:

The Direction of Maximum Variance

- Suppose we want to identify the direction \mathbf{w}_1 of maximum variance given the data matrix \mathbf{X} . We can formulate this optimization problem as follows:

$$\mathbf{w}_1 = \max_{\mathbf{w}} \mathbf{w}^T \mathbf{X}^T \mathbf{X} \mathbf{w} \dots \text{st } \|\mathbf{w}\|_2 = 1$$

The Direction of Maximum Variance

- Suppose we want to identify the direction \mathbf{w}_1 of maximum variance given the data matrix \mathbf{X} . We can formulate this optimization problem as follows:

$$\mathbf{w}_1 = \max_{\mathbf{w}} \mathbf{w}^T \mathbf{X}^T \mathbf{X} \mathbf{w} \dots \text{st } \|\mathbf{w}\|_2 = 1$$

- How can we solve this problem?

The Direction of Maximum Variance

- Let $\Sigma = \mathbf{X}^T \mathbf{X}$.

The Direction of Maximum Variance

- Let $\Sigma = \mathbf{X}^T \mathbf{X}$.
- Σ is real and symmetric, so it admits an eigendecomposition of the form:

$$\Sigma = \sum_{d=1}^p \sigma_d \mathbf{V}_d \mathbf{V}_d^T$$

The Direction of Maximum Variance

- Let $\Sigma = \mathbf{X}^T \mathbf{X}$.
- Σ is real and symmetric, so it admits an eigendecomposition of the form:

$$\Sigma = \sum_{d=1}^p \sigma_d \mathbf{V}_d \mathbf{V}_d^T$$

- $\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_p \geq 0$ are the eigenvalues of Σ .

The Direction of Maximum Variance

- Let $\Sigma = \mathbf{X}^T \mathbf{X}$.
- Σ is real and symmetric, so it admits an eigendecomposition of the form:

$$\Sigma = \sum_{d=1}^p \sigma_d \mathbf{V}_d \mathbf{V}_d^T$$

- $\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_p \geq 0$ are the eigenvalues of Σ .
- $\mathbf{V}_d \in \mathbb{R}^D$ are the eigenvectors of Σ . They satisfy:

$$\|\mathbf{V}_d\|_2 = \sqrt{\mathbf{V}_d^T \mathbf{V}_d} = 1 \dots \text{for all } d$$

$$\mathbf{V}_d^T \mathbf{V}_{d'} = 0 \dots \text{for all } d \neq d'$$

The Direction of Maximum Variance

- Using this result, we can write the optimization problem as:

$$\max_{\mathbf{w}} \mathbf{w}^T \mathbf{X}^T \mathbf{X} \mathbf{w} \dots \text{st } \|\mathbf{w}\|_2 = 1$$

The Direction of Maximum Variance

- Using this result, we can write the optimization problem as:

$$\max_{\mathbf{w}} \mathbf{w}^T \mathbf{X}^T \mathbf{X} \mathbf{w} \dots \text{st } \|\mathbf{w}\|_2 = 1$$

$$\max_{\mathbf{w}} \mathbf{w}^T \left(\sum_{d=1}^p \sigma_d \mathbf{V}_d \mathbf{V}_d^T \right) \mathbf{w} \dots \text{st } \|\mathbf{w}\|_2 = 1$$

The Direction of Maximum Variance

- Using this result, we can write the optimization problem as:

$$\max_{\mathbf{w}} \mathbf{w}^T \mathbf{X}^T \mathbf{X} \mathbf{w} \dots \text{st } \|\mathbf{w}\|_2 = 1$$

$$\max_{\mathbf{w}} \mathbf{w}^T \left(\sum_{d=1}^p \sigma_d \mathbf{V}_d \mathbf{V}_d^T \right) \mathbf{w} \dots \text{st } \|\mathbf{w}\|_2 = 1$$

$$\max_{\mathbf{w}} \sum_{d=1}^p \sigma_d (\mathbf{w}^T \mathbf{V}_d)^2 \dots \text{st } \|\mathbf{w}\|_2 = 1$$

The Direction of Maximum Variance

- \mathbf{w} can also be expressed in the orthonormal basis $\mathbf{V}_1, \dots, \mathbf{V}_p$ by letting $\mathbf{w} = \sum_{d=1}^p \omega_d \mathbf{V}_d$.

The Direction of Maximum Variance

- \mathbf{w} can also be expressed in the orthonormal basis $\mathbf{V}_1, \dots, \mathbf{V}_p$ by letting $\mathbf{w} = \sum_{d=1}^p \omega_d \mathbf{V}_d$.
- The constraint that $\|\mathbf{w}\|_2 = 1$ becomes $\sqrt{\sum_{d=1}^p \omega_d^2} = 1$.

The Direction of Maximum Variance

- \mathbf{w} can also be expressed in the orthonormal basis $\mathbf{V}_1, \dots, \mathbf{V}_p$ by letting $\mathbf{w} = \sum_{d=1}^p \omega_d \mathbf{V}_d$.
- The constraint that $\|\mathbf{w}\|_2 = 1$ becomes $\sqrt{\sum_{d=1}^p \omega_d^2} = 1$.
- This means $\sum_{d=1}^p \omega_d^2 = 1$ and $\omega_d^2 > 0$, so the ω_d^2 values act like a discrete probability distribution.

The Direction of Maximum Variance

- Plugging this back into the objective function, we have:

$$\max_{\mathbf{w}} \sum_{d=1}^p \sigma_d(\mathbf{w}^T \mathbf{V}_d)^2 \dots \text{st } \|\mathbf{w}\|_2 = 1$$

The Direction of Maximum Variance

- Plugging this back into the objective function, we have:

$$\max_{\mathbf{w}} \sum_{d=1}^p \sigma_d (\mathbf{w}^T \mathbf{V}_d)^2 \dots \text{st } \|\mathbf{w}\|_2 = 1$$

$$\max_{\omega} \sum_{d=1}^p \sigma_d \left(\sum_{d'=1}^p \omega_{d'} \mathbf{V}_{d'}^T \mathbf{V}_d \right)^2 \dots \text{st } \sum_{d=1}^p \omega_d^2 = 1$$

The Direction of Maximum Variance

- Plugging this back into the objective function, we have:

$$\max_{\mathbf{w}} \sum_{d=1}^p \sigma_d (\mathbf{w}^T \mathbf{V}_d)^2 \dots \text{st } \|\mathbf{w}\|_2 = 1$$

$$\max_{\omega} \sum_{d=1}^p \sigma_d \left(\sum_{d'=1}^p \omega_{d'} \mathbf{V}_{d'}^T \mathbf{V}_d \right)^2 \dots \text{st } \sum_{d=1}^p \omega_d^2 = 1$$

$$\max_{\omega} \sum_{d=1}^p \sigma_d \omega_d^2 \dots \text{st } \sum_{d=1}^p \omega_d^2 = 1$$

The Direction of Maximum Variance

- At this point, the solution is clear.

The Direction of Maximum Variance

- At this point, the solution is clear.
- To maximize the variance, we need to set $\omega_1 = 1$ and set $\omega_d = 0$ otherwise.

The Direction of Maximum Variance

- At this point, the solution is clear.
- To maximize the variance, we need to set $\omega_1 = 1$ and set $\omega_d = 0$ otherwise. This put's all the weight on the maximum eigenvalue of Σ , which is σ_1 by assumption.

The Direction of Maximum Variance

- At this point, the solution is clear.
- To maximize the variance, we need to set $\omega_1 = 1$ and set $\omega_d = 0$ otherwise. This put's all the weight on the maximum eigenvalue of Σ , which is σ_1 by assumption.
- Working our way back to \mathbf{w}_1 , we put all our weight on the maximum eigenvalue, so $\mathbf{w} = \sum_{d=1}^p \omega_d \mathbf{V}_d = \mathbf{V}_1$.

The Direction of Maximum Variance

- At this point, the solution is clear.
- To maximize the variance, we need to set $\omega_1 = 1$ and set $\omega_d = 0$ otherwise. This put's all the weight on the maximum eigenvalue of Σ , which is σ_1 by assumption.
- Working our way back to \mathbf{w}_1 , we put all our weight on the maximum eigenvalue, so $\mathbf{w} = \sum_{d=1}^p \omega_d \mathbf{V}_d = \mathbf{V}_1$.
- **This shows that the maximum variance direction given a data matrix \mathbf{X} is the eigenvector of $\mathbf{X}^T \mathbf{X}$ with the largest eigenvalue.**

q-Largest Directions of Variance

- Suppose instead of just the direction of maximum variance, we want the q largest directions of variance that are all mutually orthogonal.

q-Largest Directions of Variance

- Suppose instead of just the direction of maximum variance, we want the q largest directions of variance that are all mutually orthogonal.
- Finding the second-largest direction of variance corresponds to solving the problem:

$$\mathbf{w}_2 = \max_{\mathbf{w}} \sum_{d=1}^p \sigma_d (\mathbf{w}^T \mathbf{V}_d)^2 \dots \text{ st } \|\mathbf{w}\|_2 = 1 \text{ and } \mathbf{w}^T \mathbf{w}_1 = 0$$

q-Largest Directions of Variance

- Suppose instead of just the direction of maximum variance, we want the q largest directions of variance that are all mutually orthogonal.
- Finding the second-largest direction of variance corresponds to solving the problem:

$$\mathbf{w}_2 = \max_{\mathbf{w}} \sum_{d=1}^p \sigma_d (\mathbf{w}^T \mathbf{V}_d)^2 \dots \text{ st } \|\mathbf{w}\|_2 = 1 \text{ and } \mathbf{w}^T \mathbf{w}_1 = 0$$

- It's easy to see that this is going to be the eigenvector corresponding to the second largest eigenvalue.

q-Largest Directions of Variance

- Suppose instead of just the direction of maximum variance, we want the q largest directions of variance that are all mutually orthogonal.
- Finding the second-largest direction of variance corresponds to solving the problem:

$$\mathbf{w}_2 = \max_{\mathbf{w}} \sum_{d=1}^p \sigma_d (\mathbf{w}^T \mathbf{V}_d)^2 \dots \text{ st } \|\mathbf{w}\|_2 = 1 \text{ and } \mathbf{w}^T \mathbf{w}_1 = 0$$

- It's easy to see that this is going to be the eigenvector corresponding to the second largest eigenvalue.
- **In general, the top q directions of variance $\mathbf{w}_1, \dots, \mathbf{w}_q$ are given by the q eigenvectors corresponding to the q largest eigenvalues of $\mathbf{X}^T \mathbf{X}$.**

Dimensionality Reduction with PCA

- 1 Given centered data matrix $\mathbf{X} \in \mathbb{R}^{N \times p}$, compute unscaled sample covariance matrix $\Sigma = \mathbf{X}^T \mathbf{X}$.

Dimensionality Reduction with PCA

- 1 Given centered data matrix $\mathbf{X} \in \mathbb{R}^{N \times p}$, compute unscaled sample covariance matrix $\Sigma = \mathbf{X}^T \mathbf{X}$.
- 2 Compute the q leading eigenvectors w_1, \dots, w_q of Σ where $\mathbf{w}_k \in \mathbb{R}^p$.

Dimensionality Reduction with PCA

- 1 Given centered data matrix $\mathbf{X} \in \mathbb{R}^{N \times p}$, compute unscaled sample covariance matrix $\Sigma = \mathbf{X}^T \mathbf{X}$.
- 2 Compute the q leading eigenvectors w_1, \dots, w_q of Σ where $\mathbf{w}_k \in \mathbb{R}^p$.
- 3 Stack the eigenvectors together into a $p \times q$ matrix \mathbf{W} where each column k of \mathbf{W} corresponds to \mathbf{w}_k .

Dimensionality Reduction with PCA

- 1 Given centered data matrix $\mathbf{X} \in \mathbb{R}^{N \times p}$, compute unscaled sample covariance matrix $\Sigma = \mathbf{X}^T \mathbf{X}$.
- 2 Compute the q leading eigenvectors w_1, \dots, w_q of Σ where $\mathbf{w}_k \in \mathbb{R}^p$.
- 3 Stack the eigenvectors together into a $p \times q$ matrix \mathbf{W} where each column k of \mathbf{W} corresponds to \mathbf{w}_k .
- 4 Project the matrix \mathbf{X} into the rank- q sub-space of maximum variance by computing the matrix product $\mathbf{Z} = \mathbf{XW}$.

Dimensionality Reduction with PCA

- 1 Given centered data matrix $\mathbf{X} \in \mathbb{R}^{N \times p}$, compute unscaled sample covariance matrix $\Sigma = \mathbf{X}^T \mathbf{X}$.
- 2 Compute the q leading eigenvectors w_1, \dots, w_q of Σ where $\mathbf{w}_k \in \mathbb{R}^p$.
- 3 Stack the eigenvectors together into a $p \times q$ matrix \mathbf{W} where each column k of \mathbf{W} corresponds to \mathbf{w}_k .
- 4 Project the matrix \mathbf{X} into the rank- q sub-space of maximum variance by computing the matrix product $\mathbf{Z} = \mathbf{X}\mathbf{W}$.
- 5 To reconstruct \mathbf{X} given \mathbf{Z} and \mathbf{W} , we use $\hat{\mathbf{X}} = \mathbf{Z}\mathbf{W}^T$.

Outline

1 Review

2 Linear Algebra

3 PCA

4 Connection to SVD

Connection to SVD

- Last class we saw that the minimum Frobenius norm linear dimensionality reduction problem could be solved using the rank- q SVD of \mathbf{X} :

$$\|\mathbf{X} - \mathbf{U}\mathbf{D}_q\mathbf{V}_q^T\|_F^2$$

where the matrix product $\mathbf{Z} = \mathbf{U}\mathbf{D}_q$ gives the optimal rank- q representation of \mathbf{X} with respect to Frobenius norm minimization.

Connection to SVD

- If we let $q = p$ then $\mathbf{X} = \mathbf{UDV}^T$ and $\mathbf{X}^T\mathbf{X} = \mathbf{VDU}^T\mathbf{UDV}^T$.

Connection to SVD

- If we let $q = p$ then $\mathbf{X} = \mathbf{UDV}^T$ and $\mathbf{X}^T\mathbf{X} = \mathbf{VDU}^T\mathbf{UDV}^T$.
- Due to orthogonality of U this gives: $\mathbf{X}^T\mathbf{X} = \mathbf{VD}^2\mathbf{V}^T$.

Connection to SVD

- If we let $q = p$ then $\mathbf{X} = \mathbf{UDV}^T$ and $\mathbf{X}^T\mathbf{X} = \mathbf{VDU}^T\mathbf{UDV}^T$.
- Due to orthogonality of U this gives: $\mathbf{X}^T\mathbf{X} = \mathbf{VD}^2\mathbf{V}^T$.
- This means that the right singular vectors of \mathbf{X} are exactly the eigenvectors of $\mathbf{X}^T\mathbf{X}$, so SVD's \mathbf{V} and PCA's \mathbf{W} are identical (assuming \mathbf{X} is centered).

Connection to SVD

- If we let $q = p$ then $\mathbf{X} = \mathbf{UDV}^T$ and $\mathbf{X}^T\mathbf{X} = \mathbf{VDU}^T\mathbf{UDV}^T$.
- Due to orthogonality of U this gives: $\mathbf{X}^T\mathbf{X} = \mathbf{VD}^2\mathbf{V}^T$.
- This means that the right singular vectors of \mathbf{X} are exactly the eigenvectors of $\mathbf{X}^T\mathbf{X}$, so SVD's \mathbf{V} and PCA's \mathbf{W} are identical (assuming \mathbf{X} is centered).
- We can also see that the eigenvalues of $\mathbf{X}^T\mathbf{X}$ are the squares of the diagonal elements of \mathbf{D} .

Connection to SVD

- If we let $q = p$ then $\mathbf{X} = \mathbf{UDV}^T$ and $\mathbf{X}^T\mathbf{X} = \mathbf{VDU}^T\mathbf{UDV}^T$.
- Due to orthogonality of U this gives: $\mathbf{X}^T\mathbf{X} = \mathbf{VD}^2\mathbf{V}^T$.
- This means that the right singular vectors of \mathbf{X} are exactly the eigenvectors of $\mathbf{X}^T\mathbf{X}$, so SVD's \mathbf{V} and PCA's \mathbf{W} are identical (assuming \mathbf{X} is centered).
- We can also see that the eigenvalues of $\mathbf{X}^T\mathbf{X}$ are the squares of the diagonal elements of \mathbf{D} .
- This means that the q largest singular values and q largest eigenvalues correspond to the same q basis vectors.

Connection to SVD

- According to PCA, the projection operation is $\mathbf{Z} = \mathbf{XW}$.

Connection to SVD

- According to PCA, the projection operation is $\mathbf{Z} = \mathbf{XW}$.
- Using $\mathbf{X} = \mathbf{UDV}^T$ and $\mathbf{V} = \mathbf{W}$ we have:

$$\mathbf{Z} = \mathbf{XW} = (\mathbf{UDV}^T)(\mathbf{V}) = \mathbf{UD}$$

Connection to SVD

- According to PCA, the projection operation is $\mathbf{Z} = \mathbf{XW}$.
- Using $\mathbf{X} = \mathbf{UDV}^T$ and $\mathbf{V} = \mathbf{W}$ we have:

$$\mathbf{Z} = \mathbf{XW} = (\mathbf{UDV}^T)(\mathbf{V}) = \mathbf{UD}$$

- Finally, note that if the decompositions are based only on the q leading basis vectors, which are identical under both PCA and SVD, the projections $\mathbf{Z} = \mathbf{XW}$ and $\mathbf{Z} = \mathbf{UD}$ will still be identical.

Connection to SVD

- These manipulations show that PCA on $\mathbf{X}^T \mathbf{X}$ and SVD on \mathbf{X} identify exactly the same sub-space and result in exactly the same projection of the data into that sub-space.

Connection to SVD

- These manipulations show that PCA on $\mathbf{X}^T \mathbf{X}$ and SVD on \mathbf{X} identify exactly the same sub-space and result in exactly the same projection of the data into that sub-space.
- As a result, generic linear dimensionality reduction simultaneously minimizes the Frobenius norm of the reconstruction error of \mathbf{X} and maximizes the retained variance in the learned sub-space.

Connection to SVD

- These manipulations show that PCA on $\mathbf{X}^T\mathbf{X}$ and SVD on \mathbf{X} identify exactly the same sub-space and result in exactly the same projection of the data into that sub-space.
- As a result, generic linear dimensionality reduction simultaneously minimizes the Frobenius norm of the reconstruction error of \mathbf{X} and maximizes the retained variance in the learned sub-space.
- Both SVD and PCA provide the same refinement of generic linear dimensionality reduction: an orthogonal basis for exactly the same optimal linear subspace.

Demo

■ Demo