

Quantum Mechanics Technology Processing and Conclusions

Jiete XUE

October 2, 2025 - November 25, 2025

Contents

1	Quantization	1
2	Heisenberg's Breakthrough	1
3	De Broglie's & Schrödinger's Equation	2
4	Dirac's Notation	2
4.1	Bra & Ket	2
4.2	Operator	3
5	Commutator and Anticommutator	5
6	Compatible Observables	5
7	Uncertainty Relation	6
8	Position and Momentum	6
8.1	Position & Wave Function	7
8.2	Translation	7
8.3	Momentum	8
8.4	Momentum Operator in the Position Basis	8
8.5	Momentum-Space Wave Function	8
8.6	Dirac's Corresponding Rule	9
9	Schrödinger Equation	9
10	Probability Density and Probability Current	11
11	Exponential Operators and Baker-Hausdorff Lemma	12
12	Harmonic Oscillator	13
12.1	Method of Analysis (Hermite polynomials)	14
12.2	Method of Algebra	15
12.3	Further Discussion of 1D Case: Coherent State	16
12.4	High Dimensional Case	17
13	Gauge Transformations in Electromagnetism	17
13.1	Gauge Transformations	17
13.2	Landau Levels and Quantum Hall Effect	19
13.2.1	Landau Gauge	19
13.2.2	Symmetrical Gauge	20
14	Angular Momentum	21
14.1	Infinitesimal Rotations in Quantum Mechanics	21
14.2	Orbital Angular Momentum	22
14.3	Eigen Value	23

CONTENTS

14.4	Representations of the Rotation Operator	25
14.5	Spherical Harmonics	26
14.6	Schwinger's Oscillator Model of Angular Momentum	27
14.7	Addition of Angular Momenta	27
14.7.1	Clebsch-Gordan Coefficients	28
15	Discrete Symmetry	29
15.1	Wigner Theorem	29
15.2	Parity Transformation	29
15.3	Time-Reversal Symmetry	31
16	Approximation Methods	31
16.1	Time-independent Perturbation Theory: Non-degenerate Case	31

1 Quantization

$$\oint p dq = nh + C, \quad n = 1, 2, 3, \dots \quad (1.1)$$

This is not always right, but it's a good approximation.

2 Heisenberg's Breakthrough

Heisenberg use matrices to represent quantum states.¹

$$x_{n,m}(t) = x_{n,m} e^{i\omega_{n,m}t}. \quad (2.1)$$

With the **Ritz combination law**²

$$\omega_{n,m} = \omega_{n,k} + \omega_{k,m}, \quad (2.2)$$

we can calculate x^2 like

$$(x^2)_{n,m} e^{i\omega_{n,m}t} = \sum_k x_{n,k}(t) x_{k,m}(t) = \sum_k x_{n,k} x_{k,m} e^{i\omega_{n,m}t}. \quad (2.3)$$

Problem 2.1 (Harmonic Oscillator)

By solving this example, we can get how to deal with the problems in quantum situations. The equation of the harmonic oscillator in the classic mechanics is:

$$\ddot{x} + \omega_0^2 x = 0. \quad (2.4)$$

We can deduce that

$$(\omega_0^2 - \omega_{n,m}^2) x_{n,m}(t) = 0. \quad (2.5)$$

Their's a strange assumption, we may treat³ $\omega_{n,n+1}$ as ω_0 , then only for those $x_{n,n\pm 1}$ can not be zero. Then we can use (1.1) to obtain other information. (Use t as parameter to do the integration.)

Only when $n = m$, the integration of

$$-m\omega_{n,k}\omega_{k,m}x_{n,k}x_{k,m}e^{i\omega_{n,m}t} \quad (2.6)$$

in a period can not be zero.⁴ Then we obtain ⁵

$$\frac{\hbar}{2m} = \omega_0 \left[|x_{n+1,n}|^2 - |x_{n,n-1}|^2 \right]. \quad (2.7)$$

¹Here use the similar notation $x(t)$ and x , but they represent different things. In some cases, we will ignore (t) but still represent $x(t)$ for convenience.

²You can imagine a electron moving from state m to state n passing state k .

³Their must exists a $\omega_{n,m} = \omega_0$, or $x_{n,m} = 0$, or the physics phenomenon can not be detected.

⁴ $\omega_{n,n} = 0$.

⁵Here I used f-sum rule exactly. But I think there's some details need to check, about where the $\frac{1}{2}$ comes from. It appear as C in quantization condition.

3 De Broglie's & Schrödinger's Equation

De Broglie use

$$\frac{d\omega}{dk} = v_g = \frac{pc^2}{E} \quad (3.1)$$

and

$$E^2 = p^2 c^2 + m_0^2 c^4 \quad (3.2)$$

with

$$E = \hbar\omega \quad (3.3)$$

to derive that

$$p = \hbar k. \quad (3.4)$$

It is wave-like. Debye told to Schrödinger that: If it is a wave, then you can try to get a wave equation to describe it.

If we consider a plane wave, the it take the form of

$$\psi \sim e^{\frac{i}{\hbar}(px - Et)}. \quad (3.5)$$

The exponent is similar to S : the action. So we can let $\psi = e^{iS/\hbar}$, and try to explore what we can get after plugging it into the Hamilton equation.

$$\frac{\hbar^2}{2m} (\nabla\psi)^2 + (E - V(r)) \psi^2 = 0. \quad (3.6)$$

We want a linear equation. Then we let ∇ act on it, and suppose $V(r)$ is smooth at the \hbar scale. Then we get

$$\left(-\frac{\hbar^2}{2m} \nabla^2 + V(r) \right) \psi = E\psi. \quad (3.7)$$

Born gave ψ a physics meaning: The probability density for the particle that appear at position x .

Problem 3.1 (The relation between image of ψ and E)

We know that, the second order derivative can represent the curvature. So with the energy growing, the image of the wave function will twist and cross the average line more and more.

4 Dirac's Notation

4.1 Bra & Ket

P. A. M Dirac divided the word “bracket” into two parts, one is the bra, and the other is the ket. We call left vector as bra, and right vector as ket. We can denote¹ the state ψ as $|\psi\rangle$, with its dual form $\langle\psi|$. Each of these vectors contains

¹Clear and simple, its my favorite.

a complete information about the state. We have to clarify the computation law about the bras and kets.

$$a|\alpha\rangle + b|\beta\rangle \xrightarrow{DC} a^* \langle\alpha| + b^* \langle\beta|, \quad \forall a, b \in \mathbb{C}. \quad (4.1)$$

Inner product:

$$\langle\alpha|\beta\rangle := \langle\alpha| |\beta\rangle \quad (4.2)$$

is a complex number. Here we postulate two fundamental properties of inner products:

$$\langle\alpha|\beta\rangle = \langle\beta|\alpha\rangle^* \quad (4.3)$$

and

$$\langle\alpha|\alpha\rangle \geq 0. \quad (4.4)$$

The property of the second one is because we can take $|\beta\rangle = |\alpha\rangle$ in (4.3), then we can obtain $\langle\alpha|\alpha\rangle$ is a real number.

In fact, all the states forms a vector space V over \mathbb{C} .

4.2 Operator

All the operator¹ forms a unitary ring K , it can be a K -module structure acting on V . If X is an operator, then we define an operator (**Hermitian adjoint**) X^\dagger as²

$$X|\psi\rangle \xrightarrow{DC} \langle\psi| X^\dagger. \quad (4.5)$$

Notation 4.1

Since $X|\psi\rangle$ is also a ket, we denote it as $|X\psi\rangle$. And its dual form is $\langle X\psi|$.

We say X is **Hermitian** iff.

$$X^\dagger = X. \quad (4.6)$$

Easy to prove:

$$(XY)^\dagger = Y^\dagger X^\dagger, \quad (4.7)$$

$$(|\alpha\rangle \langle\beta|)^\dagger = |\beta\rangle \langle\alpha|, \quad (4.8)$$

$$(4.9)$$

We should tell difference between linear operator and anti-linear operator. They have important difference.

For linear operator A ,

$$A(\alpha|\psi\rangle + \beta|\phi\rangle) = \alpha A|\psi\rangle + \beta A|\phi\rangle. \quad (4.10)$$

¹Some where will use hat to distinguish the operator, but not always for convenience.

²When define the computing law of a Hermitian adjoint operator, we need to ensure any $\langle\psi|\phi\rangle = \langle\phi|\psi\rangle^*$. And if T is an anti-linear operator, we DO NOT define T^\dagger . And we'd better understand a expression with right combination.

$$\langle \beta | A | \alpha \rangle = \langle \alpha | A^\dagger | \beta \rangle^* . \quad (4.11)$$

But for anti-linear operator T ,

$$T(\alpha |\psi\rangle + \beta |\phi\rangle) = \alpha^* T|\psi\rangle + \beta^* T|\phi\rangle . \quad (4.12)$$

If we still want to define T^\dagger , we should satisfies the conditions in the footnote. Compare the form of the linear case, and let all the operator act on kets but not bras.

$$\langle \beta | T\alpha \rangle = \langle \alpha | T^\dagger \beta \rangle . \quad (4.13)$$

Then $|\alpha\rangle$ and $|\beta\rangle$ still anti-linear in this expression.

Theorem 4.1

The eigenvalues of a Hermitian operator A are real; the eigenkets of A corresponding to different eigenvalues are orthogonal.

Proof. We have

$$A |a_1\rangle = a_1 |a_1\rangle , \quad \langle a_2 | A = a_2^* \langle a_2 | . \quad (4.14)$$

Hence,

$$(a_1 - a_2^*) \langle a_2 | a_1 \rangle = 0 . \quad (4.15)$$

Let $a_2 = a_1$, then we obtain

$$a_1 = a_2^* . \quad (4.16)$$

So a_1 is real. If $a_1 \neq a_2$, then

$$\langle a_2 | a_1 \rangle = 0 . \quad (4.17)$$

i.e. the eigenkets are orthogonal. \square

All the observables must be real, so its operator must be Hermitian.¹ If $\{|a\rangle\}$ is complete, then

$$\sum_a |a\rangle \langle a| = 1 . \quad (4.18)$$

Let it act on a state, then we can get the state represent as the bases of $\{|a\rangle\}$. And for a particular a , we can define the **projection operator** as

$$\Lambda_a = |a\rangle \langle a| . \quad (4.19)$$

If we use (4.18) to act on both left and right of an operator X , then we can make it represent in the matrix form.

We write an observable X as

$$X = \sum_{a,b} |a\rangle \langle a| X |b\rangle \langle b| . \quad (4.20)$$

Since $\langle a | X | b \rangle$ is real,

$$X^\dagger = \sum_{a,b} |b\rangle \langle b| X |a\rangle \langle a| = X . \quad (4.21)$$

So X is Hermitian.

¹We will explain it soon.

5 Commutator and Anticommutator

For any two operators A and B , we define their **commutator** as

$$[A, B] = AB - BA. \quad (5.1)$$

And the **anticommutator** is

$$\{A, B\} = AB + BA. \quad (5.2)$$

Its easy to check

$$[AB, C] = A[B, C] + [A, C]B, \quad (5.3)$$

$$[A, BC] = [A, B]C + B[A, C], \quad (5.4)$$

$$[A, [B, C]] + [B, [A, C]] + [C, [A, B]] = 0, \quad (5.5)$$

$$[A, B]^\dagger = [B^\dagger, A^\dagger]. \quad (5.6)$$

$$\{AB, C\} = A\{B, C\} - [A, C]B, \quad (5.7)$$

$$\{A, BC\} = \{A, B\}C - B[A, C]. \quad (5.8)$$

Suppose $f(A), g(B)$ are differentiable functions of A and B , let

$$f(A) = \sum_{n=0}^{\infty} a_n A^n, \quad (5.9)$$

we have

$$[A^n, B] = \sum_{k=0}^{n-1} A^{n-k-1} [A, B] A^k. \quad (5.10)$$

If $[A, B]$ is commutable with A , then,

$$[f(A), B] = f'(A)[A, B]. \quad (5.11)$$

Similarly, if $[A, B]$ is commutable with B , then,

$$[A, g(B)] = g'(B)[A, B]. \quad (5.12)$$

6 Compatible Observables

Observables A and B are defined to be **compatible** when the corresponding operators commute,

$$[A, B] = 0. \quad (6.1)$$

Theorem 6.1

Suppose that A and B are compatible observables, and the eigen-values of A are nondegenerate. Then the matrix elements $\langle a'' | B | a' \rangle$ are all diagonal.

7 Uncertainty Relation

Define ΔA as

$$\Delta A := A - \langle A \rangle, \quad (7.1)$$

where $\langle A \rangle$ is expectation value of A . And the **dispersion** of A is

$$\langle (\Delta A)^2 \rangle = \langle (A^2 - 2A \langle A \rangle + \langle A \rangle^2) \rangle = \langle A^2 \rangle - \langle A \rangle^2. \quad (7.2)$$

Theorem 7.1

For any state, we must have the following inequality:

$$\langle (\Delta A)^2 \rangle \langle (\Delta B)^2 \rangle \geq \frac{1}{4} |\langle [A, B] \rangle|^2. \quad (7.3)$$

Proof. The Schwarz inequality¹:

$$\langle \alpha | \alpha \rangle \langle \beta | \beta \rangle \geq |\langle \alpha | \beta \rangle|^2. \quad (7.4)$$

Take²

$$|\alpha\rangle = \Delta A |\rangle, \quad |\beta\rangle = \Delta B |\rangle,$$

then,

$$\langle (\Delta A)^2 \rangle \langle (\Delta B)^2 \rangle \geq |\langle \Delta A \Delta B \rangle|^2. \quad (7.5)$$

Here we use a trick: Any operator can be divided into Hermitian part and anti-Hermitian part,

$$\Delta A \Delta B = \frac{1}{2} [\Delta A, \Delta B] + \frac{1}{2} \{\Delta A, \Delta B\}. \quad (7.6)$$

Easy to prove that the expect value of Hermitian operator is real, and the anti-Hermitian's is purely imaginary. Thus,

$$|\langle \Delta A \Delta B \rangle|^2 = \frac{1}{4} |\langle [\Delta A, \Delta B] \rangle|^2 + \frac{1}{4} |\langle \{\Delta A, \Delta B\} \rangle|^2. \quad (7.7)$$

One have $[\Delta A, \Delta B] = [A, B]$ and $|\dots|^2 \geq 0$, then we get the result. \square

8 Position and Momentum

I will follow J. J. Sakurai's steps to derive some famous results. As continuous spectra, they have following changes³:

$$\langle a' | a'' \rangle = \delta_{a', a''} \longrightarrow \langle \xi' | \xi'' \rangle = \delta(\xi' - \xi''). \quad (8.1)$$

Where $\delta(\xi' - \xi'')$ is the [Dirac delta function](#).

$$\sum_a |a\rangle \langle a| = 1 \longrightarrow \int d\xi |\xi\rangle \langle \xi| = 1. \quad (8.2)$$

¹Consider $|\alpha\rangle + \lambda |\beta\rangle$ and take $\lambda = -\langle \beta | \alpha \rangle \langle \beta | \beta \rangle$.

²Blank ket $|\rangle$ emphasizes the fact that our consideration may be applied to any ket.

³We use ξ to represent a continuous variable.

8.1 Position & Wave Function

For position, we can measure its position, which means we can get its components at the same time, they are compatible. So

$$[\hat{x}_i, \hat{x}_j] = 0. \quad (8.3)$$

$$|\mathbf{x}\rangle = |x, y, z\rangle. \quad (8.4)$$

$$\hat{x}|\mathbf{x}\rangle = x|\mathbf{x}\rangle, \hat{y}|\mathbf{x}\rangle = y|\mathbf{x}\rangle, \hat{z}|\mathbf{x}\rangle = z|\mathbf{x}\rangle. \quad (8.5)$$

We call

$$\psi_\alpha(x) = \langle x|\alpha\rangle \quad (8.6)$$

the wave function.

8.2 Translation

Consider an operator $\mathcal{J}(\mathbf{dx})$ satisfying

$$\mathcal{J}(\mathbf{dx})|\mathbf{x}\rangle = |\mathbf{x}\rangle. \quad (8.7)$$

Then

$$\mathcal{J}^\dagger(\mathbf{dx})\mathcal{J}(\mathbf{dx}) = 1 \quad (8.8)$$

needs to be guaranteed. We expect the translation does not depend on the trace, which means

$$\mathcal{J}(\mathbf{dx}_1 + \mathbf{dx}_2) = \mathcal{J}(\mathbf{dx}_1) + \mathcal{J}(\mathbf{dx}_2). \quad (8.9)$$

For the third property, suppose we consider a translation in the opposite direction; we expect the opposite-direction translation to be the same as the inverse of the original translation:

$$\mathcal{J}(-\mathbf{dx}) = \mathcal{J}^{-1}(\mathbf{dx}). \quad (8.10)$$

For the fourth property,

$$\lim_{\mathbf{dx} \rightarrow 0} \mathcal{J}(\mathbf{dx}) = 1. \quad (8.11)$$

Consider

$$\mathcal{J}(\mathbf{dx}) = 1 - i\hat{\mathbf{K}} \cdot \mathbf{dx}, \quad (8.12)$$

it let all the properties we want hold, if $\hat{\mathbf{K}}$ is Hermitian. We can check that

$$[\hat{\mathbf{x}}, \mathcal{J}(\mathbf{dx})] = \mathbf{dx}. \quad (8.13)$$

Plug in (8.12), we obtain

$$[\hat{x}_i, \hat{K}_j] = i\delta_{ij}. \quad (8.14)$$

8.3 Momentum

An infinitesimal translation in classical mechanics can be regarded as a canonical transformation,

$$\mathbf{x}_{\text{new}} = \mathbf{x} + d\mathbf{x}, \quad \mathbf{p}_{\text{new}} = \mathbf{p}. \quad (8.15)$$

obtainable from the generating function

$$F(\mathbf{x}, \mathbf{p}) = \mathbf{x} \cdot \mathbf{p}_{\text{new}} + \mathbf{p} \cdot d\mathbf{x}. \quad (8.16)$$

This equation has a striking similarity to the infinitesimal translation operator. Let

$$\mathcal{J}(d\mathbf{x}) = 1 - i\hat{\mathbf{p}} \cdot d\mathbf{x} / \hbar. \quad (8.17)$$

The commutation relation (8.14) now becomes

$$[\hat{x}_i, \hat{p}_j] = i\hbar\delta_{i,j}. \quad (8.18)$$

By (7.3), we obtain the **position-momentum uncertainty relation**:

$$\langle (\Delta x)^2 \rangle \langle (\Delta p)^2 \rangle \geq \frac{\hbar^2}{4} \quad (8.19)$$

As for finite position,

$$\mathcal{J}(\Delta\mathbf{x}) = \lim_{N \rightarrow \infty} \left(1 - \frac{i\hat{\mathbf{p}} \cdot \Delta\mathbf{x}}{N\hbar} \right)^N = \exp \left(-\frac{i\hat{\mathbf{p}} \cdot \Delta\mathbf{x}}{\hbar} \right). \quad (8.20)$$

By computation,

$$[\hat{p}_i, \hat{p}_j] = 0. \quad (8.21)$$

8.4 Momentum Operator in the Position Basis

By computing $\mathcal{J}(d\mathbf{x})|\alpha\rangle$ in the representation where the position eigenkets are used as base kets, and plug in (8.17), we obtain

$$\hat{\mathbf{p}}|\alpha\rangle = \int d\mathbf{x} |\mathbf{x}\rangle (-i\hbar\nabla \langle \mathbf{x}|\alpha\rangle). \quad (8.22)$$

That is also

$$\langle \mathbf{x}|\hat{\mathbf{p}}|\alpha\rangle = -i\hbar\nabla \langle \mathbf{x}|\alpha\rangle. \quad (8.23)$$

8.5 Momentum-Space Wave Function

The notation $\phi_\alpha(p)$ is often used

$$\langle p|\alpha\rangle = \phi_\alpha(p). \quad (8.24)$$

To transform the bases from position to momentum, we need to calculate $\langle x|p\rangle$. One have

$$\langle x|\hat{p}|p\rangle = p \langle x|p\rangle. \quad (8.25)$$

By (8.23),

$$\langle x | \hat{p} | p \rangle = -i\hbar \frac{\partial}{\partial x} \langle x | p \rangle. \quad (8.26)$$

Thus,

$$\langle x | p \rangle = A \exp\left(\frac{ipx}{\hbar}\right). \quad (8.27)$$

Do the derivation will lose a constant, but integration won't.¹

$$\delta(x - x') = \int dp \langle x | p \rangle \langle p | x' \rangle = 2\pi\hbar AA^* \delta(x - x'). \quad (8.28)$$

For convenience, we take A as a real number, then

$$\langle x | p \rangle = \frac{1}{\sqrt{2\pi\hbar}} \exp\left(\frac{ipx}{\hbar}\right). \quad (8.29)$$

For three dimension case,

$$\langle \mathbf{x} | \mathbf{p} \rangle = \left[\frac{1}{(2\pi\hbar)^{\frac{3}{2}}} \right] \exp\left(\frac{i\mathbf{p} \cdot \mathbf{x}}{\hbar}\right). \quad (8.30)$$

8.6 Dirac's Corresponding Rule

Replace classical Poission bracket by commutator as follows²

$$[\quad, \quad]_{\text{classical}} \longrightarrow \frac{[\quad, \quad]}{i\hbar}. \quad (8.31)$$

9 Schrödinger Equation

Let \mathcal{U} be the time-evolution operator, which means

$$|\psi(t)\rangle = \mathcal{U}(t, t_0) |\psi(t_0)\rangle. \quad (9.1)$$

We want it to satisfy the following conditions:

(1) \mathcal{U} should be linear:

$$\mathcal{U}(t, t_0) [c_1 |\psi_1(t_0)\rangle + c_2 |\psi_2(t_0)\rangle] = c_1 \mathcal{U}(t, t_0) |\psi_1(t_0)\rangle + c_2 \mathcal{U}(t, t_0) |\psi_2(t_0)\rangle. \quad (9.2)$$

(2) \mathcal{U} should be unitary (to make a state always normalized):

$$\mathcal{U}^\dagger \mathcal{U} = \mathcal{U} \mathcal{U}^\dagger = 1. \quad (9.3)$$

(3) Composition property:

$$\mathcal{U}(t_2, t_1) \mathcal{U}(t_1, t_0) = \mathcal{U}(t_2, t_0). \quad (9.4)$$

¹Is it an easy integration?

²I think the $i\hbar$ comes from the translation where setting a constant as $i\hbar$.

(4) Continuous and identity:

$$\lim_{dt \rightarrow 0} \mathcal{U}(t_0 + dt, t_0) = 1. \quad (9.5)$$

Similar to the previous section, we can use a Hermitian operator to construct this operator. And by the classical corresponding rule, we have

$$\mathcal{U}(t_0 + dt, t_0) = 1 - \frac{iH dt}{\hbar}. \quad (9.6)$$

Then,

$$\mathcal{U}(t + dt, t_0) = \mathcal{U}(t + dt, t) \mathcal{U}(t, t_0) = \left(1 - \frac{iH dt}{\hbar}\right) \mathcal{U}(t, t_0). \quad (9.7)$$

That is

$$i\hbar \frac{\partial}{\partial t} \mathcal{U}(t, t_0) = H \mathcal{U}(t, t_0). \quad (9.8)$$

This is the **Schrödinger equation for the time-evolution operator**. Apply this operator to a state $|\alpha, t_0; t\rangle$, we obtain

$$i\hbar \frac{\partial}{\partial t} |\alpha, t_0; t\rangle = H |\alpha, t_0; t\rangle. \quad (9.9)$$

In integration form,

$$\mathcal{U}(t, t_0) = 1 - \frac{i}{\hbar} \int_{t_0}^t dt_1 H(t_1) \mathcal{U}(t_1, t_0). \quad (9.10)$$

Then we can induction on n that

$$\mathcal{U}(t, t_0) = \sum_{n=0}^{\infty} \left(\frac{-i}{\hbar}\right)^n \int_{t_0}^t dt_1 \int_{t_0}^{t_1} dt_2 \cdots \int_{t_0}^{t_{n-1}} dt_n H(t_1) H(t_2) \cdots H(t_n). \quad (9.11)$$

Now we want to simplify this expression. The bad thing is H at different time may not commute. So we define a time-ordering operator \mathcal{T} :

$$\mathcal{T}[A(t_1)B(t_2)] = \begin{cases} B(t_2)A(t_1), & \text{if } t_1 < t_2 \\ A(t_1)B(t_2), & \text{if } t_1 \geq t_2 \end{cases}. \quad (9.12)$$

If we consider the integration is dealing on $[t_0, t]^n$, then it can be divided into $n!$ part, each part has an order of a permutation of (t_1, \dots, t_n) , but it should in order of (t_1, \dots, t_n) . Thus,

$$\begin{aligned} & \int_{t_0}^t dt_1 \int_{t_0}^{t_1} dt_2 \cdots \int_{t_0}^{t_{n-1}} dt_n H(t_1) H(t_2) \cdots H(t_n) \\ &= \frac{1}{n!} \int_{t_0}^t dt_1 \int_{t_0}^t dt_2 \cdots \int_{t_0}^t dt_n \mathcal{T}[H(t_1) H(t_2) \cdots H(t_n)] \\ &= \frac{1}{n!} \mathcal{T} \left[\left(\int_{t_0}^t dt' H(t') \right)^n \right]. \end{aligned} \quad (9.13)$$

Therefore,

$$\mathcal{U}(t, t_0) = \mathcal{T} \exp \left[-\frac{i}{\hbar} \int_{t_0}^t dt' H(t') \right]. \quad (9.14)$$

In particular, if $H(t)$ is commutable at any different time, then

$$\mathcal{U}(t, t_0) = \exp \left[-\frac{i}{\hbar} \int_{t_0}^t dt' H(t') \right]. \quad (9.15)$$

Moreover, if H is time-independent, then

$$\mathcal{U}(t, t_0) = \exp \left[-\frac{i}{\hbar} H(t - t_0) \right]. \quad (9.16)$$

10 Probability Density and Probability Current

Let ψ be the wave function, then the probability density is

$$\rho := \psi \psi^*. \quad (10.1)$$

The probability need to be conserved,

$$\frac{\partial \rho}{\partial t} + \nabla \cdot \mathbf{j} = 0. \quad (10.2)$$

Then we can deduce from Schrödinger equation that

$$\mathbf{j} = \left(\frac{\hbar}{m} \right) \text{Im}(\psi^* \nabla \psi). \quad (10.3)$$

If we write ψ into the form of

$$\psi = \sqrt{\rho} e^{\frac{iS}{\hbar}}, \quad (10.4)$$

then¹

$$\mathbf{j} = \frac{\rho \nabla S}{m}. \quad (10.5)$$

Taking the classical limit, we can see that S is the action.

$$\int \mathbf{j} d\mathbf{x} = \frac{\langle \mathbf{p} \rangle}{m}. \quad (10.6)$$

It is some thing like velocity.

¹Here is the case with charge.

11 Exponential Operators and Baker-Hausdorff Lemma

If it exists, then we denote¹

$$e^A = \sum_{n=0}^{\infty} \frac{A^n}{n!}. \quad (11.1)$$

Easy to check that if $[A, B] = 0$, then

$$\exp(A) \exp(B) = \exp(A + B). \quad (11.2)$$

Define²

$$O(\lambda) = e^{i\lambda G} A e^{-i\lambda G}, \quad (11.3)$$

where A and G are operators, and λ is a complex parameter. Then,

$$\frac{d}{d\lambda} O(\lambda) = i e^{i\lambda G} [G, A] e^{-i\lambda G} = i [G, O(\lambda)]. \quad (11.4)$$

Hence,

$$O(\lambda) = O(0) + i \int_0^\lambda d\lambda_1 [G, O(\lambda_1)]. \quad (11.5)$$

We have $O(0) = A$. And we replace $O(\lambda_1)$ similarly, also for $\lambda_2, \lambda_3, \dots$, then we obtain,

$$O(\lambda) = A + i \int_0^\lambda d\lambda_1 [G, A] + i^2 \int_0^\lambda d\lambda_1 \int_0^{\lambda_1} d\lambda_2 [G, [G, A]] + \dots \quad (11.6)$$

Easy to check

$$\int_0^\lambda d\lambda_1 \int_0^{\lambda_1} d\lambda_2 \dots \int_0^{\lambda_{n-1}} d\lambda_n = \frac{\lambda^n}{n!}. \quad (11.7)$$

Therefore³,

$$\boxed{e^{i\lambda G} A e^{-i\lambda G} = \sum_{n=0}^{\infty} \frac{(i\lambda)^n}{n!} C_G^{(n)}(A)}, \quad (11.8)$$

where⁴,

$$C_G(A) = [G, A], \quad C_G^{(n)}(A) = [G, C_G^{(n-1)}(A)]. \quad (11.9)$$

Or we can write in the form

$$e^\Omega A e^{-\Omega} = e^{\text{Ad}_\Omega}(A). \quad (11.10)$$

¹Sometimes we denote as $\exp(A)$.

²We write extra i because we often use the Hermitian conjugate, and i will give a negative sign to fit the form if G is Hermitian.

³It is easy to take the wrong signature, remember it is something like derivative and note the first term, like $[G, A]$, then it won't be wrong.

⁴This notation is not formal, but the following notation “Ad” (get from Deepseek) may be better to use in articles.

Let us consider a special case: $[A, B]$ is commutable with A and B . By Baker-Hausdorff lemma, we have

$$e^A e^B e^{-A} = e^B + [A, e^B] + \dots \quad (11.11)$$

By (5.4), we can induct on n that

$$[A, [A, B]^{n-1} e^B] = [A, B]^n e^B. \quad (11.12)$$

So,

$$e^A e^B = e^B e^A e^{[A, B]}. \quad (11.13)$$

We also have another more symmetrical form¹:

Theorem 11.1

$$[A, [A, B]] = [B, [A, B]] = 0 \Rightarrow e^A e^B = e^{A+B} e^{\frac{1}{2}[A, B]}. \quad (11.14)$$

Proof. Let

$$O(\lambda) = e^{\lambda A} e^{\lambda B}, \quad (11.15)$$

then,

$$\frac{dO}{d\lambda} = e^{\lambda A} (A + B) e^{\lambda B}. \quad (11.16)$$

We want to get a differential equation of $O(\lambda)$, so we have to take $(A + B)$ to the left of $\exp(\lambda A)$ ².

$$e^{\lambda A} B e^{\lambda B} = e^{\lambda A} B e^{-\lambda A} e^{\lambda A} e^{\lambda B}. \quad (11.17)$$

By Baker-Hausdorff lemma, we obtain

$$\frac{dO}{d\lambda} = [(A + B) + \lambda[A, B]] O. \quad (11.18)$$

The solution is

$$O = \exp \left[\lambda(A + B) + \frac{1}{2} \lambda^2 [A, B] \right]. \quad (11.19)$$

□

12 Harmonic Oscillator

We discuss the 1D case first. We have the Hamiltonian

$$\hat{H} = \frac{\hat{p}^2}{2m} + \frac{1}{2} m \omega^2 \hat{x}^2. \quad (12.1)$$

¹In fact we do not need any condition, then we can derive a general form, but that is a bit complex and unnecessary.

²Trick: If there is not we want, then create one, like $+1 - 1$, here we times 1.

12.1 Method of Analysis (Hermite polynomials)

We solve the wave function in position representation, then we obtain many information. Let $z = \frac{x}{\sqrt{\frac{\hbar}{m\omega}}}$, then

$$\left[-\frac{1}{2} \frac{d^2}{dz^2} + \frac{1}{2} z^2 \right] \psi(z) = \frac{E}{\hbar\omega} \psi(z). \quad (12.2)$$

A must to satisfy is

$$\lim_{z \rightarrow \infty} \psi(z) = 0. \quad (12.3)$$

So we first analyze the behavior of $\psi(z)$ at $z \rightarrow \pm\infty$. For $\psi'' = z^2\psi$, we try the solution¹ $\psi(z) = e^S(z)$

$$\psi' = S' \psi, \quad \psi'' = (S'' + S'^2) \psi. \quad (12.4)$$

Comparison of two expression yields

$$S'' + S'^2 = z^2. \quad (12.5)$$

The leading order leads to $s' = \pm z$, $s = \pm \frac{1}{2} z^2$. So we can try the solution

$$\psi(z) = e^{-\frac{z^2}{2}} u(z). \quad (12.6)$$

$$\frac{d^2}{dz^2} u(z) - 2z \frac{d}{dz} u(z) - (\lambda - 1) u(z) = 0. \quad (12.7)$$

We consider the series solution, let

$$u(z) = \sum_{k=0}^{+\infty} a_k z^k. \quad (12.8)$$

The we can deduce that

$$\frac{a_{k+2}}{a_k} = \frac{2k+1-\lambda}{(k+1)(k+2)}. \quad (12.9)$$

We have $u(\pm\infty) = 1$, so there must exists an $N \in \mathbb{N}$, for any $k > N$,

$$a_k = 0. \quad (12.10)$$

Therefore,

$$\lambda_n = 2n + 1, \quad n = 0, 1, 2, \dots \quad (12.11)$$

which means

$$E_n = \left(n + \frac{1}{2} \right) \hbar\omega. \quad (12.12)$$

¹This may keep the form.

The solution of (12.7) is Hermite polynomials¹ $H_n(z)$.

$$\psi(x) = e^{-\frac{1}{2}\frac{m\omega x^2}{\hbar}} \sum_{n=0}^{+\infty} \frac{c_n}{n!} H_n \left(x / \sqrt{\frac{\hbar}{m\omega}} \right). \quad (12.13)$$

And for a fixed n , the normalized wavefunction is

$$\psi_n(z) = \frac{1}{\sqrt{2^n n! \sqrt{\pi}}} e^{-\frac{z^2}{2}} H_n(z). \quad (12.14)$$

12.2 Method of Algebra

In usual computing law,

$$H = \frac{p^2}{2m} + \frac{1}{2}m\omega^2 x^2 = \frac{1}{2}m\omega^2 \left(x - \frac{ip}{m\omega} \right) \left(x + \frac{ip}{m\omega} \right). \quad (12.15)$$

Let ²

$$\hat{a}^\dagger = \sqrt{\frac{m\omega}{2\hbar}} \left(\hat{x} - \frac{i\hat{p}}{m\omega} \right), \quad \hat{a} = \sqrt{\frac{m\omega}{2\hbar}} \left(\hat{x} + \frac{i\hat{p}}{m\omega} \right). \quad (12.16)$$

Then,

$$[\hat{a}, \hat{a}^\dagger] = 1. \quad (12.17)$$

Let

$$\hat{N} = \hat{a}^\dagger \hat{a}, \quad (12.18)$$

then,

$$\hat{H} = \hbar\omega \left(\hat{N} + \frac{1}{2} \right). \quad (12.19)$$

So \hat{N} is compatible with \hat{H} . We denote an energy eigenket of \hat{N} by its eigenvalue³ n , so

$$\hat{N} |n\rangle = n |n\rangle, \quad (12.20)$$

$$\hat{H} |n\rangle = \left(n + \frac{1}{2} \right) \hbar\omega |n\rangle. \quad (12.21)$$

To appreciate the physical significance of \hat{a} , \hat{a}^\dagger , and \hat{N} , let us first note that

$$[\hat{N}, \hat{a}] = -\hat{a}, \quad [\hat{N}, \hat{a}^\dagger] = \hat{a}^\dagger. \quad (12.22)$$

As a result, we have

$$\hat{N} \hat{a}^\dagger |n\rangle = [\hat{N}, \hat{a}^\dagger] |n\rangle + \hat{a}^\dagger \hat{N} |n\rangle = (n+1) \hat{a}^\dagger |n\rangle, \quad (12.23)$$

$$\hat{N} \hat{a} |n\rangle = [\hat{N}, \hat{a}] |n\rangle + \hat{a} \hat{N} |n\rangle = (n-1) \hat{a} |n\rangle. \quad (12.24)$$

¹There's some properties about [Hermite polynomial](#).

²If we denote $l = \sqrt{\frac{\hbar}{m\omega}}$, then it will be $a = \frac{1}{\sqrt{2}} \left(\frac{x}{l} + i \frac{pl}{\hbar} \right)$. $\frac{1}{\sqrt{2}}$ seems like to normalize.

³We will later show that n must be a nonnegative integer.

We write

$$\hat{a} |n\rangle = c |n-1\rangle, \quad (12.25)$$

then,

$$\langle n | \hat{N} | n \rangle = \langle n | \hat{a}^\dagger \hat{a} | n \rangle = |c|^2. \quad (12.26)$$

Thus,

$$\hat{a} |n\rangle = \sqrt{n} |n-1\rangle. \quad (12.27)$$

Similarly, it is easy to show that¹

$$\hat{a}^\dagger |n\rangle = \sqrt{n+1} |n+1\rangle. \quad (12.28)$$

From (12.26) we know that $n \geq 0$. If we act \hat{a} on a state for many times, we can conclude that the sequence must terminate with $n = 0$, or it won't stop and n will become not real. Use

$$\hat{x} = \sqrt{\frac{\hbar}{2m\omega}} (\hat{a} + \hat{a}^\dagger), \quad \hat{p} = i\sqrt{\frac{m\hbar\omega}{2}} (-\hat{a} + \hat{a}^\dagger), \quad (12.29)$$

we can get more information. To solve the wave function, we can deduce $\langle x | \hat{a} | 0 \rangle = 0$, which leads to a differential equation for the wave function.

12.3 Further Discussion of 1D Case: Coherent State

We have seen that an energy eigenstate does not behave like the classical oscillator—in the sense of oscillating expectation values for x and p —no matter how large n may be. We may logically ask, How can we construct a superposition of energy eigenstates that most closely imitates the classical oscillator? In wave-function language, we want a wave packet that bounces back and forth without spreading in shape. It turns out that a coherent state defined by the eigenvalue equation for the non-Hermitian annihilation operator a ,

$$a |\lambda\rangle = \lambda |\lambda\rangle,$$

with, in general, a complex eigenvalue λ does the desired job.

—*Modern Quantum Mechanics*

We want it to be related to the states we have known. And the simplest one is $|0\rangle$. Let $|\lambda\rangle = A|0\rangle$. Then what we expect is $aA|0\rangle = \lambda A|0\rangle$. Or in another form:

$$(A^{-1}aA - \lambda)|0\rangle = f(a)|0\rangle,$$

where f is a arbitrary function. We want to make it easy, let us set $f(a) = a$. Then by the [Baker-Hausdorff lemma](#), the following let what we want holds

$$A = e^{\lambda a^\dagger}. \quad (12.30)$$

¹Memory trick: the square root of a larger one.

use (11.13) to do the normalization¹,

$$\langle 0 | e^{\lambda^* a} e^{\lambda a^\dagger} | 0 \rangle = e^{|\lambda|^2}. \quad (12.31)$$

Therefore²,

$$|\lambda\rangle = e^{-|\lambda|^2/2} e^{\lambda a^\dagger} |0\rangle = e^{-|\lambda|^2/2} \sum_{n=0}^{\infty} \frac{\lambda^n}{\sqrt{n!}} |n\rangle. \quad (12.32)$$

We have discussed in 8.3 that the operator $e^{-ipl/\hbar}$ is a displacement operator. Let us show we can obtain a coherent state by applying it on $|0\rangle$.

$$e^{-ip\Delta x/\hbar} |0\rangle = e^{\frac{(a^\dagger - a)\Delta x}{\sqrt{2}i}} |0\rangle \sim e^{\frac{a^\dagger \Delta x}{\sqrt{2}i}} |0\rangle. \quad (12.33)$$

12.4 High Dimensional Case

We use the Einstein summation convention. Hamilton in n -dimensional case can be written as

$$H = \frac{p^i p_i}{2m} + \frac{1}{2} m \omega^2 x^i x_i. \quad (12.34)$$

And the creation and annihilation operators are

$$a_i = \frac{1}{\sqrt{2}} \left(\frac{x_i}{l} + \frac{ip_i l}{\hbar} \right), \quad a_i^\dagger = \frac{1}{\sqrt{2}} \left(\frac{x_i}{l} - \frac{ip_i l}{\hbar} \right). \quad (12.35)$$

$$[a_i, a_j] = 0, \quad [a_i, a_j^\dagger] = \delta_{ij}. \quad (12.36)$$

If we denote $a^i a_i^\dagger$ as N , then we can written H as

$$H = \hbar \omega \left(N + \frac{n}{2} \right). \quad (12.37)$$

Or H_i^i ³ if we denote $H_i = \hbar \omega (a_i^\dagger a_i + \frac{1}{2})$.

It's easy to check that

$$[a_i^\dagger a_j, H] = 0. \quad (12.38)$$

That means the angular momentum is conserved.

13 Gauge Transformations in Electromagnetism

13.1 Gauge Transformations

In Classic mechanics, the Hamiltonian is

$$H = \frac{1}{2m} \left(\mathbf{p} - \frac{q\mathbf{A}}{c} \right)^2 + q\phi. \quad (13.1)$$

¹Expand the series we get $\exp(\lambda a) |0\rangle = |0\rangle$.

²A Poission distribution.

³It's my own notation means $\sum_i H_i$.

Where \mathbf{p} is the canonical momentum, \mathbf{A} is the vector potential, q is the charge and ϕ is the scalar potential. And now the canonical momentum \mathbf{p} is the generator of translation. And kinematical (or mechanical) momentum, denoted by:

$$\mathbf{\Pi} := \mathbf{p} - \frac{q\mathbf{A}}{c}. \quad (13.2)$$

Easy to check that

$$[\Pi_i, \Pi_j] = \frac{i\hbar q}{c} \varepsilon_{ijk} B_k. \quad (13.3)$$

The Hamiltonian:

$$H = \frac{\Pi^2}{2m} + q\phi. \quad (13.4)$$

From (13.2) we can use the substitution

$$\mathbf{p} \longrightarrow \mathbf{p} - \frac{q\mathbf{A}}{c} \quad (13.5)$$

In position representation¹,

$$\nabla \longrightarrow \nabla - \left(\frac{iq}{\hbar c} \right) \mathbf{A}. \quad (13.6)$$

to get the expressions of the electromagnetism case. For example, the probability current now is

$$\mathbf{j} = \frac{\rho}{m} \left(\nabla S - \frac{q\mathbf{A}}{c} \right). \quad (13.7)$$

$$\int \mathbf{j} d\mathbf{x} = \frac{\langle \mathbf{\Pi} \rangle}{m}. \quad (13.8)$$

\mathbf{E} and \mathbf{B} are unchanged under

$$\phi \rightarrow \phi - \frac{1}{c} \frac{\partial \Lambda}{\partial t}, \quad \mathbf{A} \rightarrow \mathbf{A} + \nabla \Lambda. \quad (13.9)$$

We consider the time-independent operator, so in the remaining part of this section, the term *gauge transformation* refers to the case $\frac{\partial \Lambda}{\partial t} = 0$.

Let $|\alpha\rangle, |\tilde{\alpha}\rangle$ be the state ket in the presence of \mathbf{A} and $\tilde{\mathbf{A}}$, where,

$$\tilde{\mathbf{A}} = \mathbf{A} + \nabla \Lambda. \quad (13.10)$$

They must have the same physical situation, at least we need to make sure

$$\langle \alpha | \alpha \rangle = \langle \tilde{\alpha} | \tilde{\alpha} \rangle, \quad \langle \alpha | \mathbf{x} | \alpha \rangle = \langle \tilde{\alpha} | \mathbf{x} | \tilde{\alpha} \rangle, \quad \langle \alpha | \mathbf{\Pi} | \alpha \rangle = \langle \tilde{\alpha} | \tilde{\mathbf{\Pi}} | \tilde{\alpha} \rangle. \quad (13.11)$$

Construct an operator g that relates $|\tilde{\alpha}\rangle$ to $|\alpha\rangle$.

$$|\tilde{\alpha}\rangle = g |\alpha\rangle. \quad (13.12)$$

¹We can use it only when the ∇ comes from \mathbf{p} .

By intuition, g is a function of Λ , and it is a function of \mathbf{x} , so it is commutable with \mathbf{x} , $g^\dagger \mathbf{x} g = \mathbf{x}$. The first and the third conditions give restrictions:

$$g^\dagger g = 1. \quad (13.13)$$

$$g^\dagger \mathbf{p} g = \mathbf{p} + \frac{q \nabla \Lambda}{c}. \quad (13.14)$$

It is easy to find

$$g = \exp \left(\frac{iq\Lambda}{\hbar c} \right) \quad (13.15)$$

satisfies the above conditions. And we have

$$g^\dagger \tilde{\Pi}^n g = \prod_{k=1}^n g^\dagger \tilde{\Pi} g = \Pi^n. \quad (13.16)$$

So, for any function f ,

$$g^\dagger f(\tilde{\Pi}) g = f(\Pi). \quad (13.17)$$

(13.14) and (13.17) tell us, mechanical momentum is gauge invariant, but the canonical momentum is not.

Form the wave function,

$$S \longrightarrow S + \frac{q\Lambda}{c}. \quad (13.18)$$

This is highly satisfactory because we see that the probability flux given by (13.7) is then gauge invariant.

DO NOT FORGET that when using the gauge transformation, not only the states but also the operators will change. Let's check the Schrödinger equation is still hold under gauge transformation.

$$\tilde{H} |\tilde{\alpha}\rangle = g g^\dagger \tilde{H} g |\alpha\rangle = g H |\alpha\rangle = i\hbar \frac{\partial}{\partial t} g |\alpha\rangle = i\hbar \frac{\partial}{\partial t} |\tilde{\alpha}\rangle. \quad (13.19)$$

Since t is only a parameter, it does not affect the commutators, so when $\frac{\partial \Lambda}{\partial t} \neq 0$, (13.17) still holds. **BUT**, $\tilde{H} \neq g^\dagger H g$, since ϕ contributes a term. So our notation is not convenient in the Λ is time-dependent case.

13.2 Landau Levels and Quantum Hall Effect

We use IS of units this subsection.

13.2.1 Landau Gauge

Let's consider a simple magnetic field,

$$\mathbf{B} = -B\hat{\mathbf{z}}. \quad (13.20)$$

We take

$$A_x = By, \quad A_y = A_z = 0. \quad (13.21)$$

We just consider 2D case, the Hamiltonian is

$$H = \frac{(p_x - qBy)^2}{2m} + \frac{p_y^2}{2m} = \frac{1}{2}m\omega_c^2 \left(y - \frac{p_x l_B^2}{\hbar} \right)^2 + \frac{p_y^2}{2m}, \quad (13.22)$$

where,

$$\omega_c = \frac{qB}{m}, \quad l_B = \sqrt{\frac{\hbar}{qB}}. \quad (13.23)$$

Since $\frac{\partial H}{\partial x} = 0$, we suppose

$$\psi_{n,k_x}(x, y) = e^{ik_x x} \psi(y). \quad (13.24)$$

Then,

$$\left[\frac{p_y^2}{2m} + \frac{1}{2}m\omega_c^2 (y - l_B^2 k_x)^2 \right] |\psi\rangle = E_n |\psi\rangle. \quad (13.25)$$

It has the same solution as the 1D harmonic oscillator, but with a displacement.

$$E_n = \hbar\omega_c \left(n + \frac{1}{2} \right). \quad (13.26)$$

13.2.2 Symmetrical Gauge

Let

$$l_B = \sqrt{\frac{\hbar}{eB}}, \quad \omega_c = \frac{eB}{m}, \quad (13.27)$$

$$a = \frac{1}{\sqrt{2}} \frac{l_B}{\hbar} (\Pi_x + i\Pi_y), \quad a^\dagger = \frac{1}{\sqrt{2}} \frac{l_B}{\hbar} (\Pi_x - i\Pi_y). \quad (13.28)$$

Then,

$$[a, a^\dagger] = 1, \quad H = \hbar\omega_c \left(a^\dagger a + \frac{1}{2} \right). \quad (13.29)$$

Then we can derivative (13.26) quickly.

To solve the wave function, we need choose a gauge, here we use symmetrical gauge, $(A_x, A_y) = \frac{B}{2}(-y, x)$. Let $w = x + iy$, $\bar{\partial} = \frac{\partial}{\partial \bar{w}} = \frac{1}{2}(\partial_x + i\partial_y)$. Then

$$\langle \mathbf{x} | a | \psi \rangle = -i\sqrt{2}l_B \left(\bar{\partial} + \frac{w}{4l_B^2} \right) \langle \mathbf{x} | \psi \rangle, \quad \langle \mathbf{x} | a^\dagger | \psi \rangle = i\sqrt{2}l_B \left(\partial - \frac{\bar{w}}{4l_B^2} \right) \langle \mathbf{x} | \psi \rangle. \quad (13.30)$$

The wave function of the ground state satisfies

$$\left(\bar{\partial} + \frac{w}{4l_B^2} \right) \psi_0(w, \bar{w}) = 0. \quad (13.31)$$

The solution is

$$\psi_0(w, \bar{w}) = e^{-\frac{|w|^2}{4l_B^2}} f(w, \bar{w}), \quad (13.32)$$

where f satisfies

$$\bar{\partial} f = 0. \quad (13.33)$$

So, we can write ψ_0 as

$$\psi_0(w, \bar{w}) = f(w) e^{-\frac{|w|^2}{4l_B^2}}. \quad (13.34)$$

Then

$$\psi_n(w, \bar{w}) = e^{-\frac{|w|^2}{4l_B^2}} \left(\partial - \frac{\bar{w}}{2l_B^2} \right)^n f(w). \quad (13.35)$$

An arbitrary $f(w)$ makes the states highly degenerate. Expand f in a series,

$$f(w) = \sum_{m \in \mathbb{N}} c_m w^m, \quad (13.36)$$

and denote

$$\psi_{0,m} \sim w^m e^{-\frac{|w|^2}{4l_B^2}}. \quad (13.37)$$

In fact, $\psi_{0,m}$ is also the eigenstates of $L_z = \hbar(\partial w - \bar{\partial} \bar{w})$, and the eigenvalues are $m\hbar$.

$|\psi_{0,m}|^2$ take the maximum when $|w| = r_m$, where

$$r_m = \sqrt{2m} l_B. \quad (13.38)$$

Easy to note that each state has a area of $2\pi l_B^2$ greater than another.

14 Angular Momentum

14.1 Infinitesimal Rotations in Quantum Mechanics

We want to construct a rotation operator \mathcal{D}^1 . The appropriate infinitesimal operators could be written as

$$U_\varepsilon = 1 - iG\varepsilon \quad (14.1)$$

with a Hermitian operator G . Define the angular-momentum operator J_k in such a way that the operator for an infinitesimal rotation around the k th axis by angle $d\varphi$ can be obtained by letting

$$G \rightarrow \frac{J_k}{\hbar}, \quad \varepsilon \rightarrow d\phi. \quad (14.2)$$

¹The symbol \mathcal{D} stems from the German word Drehung, meaning “rotation.”

With J_k taken to be Hermitian, the infinitesimal-rotation operator is guaranteed to be unitary and reduces to the identity operator in the limit $d\phi \rightarrow 0$. More generally, we have

$$\mathcal{D}(\hat{\mathbf{n}}, d\phi) = 1 - i \left(\frac{\mathbf{J} \cdot \hat{\mathbf{n}}}{\hbar} \right) d\phi \quad (14.3)$$

for a rotation about the direction characterized by a unit vector $\hat{\mathbf{n}}$ by an infinitesimal angle $d\phi$. For instance, if we are interested in a finite rotation about the $\hat{\mathbf{n}}$ -axis by angle ϕ , we consider

$$\mathcal{D}_{\hat{\mathbf{n}}}(\phi) = \lim_{N \rightarrow +\infty} \left[1 - i \left(\frac{\mathbf{J} \cdot \hat{\mathbf{n}}}{\hbar} \right) \frac{\phi}{N} \right]^N = \exp \left[-i \left(\frac{\mathbf{J} \cdot \hat{\mathbf{n}}}{\hbar} \right) \phi \right]. \quad (14.4)$$

Taking $\phi = \varepsilon$, the first three term is

$$\mathcal{D}_{\hat{\mathbf{n}}}(\varepsilon) \approx 1 - i \left(\frac{\mathbf{J} \cdot \hat{\mathbf{n}}}{\hbar} \right) \varepsilon - \frac{1}{2!} \left(\frac{\mathbf{J} \cdot \hat{\mathbf{n}}}{\hbar} \right)^2 \varepsilon^2. \quad (14.5)$$

Let $\hat{\mathbf{n}}$ be $\hat{\mathbf{x}}$ and $\hat{\mathbf{y}}$, by the properties of rotation, \mathcal{D} need to satisfy

$$[\mathcal{D}_x(\varepsilon), \mathcal{D}_y(\varepsilon)] = \mathcal{D}_z(\varepsilon^2) - 1. \quad (14.6)$$

So,

$$[J_x, J_y] = i\hbar J_z. \quad (14.7)$$

In general,

$$[J_i, J_k] = i\hbar \varepsilon_{ijk} J_k. \quad (14.8)$$

By the meaning of rotation, we can also know that for any scalar a and vector v_i ,

$$[J_i, a] = 0, \quad [J_i, v_j] = i\hbar \varepsilon_{ijk} v_k. \quad (14.9)$$

14.2 Orbital Angular Momentum

Orbital angular momentum is defined as

$$\mathbf{L} = \mathbf{x} \times \mathbf{p}. \quad (14.10)$$

Easy to check that \mathbf{L} satisfies (14.7). Expand \mathbf{L}^2 :

$$\begin{aligned} \mathbf{L}^2 &= \sum_{ijklmk} \varepsilon_{ijk} x_i p_j \varepsilon_{lmk} x_l p_m \\ &= \sum_{ijlm} (\delta_{il} \delta_{jm} - \delta_{im} \delta_{jl}) x_i p_j x_l p_m \\ &= \sum_{ijlm} [\delta_{il} \delta_{jm} x_i (x_l p_j - i\hbar \delta_{jl}) p_m - \delta_{im} \delta_{jl} x_i p_j (p_m x_l + i\hbar \delta_{lm})] \\ &= \mathbf{x}^2 \mathbf{p}^2 - i\hbar \mathbf{x} \cdot \mathbf{p} - \sum_{ijlm} \delta_{im} \delta_{jl} [x_i p_m (x_l p_j - i\hbar \delta_{jl}) + i\hbar \delta_{lm} x_i p_j] \\ &= \mathbf{x}^2 \mathbf{p}^2 - (\mathbf{x} \cdot \mathbf{p})^2 + i\hbar \mathbf{x} \cdot \mathbf{p}. \end{aligned} \quad (14.11)$$

We have

$$[L_i, H] = 0, \quad [L^i L_i, H] = 0, \quad [L^i L_i, L_j] = 0. \quad (14.12)$$

So we can get the eigen value of L_z, \mathbf{L}^2, H at the same time.

$$\langle \mathbf{x} | \mathbf{L} | \alpha \rangle = -i\hbar \mathbf{x} \times \nabla \langle \mathbf{x} | \alpha \rangle. \quad (14.13)$$

Under spherical coordinates, the nabla operator can be written as¹

$$\nabla = \frac{\partial}{\partial r} \hat{\mathbf{r}} + \frac{1}{r} \frac{\partial}{\partial \theta} \hat{\theta} + \frac{1}{r \sin \theta} \frac{\partial}{\partial \phi} \hat{\phi}. \quad (14.14)$$

One has,

$$\mathbf{x} = r \hat{\mathbf{r}}, \quad (14.15)$$

so,

$$\mathbf{x} \times \nabla = \frac{\partial}{\partial \theta} \hat{\phi} - \frac{1}{\sin \theta} \frac{\partial}{\partial \phi} \hat{\theta}. \quad (14.16)$$

Then, project on $\hat{\mathbf{x}}, \hat{\mathbf{y}}, \hat{\mathbf{z}}$, we obtain

$$\langle \mathbf{x} | L_z | \alpha \rangle = -i\hbar \frac{\partial}{\partial \phi} \langle \mathbf{x} | \alpha \rangle, \quad (14.17)$$

$$\langle \mathbf{x} | L_x | \alpha \rangle = -i\hbar \left(-\sin \phi \frac{\partial}{\partial \theta} - \cot \theta \cos \phi \frac{\partial}{\partial \phi} \right) \langle \mathbf{x} | \alpha \rangle, \quad (14.18)$$

$$\langle \mathbf{x} | L_y | \alpha \rangle = -i\hbar \left(\cos \phi \frac{\partial}{\partial \theta} - \cot \theta \sin \phi \frac{\partial}{\partial \phi} \right) \langle \mathbf{x} | \alpha \rangle, \quad (14.19)$$

$$\langle \mathbf{x} | L_{\pm} | \alpha \rangle = -i\hbar e^{\pm i\phi} \left(\pm i \frac{\partial}{\partial \theta} - \cot \theta \frac{\partial}{\partial \phi} \right) \langle \mathbf{x} | \alpha \rangle. \quad (14.20)$$

Where, $L_{\pm} = L_x \pm iL_y$. (14.11) gives the fact that

$$\langle \mathbf{x} | \mathbf{L}^2 | \alpha \rangle = -r^2 \hbar^2 \nabla_{\theta, \phi}^2 \langle \mathbf{x} | \alpha \rangle. \quad (14.21)$$

$\nabla_{\theta, \phi}$ is the laplacian in spherical coordinates, but without the term² of r .

14.3 Eigen Value

We now look for the simultaneous eigenkets of \mathbf{J}^2 and J_z . We denote the eigenvalues of \mathbf{J}^2 and J_z by a and b , respectively:

$$\mathbf{J}^2 |a, b\rangle = a |a, b\rangle, \quad J_z |a, b\rangle = b |a, b\rangle. \quad (14.22)$$

Consider

$$J_{\pm} := J_x \pm iJ_y, \quad (14.23)$$

¹ \mathbf{p} represent a displacement, it always acts like a gradient.

²Term of r is $\frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial}{\partial r} \cdot \right)$. This is equivalent to $\frac{1}{r} \frac{\partial^2}{\partial r^2} (r \cdot)$

then,

$$J_- = J_+^\dagger, \quad \mathbf{J}^2 = J_z^2 + \frac{1}{2}\{J_+, J_-\}. \quad J_- J_+ = \mathbf{J}^2 - J_z^2 - \hbar J_z, \quad (14.24)$$

$$[J_+, J_-] = 2\hbar J_z, \quad [J_z, J_\pm] = \pm\hbar J_\pm, \quad [\mathbf{J}^2, J_\pm] = 0. \quad (14.25)$$

$$J_z (J_\pm |a, b\rangle) = (b \pm \hbar) (J_\pm |a, b\rangle). \quad (14.26)$$

So, we call J_\pm the **ladder operators**.

$$\mathbf{J}^2 (J_\pm |a, b\rangle) = J_\pm \mathbf{J}^2 |a, b\rangle = a (J_\pm |a, b\rangle). \quad (14.27)$$

So,

$$J_\pm |a, b\rangle = c_\pm |a, b \pm \hbar\rangle. \quad (14.28)$$

Then there must exist b_{\max} and b_{\min} , such that¹

$$a = b_{\max} (b_{\max} + \hbar) = b_{\min} (b_{\min} - \hbar). \quad (14.29)$$

We can infer that

$$-b_{\max} = b_{\min}, \quad (14.30)$$

with b_{\max} is positive, and that the allowed values of b lie within

$$b_{\min} \leq b \leq b_{\max}. \quad (14.31)$$

$$b_{\max} = \frac{n\hbar}{2} \equiv j\hbar, \quad (14.32)$$

where j is either an integer or a half-integer², then

$$a = \hbar^2 j(j+1). \quad (14.33)$$

Let us also define m such that

$$b \equiv m\hbar. \quad (14.34)$$

The allowed m -values for a given j are

$$m = \underbrace{-j, -j+1, \dots, j-1, j}_{2j+1 \text{ states}}. \quad (14.35)$$

Instead of $|a, b\rangle$, it is more convenient to denote a simultaneous eigenket of \mathbf{J}^2 and J_z by $|j, m\rangle$.

$$\mathbf{J}^2 |j, m\rangle = \hbar^2 j(j+1) |j, m\rangle, \quad J_z |j, m\rangle = m\hbar |j, m\rangle. \quad (14.36)$$

Then we can deduce that

$$J_\pm |j, m\rangle = \sqrt{(j \mp m)(j \pm m + 1)} \hbar |j, m \pm 1\rangle, \quad (14.37)$$

Important result: If $[K, \mathbf{J}] = 0$, then $K |j, m\rangle$ is also an eigenket of \mathbf{J}^2 and J_z . So,

$$\langle j', m' | K |j, m\rangle = \delta_{j'j} \delta_{m'm} f(j, m). \quad (14.38)$$

By using \mathbf{J}_\pm , we can prove that

$$\langle j, m+1 | K |j, m+1\rangle = \langle j, m | K |j, m\rangle. \quad (14.39)$$

¹By (14.24)

²If it represent the orbital angular momentum, we will denote j as l , and l can only take integer number, since the following will show the factor $e^{il\phi}$, and it must admit 2π as a period.

14.4 Representations of the Rotation Operator

If an arbitrary y -axis is rotated around the arbitrary z -axis by α to become y' , then for any angle β , then¹

$$R_{y'}(\beta) = R_z(\alpha)R_y(\beta)R_z^{-1}(\alpha). \quad (14.40)$$

If we use the Euler Rotations², then

$$R(\alpha, \beta, \gamma) \equiv R_{z'}(\gamma)R_{y'}(\beta)R_z(\alpha) = R_z(\alpha)R_y(\beta)R_z(\gamma). \quad (14.41)$$




FIGURE 3.4 Euler rotations.

Figure 14.1: Euler Rotations

Having obtained the matrix elements of J_z and J_{\pm} , we are now in a position to study the matrix elements of the rotation operator $\mathcal{D}(R)$. If a rotation R is specified by $\hat{\mathbf{n}}$ and ϕ , we can define its matrix elements by

$$\mathcal{D}_{m'm}^{(j)}(R) = \langle j, m' | \exp \left(\frac{-i\mathbf{J} \cdot \hat{\mathbf{n}}\phi}{\hbar} \right) | j, m \rangle. \quad (14.42)$$

These matrix elements are sometimes called **Wigner functions**. In Euler Rotations, the rotation operator is defined by

$$\begin{aligned} \mathcal{D}_{m'm}^{(j)}(\alpha, \beta, \gamma) &= \langle j, m' | \exp \left(\frac{-iJ_z\alpha}{\hbar} \right) \exp \left(\frac{iJ_y\beta}{\hbar} \right) \exp \left(\frac{-iJ_z\gamma}{\hbar} \right) | j, m \rangle \\ &= e^{-i(m'\alpha+m\gamma)} \langle j, m' | \exp \left(\frac{-iJ_y\beta}{\hbar} \right) | j, m \rangle. \end{aligned} \quad (14.43)$$

¹ R also means rotation.

²This figure is from *Modern Quantum Mechanics*

We denote nontrivial part as

$$d_{m'm}^{(j)}(\beta) = \langle j, m' | \exp\left(\frac{-iJ_y\beta}{\hbar}\right) | j, m \rangle. \quad (14.44)$$

Since

$$[\mathbf{J}^2, \mathcal{D}(R)] = 0, \quad (14.45)$$

any state $|j, m\rangle$ after a rotation R is also an eigenket of \mathbf{J}^2 , and its eigenvalue is still $j(j+1)\hbar^2$.

To find the amplitude for being found in $|j, m\rangle$, we simply expand the rotated state as follows:

$$\mathcal{D}(R) |j, m\rangle = \sum_{m'} |j, m'\rangle \langle j, m' | \mathcal{D}(R) |j, m\rangle = \sum_{m'} |j, m'\rangle \mathcal{D}_{m'm}^{(j)}(R). \quad (14.46)$$

Theorem 14.1 (Wigner)

Given an operator R , if for any state $|\psi\rangle, |\phi\rangle$, $|\langle R\psi | R\phi \rangle|$ is held, then R is unitary or anti-unitary.

14.5 Spherical Harmonics

¹ Let us consider a problem with spherical symmetry, we can consider

$$\langle \hat{\mathbf{n}} | l, m \rangle = Y_l^m(\hat{\mathbf{n}}) = Y_l^m(\theta, \phi), \quad (14.47)$$

where we have defined a **direction eigenket** $|\hat{\mathbf{n}}\rangle$.

By (14.36), we have

$$-i\hbar \frac{\partial}{\partial \phi} Y_l^m(\theta, \phi) = m\hbar Y_l^m(\theta, \phi), \quad (14.48)$$

$$\left[\frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial}{\partial \theta} \right) + \frac{1}{\sin^2 \theta} \frac{\partial^2}{\partial \phi^2} + l(l+1) \right] Y_l^m = 0. \quad (14.49)$$

By

$$\langle l', m' | l, m \rangle = \delta_{ll'} \delta_{mm'}, \quad (14.50)$$

we obtain,

$$\int_0^{2\pi} d\phi \int_{-1}^1 d(\cos \theta) Y_{l'}^{m'*}(\theta, \phi) Y_l^m(\theta, \phi) = \delta_{ll'} \delta_{mm'}. \quad (14.51)$$

(14.48) implies that the ϕ -independence $Y_l^m(\theta, \phi)$ must behave like $e^{im\phi}$. By (14.20),

$$\left[i \frac{\partial}{\partial \theta} - \cot \theta \frac{\partial}{\partial \phi} \right] \langle \hat{\mathbf{n}} | l, l \rangle = 0. \quad (14.52)$$

¹Spherical Functions.

So,

$$Y_l^l(\theta, \phi) = c_l e^{il\phi} \sin^l \theta. \quad (14.53)$$

Use (14.51) to normalize,

$$\int_0^\pi \sin^{2l} \theta \sin \theta d\theta = B \left(l+1, \frac{1}{2} \right) = \frac{\Gamma(l+1) \Gamma(\frac{1}{2})}{\Gamma(l+\frac{3}{2})}. \quad (14.54)$$

Then we can use the ladder operator to get Y_l^m . Check details in the reference.

14.6 Schwinger's Oscillator Model of Angular Momentum

Consider a 2D harmonic oscillator $H = \hbar\omega(a_1^\dagger a_1 + a_2^\dagger a_2)$ exhibiting the SU(2) symmetry. We define,

$$J_\mu = \frac{1}{2} a_\alpha^\dagger (\sigma^\mu)^\alpha_\beta a^\beta, \quad (14.55)$$

where σ^μ is one of the [Pauli matrix](#) with $\mu = 1, 2, 3$, $\{\alpha, \beta\} \subseteq \{1, 2\}$. We can prove that

$$[J_\mu, J_\nu] = i\varepsilon_{\mu\nu\rho} J_\rho. \quad (14.56)$$

If we let $2J = a_\alpha^\dagger a^\alpha$, then

$$J^\mu J_\mu = J(J+1). \quad (14.57)$$

And the state can be represented as:

$$|j, m\rangle = \frac{(a_1^\dagger)^{j+m}}{\sqrt{(j+m)!}} \frac{(a_2^\dagger)^{j-m}}{\sqrt{(j-m)!}} |0, 0\rangle. \quad (14.58)$$

It satisfies (14.36) and (14.37).

14.7 Addition of Angular Momenta

Consider two independent state $|\alpha\rangle, |\beta\rangle$, we can see them as a state $|\alpha, \beta\rangle$ ¹,

$$|\alpha, \beta\rangle = |\alpha\rangle \otimes |\beta\rangle, \quad (14.59)$$

Consider the rotation

$$\mathcal{D}(R) = \mathcal{D}_1(R) \otimes \mathcal{D}_2(R) = \exp\left(\frac{-i\mathbf{J}_1 \cdot \hat{\mathbf{n}}\phi}{\hbar}\right) \otimes \exp\left(\frac{-i\mathbf{J}_2 \cdot \hat{\mathbf{n}}\phi}{\hbar}\right). \quad (14.60)$$

Let $\phi = \varepsilon$, and $\mathcal{D}(R) = \exp(-i\mathbf{J} \cdot \hat{\mathbf{n}}\phi/\hbar)$ we can get

$$\mathbf{J} = \mathbf{J}_1 \otimes 1 + 1 \otimes \mathbf{J}_2. \quad (14.61)$$

When we are talking about²

$$\mathbf{J} = \mathbf{J}_1 + \mathbf{J}_2, \quad (14.62)$$

¹The following is tensor product, but in many physics books they mess up with direct product.

²The computing law are very similar, so many other places always write like this.

we should understand it as (14.61). Easy to check that (14.62) satisfies the commutation relation of angular momentum. But neither $\mathbf{J}_1 + \mu\mathbf{J}_2$ nor $\lambda(\mathbf{J}_1 + \mathbf{J}_2)$ ($\mu, \lambda \neq 1$).

Also, we define

$$J_x = J_{1x} \otimes 1 + 1 \otimes J_{2x}, \quad J_y = J_{1y} \otimes 1 + 1 \otimes J_{2y}, \quad J_z = J_{1z} \otimes 1 + 1 \otimes J_{2z}, \quad (14.63)$$

$$J_+ = J_{1+} \otimes 1 + 1 \otimes J_{2+}, \quad J_- = J_{1-} \otimes 1 + 1 \otimes J_{2-}, \quad (14.64)$$

$$\mathbf{J}^2 = J_x^2 + J_y^2 + J_z^2. \quad (14.65)$$

A calculate example¹:

$$(A \otimes B)(C \otimes D) = (AC) \otimes (BD). \quad (14.66)$$

$$J_z^2 = J_{1z}^2 \otimes 1 + 1 \otimes J_{2z}^2 + 2J_{1z} \otimes J_{2z}. \quad (14.67)$$

So we can also write \mathbf{J}^2 into the form of

$$\mathbf{J}^2 = \mathbf{J}_1^2 \otimes 1 + 1 \otimes \mathbf{J}_2^2 + 2J_{1z} \otimes J_{2z} + J_{1+} \otimes J_{2-} + J_{1-} \otimes J_{2+}. \quad (14.68)$$

14.7.1 Clebsch-Gordan Coefficients

As for the choice of base kets, since $[\mathbf{J}^2, J_{1,2z}] \neq 0$, we have two options:

$$|j_1 j_2; j m\rangle \text{ and } |j_1 j_2; m_1 m_2\rangle.$$

We connect them with Clebsch-Gordan coefficients $\langle j_1 j_2 m_1 m_2 | j_1 j_2; j m \rangle$:

$$|j_1 j_2; j m\rangle = \sum_{m_1, m_2} |j_1 j_2; m_1 m_2\rangle \langle j_1 j_2 m_1 m_2 | j_1 j_2; j m \rangle. \quad (14.69)$$

We figure out some properties about CG coefficients:

(1) CG coefficients vanish unless

$$m = m_1 + m_2, \quad (14.70)$$

because

$$\begin{aligned} 0 &= \langle j_1 j_2; m_1 m_2 | (J_z - J_{1z} - J_{2z}) | j_1 j_2; j m \rangle \\ &= (m - m_1 - m_2) \langle j_1 j_2; m_1 m_2 | j_1 j_2; j m \rangle. \end{aligned} \quad (14.71)$$

(2) The coefficients vanish unless

$$|j_1 - j_2| \leq j \leq j_1 + j_2. \quad (14.72)$$

This is based on our choice of base kets, just for convenience, it has already provided a complete bases². In addition, we even choose CG coefficients to be real numbers.

(3) Recursive relations

¹Attention: $A \otimes B + C \otimes D \stackrel{?}{=} (A+C) \otimes (B+D)$. But this is true: $A \otimes C + B \otimes C = (A+B) \otimes C$.

²Check the dimension yourself.

15 Discrete Symmetry

15.1 Wigner Theorem

Theorem 15.1 (Wigner)

Given an operator R , if for any state $|\psi\rangle, |\phi\rangle$, $|\langle R\psi|R\phi\rangle|$ is held, then R is unitary or anti-unitary¹.

Let \mathbb{H} denote the Hilbert space describing a system, with its inner product denoted by $\langle \cdot, \cdot \rangle$, and let \mathbb{PH} denote the corresponding projective space (the set of all one-dimensional subspaces of \mathbb{H}). The reason for using \mathbb{PH} is that its points correspond bijective to physical states. The inner product $\langle \cdot, \cdot \rangle$ on \mathbb{H} induces a metric d on \mathbb{PH} (the Fubini–Study metric). It is easy to see that a unitary or anti-unitary operator $K : \mathbb{H} \rightarrow \mathbb{H}$ (where anti-unitary means $\langle Kv, Kw \rangle = \langle w, v \rangle$) induces a homeomorphism from \mathbb{PH} to \mathbb{PH} that preserves the metric d , meaning $d(Kx, Ky) = d(x, y)$. Wigner’s theorem states that any homeomorphism of \mathbb{PH} onto itself that preserves the metric d is necessarily induced by some unitary or anti-unitary operator on \mathbb{H} .

Theorem (Wigner): If the dimension of \mathbb{H} is greater than or equal to 2, and $U : \mathbb{PH} \rightarrow \mathbb{PH}$ is a homeomorphism preserving the metric d , i.e., $d(Ux, Uy) = d(x, y)$, then there exists a unitary or anti-unitary operator \mathcal{U} on \mathbb{H} such that \mathcal{U} induces U . If \mathcal{V} also induces U , then $\mathcal{V} = e^{i\theta}\mathcal{U}$, where $\theta \in \mathbb{R}$. —Reference

15.2 Parity Transformation

Parity operator π will changes a right-handed system into a left-handed system. Somewhere, we denote any ket $|\alpha\rangle$ after action by π as $|\alpha^\pi\rangle$, but in most of my habits, I will just write $|\pi\alpha\rangle$.

$$|\pi\alpha\rangle := \pi|\alpha\rangle \quad (15.1)$$

Consider the physical meaning, we require

$$\langle \pi\alpha | \pi\alpha \rangle = \langle \alpha | \alpha \rangle, \quad (15.2)$$

$$\langle \pi\alpha | \mathbf{x} | \pi\alpha \rangle = -\langle \alpha | \mathbf{x} | \alpha \rangle. \quad (15.3)$$

15.2 implies that π is whether unitary or anti-unitary.

Since the expectation value is a real number, and π keep the module invariant,

$$\langle \pi\alpha | \mathbf{x} | \pi\alpha \rangle = \langle \alpha | \pi^{-1} \mathbf{x} \pi | \alpha \rangle. \quad (15.4)$$

Then we will use $\langle \alpha | \pi^{-1} \mathbf{x} \pi | \alpha \rangle = -\langle \alpha | \mathbf{x} | \alpha \rangle$ to prove $\pi^{-1} \mathbf{x} \pi = -\mathbf{x}$. Since $|\alpha\rangle$ is arbitrary, let $|\alpha\rangle = |\psi\rangle + |\phi\rangle$. Then, we can deduce that

$$\text{Re}(\langle \psi | \pi^{-1} \mathbf{x} \pi | \phi \rangle) = -\text{Re}(\langle \psi | \mathbf{x} | \phi \rangle). \quad (15.5)$$

¹Review 4.2 first.

Let $|\alpha\rangle = |\psi\rangle + i|\phi\rangle$, then,

$$\text{Im}(\langle\psi|\pi^{-1}\mathbf{x}\pi|\phi\rangle) = -\text{Im}(\langle\psi|\mathbf{x}|\phi\rangle). \quad (15.6)$$

So, for any $|\psi\rangle, |\phi\rangle$,

$$\langle\psi|\pi^{-1}\mathbf{x}\pi|\phi\rangle = -\langle\psi|\mathbf{x}|\phi\rangle. \quad (15.7)$$

Which means

$$\pi^{-1}\mathbf{x}\pi = -\mathbf{x}, \quad \mathbf{x}\pi = -\pi\mathbf{x}. \quad (15.8)$$

Then we derive the effect π act on momentum p . Let us start from position translation. We have

$$\pi\mathcal{T}(\mathbf{d}\mathbf{x}') = \mathcal{T}(-\mathbf{d}\mathbf{x}')\pi, \quad (15.9)$$

so,

$$\pi^{-1}\mathbf{p}\pi = -\mathbf{p}, \quad \{\pi, \mathbf{p}\} = 0. \quad (15.10)$$

Easy to check that

$$\pi^{-1}i\hbar\pi = \pi^{-1}[\mathbf{x}, \mathbf{p}]\pi = [\mathbf{x}, \mathbf{p}] = i\hbar. \quad (15.11)$$

So, π is unitary. Then we can define π^\dagger , and $\pi^\dagger\pi = 1$.

Note that

$$\mathbf{x}\pi|\mathbf{x}'\rangle = -\pi\mathbf{x}|\mathbf{x}'\rangle = (-\mathbf{x}')\pi|\mathbf{x}'\rangle. \quad (15.12)$$

So $\pi|\mathbf{x}'\rangle = e^{i\delta}|\mathbf{x}'\rangle$. It is customary to take $e^{i\delta} = 1$ by convention. Then $\pi^2 = 1$. Hence,

$$\pi^{-1} = \pi^\dagger = \pi. \quad (15.13)$$

Its eigen-value can only be ± 1 .

In geometry, reflection is commutable with rotation. In the language of operators,

$$\pi\mathcal{D}(R) = \mathcal{D}(R)\pi. \quad (15.14)$$

So,

$$\pi^\dagger\mathbf{J}\pi = \mathbf{J}, \quad [\pi, \mathbf{J}] = 0. \quad (15.15)$$

Now, we consider the wave-function under parity transformation. For any state $|\psi\rangle$, let the its state after parity transformation be $|\psi'\rangle$, then we have

$$\psi'(\mathbf{x}) = \langle\mathbf{x}|\psi'\rangle = \langle\mathbf{x}|\pi|\psi\rangle = \langle-\mathbf{x}|\psi\rangle = \psi(-\mathbf{x}). \quad (15.16)$$

If $|\psi\rangle$ is an eigen-state of π , $\pi|\psi\rangle = \pm|\psi\rangle$, then

$$\psi(\mathbf{x}) = \pm\psi(-\mathbf{x}) \begin{cases} \text{even parity,} \\ \text{odd parity.} \end{cases} \quad (15.17)$$

So for 1D harmonic oscillator,

$$\psi_n(-x) = (-1)^n\psi_n(x). \quad (15.18)$$

For angular momentum eigenstates $Y_{l,m}(\hat{\mathbf{n}})$

$$Y_{lm}(-\hat{\mathbf{n}}) = Y_{lm}(\pi - \theta, \pi + \varphi) = (-1)^l Y_{lm}(\theta, \varphi). \quad (15.19)$$

Theorem 15.2

Suppose

$$[H, \pi] = 0$$

and $|n\rangle$ is a nondegenerate eigenket of H with eigenvalue E_n :

$$H|n\rangle = E_n|n\rangle;$$

then $|n\rangle$ is also a parity eigenket.

Proof. Note that $\frac{1}{2}(1 \pm \pi)|n\rangle$ is a parity eigenket with eigenvalues ± 1 . But this is also an energy eigenket with eigenvalue E_n , which means it represents the same state as $|n\rangle$. \square

15.3 Time-Reversal Symmetry

First we claim that any anti-unitary operator can be written as

$$\theta = UK, \tag{15.20}$$

where U is a unitary operator, and K is the complex-conjugate operator that forms the complex conjugate of any coefficient that multiplies a ket.

16 Approximation Methods**16.1 Time-independent Perturbation Theory: Non-degenerate Case**

Consider

$$H = H_0 + \lambda V, \tag{16.1}$$

where $\lambda \ll 1$, and we have known the eigenvalues and eigenkets of H_0 . Then we want to know the eigenvalues and eigenkets of H . Let

$$\Delta_n = E_n - E_n^{(0)}, \tag{16.2}$$

then Schrödinger equation gives

$$(E_n^{(0)} - H_0)|n\rangle = (\lambda V - \Delta_n)|n\rangle. \tag{16.3}$$

For convenience, we define the projection operator and an operator like inverse of $(E_n^{(0)} - H_0)$:

$$\phi_n := 1 - |n^{(0)}\rangle \langle n^{(0)}| = \sum_{m \neq n} |m^{(0)}\rangle \langle m^{(0)}|, \tag{16.4}$$

$$\frac{\phi_n}{E_n^{(0)} - H_0} := \sum_{m \neq n} \frac{|m^{(0)}\rangle \langle m^{(0)}|}{E_n^{(0)} - E_m^{(0)}}. \tag{16.5}$$

We define ϕ_n aim at projecting out the component of $|n^{(0)}\rangle$, eliminating the singularity of $(E_n^{(0)} - H_0)^{-1}$.

Now, we have

$$|n\rangle = \frac{\phi_n}{E_n^{(0)} - H_0} (\lambda V - \Delta_n) |n\rangle + |n^{(0)}\rangle \langle n^{(0)} | n \rangle. \quad (16.6)$$

Note that

$$\langle n^{(0)} | (E_n^{(0)} - H_0) | n \rangle = 0. \quad (16.7)$$

So,

$$\Delta_n \langle n^{(0)} | n \rangle = \lambda \langle n^{(0)} | V | n \rangle. \quad (16.8)$$

Since there's just a common multiplicative factor, we can normalize $|n\rangle$ later, and now we let $\langle n^{(0)} | n \rangle = 1$. Then, we can set

$$|n\rangle = |n^{(0)}\rangle + \lambda |n^{(1)}\rangle + \lambda^2 |n^{(2)}\rangle + \dots, \quad (16.9)$$

$$\Delta_n = \lambda E_n^{(1)} + \lambda^2 E_n^{(2)} + \dots. \quad (16.10)$$

equating the coefficient of various powers of λ , we obtain

$$\Delta_n^{(k)} = \langle n^{(0)} | V | n^{(k-1)} \rangle, \quad k = 1, 2, \dots, \quad (16.11)$$

Also,

$$|n^{(k)}\rangle = \frac{\phi_n}{E_n^{(0)} - H_0} \left[\left(V - \Delta_n^{(1)} \right) |n^{(k-1)}\rangle - \sum_{j=2}^k \Delta_n^{(j)} |n^{(k-j)}\rangle \right], \quad k = 1, 2, \dots \quad (16.12)$$

Let

$$V_{nk} := \langle n^{(0)} | V | k^{(0)} \rangle, \quad (16.13)$$

let us simplify the first and second order corrections:

$$\Delta_n^{(1)} = V_{nn}, \quad (16.14)$$

So,

$$|n^{(1)}\rangle = \sum_{m \neq n} \frac{V_{mn}}{E_n^{(0)} - E_m^{(0)}} |m^{(0)}\rangle, \quad (16.15)$$

$$\Delta_n^{(2)} = \sum_{m \neq n} \frac{|V_{mn}|^2}{E_n^{(0)} - E_m^{(0)}}. \quad (16.16)$$

$$|n^{(2)}\rangle = \sum_{m \neq n} \left[\sum_{l \neq n} \frac{V_{ml} V_{ln}}{(E_n^{(0)} - E_m^{(0)})(E_n^{(0)} - E_l^{(0)})} - \frac{V_{nn} V_{mn}}{(E_n^{(0)} - E_m^{(0)})^2} \right] |m^{(0)}\rangle. \quad (16.17)$$

Thus, up to second order in λ , we have

$$E_n \approx E_n^{(0)} + \lambda V_{nn} + \lambda^2 \sum_{m \neq n} \frac{|V_{mn}|^2}{E_n^{(0)} - E_m^{(0)}}, \quad (16.18)$$

$$|n\rangle \approx |n^{(0)}\rangle + \lambda \sum_{m \neq n} \frac{V_{mn}}{E_n^{(0)} - E_m^{(0)}} |m^{(0)}\rangle. \quad (16.19)$$

Second order correction of energy at ground state always show minus sign.

It is time to normalize $|n\rangle$. We define

$$|n\rangle_N = Z_n^{1/2} |n\rangle, \quad Z_n^{1/2} = \langle n^{(0)} | n \rangle_N. \quad (16.20)$$

Since $\langle n^{(0)} | n^{(k)} \rangle = 0$, if $k \geq 1$,

$$Z_n^{-1} = 1 + \lambda^2 \langle n^{(1)} | n^{(1)} \rangle + \mathcal{O}(\lambda^3) \quad (16.21)$$

$$= 1 + \lambda^2 \sum_{m \neq n} \frac{|V_{mn}|^2}{(E_n^{(0)} - E_m^{(0)})^2} + \mathcal{O}(\lambda^3). \quad (16.22)$$

$$Z_n = 1 - \lambda^2 \sum_{m \neq n} \frac{|V_{mn}|^2}{(E_n^{(0)} - E_m^{(0)})^2} + \mathcal{O}(\lambda^3). \quad (16.23)$$

The second term in (16.23) is to be understood as the probability for “leakage” to states other than $|n^{(0)}\rangle$. Notice that Z_n is less than 1, as expected on the basis of the probability interpretation for Z .

In fact, we have (not only up to second order)

$$Z_n = \frac{\partial E_n}{\partial E_n^{(0)}}. \quad (16.24)$$