

# Quantum Mechanics 2025 HW10

Due 12/16 in Class  
December 16, 2025

## Problem 1.

A one-dimensional simple harmonic oscillator of angular frequency  $\omega$  is acted upon by a spatially uniform but time-dependent force (*not* potential)

$$F(t) = \frac{(F_0\tau/\omega)}{\tau^2 + t^2}, \quad -\infty < t < \infty.$$

At  $t = -\infty$ , the oscillator is known to be in the ground state. Using the time-dependent perturbation theory to first order, calculate the probability that the oscillator is found in the first excited state at  $t = +\infty$ .

Bonus question:  $F(t)$  is so normalized that the impulse

$$\int F(t) dt$$

imparted to the oscillator is always the same—that is, independent of  $\tau$ ; yet for  $\tau \gg 1/\omega$ , the probability for excitation is essentially negligible. Is this reasonable? [Matrix element of  $x$ :  $\langle n'|x|n \rangle = (\hbar/2m\omega)^{1/2} (\sqrt{n}\delta_{n',n-1} + \sqrt{n+1}\delta_{n',n+1})$ .]

## Problem 2.

A hydrogen atom in its ground state  $[(n, l, m) = (1, 0, 0)]$  is placed between the plates of a capacitor. A time-dependent but spatial uniform electric field (not potential!) is applied as follows:

$$\mathbf{E} = \begin{cases} 0 & \text{for } t < 0, \\ \mathbf{E}_0 e^{-t/\tau} & \text{for } t > 0 \end{cases} \quad (\mathbf{E}_0 \text{ is the positive } z\text{-direction})$$

Using first-order time-dependent perturbation theory, compute the probability for the atom to be found at  $t \gg \tau$  in each of the three  $2p$  states:  $(n, l, m) = (2, 1, \pm 1 \text{ or } 0)$ . Repeat the problem for the  $2s$  state:  $(n, l, m) = (2, 0, 0)$ . You need not attempt to evaluate radial integrals, but perform all other integrations (with respect to angles and time).

### Problem 3.

Consider a two-level system with  $E_1 < E_2$ . There is a time-dependent potential that connects the two levels as follows:

$$V_{11} = V_{22} = 0, \quad V_{12} = \gamma e^{i\omega t}, \quad V_{21} = \gamma e^{-i\omega t} \quad (\gamma \text{ real}).$$

At  $t = 0$ , it is known that only the lower level is populated—that is,  $c_1(0) = 1$ ,  $c_2(0) = 0$ .

(a) Find  $|c_1(t)|^2$  and  $|c_2(t)|^2$  for  $t > 0$  by *exactly* solving the coupled differential equation

$$i\hbar\dot{c}_k = \sum_{n=1}^2 V_{kn}(t)e^{i\omega_{kn}t}c_n, \quad (k = 1, 2).$$

(b) Do the same problem using time-dependent perturbation theory to lowest nonvanishing order. Compare the two approaches for small values of  $\gamma$ . Treat the following two cases separately: (i)  $\omega$  very different from  $\omega_{21}$  and (ii)  $\omega$  close to  $\omega_{21}$ . Answer for (a): (Rabi's formula)

$$|c_2(t)|^2 = \frac{\gamma^2/\hbar^2}{\gamma^2/\hbar^2 + (\omega - \omega_{21})^2/4} \sin^2 \left[ \frac{\gamma^2}{\hbar^2} + \frac{(\omega - \omega_{21})^2}{4} \right]^{1/2} t,$$

$$|c_1(t)|^2 = 1 - |c_2(t)|^2.$$

### Problem 4.

An atom is irradiated by right-hand circularly polarized light propagating along the  $z$ -direction. The atom transitions from energy level  $E_{nl}$  to  $E_{n'l'}$ . Find the selection rules.

### Problem 5.

Assuming spin is neglected, the electronic state in an atom is expressed as

$$\psi_{nlm}(r, \theta, \varphi) = R_n(r)Y_l^m(\theta, \varphi).$$

(a) For an electric dipole spontaneous transition from an initial  $s$ -state (with energy  $E_{nl}$ ,  $l = 0$ ) to a final  $p$ -state (with energy  $E_{n'l'}$ ,  $l' = 1$ ), find the branching ratios for the final states with  $m = 1, 0, -1$  respectively.

(b) Let the initial state be  $nlm$  and final state  $n'l'm'$ . (1) For a fixed final state  $n'l'$ , find the branching ratios for transitions to different  $m'$  states. (2) Find the branching ratios for transitions to different  $l'$  levels (neglecting the differences in the radial wave functions).