

The Complete Genome Sequence of Shope (Rabbit) Fibroma Virus

David O. Willer,* Grant McFadden,† and David H. Evans*¹

*Department of Molecular Biology and Genetics, The University of Guelph, Guelph, Ontario N1G 2W1, Canada; and †Department of Microbiology and Roberts Research Institute, The University of Western Ontario, London, Ontario N6G 2V4, Canada

Received July 27, 1999; returned to author for revision August 18, 1999; accepted September 10, 1999

We have determined the complete DNA sequence of the Leporipoxvirus Shope fibroma virus (SFV). The SFV genome spans 159.8 kb and encodes 165 putative genes of which 13 are duplicated in the 12.4-kb terminal inverted repeats. Although most SFV genes have homologs encoded by other *Chordopoxvirinae*, the SFV genome lacks a key gene required for the production of extracellular enveloped virus. SFV also encodes only the smaller ribonucleotide reductase subunit and has a limited nucleotide biosynthetic capacity. SFV preserves the *Chordopoxvirinae* gene order from S012L near the left end of the chromosome through to S142R (homologs of vaccinia F2L and B1R, respectively). The unique right end of SFV appears to be genetically unstable because when the sequence is compared with that of myxoma virus, five myxoma homologs have been deleted (C. Cameron, S. Hota-Mitchell, L. Chen, J. Barrett, J.-X. Cao, C. Macaulay, D. Willer, D. Evans, and G. McFadden, 1999, *Virology* 264, 298–318). Most other differences between these two Leporipoxviruses are located in the telomeres. Leporipoxviruses encode several genes not found in other poxviruses including four small hydrophobic proteins of unknown function (S023R, S119L, S125R, and S132L), an α 2,3-sialyltransferase (S143R), a protein belonging to the Ig-like protein superfamily (S141R), and a protein resembling the DNA-binding domain of proteins belonging to the HIN-200 protein family (S013L). SFV also encodes a type II DNA photolyase (S127L). *Melanoplus sanguinipes* entomopoxvirus encodes a similar protein, but SFV is the first mammalian virus potentially capable of photoreactivating ultraviolet DNA damage. © 1999 Academic Press

INTRODUCTION

The Poxviridae comprise of a large family of double-stranded DNA viruses that infect vertebrates (Chordopoxvirinae) and insects (Entomopoxvirinae). These viruses are characterized by a complex capsid architecture, replicate in the cytoplasm of infected cells, and are of particular interest for their remarkable capacity to disarm or delay many different host antiviral responses. Poxviruses are endowed with some of the largest known viral genomes (139–260 kb), and thus few complete genomic DNA sequences have been reported to date. Those viral DNA sequences that have been completely elucidated include that of a single viral member of the Entomopoxvirinae family [*Melanoplus sanguinipes* entomopoxvirus, MSV (Afonso *et al.*, 1999)] as well as those of several Chordopoxvirinae. These latter viruses include the *Orthopoxviruses* vaccinia [VAC, (Goebel *et al.*, 1990; Antoine *et al.*, 1998)] and variola [VAR, (Massung *et al.*, 1994; Shchelkunov *et al.*, 1995)], as well as the *Molluscipoxvirus* mollusci contagiosum virus [MCV, (Senkevich *et al.*, 1997)].

Shope (or rabbit) fibroma virus (SFV) belongs to the *Leporipoxvirus* genus, a family of viruses whose host range includes rabbits, hares, and squirrels (reviewed in Digiacomo and Mare, 1994). SFV was originally shown to

cause fibroxanthosarcoma-like tumours in its natural host, the eastern cottontail rabbit *Sylvilagus floridanus* (Shope, 1932b). Leporipoxviruses appear to be transmitted from rabbit to rabbit by biting insects, and the widespread prevalence of antibodies to the virus suggests that SFV infections may be endemic throughout US and Canadian rabbit populations (Digiacomo and Mare, 1994). Superficially similar disease symptoms and virus particles also have been reported in the African hare *Lepus capensis*, suggesting that the range of Leporipoxviruses may extend as far as African rabbit populations (Karstad *et al.*, 1977). Healthy adult rabbits mount an effective cell-mediated immune response, which typically starts to induce tumour regression within 10–12 days post infection (Scott *et al.*, 1981; Sell and Scott, 1981). However, SFV sometimes can cause a lethal disseminated infection in newborn rabbits and immunocompromised adults. Immunological studies (Shope, 1932a) and subsequent DNA sequence analyses have shown that SFV is closely related to myxoma virus (MYX), the causative agent of rabbit myxomatosis. The two kinds of viral infections are reportedly hard to differentiate in cottontail rabbits, but whereas SFV-induced disease in European rabbits (*Oryctolagus cuniculus*) closely resembles the disease in *S. floridanus*, myxoma causes a rapidly lethal disseminated infection in European rabbits that is strikingly different from the benign fibromas caused by SFV. This phenotype is partially explicable by the fact that myxoma virus, unlike SFV, can inhibit an

¹ To whom reprint requests should be addressed. Fax: (519) 837-2075. E-mail: dhevans@uoguelph.ca.

apoptotic response in rabbit T lymphocytes and thus is able to replicate and spread through the lymphatic system (Macen *et al.*, 1996).

A number of studies have been directed at mapping and sequencing the genomes of SFV and myxoma viruses as well as what appears to be a natural recombinant called malignant rabbit fibroma virus (Strayer *et al.*, 1983; Block *et al.*, 1985; Upton *et al.*, 1988). Restriction sites are highly conserved amongst SFV strains and a detailed restriction map suggested a total genome size of ~160 kb with 12-kb terminal inverted repeats (TIR) (Wills *et al.*, 1983; Delange *et al.*, 1984; Cabirac *et al.*, 1985). DNA sequencing studies have shown that the SFV genome bears the hairpin tips characteristic of poxviruses (DeLange *et al.*, 1986) and encodes an array of genes thought to be required either for growth (Stuart *et al.*, 1993) or pathogenesis (Chang *et al.*, 1987, 1990; Upton *et al.*, 1987; Smith *et al.*, 1991a; Strayer *et al.*, 1991; Opgenorth *et al.*, 1993; Mossman *et al.*, 1995; Brick *et al.*, 1998). DNA transfection methods have been used to show that SFV catalyzes an efficient nonspecific DNA replication reaction (DeLange and McFadden, 1986), and this reaction has been exploited to study the mechanics of both SFV telomere resolution (DeLange *et al.*, 1986) and poxvirus genetic recombination (Evans *et al.*, 1988; Fisher *et al.*, 1991; Parks and Evans, 1991a,b). Biochemical and genetic studies also have characterized the reactions catalyzed by some of the most highly conserved of the replicative genes encoded by poxviruses. These include SFV topoisomerase (Upton *et al.*, 1990b; Palaniyar *et al.*, 1996, 1999), DNA ligase (Parks *et al.*, 1994, 1998), thymidine kinase (Upton and McFadden, 1986), and uracil-*N*-glycosylase (Upton *et al.*, 1993). However, these and other studies have yet to explain how or why SFV-infected cells catalyze such high levels of genetic recombination or *trans*-replication.

As a way of gaining further insights into these varied aspects of SFV biology, we have determined the complete genomic DNA sequence of the virus using a combination of random sequence analysis and primer walking. Together with a separate report on the genome sequence of myxoma virus (Cameron *et al.*, 1999) our work provides the first insights into the complete gene complement of *Leporipoxviruses*.

RESULTS AND DISCUSSION

Overview of the SFV genome

The SFV genome is 159,857 bp in length and encodes 60.5% (A+T) base pairs. The calculated *Bam*HI restriction map was in exact concordance with that determined experimentally (Delange *et al.*, 1984). The gene arrangement and direction of transcription closely resembles that seen in other Chordopoxviridae (Fig. 1) in that there is a pronounced tendency for genes located near the ends of the virus to be transcribed toward the telomeres.

This can be demonstrated by arbitrarily dividing the genome into four 40,000-nt quarter sections. Within the left- and right-hand portions of the genome, only six genes in each of these two quarter sections disregard this rule. Throughout the rest of the genome, genes also tend to be arranged in transcriptional clusters with the occasional single gene periodically interrupting clusters of genes all transcribed in a common direction. The genes were numbered starting from the left end of the virus using a scheme similar to that used for molluscum contagiosum and *Melanoplus sanguinipes* viruses (Table 1) (Senkevich *et al.*, 1996; Afonso *et al.*, 1999). Several exceptions to the sequential number system were employed to maintain continuity with the earlier literature and with the gene order in myxoma virus. For example, we have assigned decimals to the genes previously called T-3A (S003.1L) and T-3C (S003.2L) to ensure that other telomeric genes (e.g., T-6 = S006L) retain their original numbering. Other features of the gene nomenclature are discussed in the accompanying paper (Cameron *et al.*, 1999).

The SFV gene order also closely matches that of the other sequenced Chordopoxviridae. Within the central core of the virus only a small number of genes have been deleted or replaced, and we can readily align nearly every SFV gene stretching from S019L to S126R with VAC and MCV homologs running from VAC F9L/MC016L to VAC A37R/MC149R. Thereafter the alignment can still be followed in vaccinia virus but becomes less obvious due to the absence or substitution of many genes. Nevertheless, it can still be traced out to S012L on the left end of the virus [which encodes a homolog of VAC F2L (dUT-Pase)] and out to S142R on the right side of the genome (which encodes a homolog of VAC B1R protein kinase). The rightmost portion of SFV also encodes a few genes found to the left of F2L in vaccinia virus. These include S144R, S146R, and S154R, which appear to be homologs of VAC C3L, N1L, and VAC M2L, respectively, and the different locations suggest that a translocation may have occurred sometime after the genetic isolation of Leporipoxviruses and Orthopoxviruses. SFV, like other Chordopoxviridae, shares no obvious conservation of gene order with the one sequenced Entomopoxvirus.

Terminal inverted repeats

The SFV TIRs have been previously shown to span 12,397 nt. They are most notable for the complete absence of any repetitive DNA and a high density of encoded genes. An odd feature of the terminal inverted repeats is that the TIR junction falls within the boundaries of genes designated S009L (previously called T-9) and S155R. Because the C termini of these two proteins are located within the TIR, they share 443 amino acids of perfect amino acid identity. However, S009L encodes 67 more amino acids than does S155R, in a portion of the




FIG. 1. SFV genetic map. Rectangles indicate open reading frames, and the direction of transcription is indicated by the box placement. The regions corresponding to each of the known *Bam*HI restriction fragments also have been located on the genetic map in white lettering. Where genes are transcribed in a leftward or a rightward direction, they have been drawn below or above the line, respectively. There is a tendency for genes to be transcribed in commonly oriented blocks, interrupted by single-isolated genes. A selected subset of SFV genes and genome features also have been identified to illustrate points raised in the Discussion and to facilitate comparison with other poxvirus genetic maps.

TABLE 1
SFV Gene Products

ORF ^a	Translation		Size (aa)	MW (kDa) ^c	Prom. type ^d	Function/feature(s)	Homolog(s) ^e	BLASTP score ^f	Amino acid identity ^g	Reference(s) ^h
	Start	Stop ^b								
S001L/R	1251	475	258	28.4	E	Chemokine binding protein	VAC C23L/B29R	4E-32	74/164 (45)	Graham <i>et al.</i> (1997); Patel <i>et al.</i> (1990)
S002L/R	158607	159383	325	35.1	E	TNF- α Receptor	VAC C22L/B28R	5E-25	154/320 (48)	Smith <i>et al.</i> (1990, 1991a)
S003.1L/R	2325	1348	325	35.1	E	TNF- α Receptor	VAC C22L/B28R	5E-25	154/320 (48)	Smith <i>et al.</i> (1990, 1991a)
S003.2L/R	157533	158510	325	35.1	E	TNF- α Receptor	VAC C22L/B28R	5E-25	154/320 (48)	Smith <i>et al.</i> (1990, 1991a)
S004L/R	2861	2406	151	17.7	E	Unknown	VAC B15R	4E-08	37/120 (30)	
S005L/R	156997	157452	151	17.7	E	Unknown	VAC B15R	4E-08	37/120 (30)	
S006L/R	3306	2929	125	14.4	?	Unknown	SHP T3C	2E-17	44/104 (42)	Gershon and Black (1989)
S007L/R	156552	156929	125	14.4	?	Unknown	SHP T3C	2E-17	44/104 (42)	Gershon and Black (1989)
S008L/R	4211	3534	225	25.1	E	Virulence factor (RDEL motif)	VAC B9R	0.002	23/69 (33)	Barry <i>et al.</i> (1997)
S009L	155647	156324	225	25.1	E	Virulence factor (RDEL motif)	VAC B9R	0.002	23/69 (33)	Barry <i>et al.</i> (1997)
S010L	4493	4251	80	9.2	L	Unknown	SPV C2L	4E-17	44/92 (47)	Massung <i>et al.</i> (1993)
S011L	155365	155607	80	9.2	L	Unknown	SPV C2L	4E-17	44/92 (47)	Massung <i>et al.</i> (1993)
S012L	5967	4513	484	55.8	E	Virulence factor (ankyrin-like protein)	VAC B4R	3E-10	90/427 (21)	Howard <i>et al.</i> (1991) Mossman <i>et al.</i> (1996a)
S013L	153891	155345	484	55.8	E	Virulence factor (ankyrin-like protein)	VAC B4R	3E-10	90/427 (21)	Howard <i>et al.</i> (1991) Mossman <i>et al.</i> (1996a)
S014L	7530	6004	508	58.2	E	Kelch repeat	VAC A55R	8E-32	124/506 (24)	
S015L	152328	153854	508	58.2	E	Kelch repeat	VAC A55R	8E-32	124/506 (24)	
S016L	8360	7572	262	30.6	E	IFN- γ receptor mimic	VAC B8R	7E-12	48/171 (28)	Upton <i>et al.</i> (1992) Mossman <i>et al.</i> (1996b)
S017L	151498	152286	262	30.6	E	IFN- γ receptor mimic	VAC B8R	7E-12	48/171 (28)	Upton <i>et al.</i> (1992) Mossman <i>et al.</i> (1996b)
S018L	9959	8415	514	59.4	E	Kelch repeat	VAC A55R	1E-30	120/502 (23)	
S019L	149899	151443	514	59.4	E	Kelch repeat	VAC A55R	1E-30	120/502 (23)	
S020L	10567	10397	56	6.3	—	Serpin fragment ("SERP-1")	MYX MAP1	0.38	19/37 (51)	Upton <i>et al.</i> (1990a)
S021L	149291	149461	56	6.3	—	Serpin fragment ("SERP-1")	MYX MAP1	0.38	19/37 (51)	Upton <i>et al.</i> (1990a)
S022L	10842	10606	78	9.1	E	Interferon resistance	VAC K3L	0.014	18/60 (30)	Beattie <i>et al.</i> (1991) Davies <i>et al.</i> (1992)
S023R	149016	149252	78	9.1	E	Interferon resistance	VAC K3L	0.014	18/60 (30)	Beattie <i>et al.</i> (1991) Davies <i>et al.</i> (1992)
S024L	12548	11016	510	58.3	E	Kelch repeat	VAC A55R	6E-27	125/511 (24)	Goebel <i>et al.</i> (1990)
S025L	12936	12694	80	9.2	L	EGF-like growth factor	VAC C11R	0.22	15/51 (29)	Chang <i>et al.</i> (1987)
S026L	12936	12694	80	9.2	L	EGF-like growth factor	VAC C11R	0.22	15/51 (29)	Chang <i>et al.</i> (1987)
S027L	13409	12918	163	18.5	E	Membrane virulence factor	SPV C10L (?)	2.5	20/78 (25)	Graham <i>et al.</i> (1992) Massung <i>et al.</i> (1993)
S028L	13856	13425	143	15.4	E	dUTPase	VAC F2L	5E-40	82/138 (59)	Roseman and Slabaugh (1990) Roseman <i>et al.</i> (1996)
S029L	14217	13894	107	12.7	E	Resembles <i>M.</i> <i>musculus</i> interferon- inducible protein	IFN-203	0.007	22/83 (26)	Gribaudo <i>et al.</i> (1997)
S030L	14217	13894	107	12.7	E	Resembles <i>M.</i> <i>musculus</i> interferon- inducible protein	IFN-203	0.007	22/83 (26)	Gribaudo <i>et al.</i> (1997)
S031L	15802	14252	516	58.7	E	Kelch repeat	VAC F3L	2E-41	135/511 (26)	Roseman and Slabaugh (1990)
S032L	16824	15856	322	37.5	E	Ribonucleotide reductase (small subunit)	VAC F4L	1E-143	243/319 (76)	Slabaugh <i>et al.</i> (1988) Roseman and Slabaugh (1990)
S033L	15802	14252	516	58.7	E	Kelch repeat	VAC F3L	2E-41	135/511 (26)	Roseman and Slabaugh (1990)
S034L	16824	15856	322	37.5	E	Ribonucleotide reductase (small subunit)	VAC F4L	1E-143	243/319 (76)	Slabaugh <i>et al.</i> (1988) Roseman and Slabaugh (1990)
S035L	17101	16853	82	9.4	?	Unknown	SPV C15L	4E-9	29/74 (39)	Massung <i>et al.</i> (1993)
S036L	17337	17104	77	8.8	E	Unknown	SPV C15L	4E-9	29/74 (39)	Massung <i>et al.</i> (1993)
S037L	17726	17526	66	7.5	E	Unknown	VAC F8L	0.043	19/44 (43)	
S038L	18427	17780	215	24.5	L	Unknown	VAC F9L	2E-49	100/217 (46)	
S039L	18427	17780	215	24.5	L	Unknown	MC016L	8E-36	78/214 (36)	
S040L	19742	18405	445	52.5	L	S/T/Y protein kinase 2	VAC F10L	0.0	300/440 (68)	Lin and Broyles (1994) Derrien <i>et al.</i> (1999)
S041L	21728	19845	627	72.5	E	EEV maturation (fowlpox)	MC017L	1E-165	264/439 (60)	Ogawa <i>et al.</i> (1993)
S042L	21728	19845	627	72.5	E	EEV maturation (fowlpox)	VAC F12L	1E-101	215/618 (34)	Ogawa <i>et al.</i> (1993)
S043L	22876	21764	370	41.5	L	Envelope antigen	MC019L	2E-66	176/620 (28)	
S044L	22876	21764	370	41.5	L	Envelope antigen	VAC F13L	1E-115	201/369 (54)	Jackson and Hall (1998)
S045L	22876	21764	370	41.5	L	Envelope antigen	MC021L	6E-84	158/364 (43)	Hirt <i>et al.</i> (1986)
S046L	22958	23064	35	3.8	L	Unknown	VAC F15L	1E-36	65/148 (43)	
S047L	23488	23042	148	17	E	Unknown	MC025L	6E-35	63/147 (42)	
S048L	23488	23042	148	17	E	Unknown	MC025L	6E-35	63/147 (42)	
S049L	24181	23552	209	24.5	E	Unknown	VAC F16L	3E-30	77/223 (34)	
S050L	24181	23552	209	24.5	E	Unknown	MC029L	1E-12	53/229 (23)	

TABLE 1—Continued

ORF ^a	Translation		Size (aa)	MW (kDa) ^c	Prom. type ^d	Function/feature(s)	Homolog(s) ^e	BLASTP score ^f	Amino acid identity ^g	Reference(s) ^h
	Start	Stop ^b								
S026R	24220	24525	101	11.2	L	DNA-binding phosphoprotein (VP11)	VAC F17R MC030R	7E-28 5E-17	63/104 (60) 44/94 (46)	Kao <i>et al.</i> (1981) Jackson and Hall (1998)
S027L	25934	24522	470	53.8	?	Poly (A) polymerase catalytic subunit	VAC E1L MC031L	1E-175 1E-138	290/462 (62) 249/471 (52)	Gershon <i>et al.</i> (1991)
S028L	28906	25931	721	83.9	L	Unknown	VAC E2L MC032L	1E-151 1E-101	286/740 (38) 226/749 (30)	
S029L	28485	28135	116	13.1	E	dsRNA-dependent protein kinase inhibitor	VAC E3L	6E-15	35/90 (38)	Chang <i>et al.</i> (1995)
S030L	29215	28547	222	25.8	E	RNA polymerase subunit (rpo30, VITF-1)	VAC E4L MC034L	6E-74 3E-58	130/192 (67) 102/180 (56)	Ahn <i>et al.</i> (1990) Broyles and Pennington (1990)
S031R	29315	30493	392	45.7	E	Unknown	VAC E5R	2E-12	79/314 (25)	
S032R	30503	32200	565	66.7	L	Unknown	VAC E6R MC037R	0.0 0.0	333/567 (58) 283/565 (50)	
S033R	32203	33027	274	31.8	E	Unknown	VAC E8R MC038R	1E-102 1E-89	177/264 (67) 160/269 (59)	
S034L	36041	33024	1005	117.1	E	DNA polymerase	VAC E9L MC039L	0.0 0.0	641/1008 (63) 490/1009 (48)	Earl <i>et al.</i> (1986) Traktman <i>et al.</i> (1984)
S035R	36075	36365	96	11.1	?	Unknown	VAC E10R MC040R	3E-38 3E-30	70/95 (73) 55/95 (57)	
S036L	38431	36386	681	78.2	E	Unknown	VAC O1L	1E-111	231/677 (34)	
S037L	38565	38467	32	3.8	L	Unknown	MC042L FPV O3L VAC O3L(?) MC043.1L(?)	5E-48 0.32 >1 >1	113/488 (23) 13/27 (48)	
S038L	39517	38576	313	36.4	L	DNA-binding, late stage virion morphogenesis	VAC I1L MC044L	1E-117 2E-94	208/313 (66) 165/308 (53)	Klemperer <i>et al.</i> (1997)
S039L	39742	39518	74	8.6	L	Unknown	VAC I2L MC045L	1E-12 7E-9	35/71 (49) 29/73 (39)	
S040L	40555	39743	270	30.3	E	DNA-binding phosphoprotein	VAC I3L MC046L	1E-65 4E-49	126/228 (55) 102/237 (43)	Davis and Mathews (1993)
S041L	40859	40623	78	8.9	?	Structural protein (VP13K)	MCO47L	2E-10	29/76 (38)	Takahashi <i>et al.</i> (1994)
S042L	42040	40880	386	44.7	L	Unknown	VAC I5L VAC I6L MC048L	8E-9 1E-104 6E-53	25/69 (36) 187/386 (48) 111/363 (30)	
S043L	43322	42033	429	49.9	L	Virion morphogenesis, cysteine protease?	VAC I7L MC049L	1E-163 1E-129	270/425 (63) 225/424 (53)	Kane and Shuman (1993) Li and Hochstrasser (1999)
S044R	43328	45364	678	78.2	?	NPH-II, RNA helicase	VAC I8R MC050L	0.0 1E-171	381/676 (56) 304/683 (44)	Koonin and Senkevich (1992) Shuman (1992)
S045L	47133	45361	590	68.4	L	Metalloendo-proteinase?, virion morphogenesis	VAC G1L MC056L	0.0 1E-156	322/592 (54) 259/591 (43)	Whitehead and Hruby (1994)
S046L	47465	47130	111	12.8	E/L	Unknown	VAC G3L MC057L	4E-11 4.0	36/89 (40) 26/87 (29)	
S047R	47459	48136	225	26.2	E	IBT-dependent protein, late elongation factor (VLTF)	VAC G2R MC058R	5E-53 2E-25	102/225 (45) 62/235 (26)	Meis and Condit (1991) Black <i>et al.</i> (1998)
S048L	48480	48094	128	14.7	L	Glutaredoxin 2, membrane protein	VAC G4L MC059L	5E-28 2E-13	56/121 (46) 36/121 (29)	Gvakharia <i>et al.</i> (1996) Jensen <i>et al.</i> (1996)
S049R	48483	49781	432	49.8	?	Unknown	VAC G5R MC060R	9E-74 2E-54	163/391 (41) 127/395 (32)	

TABLE 1—Continued

ORF ^a	Translation		Size (aa)	MW (kDa) ^c	Prom. type ^d	Function/feature(s)	Homolog(s) ^e	BLASTP score ^f	Amino acid identity ^g	Reference(s) ^h
	Start	Stop ^b								
S050R	49785	49976	63	7.1	E/L	RNA polymerase subunit (rpo7)	VAC G5.5R MC061R	3E-23 4E-18	49/63 (77) 40/63 (63)	Meis and Condit (1991) Amegadzie <i>et al.</i> (1992) Johnson <i>et al.</i> (1993)
S051R	49976	50500	174	20	?	Unknown	VAC G6R MC062R	2E-36 2E-18	75/164 (45) 58/171 (33)	
S052L	51521	50469	350	39.4	L	Structural protein (VP16K)	VAC G7L MC065L	8E-91 1E-52	175/372 (47) 108/250 (43)	Takahashi <i>et al.</i> (1994)
S053R	51551	52333	260	29.9	I	Late transcription factor (VLTF-1)	VAC G8R MC067R	1E-125 1E-107	214/260 (82) 186/260 (71)	Keck <i>et al.</i> (1990) Wright <i>et al.</i> (1991)
S054R	52354	53352	332	38.6	L	Myristylprotein	VAC G9R MC068R	7E-78 7E-41	143/324 (44) 102/322 (31)	Martin <i>et al.</i> (1997)
S055R	53353	54081	242	26.2	L	Myristylprotein, IMV virion protein	VAC L1R MC069R	3E-89 5E-80	153/239 (64) 141/242 (58)	Franke <i>et al.</i> (1990) Martin <i>et al.</i> (1997)
S056R	54137	54436	99	11.8	E	Unknown	VAC L2R MC070R	2E-8 2.0	26/68 (38) 19/80 (23)	
S057L	55348	54386	320	37.5	L	Unknown	VAC L3L MC072L	7E-86 5E-63	167/328 (50) 122/286 (42)	
S058R	55373	56128	251	28.6	L	DNA binding structural protein precursor (VP8)	VAC L4R MC073R	4E-87 6E-65	151/250 (60) 129/254 (50)	Yang and Bauer (1988) Bayliss and Smith (1997)
S059R	56147	56536	129	15	L	Unknown	VAC L5R MC074R	2E-19 0.74	50/123 (40) 24/111 (21)	
S060R	56490	56936	148	17.1	L	Dimeric virion protein	VAC J1R MC075R	5E-35 5E-32	72/150 (48) 69/147 (46)	cited in Antoine <i>et al.</i> (1998)
S061R	56969	57505	178	19.9	E	Thymidine kinase	VAC J2R	1E-62	116/175 (66)	Upton and McFadden (1986)
S062R	57587	58063	158	18.5	E	Host range, triplicated	VAC C7L	2E-14	48/148 (32)	Perkus <i>et al.</i> (1990)
S063R	58120	58728	202	22.8	E	Host range, triplicated	VAC C7L (?)	0.64	38/156 (24)	Perkus <i>et al.</i> (1990)
S064R	58770	59366	198	23.3	E	Host range, triplicated	VAC C7L	3E-17	42/138 (30)	Perkus <i>et al.</i> (1990)
S065R	59468	60484	328	39.7	E/L	Poly (A) polymerase regulatory subunit (VP39)	VAC J3R MC076R	1E-129 8E-89	217/333 (65) 165/342 (48)	Jackson and Bults (1990) Schnierle <i>et al.</i> (1992)
S066R	60384	60941	185	21	?	RNA polymerase subunit (rpo22)	VAC J4R MC077R	2E-68 5E-59	123/185 (66) 107/187 (57)	Broyles and Moss (1986) Jackson and Bults (1990)
S067L	61345	60944	133	15.4	?	Unknown (essential)	VAC J5L MC078L	7E-45 2E-28	79/131 (60) 52/114 (45)	Zajac <i>et al.</i> (1995)
S068R	61416	65276	1286	147.5	E/L	RNA polymerase subunit (rpo147)	VAC J6R MC079R	0.0 0.0	1019/1288 (79) 918/1289 (71)	Broyles and Moss (1986)
S069L	65803	65282	173	20.1	L	S/Y phosphatase	VAC H1L MC082L	7E-58 3E-48	108/172 (62) 88/171 (51)	Mossman <i>et al.</i> (1995)
S070R	65819	66391	190	21.6	L	Unknown	VAC H2R MC083R	9E-74 1E-59	124/188 (65) 96/181 (53)	
S071L	67368	66394	324	37.2	L	IMV envelope protein	VAC H3L MC084L	9E-52 9E-20	109/326 (33) 83/315 (26)	Chertov <i>et al.</i> (1991)
S072L	69765	67369	798	94.3	L	RNA polymerase-associated TF (RAP 94)	VAC H4L MC085L	0.0 0.0	504/800 (63) 440/800 (55)	Ahn and Moss (1992b)
S073R	69894	70490	198	22.3	E/L	Late transcription factor (VLTF-4)	VAC H5R MC086R	6E-9 0.003	29/60 (48) 21/59 (35)	Kovacs and Moss (1996)
S074R	70518	71462	314	37	L	Topoisomerase I, (telomere resolvase)	VAC H6R MC087R	1E-113 6E-93	193/312 (61) 164/310 (52)	Upton <i>et al.</i> (1990b) Palaniyar <i>et al.</i> (1999)
S075R	71462	71914	150	17.6	I	Unknown	VAC H7R MC088R	6E-26 1E-14	58/145 (40) 47/148 (31)	

TABLE 1—Continued

ORF ^a	Translation		Size (aa)	MW (kDa) ^c	Prom. type ^d	Function/feature(s)	Homolog(s) ^e	BLASTP score ^f	Amino acid identity ^g	Reference(s) ^h
	Start	Stop ^b								
S076R	71907	74417	836	97	E	mRNA capping enzyme, large subunit	VAC D1R MC090R	0.0 0.0	540/832 (64) 467/819 (57)	Upton <i>et al.</i> (1991)
S077L	74810	74379	143	16.6	?	Virion component	VAC D2L MC091L	9E-20 3E-13	51/150 (34) 46/161 (28)	Upton <i>et al.</i> (1991) Dyster and Niles (1991)
S078R	74816	75541	241	28	?	Structural protein	VAC D3R MC092R	2E-22 5E-8	79/247 (31) 49/218 (22)	Dyster and Niles (1991)
S079R	75538	76194	218	25.6	E	Uracil DNA glycosylase	VAC D4R MC093R	3E-93 2E-65	150/218 (68) 112/222 (50)	Upton <i>et al.</i> (1993)
S080R	76228	78588	786	90.9	E/L	Replication/recomb. NTPase	VAC D5R MC094R	0.0 0.0	498/785 (63) 419/787 (53)	Evans <i>et al.</i> (1995)
S081R	78585	80492	635	73.2	L	Early transcription factor (VETF-1)	VAC D6R MC095R	0.0 0.0	514/637 (80) 493/631 (78)	Strayer <i>et al.</i> (1991) Broyles and Fesler (1990) Gershon and Moss (1990)
S082R	80523	81014	163	18.4	E/L	RNA polymerase subunit (rpo18)	VAC D7R MC097R	6E-50 2E-44	97/161 (60) 88/158 (55)	Strayer <i>et al.</i> (1991) Ahn <i>et al.</i> (1990) Quick and Broyles (1990)
S083L	81842	80985	285	32.4	?	Carbonic anhydrase-like structural protein	VAC D8L	3E-28	82/224 (36)	Niles <i>et al.</i> (1986); Niles and Seto (1988) Strayer <i>et al.</i> (1991)
S084R	81888	82508	206	24	E	Mut-T like protein	VAC D9R MC098R	1E-58 4E-45	105/203 (51) 90/209 (43)	Niles <i>et al.</i> (1986) Koonin <i>et al.</i> (1993)
S085R	82505	83287	260	30.4	E/L	Mut-T like protein, neg. regulates gene expression	VAC D10R MC099R	5E-56 1E-41	111/244 (45) 90/211 (42)	Niles <i>et al.</i> (1986) Shors <i>et al.</i> (1999)
S086L	85182	83284	632	72.8	L	NTPase I, DNA helicase	VAC D11L MC100L	0.0 0.0	433/630 (68) 365/626 (58)	Christen <i>et al.</i> (1998) Koonin and Senkevich (1992)
S087L	86063	85200	287	33.2	?	mRNA capping enzyme small subunit, transcrip. initiation factor (VITF)	VAC D12L MC101L	1E-125 1E-106	207/287 (72) 176/284 (61)	Niles <i>et al.</i> (1989) Vos <i>et al.</i> (1991) Weinrich and Hruby (1986)
S088L	87750	86092	552	62.3	L	Rifampicin resistance IMV protein	VAC D13L MC102L	0.0 0.0	378/550 (68) 318/549 (57)	Tartaglia and Paoletti (1985) Sodeik <i>et al.</i> (1994)
S089L	88226	87777	149	16.8	I	Trans-activator protein (VLTF-2)	VAC A1L MC103L	1E-52 9E-42	98/150 (65) 77/147 (52)	Keck <i>et al.</i> (1990) Keck <i>et al.</i> (1993)
S090L	88937	88263	224	26.2	I	Trans-activator protein (VLTF-3)	VAC A2L MC104L	1E-112 2E-97	190/224 (84) 165/222 (74)	Keck <i>et al.</i> (1990) Keck <i>et al.</i> (1993)
S091L	89161	88934	75	8.8	L	Unknown	VAC (MVA) 8.9K MC105L	9E-19 6E-7	42/76 (55) 24/65 (36)	Antoine <i>et al.</i> (1998)
S092L	91131	89170	653	74.3	L	Major core protein P4b precursor	VAC A3L MC106L	0.0 0.0	398/652 (61) 368/667 (55)	Rosel and Moss (1985)
S093L	91627	91166	153	17.7	L	Core protein?	MC107L VAC A4L (?)	6E-4 >1	25/73 (34) 19/64 (29)	Demkowicz <i>et al.</i> (1992) Cudmore <i>et al.</i> (1996)
S094R	91667	92161	164	18.8	L	RNA polymerase subunit (rpo19)	VAC A5R MC108R	2E-36 2E-35	78/126 (61) 76/129 (58)	Ahn <i>et al.</i> (1992)
S095L	93279	92158	373	43.4	I	Unknown	VAC A6L MC109L	1E-114 1E-95	201/372 (54) 162/372 (43)	
S096L	95444	93309	711	82.6	L	Early transcription factor large subunit (VETF)	VAC A7L MC110L	0.0 0.0	495/711 (69) 449/708 (63)	Gershon and Moss (1990)
S097R	95491	96351	286	33.8	E	Intermediate transcription factor (VITF-3)	VAC A8R MC111R	1E-98 3E-68	175/286 (61) 129/291 (44)	Sanz and Moss (1999)
S098L	96581	96348	77	8.7	E/L	Unknown	VAC A9L MC112L	2E-20 1E-12	43/58 (74) 35/58 (60)	

TABLE 1—Continued

ORF ^a	Translation		Size (aa)	MW (kDa) ^c	Prom. type ^d	Function/feature(s)	Homolog(s) ^e	BLASTP score ^f	Amino acid identity ^g	Reference(s) ^h
	Start	Stop ^b								
S099L	99290	96582	902	103.7	L	Major core protein P4a precursor	VAC A10L MC113L	0.0 0.0	433/907 (47) 385/908 (42)	Van Meir and Wittek (1988) Vanslyke <i>et al.</i> (1991)
S100R	99305	100249	314	35.5	L	Unknown	VAC A11R MC114R	2E-81 4E-66	161/323 (49) 140/325 (43)	
S101L	100716	100246	156	16.9	L	Structural protein	MC115L VAC A12L	6E-30 1E-27	75/151 (49) 78/182 (42)	Takahashi <i>et al.</i> (1994)
S102L	100955	100749	68	7.8	L	Structural protein, IMV membrane protein p8	VAC A13L	0.00067	25/69 (36)	Takahashi <i>et al.</i> (1994) Jensen <i>et al.</i> (1996)
S103L	101296	101009	95	10.4	L	IMV membrane protein, assembly factor	VAC A14L MC118L	2E-24 2E-17	52/93 (55) 40/95 (42)	Jensen <i>et al.</i> (1996) Rodriguez <i>et al.</i> (1998)
S104L	101474	101313	53	6.3	L	Unknown	VAC A14.5L ⁱ MC119L	2E-14 ^j 9E-9	34/51 (66) 24/53 (45)	
S105L	101748	101464	94	10.9	L	Unknown	VAC A15L MC120L	9E-20 6E-8	43/93 (46) 33/94 (35)	
S106L	102862	101732	376	43.9	L	35K myristylprotein	VAC A16L MC121L	1E-110 4E-85	185/381 (48) 148/359 (41)	Martin <i>et al.</i> (1997)
S107L	103478	102879	199	22.4	L	IMV membrane protein, morphogenesis factor	VAC A17L MC122L	5E-32 5E-28	76/170 (44) 76/190 (40)	Rodriguez <i>et al.</i> (1995) Krijnse-Locker <i>et al.</i> (1996)
S108R	103493	104929	478	55.2	I	Transcript release factor, ATP-dep. DNA helicase	VAC A18R MC123R	1E-162 1E-128	280/490 (57) 220/459 (47)	Bayliss and Condit (1995) Xiang <i>et al.</i> (1998)
S109L	105131	104910	73	8	L	Unknown	VAC A19L MC124L	9E-17 4E-10	44/72 (61) 35/75 (46)	
S110L	105473	105132	113	13.3	?	Unknown	VAC A21L MC125L	3E-32 3E-19	66/117 (56) 48/111 (43)	
S111R	105472	106770	432	50.8	E	DNA polymerase processivity factor	VAC A20R MC126R	2E-79 1E-45	154/410 (37) 109/383 (28)	McDonald <i>et al.</i> (1997)
S112R	106733	107215	160	18.8	?	Unknown	VAC A22R MC127R	1E-52 1E-33	97/154 (62) 71/142 (50)	
S113R	107243	108400	385	44.9	E	Intermediate transcrip. factor (VITF-3)	VAC A23R MC128R	1E-120 1E-95	221/386 (57) 184/385 (47)	Sanz and Moss (1999)
S114R	108426	111893	1155	132.5	E/L	RNA polymerase subunit (rpo132)	VAC A24R MC129R	0.0 0.0	944/1155 (81) 862/1156 (74)	Hooda-Dhingra <i>et al.</i> (1990) Amegadzie <i>et al.</i> (1991b)
S115L	112448	111909	179	20.6	L	Fusion protein	VAC A27L MC131L/MC133L	2E-4 0.18	28/88 (31) 22/68 (32)	Rodriguez and Esteban (1987) Gong <i>et al.</i> (1990) Rodriguez and Smith (1990)
S116L	112871	112449	140	16.1	L	Unknown	VAC A28L MC134L	1E-30 5E-29	71/148 (47) 59/141 (41)	
S117L	113785	112877	302	35	E	RNA polymerase subunit (rpo35)	VAC A29L MC135L	1E-102 9E-81	172/300 (57) 146/298 (48)	Amegadzie <i>et al.</i> (1991a)
S118L	113984	113754	76	8.5	L	Unknown	VAC A30L MC136L	4E-12 2E-9	35/60 (58) 35/69 (50)	
S119L	114122	114003	39	4.4	L	Unknown				
S120L	114924	114157	255	29.1	E/L	Putative ATPase	MC140L VAC A32L	4E-90 1E-84	158/249 (63) 151/255 (59)	Koonin <i>et al.</i> (1993)
S121R	115011	115529	172	19.2	E	EEV glycoprotein	VAC A33R MC142R	4E-18 0.027	57/190 (30) 33/131 (25)	Roper <i>et al.</i> (1996)
S122R	115539	116054	171	19.8	L	Lectin-like EEV glycoprotein	VAC A34R MC143R	2E-40 8E-4	76/172 (44) 40/173 (23)	Duncan and Smith (1992) McIntosh and Smith (1996)
S123R	116088	116627	179	20.4	E	Unknown	VAC A35R MC145R	2E-21 1E-11	53/172 (30) 43/170 (25)	
S124R	116664	117524	286	32.6	E/L	Unknown	MC144R	9E-12	52/221 (23)	
S125R	117533	118027	164	18.9	?	Unknown	SHP HM1 (?)	0.64	33/143 (23)	Gershon <i>et al.</i> (1989)

TABLE 1—Continued

ORF ^a	Translation		Size (aa)	MW (kDa) ^c	Prom. type ^d	Function/feature(s)	Homolog(s) ^e	BLASTP score ^f	Amino acid identity ^g	Reference(s) ^h
	Start	Stop ^b								
S126R	118067	118882	271	31.5	E	Unknown	VAC A37R MC149R	1E-21 4E-11	69/262 (26) 46/191 (24)	
S127L	120206	118869	445	52.5	?	Type II CPD photolyase (PL)	<i>M. domestica</i> PL MSV235 PL	1E-125 1E-124	212/435 (48) 211/451 (46)	Kato <i>et al.</i> (1994) Afonso <i>et al.</i> (1999)
S128L	121081	120209	290	33	?	CD47-like transmemb. integrin assoc. protein	Human OA3-312 VAC A38L (?)	3E-4 7.0	53/268 (19) 24/99 (24)	Campbell <i>et al.</i> (1992) Sanderson <i>et al.</i> (1996)
S129R	121224	121460	78	8.8	E	Myristylprotein	VAC E7R (?)	0.79	13/50 (26)	Martin <i>et al.</i> (1997)
S130R	121545	121859	104	11.6	?	Unknown				
S131R	121931	122422	163	17.8	L	Cu-Zn superoxide dismutase (SOD)	<i>O. cuniculus</i> SOD VAC A45R	2E-37 4E-06	80/150 (53) 35/199 (29)	Reinecke <i>et al.</i> (1988) Smith <i>et al.</i> (1991b)
S132L	122933	122406	175	20.6	E	Unknown	MC085L (?)	2.1	30/91 (32)	
S133R	123023	124702	559	63.2	L	ATP-dependent DNA ligase	VAC A50R	1E-162	283/557 (50)	Parks <i>et al.</i> (1994)
S134R "S135R"	124813	130632	1939	219.8	E	Unknown (deleted in SFV, retained by MYX)	MC035R	0.0	411/944 (43)	Cameron <i>et al.</i> (1999)
"S136R"						(deleted in SFV, retained by MYX)				Cameron <i>et al.</i> (1999)
S137R	131457	132386	309	35.8	L	Unknown	VAC A51R	8E-34	89/263 (33)	
S138L	133285	132413	290	33.7	E	α 2,3- sialyltransferase (ST)	MYX MST3N <i>M. musculus</i> ST	1E-132 3E-49	225/290 (77) 105/243 (43)	Jackson <i>et al.</i> (1999)
"S139R"						(deleted in SFV, retained by MYX)				Cameron <i>et al.</i> (1999)
S140R	133842	135503	553	63.1	?	Kelch repeat	VAC A55R	7E-65	166/499 (33)	
S141R	135538	136209	223	24.4	E	Fam. of surface antigen/receptor proteins	MYX MA56 <i>X. laevis</i> N-CAM	2E-76 0.029	156/224 (69) 16/47 (34)	Jackson <i>et al.</i> (1999) Tonissen and Krieg (1993)
S142R	136203	137123	306	35.5	E	S/T protein kinase	VAC B1R	1E-75	138/290 (47)	Banham and Smith (1992) Lin <i>et al.</i> (1992)
S143R	137135	137839	234	28.1	L	RING finger protein, localizes to viral factories	CPX D7R	1E-19	60/218 (27)	Brick <i>et al.</i> (1998); Upton <i>et al.</i> (1994) Safronov <i>et al.</i> (1996)
S144R	137878	138777	299	34.7	?	Complement control protein (VCP) precursor (omitted during sequence anotation)	VAC C3L	1E-17	51/160 (31)	Kotwal and Moss (1998b)
S145R										
S146R	138784	139110	108	13.7	E	Virulence factor, secreted protein	VAC N1L	0.025	31/101 (30)	Kotwal and Moss (1988a) Kotwal <i>et al.</i> (1989)
S147R	139169	140032	287	33.5	E	S/T protein kinase?	FPV protein kinase	1E-18	66/259 (25)	Hertig <i>et al.</i> (1997)
S148R	140100	142121	673	77.2	E	Host range, ankyrin-like	VAC B18R	2E-12	119/551 (21)	Mossman <i>et al.</i> (1996a) Smith <i>et al.</i> (1991b)
S149R	142123	143595	490	56.7	E	Host range, ankyrin-like	VAC B4R	8E-12	79/392 (20)	Howard <i>et al.</i> (1991) Mossmann <i>et al.</i> (1996a) Smith <i>et al.</i> (1991b)
"S150R"						(deleted in SFV, retained by MYX)				Cameron <i>et al.</i> (1999)
S151R	144598	145599	333	38.3	E	Serpin ("SERP-2")	VAC C12L	5E-40	114/356 (32)	Petit <i>et al.</i> (1996)

TABLE 1—Continued

ORF ^a	Translation		Size (aa)	MW (kDa) ^c	Prom. type ^d	Function/feature(s)	Homolog(s) ^e	BLASTP score ^f	Amino acid identity ^g	Reference(s) ^h
	Start	Stop ^b								
"S152R"						(deleted in SFV, retained by MYX)				Cameron <i>et al.</i> (1999)
S153R	146088	146693	187	23.1	E	Unknown, RGD motif, may bind metals (see text)	SPV C7L BHV4-IE1	2E-19 3E-10	47/158 (29) 27/82 (32)	Massung <i>et al.</i> (1993) Cameron <i>et al.</i> (1999)
S154L	147428	146784	214	24.2	E	Unknown	VAC M2L	1E-56	109/218 (50)	
S155R	147511	148842	443	50.8	—	Truncated duplicate of S009L, probably inactive	VAC C2L	5E-25	90/376 (23)	

Note. CPX, cowpox virus; FPV, fowlpox virus; MC, molluscum contagiosum virus (subtype-1); MSV, *Melanoplus sanguinipes* entomopoxvirus; MYX, myxoma virus; SHP, sheeppox virus; SPV, swinepox virus; VAC, vaccinia virus (strain Copenhagen unless otherwise noted).

^a The open reading frame numbering system is described in the text. Several numbers were not used to maintain congruence with the myxoma gene notation.

^b Including the last base of the stop codon.

^c Predicted molecular weight for an unmodified polypeptide.

^d E, early; I, intermediate; L, late.

^e Although many poxviral homologs were often detected, normally only the vaccinia and molluscum contagiosum gene homologs are described where such genes exist. When no vaccinia and/or molluscum contagiosum homolog(s) were found, the closest alternative poxvirus and/or cellular homolog was identified. In cases where BLASTP scores exceed 0.1, synteny and amino acid identity were also used to identify homologs.

^f BLAST expectation scores depend upon the size of the sequence database. These BLASTP scores were determined on July 21–23, 1999.

^g Percentages shown in parentheses.

^h See Goebel *et al.* (1990) and Senkevich *et al.* (1996) for a description of the vaccinia and molluscum contagiosum virus genes listed in Table 1.

ⁱ A 14.5L is a small vaccinia gene (53 aa, nucleotides 127,154–127,306) not annotated in M35027. A late promoter motif precedes A14.5L and homologs are also encoded by variola (data not shown).

^j Tblastn score determined August 25–27, 1999.

gene extending beyond the TIR junction and clearly preceded by an early poxviral promoter. Close examination of the two ORFs suggests that S155R is probably a pseudogene in which the N terminus has been disrupted by mutation. This is evidenced by the fact that the identity with S009L can be traced back from the putative ATG start codon to the TIR junction, where it disappears without encountering an in-frame coding element or an upstream promoter. Moreover, the missing portion of S155R encodes only part of the BTB/POZ domain encoded by S009L. Because this domain mediates protein–protein interactions (Bardwell and Treisman, 1994), it seems unlikely that the encoded protein would be functional were it to be expressed.

Repetitive DNA

SFV contains only a single example of simple repetitive DNA. This is ~150 bp of DNA located immediately 3' of the S149R gene and encoding ~30 copies of the pentanucleotide repeat 5' -ATAGA- 3' (nt. 143,592–143,746). The repeat is located within a 1-kb ORF-free gap in the coding sequence from which the SFV homolog of the myxoma M150R gene also appears to have been

deleted (see below). The function of the ATAGA repeat is unknown, although BLASTN searches also detected identical repeats in clones of varied metazoan origin. The repeat is not detected in myxoma virus or in any other poxviruses.

Transcriptional regulation

Conserved sequence motifs were found immediately upstream of many SFV genes, which closely resembled poxviral early and/or late promoter elements. Figure 2 presents histograms illustrating the two conserved motifs, and the relative probability of encountering particular bases at different positions within these two sequence motifs. Although it is not shown in Fig. 2, the TATAAATG late motif also was preceded by a highly A/T rich upstream consensus motif again characteristic of Chordopoxviral late promoters. Using these consensus sequences, we can tentatively identify those genes that are most likely regulated by early, late, or hybrid early/late promoters (Table 1). Intermediate promoters are more difficult to identify due to the limited number encoded by Chordopoxviruses. We do find sequences resembling an intermediate promoter immediately upstream of the




FIG. 2. SFV promoter motifs. Three types of promoter motifs were found immediately upstream of many SFV open reading frames along with a fourth, AT-rich pattern, usually found upstream of the TATAAATG late motif (not shown). The histograms illustrate the frequency of occurrence of each of the four bases within these three conserved motifs. The early and late promoter consensus patterns are found upstream of ~70 and 85 genes, respectively, and the statistics derive from an analysis of these two gene families. The intermediate motif was discovered through an analysis of seven SFV genes whose VAC and MCV gene homologs are thought to be under intermediate promoter control.

seven SFV genes that are homologs of known or putative VAC and MCV intermediate genes [S053R (VAC G8R), S075R (VAC H7R), S087L (VAC D12L), S089L (VAC A1L), S090L (VAC A2L), S095L (VAC A6L), S108R (VAC A18R)] (Fig. 2). However, the consensus is not sufficiently robust to identify other examples with any degree of confidence.

We also noted that a T₅NT motif is usually found near the 3' end of putative SFV early genes. The same T₅NT early-transcription-termination motif appears to be randomly distributed throughout the coding regions of putative late genes. This distribution pattern was first noted in vaccinia virus (Yuen and Moss, 1987) and has been used as a sequence clue that helps to differentiate early and late poxvirus genes. However, we noted numerous exceptions to the rule that T₅NT motifs are not expected near the 5' ends of early genes. The S033R, S034L, S064R, S065R, S068R, S079R, S080R, S085R, S111R, S114R, and S117R genes all encode early or hybrid early/late promoters (Fig. 2) in addition to a "precocious" T₅NT motif. In most such cases, the motif is located a few hundred nucleotides downstream of the probable transcription initiation site. There are no other obvious associated genetic features, and it seems unlikely that we might have erroneously assigned the promoters, since most of these genes encode typical poxvirus early pro-

teins (replicative enzymes and RNA polymerase components). It is difficult to say whether these genes are subject to a special form of transcription regulation or whether the elements are inactive due to sequence context effects. However, it is apparent that one cannot use the distribution of T₅NT motifs to predict Leporipoxvirus gene expression kinetics with confidence.

Similarity among SFV, vaccinia, and molluscum contagiosum proteins

When compared with other fully sequenced non-Leporipoxviral genomes, the individual genes of SFV most closely resemble genes encoded by vaccinia and variola viruses. When these same gene products are compared with those encoded by MCV and MSV, the SFV proteins produced alignments displaying consistently lower alignment scores. This point is illustrated in Fig. 3 where we summarize those SFV genes which generated the 20 highest BLASTP alignment scores. The preponderance of putative transcription and replicative proteins amongst this selected set of SFV genes is also noteworthy, as are the many proteins required for viral assembly. Figure 3 also illustrates a singular exception to the rule that highly conserved SFV genes most closely resemble their Or-




FIG. 3. Conservation of amino acid sequence between SFV, VAC, MCV, and MSV gene products. BLASTP alignment scores were calculated for all of the proteins encoded by SFV and used to identify the 20 gene products showing the highest conservation of amino acid sequence over the greatest peptide distance. The identity of the vaccinia virus gene homolog also is listed to aid in the identification of each SFV gene product. No MSV homologs of S028L, S133R, S020L, or S134R can be detected, or a MCV homolog of S133R (DNA ligase). Most VAR alignment scores are essentially identical to VAC scores and have been omitted for brevity. However, S134R encodes a homolog of the VAR B22R and MC035R gene products, a gene not encoded by vaccinia virus.

thopoxviral homologs. The exception is S134R, which encodes the largest protein in the SFV genome (220 kDa). The S134R gene encodes a transmembrane protein of unknown function and more closely resembles the MC035R rather than the VAR B22R gene product. Moreover there is no VAC homolog of this gene.

The closer resemblance to Orthopoxviridae also can be illustrated by a more comprehensive comparison of the similarity between less highly conserved genes. Figure 4 illustrates a pairwise comparison of BLASTP expectation scores (E-values) obtained through alignments of SFV, VAC, MCV, and MSV gene homologs. A clear trend is noted wherein the logarithm of the expectation scores more closely favours vaccinia over MCV or MSV gene products. Parenthetically this approach provides a method of determining whether viruses such as MCV or MSV really are missing certain genes or whether the presence of the homolog has been obscured by gene-specific patterns of amino acid sequence divergence. We note that points lying along the abscissa indicate SFV gene homologs encoded by vaccinia virus, but apparently absent in MCV and/or MSV. Clearly the greater the distance from the regression line, the greater the statistical likelihood that the gene is truly absent. A more formal analysis of this methodology will be detailed in a later communication.

Vaccinia, VAR, and MCV genes absent in SFV

Most of the SFV genes located within the genomic core interval bounded by S012L and S142R appear to

have syntenic VAC, VAR, and/or MCV homologs. However, there are a few "common" genes whose absence from the SFV genome serves to further delimit the minimal gene complement of Chordopoxviruses (Table 2). One of these genes has recently been shown to be mutable in modified vaccinia strain Ankara (Antoine *et al.*, 1998) and thus the absence of F11L/C15L/MC018L gene homologs is not entirely surprising. SFV also lacks a homolog of the gene encoding a steroid dehydrogenase/isomerase (A44L/A47L/MC152R), although this conserved virulence factor is also not required for growth in culture (Moore and Smith, 1992).

Several absent genes suggest that SFV may not utilize the same infection route(s) as do other Chordopoxviruses. SFV appears to lack homologs of the A-type inclusion proteins. These genes have been shown to be mutable in a variety of Orthopoxviruses, but most viruses produce at least a truncated form of the protein that may promote infection of phagocytic cells by intracellular mature virions (IMV) (Ulaeto *et al.*, 1996). The complete absence of any SFV gene homolog is curious and predicts that Leporipoxvirus infections might produce relatively small amounts of IMV. It is also noteworthy that SFV lacks an A36R/A38R/MC146R gene homolog. Even though the gene is not essential for growth, disruption of VAC A36R creates a small plaque phenotype and reduces extracellular enveloped virus production (EEV) three- to fivefold (Parkinson and Smith, 1994). Recent studies have shown that A36R gene products are found on the surface of intracellular enveloped viruses (IEV)




FIG. 4. Similarities between SFV, VAC, MCV, and MSV genomes. A BLASTP score was calculated for each SFV gene vs its putative VAC-, MCV-, and/or MSV-encoded gene homolog. The negative logarithm of the resulting SFV-VAC score (E-value) then was plotted on the abscissa vs the corresponding MCV (■) or the MSV (◇) score. The data points would be expected to distribute along the line where $\log_{10}(x) = \log_{10}(y)$ were SFV no more closely related to vaccinia than to MCV or MSV. However, most comparisons favour VAC over MCV (i.e., SFV genes more closely resemble vaccinia genes) and greatly favour VAC over MSV. The SFV DNA photolyase gene, S127L, is an obvious exception to this trend. MSV235 encodes a homolog of S127L (Fig. 5), but no such gene seems to be encoded by VAC or MCV.

where they serve to nucleate actin and thus facilitate EEV dissemination. Because SFV encodes homologs of most of the other proteins thought to be required for vaccinia IEV assembly (S022L/F13L, S115L/A27L, S121R/A33R, S122R/A34R, but probably not B5R) (Vazquez *et al.*, 1998; Rottger *et al.*, 1999), it seems likely that although SFV can form intracellular enveloped virions, the virus is not subsequently spread by a route dependent upon actin-driven EEV production. This is certainly consistent with the very small plaques produced by the virus. Experience suggests that most infectious SFV particles are cell associated, but it is presently unknown whether these are IEV, or cell-associated enveloped viruses.

Two additional genes are conserved among VAC, VAR, and MCV even though there appear to be no SFV-encoded homologs. One is A31R/A34R/MC138R and no mutations have yet been shown to map to the gene. Synteny predicts that S119L might encode a homolog of these genes, but the SFV peptide is too small (39 vs 124 amino acids) and shows little resemblance to the Orthopoxviral and Molluscipoxviral proteins. SFV also lacks a homolog of the E11L/E11L/MC041L gene. Vaccinia

E11L encodes a late protein that is encapsidated within the vaccinia virus core, and a conditionally lethal temperature-sensitive mutation has been mapped to the gene (Wang and Shuman, 1996). What biological function is served by VAC E11L and its homologs is uncertain.

Another peculiar feature of the SFV genome is the partial absence of most of the genes involved in nucleotide biosynthesis. In this respect the virus lies midway between vaccinia and variola viruses, which encode several such genes, and MCV, which encodes none at all. For example, SFV seemingly lacks thymidylate and guanylate kinases (VAC A48R and VAC A57R, respectively) while still encoding a thymidine kinase (S061R) (Upton and McFadden, 1986). It also lacks the putative cytidine kinase encoded by fowlpox virus (Koonin and Senkevich, 1993). SFV also encodes only the smaller R2 subunit of the ribonucleotide reductase (S015L = VAC F4L), presumably rendering it dependent upon host cell functions for the synthesis of DNA biosynthetic precursors. A similar situation is observed in human cytomegalovirus except that this virus encodes only the larger R1 subunit (Bankier *et al.*, 1991). Why SFV should retain only one-half of an $\alpha_2\beta_2$ protein is unclear. It has been observed that heterodimers composed of a vaccinia R1 subunit and mouse R2 subunit are catalytically active (Hendricks and Mathews, 1998), and it is quite possible that SFV has

TABLE 2
Selected VAC, VAR, and MCV Genes Missing
from the SFV Chromosome

Virus			Predicted function or properties
VAC	VAR	MCV	
F11L	C15L	MC018L	Non-essential (Antoine <i>et al.</i> , 1998)
E11L	E11L	MC041L	Essential virion component (Wang and Shuman, 1996)
O2L	Q2L	—	Glutaredoxin 1 (Ahn and Moss, 1992a)
I4L	K4L	—	Ribonucleotide reductase (large subunit) (Tengelsen <i>et al.</i> , 1988)
—	—	MC066L	Glutathione peroxidase (Shisler <i>et al.</i> , 1998)
A25L	A26L	MC130L	A-type inclusion protein (Funahashi <i>et al.</i> , 1998)
A26L	A30L	MC131L	A-type inclusion protein (Funahashi <i>et al.</i> , 1988)
A31R	A34R	MC138R	Function unknown
A36R	A38R	MC146R	Intracellular enveloped virion protein (Rottger <i>et al.</i> , 1999)
A44L	A47L	MC152R	Steroid dehydrogenase/isomerase (Moore and Smith, 1992)
A48R	J2R	—	Thymidylate kinase (Smith <i>et al.</i> , 1991b)
A57R	J8R	—	Guanylate kinase (Smith <i>et al.</i> , 1991b)

TABLE 3

SFV Genes Exhibiting Shared Structural Motifs

Family	Genes	Comment ^a
Ankyrin-like repeat	S005L/R	5 ankyrin-like repeats
	S148R	5 ankyrin-like repeats
	S149R	4 ankyrin-like repeats
Kelch repeat	S006L/R	4 kelch motifs plus BTB/POZ domain
	S008L/R	4 kelch motifs plus BTB/POZ domain
	S009L	4 kelch motifs plus BTB/POZ domain
	S0014L	4 kelch motifs plus BTB/POZ domain
	S140R	6 kelch motifs plus BTB/POZ domain
	S155R	4 kelch motifs plus a portion of BTB/POZ domain
C7L-like proteins	S062R	VAC C7L homolog
	S063R	VAC C7L homolog (distant)
	S064R	VAC C7L homolog

^a The number of repeat motifs varies depending upon cut-off score. We set E = 10 when searching the Pfam HMM (v. 4.1) database.

evolved to exploit the opposite arrangement consisting of the SFV R2 and host R1 subunits (C. Mathews, personal communication). If this was the case, one would expect the small subunit to be the one retained by SFV because this is the subunit that, in vaccinia virus, permits a virus-specific interaction with the replicative single-strand DNA binding protein (Davis and Mathews, 1993). The DNA sequence suggests that S015L encodes a catalytically active protein because the gene still encodes the residues required to coordinate a diferric iron centre as well as the tyrosine involved in stable free radical formation (Y₁₁₀) and all, except the very last, of the residues postulated to transport electrons from the centre of the R2 dimer to the nucleotide-binding R1 dimer (Jordan and Reichard, 1998). If S015L does encode an active protein, the electron donor remains to be identified because SFV lacks a gene homologous to VAC O2L. The O2L-encoded glutaredoxin 1 protein is thought to donate electrons to VAC ribonucleotide reductase (Rajagopal *et al.*, 1995) and yet the only glutaredoxin encoded by SFV (S048L) most closely resembles the glutaredoxin 2 protein encoded by VAC G4L.

SFV gene families

Several SFV genes share common structural features that permit their assignment to viral gene families. These are summarized in Table 3. Perhaps most striking are the number of encoded proteins displaying ankyrin-like and kelch motif repeats. Also noteworthy are the three homologs of VAC C7L encoded by SFV genes S062R, S063R, and S064R. Nothing is known concerning the

function of these SFV genes, but the C7L gene product is required for vaccinia replication in certain cell types and therefore has been categorized as a host-range function (Oguiura *et al.*, 1993). SFV has a very limited host range, growing only on rabbit and a few selected primate cells, and it is thus somewhat surprising that it encodes such an abundance of putative host-range genes.

Leporipoxvirus genes not encoded by other poxviruses

A Leporipoxvirus-specific SFV gene is difficult to define because novelty will depend upon the size of the poxvirus sequence database and upon the cut-off point for alignment scores. Furthermore, alignment scores are dependent upon protein size, and one can miss homologous genes if they happen to be very small. [This point is clearly illustrated by comparing BLASTP and amino acid identity scores for ORFs S008L and S008.1L (Table 1)]. We have taken a conservative approach that uses a combination of semiquantitative factors to decide whether a gene should be considered unique to Leporipoxviruses. These factors include the BLASTP expectation score (which should normally exceed 0.01–0.1 except in the case of very small proteins) and no evidence of a possible syntenic relationship between the SFV gene and other poxvirus genes. The degree of amino acid identity also was taken into consideration, with amino acid identities exceeding 20–25% being considered probable evidence of homology. By these criteria SFV and myxoma encode a number of genes seemingly unique to Leporipoxviruses that are summarized in Table 4. Several small hypothetical genes were discovered whose size renders similarity searches problematical (Table 4). A disproportionate number of these new proteins are predicted to encode transmembrane helices including S023R, S119L, S125R, and S132L, but little else is known of their function. S130R encodes an 11.6-kDa

TABLE 4
Novel Leporipoxvirus Genes

SFV gene	Predicted or known encoded function
S013L	Resembles DNA binding domain of HIN-200 protein family
S017L	Possibly a homolog of VAC F7L
S023R	Unknown, predicted transmembrane helix
S119L	Unknown, predicted transmembrane helix
S125R	Unknown, predicted transmembrane helix
S127L	Type II CPD photolyase
S130R	Unknown
S132L	Unknown, predicted transmembrane helices
S138L	α 2,3-sialyltransferase (Jackson, Hall, and Kerr, 1999)
S141R	Ig-like protein superfamily, predicted transmembrane helices

[illegible]

FIG. 5. ClustalW alignment of the S127L-encoded protein with other Type II CPD photolyases. GenBank protein sequence IDs were: g4049783 (MSV235), g1079459 (*Monodelphis domestica*), g1401039 (*Potorous tridactylus*), g464378 (*Carassius auratus*), g1401038 (*Oryzias latipes*), g1079066 (*Drosophila melanogaster*), g1617219 (*Arabidopsis thaliana*), g1197525 (*Myxococcus xanthus*), and g639674 (*Methanobacterium thermoautotrophicum*). Black boxes indicate identical residues found at ≥ 9 sites amongst the 10 aligned proteins.

protein displaying no significant similarity to any other protein in the sequence databases.

The S013L gene encodes a small protein (12,729 Da, $pI = 7.91$) whose gene expression appears to be regulated by a canonical poxvirus early promoter. The gene product resembles the N-terminal DNA-binding portion of proteins belonging to the hemopoietic IFN-inducible nuclear 200-amino acid repeat (HIN-200) protein family (Dawson and Trapani, 1995, 1996). Expression of these proteins can be induced by interferons α and γ , and one protein family member, IFI16, has been shown to be a transcriptional repressor that may regulate myelopoiesis (Johnstone *et al.*, 1998). An interesting feature of S013L is that it appears to have lost the 200 amino acid repeats that are required if IFI16 is to repress transcription but still encodes a putative SV40 T-antigen-like nuclear localization signal (PKKQRNR) near the C terminus (Mattaj and Englmeier, 1998). Although the function of S013L is uncertain, its structure suggests that the encoded protein could compete with the activity of certain transcriptional repressors and perhaps interfere in the differentiation of hemopoietic cells.

Perhaps the most unusual protein encoded by Lepo-

riporipoxviruses is the S127L gene product. S127L encodes a 52.5-kDa protein that is clearly homologous to proteins belonging to the family of type II cyclobutane pyrimidine dimer (CPD) photolyases (Fig. 5). Photolyases catalyze a light-dependent reaction that returns CPD adducts back to their component pyrimidines and thus reverses the DNA damage caused by ultraviolet light. Although a similar gene is encoded by MSV (Afonso *et al.*, 1999), this is the first DNA repair gene of its kind found to be encoded by a Chordopoxvirus or, indeed, any mammalian virus. The gene is particularly curious in that whilst marsupials such as the common opossum (*Monodelphis domestica*) are known to encode photolyases (Yasui *et al.*, 1994), no placental mammal has yet been shown to have retained the gene. Because it is generally believed that poxviruses originally acquired most genes from their hosts, this discovery raises interesting questions concerning the evolutionary history of a New World virus now thought to infect only rabbits. Preliminary experiments show that the gene can complement an *Escherichia coli phr* (photolyase deficient) mutation *in vivo* (D. Willier, M. Moon, and D. Evans, unpublished data) and imply that Leporipoxviruses can repair the damage

SFV	1	MNNRACPRYYEFNKVNYKAVCMIRGND-IRGVIHFQEQLQNDIVVVFQVILGLTEGFHGIN
BOV	1	-----MATKAVCVLKGDPVOGTIHFQAKG-DIVVVTGSIITGLTEGDHGFH
VAC	1	-----MAVCIIDHDN-IRGVIYFEPVH-GKDKVLGSGVIGLKSCTYSLI
SFV	60	VHEFGDACG--NELGEHYNPEKSSHCSPSDVHRHVGDGLGNIHASKYGVSVVHIIDGKITL
BOV	46	VHOFGDNTQGCTSAQPHFNPLSKKHGGPKDEERHVGDGLGNVTADKNGVAIVDIVDPLISL
VAC	42	IHRYGDISQG-----CDSTGSP--E--I-FIGNIFVNRVGVAVV-YLDTDVNI
SFV	118	TCDRSIIGRSLAVHEREDDLGRGPNVDSSIHGNSGNVIAWGIIGVA-
BOV	106	SGEYSIIGRTMVVHEKPDLLGRGNGNEESTKTGNAGSRLACGVIGTAK
VAC	84	S---TIIGKALSISKNDQRLACGVIGISYINEKIIHFLTINENGV--

FIG. 6. ClustalW alignment of the S131R-encoded protein with bovine and vaccinia superoxide dismutases. GenBank protein sequence IDs were: g134601 (bovine Cu/Zn SOD) and g134630 (VAC A45R gene product). The bovine SOD active site arginine is residue R₁₄₂.

caused by ultraviolet light exposure either before and/or during viral infection.

Two additional SFV genes are homologous to previously identified myxoma virus genes, but Leporipoxviruses otherwise appear to be the only poxviruses encoding such genes. The S138L gene is predicted to encode a 34-kDa α 2,3-sialyltransferase nearly identical to the myxoma enzyme recently characterized by Jackson and colleagues (Jackson *et al.*, 1999). SFV sialyltransferase presumably catalyzes an identical reaction and likewise promotes viral virulence. The S141R gene product also has a previously identified myxoma gene homolog (MA56) (Jackson *et al.*, 1999). S141R encodes a 24-kDa protein that is predicted to contain two transmembrane helices and resembles the immunoglobulin-like domains of a large family of eukaryotic cell-surface proteins. Further commentary on the possible function of these two Leporipoxvirus gene products can be found in the accompanying paper (Cameron *et al.*, 1999).

Other unusual genes—SFV superoxide dismutase (S131R) and S153R

Although all of the sequenced Chordopoxviruses encode genes resembling some type of superoxide dismutase (SOD), the SFV-encoded superoxide dismutase homolog is peculiar in just how closely it still resembles mammalian cytoplasmic Cu/Zn superoxide dismutases. Figure 6 shows an alignment of the SFV gene product along with the proteins encoded by vaccinia virus (Smith *et al.*, 1991b) and by cattle (Steinman *et al.*, 1974). A 1.9 Å high-resolution X-ray crystal structure of bovine erythrocyte SOD provides insights into the structure of these proteins (Rypniewski *et al.*, 1995). The SFV gene encodes a nearly intact SOD, whereas the vaccinia A45R gene product has lost portions of the protein encoding all of the bovine protein's Zn-binding domain plus several small surface peptide loops (Fig. 6). Still retained by both proteins are many of the glycine residues which, in the crystal structure, form the SOD dimer interface. Thus the poxvirus proteins are potentially capable of forming heterodimers with cellular SOD or of complexing with a

cellular copper chaperone (Casareno *et al.*, 1998). Closer examination of the SFV gene suggests that while SFV SOD must have some biological function, the protein is unlikely to still catalyze superoxide dismutation. This is suggested by an R-to-V substitution at the SOD active site (residue V₁₅₄ in the SFV gene), and by preliminary experiments which have thus far failed to detect any residual superoxide dismutase activity in either vaccinia or SFV recombinant proteins (A. Winter, M. Moon, and D. Evans, unpublished results). Why so many poxviruses encode seemingly "inactive" superoxide dismutase homologs remains an intriguing and yet elusive question.

S153R encodes a 23-kDa gene that resembles swinepox C7L and a portion of some gamma herpes virus IE1 proteins (Table 1). The protein encodes a highly acidic tail (pI = 4.5) as well as an RGD cell-adhesion motif. The alignment between S153R and C7L spans most of the two genes, although the RGD motif and C-terminal acidic tail are unique features of S153R (data not shown). A common property of S153R and C7L is that both may be metal-binding proteins. For example S153R encodes an imperfect zinc-finger motif (Prosite PDOC00449), and it is this region of the protein that aligns with a zinc-finger motif encoded by a family of alternatively spliced herpes virus immediate-early proteins. C7L encodes a similar cysteine-rich motif characteristic of metallothioneins (Prosite PDOC00180). What function might be served by S153R is uncertain.

Similarities and differences between Shope fibroma and myxoma viruses

A later publication will focus on the relationship between these two viruses in greater detail. At this point, we will simply highlight some of the more obvious genetic rearrangements and speculate as to the function of some of the SFV genes that have been subjected to these mutational processes. The probable function of genes found in myxoma virus, but seemingly mutated in SFV, is more appropriately discussed in the accompanying paper (Cameron *et al.*, 1999).

SFV and myxoma viruses encode a nearly identical

TABLE 5
Genetic Differences between SFV and Myxoma Viruses

SFV gene	Myxoma gene	SFV arrangement	Myxoma gene properties ^a
—	M000.5L/R	No trace of ORF	—
S008.1L/R	M008.1L/R	Fragmentary remains	—
S009L/S155R	M009L	Partially duplicated	Single gene
—	M135R	Fragmentary remains	—
—	M136R	Fragmentary remains	—
—	M139R	Fragmentary remains	—
—	M150R	Fragmentary remains	—
—	M152R	Fragmentary remains	—
S153R	M153R/M153.1R	SPV C7L homolog	Probably disrupted by frameshift
S008.2L/R	M156R	Duplicated gene	Single gene

^a Myxoma gene properties are described in the accompanying paper (Cameron *et al.*, 1999).

complement of homologous genes. The similarity is sufficiently high that it was used in our analysis of the two viruses to differentiate start codons from fortuitous in-frame ATG codons, identify frameshifting sequence errors, and detect genes encoding very small peptides. It also permitted using a single common system of gene nomenclature. Despite this high degree of sequence conservation, SFV and myxoma are still clearly distinct viruses as judged by the criteria currently used to categorize poxvirus "species" (Douglas and Dumbell, 1996). This can be demonstrated by examining phylogenetic trees constructed using the sequence of such widely distributed poxvirus proteins as the thymidine kinase. Such phylogenies show that there is a greater degree of sequence divergence between SFV and myxoma than between most Orthopoxviruses (data not shown).

Closer inspection of the two genomes detects a few gene rearrangements and a number of genes deleted in SFV relative to myxoma (Table 5). The terminal inverted repeats (TIR) encode several of these changes. First, SFV lacks a homolog of a small gene encoded at the very ends of the myxoma genome about which nothing is yet known (M000.5L/R). Myxoma also encodes a 41,557-Da serpin (M008.1L/R, SERP-1) (Upton *et al.*, 1990a) of which only 6356 Da of fragmentary remains can still be detected in SFV (S008.1L/R). Both SFV and myxoma still encode divergent homologs of the VAC K3L gene (S008.2L/R and M156R, respectively), but the SFV gene is duplicated within the TIR whereas myxoma (like vaccinia) encodes only a single copy of the gene located immediately adjacent to the boundary with the myxoma right TIR (Cameron *et al.*, 1999). The function of these Leporipoxvirus K3L gene homologs is unknown. However, the VAC protein resembles the N-terminal portion of eIF-2 α and inhibits interferon activity by preventing phosphorylation of eIF-2 α (Beattie *et al.*, 1991; Carroll *et al.*, 1993). A somewhat similar rearrangement also pertains to S009L/S155R. These genes are partially duplicated in SFV, as discussed previously, whereas myxoma encodes

only a single gene homolog located immediately adjacent to the left TIR border (M009L) (Cameron *et al.*, 1999).

Although these telomeric rearrangements appear complex, they can be fairly easily rationalized. We suggest that the TIRs of a hypothetical "ancestral" Leporipoxvirus once most closely resembled that of SFV, except that these terminal inverted repeats were longer than modern SFV TIRs and still encoded two complete copies of the genes now called S/M009L, S008.2L/R, and S/M008.1L/R. SFV, strain Kasza, has likely undergone a partial deletion of the hypothetical ancestral "S009R" gene (leaving a pseudogene now called S155R) and a second excision of most of the S/M008.1L/R gene homolog by a well documented, but still poorly understood, mirror-deletion mechanism (McFadden and Dales, 1979). Myxoma, strain Lausanne, has probably suffered a complete deletion of the ancestral "M009R" gene, followed by a deletion of the ancestor of the gene now called S008.2L from the left TIR. The resulting myxoma TIR now encodes only one copy of M009L, and M156R actually derives from a gene originally syntenic with S008.2R. These genetic models are outlined in Fig. 7.

Elsewhere in the SFV genome we find deletions that seem to have eradicated a further five genes relative to myxoma virus and a frameshift in myxoma that seems to have disrupted that virus's homolog of an SFV gene. Most of these rearrangements are located toward the right end of the virus, which seems to be an unusually dynamic poxvirus genetic region. The missing genes are the SFV homologs of myxoma M135R, M136R, M139R, M150R, and M152R. That these are SFV deletions, and not myxoma gene insertions, is suggested by the presence of DNA fragments in SFV that still exhibit patchy DNA homology to the genes retained by myxoma virus. No simple pattern of repeats or other sequence motifs can be discerned at the putative break points in the myxoma genome, but sequence drift during the course of SFV and myxoma evolution may have long since erased such features (data not shown). This is in contrast to the




FIG. 7. Hypothetical rearrangements underlying the structure of SFV and myxoma telomeres. SFV and myxoma are proposed to have derived from a common ancestor strain that once encoded a longer TIR (arrows) and a greater complement of duplicated genes. Sequential deletions coupled with sequence drift have obscured the hypothetical common origin of the genes now called M156R and S008.2L/R (hatched boxes). The precise order in which these events might have taken place cannot be predicted with certainty.

small deletions observed in vaccinia and variola that sometimes still show evidence of having occurred between small directly repeated sequences (Shchelkunov and Totmenin, 1995). As described above, the S153R gene encodes a protein of unknown function that resembles swinepox virus C7L as well as a portion of genes encoded by some herpes viruses. A frameshift appears to have split the myxoma homolog of S153R into two open-reading frames, somewhere near the middle of the gene.

Inspection of the genes altered in SFV and myxoma viruses provides only limited insights into the altered virulence of SFV compared with myxoma. The most obvious deficiency is that SFV seems to be missing the two serpin-like genes encoded by M008.1L/R and M152R, one of which (M008.1) is known to be important for virulence (Upton *et al.*, 1990a; Cameron *et al.*, 1999). The

biological and genetic properties of the other missing genes are less obvious. One of the mutated genes ("S135R") resembled a portion of the VAC B19R putative interleukin-1 receptor (Smith and Chan, 1991), two others ("S136R" and "S139R") resembled VAC A52R, and "S150R" appears to have once encoded several ankyrin-like repeats as do three other genes still retained by SFV (Table 3) (Cameron *et al.*, 1999). The impact these genetic alterations might have on Leporipoxvirus virulence is discussed further in the accompanying paper (Cameron *et al.*, 1999).

In conclusion, the DNA sequence of SFV provides further insights into the evolutionary history and minimal essential gene complement of Chordopoxviruses. Through comparative analysis of the genome sequences of myxoma and Shope fibroma viruses we expect to gain insights into how the two viruses can cause such dis-

tinctly different diseases. Furthermore this work has identified all of the genes that are collectively responsible for catalyzing high-frequency recombination and *trans*-replication in SFV-infected cells. The discovery of a novel DNA repair gene and a curious new superoxide dismutase homolog also will provide fascinating insights into how poxviruses avoid the genotoxic effects of ultraviolet light and protect themselves from the DNA, lipid, and protein-damaging effects of reactive oxygen species.

MATERIALS AND METHODS

SFV, myxoma virus, and other genome sequences

The completed nucleotide sequence of SFV has been submitted to GenBank and assigned Accession No. AF170722. The sequence of myxoma virus can be retrieved from GenBank using Accession No. AF170726 (Cameron *et al.*, 1999). Where we have compared gene sequences "vaccinia" refers to the Copenhagen strain (GenBank ID M35027), "variola" refers to variola major strain Bangladesh-75 (GenBank ID L22579), MCV refers to Molluscum contagiosum virus subtype 1 (GenBank ID U60315), and MSV refers to Melanoplus sanguinipes entomopoxvirus (GenBank ID AF063866), unless otherwise indicated.

Materials, cell, and virus culture

SIRC cells and the Kasza strain of SFV (Kasza, 1974) were originally obtained from the American Type Culture Collection. SIRC cells were grown at 37° in Dulbecco's modified Eagle's medium (DMEM) (Gibco BRL) supplemented with 1% nonessential amino acids and 10% fetal bovine serum (CanSera) in a 5% CO₂ atmosphere.

Virus purification and DNA extraction

SFV virions were isolated essentially as described (Dales and Mosbach, 1968). Briefly, SIRC cells were infected with virus at a m.o.i. of 1 and incubated for 3 days at 37°C, and cell-associated virus particles ($\sim 1.5 \times 10^{10}$ plaque forming units) were extracted by Dounce homogenization at 4°C in 80 mL of 1:10 diluted phosphate-buffered saline. The cell debris were removed by low-speed centrifugation (1000 *g* for 10 min at 4°C), resuspended in 40 mL of serum-free DMEM, sonicated, and recentrifuged, and the two supernatants pooled. The virus was recovered by centrifugation (7000 *g* for 10 min at 4°C), treated with trypsin (15 min at 37°C), diluted to 10 mL with serum-containing DMEM, pelleted again, and resuspended in 2 mL of buffer containing 0.25M sucrose and 10 mM Tris—HCl pH 8.5. Viral particles then were applied to the top of a 20–50% (w/v) potassium tartrate gradient, and centrifuged at 32,000 *g* for 2.5 h at 4°C. The virus band was retrieved by side puncture, diluted two-fold with water, centrifuged at 13,000 *g* for 30 min at 4°C,

and the viral pellet resuspended in 40 mL of buffer (10 mM Tris—HCl, 1 mM EDTA, pH 8.5). Virion proteins were eliminated by overnight digestion with 0.6 mg/mL proteinase K at 37°C in the presence of 0.1% (w/v) sodium dodecyl sulfate. High-molecular-weight viral DNA was recovered by phenol-chloroform extraction followed by precipitation with ethanol.

Acquisition of primary sequence data and sequence assembly

Random DNAs were produced by fragmenting virus DNA on ice with a probe sonicator. The ends of the sheared DNA fragments were repaired using T4 and Klenow polymerases, and then fragments ranging in size from 1.5 to 3.0 kb were separated by electrophoresis and recovered. A pBlueScript II (KS+) plasmid vector (Stratagene) was digested with *EcoRV* and treated with shrimp alkaline phosphatase, ligated to the SFV DNA fragments using T4 ligase, and transformed into *E. coli* JM105 cells (Promega). Suitably sized plasmid clones were identified using the PCR and purified using a "High Pure" plasmid kit (Roche), and the double-stranded templates were sequenced from both ends of the insert using T3 and T7 primers. A cycle-sequencing protocol, AmpliTaq DNA polymerase (Perkin-Elmer), dye terminators, and an Applied Biosystems PRISM 377 sequencer that had been modified to provide extended sequence reads were used to obtain all of the raw sequence data. Data were collected using ABI data collection (v 1.1) and DNA sequencing software (v 3.3). As contigs started to assemble, we switched to a more efficient primer walking strategy using, as templates, a library of *Bam*HI restriction fragments previously cloned from SFV strain Kasza (Wills *et al.*, 1983; Delange *et al.*, 1984) plus custom sequencing primers (Gibco BRL). Primers were designed using RightPrimer (v 1.2.5).

Sequence traces were manually edited to correct errors in automated base calls and to trim the ill-defined sequences found at the 5' and 3' ends of each fragment. The final contig then was assembled from these fragmentary elements using the program Sequencer (v 3.1). After a completed sequence had been acquired, each base call within the assembly was manually inspected, and several portions of the genome were resequenced on the opposite strand, using new sequencing primers, to confirm the integrity of the sequence. This final sequence was overdetermined with about threefold redundancy at each base pair from 565 separate sequence fragments. Although portions of the SFV genome have been previously sequenced, including the TIRs, TIR junctions, and several internal *Bam*HI restriction fragments, we nevertheless resequenced all of these regions to confirm the accuracy of the final assembly. We did not resequence the very end of the virus. As defined by the an *Afl*II site located at the hairpin centre of inverted

repeat symmetry (DeLange *et al.*, 1986), the first base of our contig is located either 34 or 40 bases from the end of the genome (nt 125 or 198 in GenBank entry M14003.1). The exact distance depends upon the number of intervening extrahelical bases.

Sequence analysis and assignment of open-reading frames (ORFs)

The SFV genome is generally syntenic with the genomes of Orthopoxviridae so the open reading frames were easily identified using a combination of BLAST analyses (Altschul *et al.*, 1997) and visual inspection. In some locations, we encountered gaps between genes that were too large to be occupied by the accompanying poxviral promoters (30–40 bp), and this was taken to be indicative of the possible presence of a small ORF. Where such ORFs were found on both SFV and myxoma viruses, they were provided with gene designations. The gene numbering starts from the revised left end of the virus [the right end of the virus in the original restriction map (DeLange *et al.*, 1984)] and throughout most of the genome was incremented in steps of one. Where SFV clearly lacks a gene found in myxoma virus at the syntenic position (e.g., "S139") we have skipped that number to maintain consistency in the naming of homologous genes (Table 1).

Putative early and late promoters were identified using internet implementations of MEME version 2.2 (Bailey and Elkan, 1994) and MAST version 2.2 (Bailey and Gribskov, 1998). An iterative process was used whereby small collections of putative early and late promoter elements were used as training sets in a search for additional examples of conserved sequence motifs. Once genes had been roughly categorized as encoding either or both promoter motifs, histograms were assembled illustrating the likelihood of encountering particular bases at given positions within each conserved motif. BLAST scores (Altschul *et al.*, 1997) were calculated using NCBI internet resources and sequence alignments prepared using Macintosh or internet versions of MACAW (Schuler *et al.*, 1991) and Clustal W (Thompson *et al.*, 1994). Phylogenies were calculated using Felsenstein's PHYLIP package of software (Felsenstein, 1989). Protein motifs were identified by searches against the Pfam HMM database.

ACKNOWLEDGMENTS

We especially thank Ms. Angela Hollis for performing the hundreds of DNA sequencing reactions upon which this work depended. We also thank J. Barrett, C. Cameron, L. Chen, J. Cao, S. Hota-Mitchell, C. Macaulay, and J. Tyrone for advice, assistance, and commentary on this work during the process of sequence assembly and annotation, and C. Mathews for providing helpful insights into aspects of this work. This study was supported by operating grants awarded to D.E. and G.M. from the Medical Research Council of Canada and to G.M. from the National Cancer Institute of Canada. G.M. is a Medical Research

Council of Canada Senior Scientist and D.W. is a recipient of an Ontario Graduate Scholarship in Science and Technology. The College of Biological Sciences DNA sequencing facility at the University of Guelph is partially supported by National Sciences and Engineering Research Council, ORDCF, and CFI funds.

REFERENCES

- Afonso, C. L., Tulman, E. R., Lu, Z., Oma, E., Kutish, G. F., and Rock, D. L. (1999). The genome of *Melanoplus sanguinipes* entomopoxvirus. *J. Virol* **73**, 533–552.
- Ahn, B. Y., Gershon, P. D., Jones, E. V., and Moss, B. (1990a). Identification of rpo30, a vaccinia virus RNA polymerase gene with structural similarity to a eucaryotic transcription elongation factor. *Mol. Cell. Biol.* **10**, 5433–5441.
- Ahn, B. Y., Jones, E. V., and Moss, B. (1990b). Identification of the vaccinia virus gene encoding an 18-kilodalton subunit of RNA polymerase and demonstration of a 5' poly(A) leader on its early transcript. *J. Virol* **64**, 3019–3024.
- Ahn, B. Y., and Moss, B. (1992a). Glutaredoxin homolog encoded by vaccinia virus is a virion-associated enzyme with thioltransferase and dehydroascorbate reductase activities. *Proc. Natl. Acad. Sci. USA* **89**, 7060–7064.
- Ahn, B. Y., and Moss, B. (1992b). RNA polymerase-associated transcription specificity factor encoded by vaccinia virus. *Proc. Natl. Acad. Sci. USA* **89**, 3536–3540.
- Ahn, B. Y., Rosel, J., Cole, N. B., and Moss, B. (1992). Identification and expression of rpo19, a vaccinia virus gene encoding a 19-kilodalton DNA-dependent RNA polymerase subunit. *J. Virol* **66**, 971–982.
- Altschul, S. F., Madden, T. L., Schaffer, A. A., Zhang, J., Zhang, Z., Miller, W., and Lipman, D. J. (1997). Gapped BLAST and PSI-BLAST: A new generation of protein database search programs. *Nucleic Acids Res.* **25**, 3389–4302.
- Amegadzie, B. Y., Ahn, B. Y., and Moss, B. (1991a). Identification, sequence, and expression of the gene encoding a Mr 35000 subunit of the vaccinia virus DNA-dependent RNA polymerase. *J. Biol. Chem.* **266**, 13712–13718.
- Amegadzie, B. Y., Ahn, B. Y., and Moss, B. (1992). Characterization of a 7-kilodalton subunit of vaccinia virus DNA-dependent RNA polymerase with structural similarities to the smallest subunit of eukaryotic RNA polymerase II. *J. Virol* **66**, 3003–3010.
- Amegadzie, B. Y., Holmes, M. H., Cole, N. B., Jones, E. V., Earl, P. L., and Moss, B. (1991b). Identification, sequence, and expression of the gene encoding the second-largest subunit of the vaccinia virus DNA-dependent RNA polymerase. *Virology* **180**, 88–98.
- Antoine, G., Scheifflinger, F., Dorner, F., and Falkner, F. G. (1998). The complete genomic sequence of the modified vaccinia Ankara strain: Comparison with other orthopoxviruses. *Virology* **244**, 365–396.
- Bailey, T. L., and Elkan, C. (1994). "Fitting a mixture model by expectation maximization to discover motifs in biopolymers." Proceedings of the Second International Conference on Intelligent Systems for Molecular Biology. AAAI Press, Menlo Park.
- Bailey, T. L., and Gribskov, M. (1998). Combining evidence using p-values: Application to sequence homology searches. *Bioinformatics* **14**, 48–54.
- Banham, A. H., and Smith, G. L. (1992). Vaccinia virus gene B1R encodes a 34-kDa serine/threonine protein kinase that localizes in cytoplasmic factories and is packaged into virions. *Virology* **191**, 803–812.
- Bankier, A. T., Beck, S., Bohni, R., Brown, C. M., Cerny, R., Chee, M. S., Hutchison, C. A. d., Kouzarides, T., Martignetti, J. A., Preddie, E., et al. (1991). The DNA sequence of the human cytomegalovirus genome. *DNA Seq.* **2**, 1–12.
- Bardwell, V. J., and Treisman, R. (1994). The POZ domain: A conserved protein-protein interaction motif. *Genes Dev.* **8**, 1664–1677.
- Barry, M., Hnatiuk, S., Mossman, K., Lee, S. F., Boshkov, L., and McFadden, G. (1997). The myxoma virus M-T4 gene encodes a novel

- RDEL-containing protein that is retained within the endoplasmic reticulum and is important for the productive infection of lymphocytes. *Virology* **239**, 360–377.
- Bayliss, C. D., and Condit, R. C. (1995). The vaccinia virus A18R gene product is a DNA-dependent ATPase. *J. Biol. Chem.* **270**, 1550–1556.
- Bayliss, C. D., and Smith, G. L. (1997). Vaccinia virion protein VP8, the 25 kDa product of the L4R gene, binds single-stranded DNA and RNA with similar affinity. *Nucleic Acids Res.* **25**, 3984–3990.
- Beattie, E., Tartaglia, J., and Paoletti, E. (1991). Vaccinia virus-encoded eIF-2 alpha homolog abrogates the antiviral effect of interferon. *Virology* **183**, 419–422.
- Black, E. P., Moussatche, N., and Condit, R. C. (1998). Characterization of the interactions among vaccinia virus transcription factors G2R, A18R, and H5R. *Virology* **245**, 313–322.
- Block, W., Upton, C., and McFadden, G. (1985). Tumorigenic poxviruses: Genomic organization of malignant rabbit virus, a recombinant between Shope fibroma virus and myxoma virus. *Virology* **140**, 113–124.
- Brick, D. J., Burke, R. D., Schiff, L., and Upton, C. (1998). Shope fibroma virus RING finger protein N1R binds DNA and inhibits apoptosis. *Virology* **249**, 42–51.
- Broyles, S. S., and Fesler, B. S. (1990). Vaccinia virus gene encoding a component of the viral early transcription factor. *J. Virol.* **64**, 1523–1529.
- Broyles, S. S., and Moss, B. (1986). Homology between RNA polymerases of poxviruses, prokaryotes, and eukaryotes: Nucleotide sequence and transcriptional analysis of vaccinia virus genes encoding 147-kDa and 22-kDa subunits. *Proc. Natl. Acad. Sci. USA* **83**, 3141–3145.
- Broyles, S. S., and Pennington, M. J. (1990). Vaccinia virus gene encoding a 30-kilodalton subunit of the viral DNA-dependent RNA polymerase. *J. Virol.* **64**, 5376–5382.
- Cabirac, G. F., Strayer, D. S., Sell, S., and Leibowitz, J. L. (1985). Characterization, molecular cloning, and physical mapping of the Shope fibroma virus genome. *Virology* **143**, 663–670.
- Cameron, C., Macaulay, C., Cao, J., Barrett, J., Hota-Mitchell, S., Willer, D. O., Evans, D. H., and McFadden, G. (In press). The complete genome sequence of myxoma virus. *Virology*
- Campbell, I. G., Freemont, P. S., Foulkes, W., and Trowsdale, J. (1992). An ovarian tumor marker with homology to vaccinia virus contains an IgV-like region and multiple transmembrane domains. *Cancer Res* **52**, 5416–5420.
- Carroll, K., Elroy-Stein, O., Moss, B., and Jagus, R. (1993). Recombinant vaccinia virus K3L gene product prevents activation of double-stranded RNA-dependent, initiation factor 2 alpha-specific protein kinase. *J. Biol. Chem.* **268**, 12837–12842.
- Casareno, R. L., Waggoner, D., and Gitlin, J. D. (1998). The copper chaperone CCS directly interacts with copper/zinc superoxide dismutase. *J. Biol. Chem.* **273**, 23625–23628.
- Chang, H. W., Uribe, L. H., and Jacobs, B. L. (1995). Rescue of vaccinia virus lacking the E3L gene by mutants of E3L. *J. Virol.* **69**, 6605–6608.
- Chang, W., Macaulay, C., Hu, S. L., Tam, J. P., and McFadden, G. (1990). Tumorigenic poxviruses: Characterization of the expression of an epidermal growth factor related gene in Shope fibroma virus. *Virology* **179**, 926–930.
- Chang, W., Upton, C., Hu, S. L., Purchio, A. F., and McFadden, G. (1987). The genome of Shope fibroma virus, a tumorigenic poxvirus, contains a growth factor gene with sequence similarity to those encoding epidermal growth factor and transforming growth factor alpha. *Mol. Cell. Biol.* **7**, 535–540.
- Chertov, O., Telezhinskaya, I. N., Zaitseva, E. V., Golubeva, T. B., Zinov'ev, V. V., Ovechkina, L. G., Mazkova, L. B., and Malygin, E. G. (1991). Amino acid sequence determination of vaccinia virus immunodominant protein p35 and identification of the gene. *Biomed. Sci.* **2**, 151–154.
- Christen, L. M., Sanders, M., Wiler, C., and Niles, E. G. (1998). Vaccinia virus nucleoside triphosphate phosphohydrolase I is an essential viral early gene transcription termination factor. *Virology* **245**, 360–371.
- Cudmore, S., Blasco, R., Vincentelli, R., Esteban, M., Sodeik, B., Griffiths, G., and Krijnse Locker, J. (1996). A vaccinia virus core protein, p39, is membrane associated. *J. Virol.* **70**, 6909–6921.
- Dales, S., and Mosbach, E. H. (1968). Vaccinia as a model for membrane biogenesis. *Virology* **35**, 564–583.
- Davies, M. V., Elroy-Stein, O., Jagus, R., Moss, B., and Kaufman, R. J. (1992). The vaccinia virus K3L gene product potentiates translation by inhibiting double-stranded-RNA-activated protein kinase and phosphorylation of the alpha subunit of eukaryotic initiation factor 2. *J. Virol.* **66**, 1943–1950.
- Davis, R. E., and Mathews, C. K. (1993). Acidic C terminus of vaccinia virus DNA-binding protein interacts with ribonucleotide reductase. *Proc. Natl. Acad. Sci. USA* **90**, 745–749.
- Dawson, M. J., and Trapani, J. A. (1995). The interferon-inducible autoantigen, IFI 16: Localization to the nucleolus and identification of a DNA-binding domain. *Biochem. Biophys. Res. Commun.* **214**, 152–162.
- Dawson, M. J., and Trapani, J. A. (1996). HIN-200: A novel family of IFN-inducible nuclear proteins expressed in leukocytes. *J. Leukocyte Biol.* **60**, 310–316.
- Delange, A. M., Macaulay, C., Block, W., Mueller, T., and McFadden, G. (1984). Tumorigenic poxviruses: Construction of the composite physical map of the Shope fibroma virus genome. *J. Virol.* **50**, 408–416.
- DeLange, A. M., and McFadden, G. (1986). Sequence-nonspecific replication of transfected plasmid DNA in poxvirus-infected cells. *Proc. Natl. Acad. Sci. USA* **83**, 614–618.
- DeLange, A. M., Reddy, M., Scraba, D., Upton, C., and McFadden, G. (1986). Replication and resolution of cloned poxvirus telomeres in vivo generates linear minichromosomes with intact viral hairpin termini. *J. Virol.* **59**, 249–259.
- Demkowicz, W. E., Maa, J. S., and Esteban, M. (1992). Identification and characterization of vaccinia virus genes encoding proteins that are highly antigenic in animals and are immunodominant in vaccinated humans. *J. Virol.* **66**, 386–398.
- Derrien, M., Punjabi, A., Khanna, M., Grubisha, O., and Traktman, P. (1999). Tyrosine phosphorylation of A17 during vaccinia virus infection: Involvement of the H1 phosphatase and the F10 kinase. *J. Virol.* **73**, 7287–7296.
- Digiorgio, R. F., and Mare, C. J. (1994). Viral disease. In "The Biology of the Laboratory Rabbit" (P. J. E. A. Manning, Ed.), 2nd Ed., pp. 171–197. Academic Press, San Diego.
- Douglas, N. J., and Dumbell, K. R. (1996). DNA sequence variation as a clue to the phylogenesis of orthopoxviruses. *J. Gen. Virol.* **77**, 947–951.
- Duncan, S. A., and Smith, G. L. (1992). Identification and characterization of an extracellular envelope glycoprotein affecting vaccinia virus egress. *J. Virol.* **66**, 1610–1621.
- Dyster, L. M., and Niles, E. G. (1991). Genetic and biochemical characterization of vaccinia virus genes D2L and D3R which encode virion structural proteins. *Virology* **182**, 455–467.
- Earl, P. L., Jones, E. V., and Moss, B. (1986). Homology between DNA polymerases of poxviruses, herpesviruses, and adenoviruses: Nucleotide sequence of the vaccinia virus DNA polymerase gene. *Proc. Natl. Acad. Sci. USA* **83**, 3659–3663.
- Evans, D. H., Stuart, D., and McFadden, G. (1988). High levels of genetic recombination among cotransfected plasmid DNAs in poxvirus-infected mammalian cells. *J. Virol.* **62**, 367–375.
- Evans, E., Klemperer, N., Ghosh, R., and Traktman, P. (1995). The vaccinia virus D5 protein, which is required for DNA replication, is a nucleic acid-independent nucleoside triphosphatase. *J. Virol.* **69**, 5353–5361.
- Felsenstein, J. (1989). PHYLIP—Phylogeny Inference Package (Version 3.2). *Cladistics* **5**, 164–166.
- Fisher, C., Parks, R. J., Lauzon, M. L., and Evans, D. H. (1991). Hetero-

- duplex DNA formation is associated with replication and recombination in poxvirus-infected cells. *Genetics* **129**, 7–18.
- Franke, C. A., Wilson, E. M., and Hruby, D. E. (1990). Use of a cell-free system to identify the vaccinia virus L1R gene product as the major late myristylated virion protein M25. *J. Virol.* **64**, 5988–5996.
- Funahashi, S., Sato, T., and Shida, H. (1988). Cloning and characterization of the gene encoding the major protein of the A-type inclusion body of cowpox virus. *J. Gen. Virol.* **69**, 35–47.
- Gershon, P. D., Ahn, B. Y., Garfield, M., and Moss, B. (1991). Poly(A) polymerase and a dissociable polyadenylation stimulatory factor encoded by vaccinia virus. *Cell* **66**, 1269–1278.
- Gershon, P. D., Ansell, D. M., and Black, D. N. (1989). A comparison of the genome organization of capripoxvirus with that of the orthopoxviruses. *J. Virol.* **63**, 4703–4708.
- Gershon, P. D., and Black, D. N. (1989). A capripoxvirus pseudogene whose only intact homologs are in other poxvirus genomes. *Virology* **172**, 350–354.
- Gershon, P. D., and Moss, B. (1990). Early transcription factor subunits are encoded by vaccinia virus late genes. *Proc. Natl. Acad. Sci. USA* **87**, 4401–4405.
- Goebel, S. J., Johnson, G. P., Perkus, M. E., Davis, S. W., Winslow, J. P., and Paoletti, E. (1990). The complete DNA sequence of vaccinia virus. *Virology* **179**, 247–266, 517–563.
- Gong, S. C., Lai, C. F., and Esteban, M. (1990). Vaccinia virus induces cell fusion at acid pH and this activity is mediated by the N-terminus of the 14-kDa virus envelope protein. *Virology* **178**, 81–91.
- Graham, K. A., Lalani, A. S., Macen, J. L., Ness, T. L., Barry, M., Liu, L. Y., Lucas, A., Clark-Lewis, I., Moyer, R. W., and McFadden, G. (1997). The T1/35kDa family of poxvirus-secreted proteins bind chemokines and modulate leukocyte influx into virus-infected tissues. *Virology* **229**, 12–24.
- Graham, K. A., Opgenorth, A., Upton, C., and McFadden, G. (1992). Myxoma virus M11L ORF encodes a protein for which cell surface localization is critical in manifestation of viral virulence. *Virology* **191**, 112–124.
- Gribaudo, G., Ravaglia, S., Guandalini, L., Riera, L., Gariglio, M., and Landolfo, S. (1997). Molecular cloning and expression of an interferon-inducible protein encoded by gene 203 from the gene 200 cluster. *Eur. J. Biochem.* **249**, 258–264.
- Gvakharia, B. O., Koonin, E. K., and Mathews, C. K. (1996). Vaccinia virus G4L gene encodes a second glutaredoxin. *Virology* **226**, 408–411.
- Hendricks, S. P., and Mathews, C. K. (1998). Allosteric regulation of vaccinia virus ribonucleotide reductase, analyzed by simultaneous monitoring of its four activities. *J. Biol. Chem.* **273**, 29512–29518.
- Hertig, C., Coupar, B. E., Gould, A. R., and Boyle, D. B. (1997). Field and vaccine strains of fowlpox virus carry integrated sequences from the avian retrovirus, reticuloendotheliosis virus. *Virology* **235**, 367–376.
- Hirt, P., Hiller, G., and Wittek, R. (1986). Localization and fine structure of a vaccinia virus gene encoding an envelope antigen. *J. Virol.* **58**, 757–764.
- Hooda-Dhingra, U., Patel, D. D., Pickup, D. J., and Condit, R. C. (1990). Fine structure mapping and phenotypic analysis of five temperature-sensitive mutations in the second largest subunit of vaccinia virus DNA-dependent RNA polymerase. *Virology* **174**, 60–69.
- Howard, S. T., Chan, Y. S., and Smith, G. L. (1991). Vaccinia virus homologues of the Shope fibroma virus inverted terminal repeat proteins and a discontinuous ORF related to the tumor necrosis factor receptor family. *Virology* **180**, 633–647.
- Jackson, R. J., and Bults, H. G. (1990). A myxoma virus nucleotide sequence with homology to the vaccinia virus RNA polymerase 22-kDa subunit gene. *Nucleic Acids Res.* **18**, 5290.
- Jackson, R. J., and Hall, D. F. (1998). The myxoma virus EcoRI-O fragment encodes the DNA binding core protein and the major envelope protein of extracellular poxvirus. *Virus Genes* **17**, 55–62.
- Jackson, R. J., Hall, D. F., and Kerr, P. J. (1999). Myxoma virus encodes an alpha2,3-sialyltransferase that enhances virulence. *J. Virol.* **73**, 2376–2384.
- Jensen, O. N., Houthaeve, T., Shevchenko, A., Cudmore, S., Ashford, T., Mann, M., Griffiths, G., and Krijnse Locker, J. (1996). Identification of the major membrane and core proteins of vaccinia virus by two-dimensional electrophoresis. *J. Virol.* **70**, 7485–7497.
- Johnson, G. P., Goebel, S. J., and Paoletti, E. (1993). An update on the vaccinia virus genome. *Virology* **196**, 381–401.
- Johnstone, R. W., Kerry, J. A., and Trapani, J. A. (1998). The human interferon-inducible protein, IFI 16, is a repressor of transcription. *J. Biol. Chem.* **273**, 17172–17177.
- Jordan, A., and Reichard, P. (1998). Ribonucleotide reductases. *Annu. Rev. Biochem.* **67**, 71–98.
- Kane, E. M., and Shuman, S. (1993). Vaccinia virus morphogenesis is blocked by a temperature-sensitive mutation in the I7 gene that encodes a virion component. *J. Virol.* **67**, 2689–2698.
- Kao, S. Y., Rensner, E., Kates, J., and Bauer, W. R. (1981). Purification and characterization of a superhelix binding protein from vaccinia virus. *Virology* **111**, 500–508.
- Karstad, L., Thorsen, J., Davies, G., and Kaminjolo, J. S. (1977). Poxvirus fibromas on African hares. *J. Wildl. Dis.* **13**, 245–247.
- Kasza, L. (1974). Isolation and characterization of a rabbit fibroma virus from a naturally occurring tumor. *Am. J. Vet. Res.* **35**, 87–89.
- Kato, T., Jr., Todo, T., Ayaki, H., Ishizaki, K., Morita, T., Mitra, S., and Ikenaga, M. (1994). Cloning of a marsupial DNA photolyase gene and the lack of related nucleotide sequences in placental mammals. *Nucleic Acids Res.* **22**, 4119–4124.
- Keck, J. G., Baldick, C. J., Jr., and Moss, B. (1990). Role of DNA replication in vaccinia virus gene expression: A naked template is required for transcription of three late trans-activator genes. *Cell* **61**, 801–809.
- Keck, J. G., Kovacs, G. R., and Moss, B. (1993). Overexpression, purification, and late transcription factor activity of the 17-kilodalton protein encoded by the vaccinia virus A1L gene. *J. Virol.* **67**, 5740–5748.
- Klemperer, N., Ward, J., Evans, E., and Traktman, P. (1997). The vaccinia virus I1 protein is essential for the assembly of mature virions. *J. Virol.* **71**, 9285–9294.
- Koonin, E. V., and Senkevich, T. G. (1992). Vaccinia virus encodes four putative DNA and/or RNA helicases distantly related to each other. *J. Gen. Virol.* **73**, 989–993.
- Koonin, E. V., and Senkevich, T. G. (1993). Fowlpox virus encodes a protein related to human deoxycytidine kinase: Further evidence for independent acquisition of genes for enzymes of nucleotide metabolism by different viruses. *Virus Genes* **7**, 289–295.
- Koonin, E. V., Senkevich, T. G., and Chernos, V. I. (1993). Gene A32 product of vaccinia virus may be an ATPase involved in viral DNA packaging as indicated by sequence comparisons with other putative viral ATPases. *Virus Genes* **7**, 89–94.
- Kotwal, G. J., Hugin, A. W., and Moss, B. (1989). Mapping and insertional mutagenesis of a vaccinia virus gene encoding a 13800-Da secreted protein. *Virology* **171**, 579–587.
- Kotwal, G. J., and Moss, B. (1988a). Analysis of a large cluster of nonessential genes deleted from a vaccinia virus terminal transposition mutant. *Virology* **167**, 524–537.
- Kotwal, G. J., and Moss, B. (1988b). Vaccinia virus encodes a secretory polypeptide structurally related to complement control proteins. *Nature* **335**, 176–178.
- Kovacs, G. R., and Moss, B. (1996). The vaccinia virus H5R gene encodes late gene transcription factor 4: Purification, cloning, and overexpression. *J. Virol.* **70**, 6796–6802.
- Krijnse-Locker, J., Schleich, S., Rodriguez, D., Goud, B., Snijder, E. J., and Griffiths, G. (1996). The role of a 21-kDa viral membrane protein in the assembly of vaccinia virus from the intermediate compartment. *J. Biol. Chem.* **271**, 14950–14958.
- Li, S. J., and Hochstrasser, M. (1999). A new protease required for cell-cycle progression in yeast. *Nature* **398**, 246–251.
- Lin, S., and Broyles, S. S. (1994). Vaccinia protein kinase 2: A second essential serine/threonine protein kinase encoded by vaccinia virus. *Proc. Natl. Acad. Sci. USA* **91**, 7653–7657.

- Lin, S., Chen, W., and Broyles, S. S. (1992). The vaccinia virus B1R gene product is a serine/threonine protein kinase. *J. Virol.* **66**, 2717–2723.
- Macen, J. L., Graham, K. A., Lee, S. F., Schreiber, M., Boshkov, L. K., and McFadden, G. (1996). Expression of the myxoma virus tumor necrosis factor receptor homologue and M11L genes is required to prevent virus-induced apoptosis in infected rabbit T lymphocytes. *Virology* **218**, 232–237.
- Martin, K. H., Grosenbach, D. W., Franke, C. A., and Hruby, D. E. (1997). Identification and analysis of three myristylated vaccinia virus late proteins. *J. Virol.* **71**, 5218–5226.
- Massung, R. F., Jayarama, V., and Moyer, R. W. (1993). DNA sequence analysis of conserved and unique regions of swinepox virus: Identification of genetic elements supporting phenotypic observations including a novel G protein-coupled receptor homologue. *Virology* **197**, 511–528.
- Massung, R. F., Liu, L. I., Qi, J., Knight, J. C., Yuran, T. E., Kerlavage, A. R., Parsons, J. M., Venter, J. C., and Esposito, J. J. (1994). Analysis of the complete genome of smallpox variola major virus strain Bangladesh-1975. *Virology* **201**, 215–240.
- Mattaj, J. W., and Englmeier, L. (1998). Nucleocytoplasmic transport: The soluble phase. *Annu. Rev. Biochem.* **67**, 265–306.
- McDonald, W. F., Klemperer, N., and Traktman, P. (1997). Characterization of a processive form of the vaccinia virus DNA polymerase. *Virology* **234**, 168–175.
- McFadden, G., and Dales, S. (1979). Biogenesis of poxviruses: Mirror-image deletions in vaccinia virus DNA. *Cell* **18**, 101–108.
- McIntosh, A. A., and Smith, G. L. (1996). Vaccinia virus glycoprotein A34R is required for infectivity of extracellular enveloped virus. *J. Virol.* **70**, 272–281.
- Meis, R. J., and Condit, R. C. (1991). Genetic and molecular biological characterization of a vaccinia virus gene which renders the virus dependent on isatin-beta-thiosemicarbazone (IBT). *Virology* **182**, 442–454.
- Moore, J. B., and Smith, G. L. (1992). Steroid hormone synthesis by a vaccinia enzyme: A new type of virus virulence factor. *EMBO J.* **11**, 1973–1980.
- Mossman, K., Lee, S. F., Barry, M., Boshkov, L., and McFadden, G. (1996a). Disruption of M-T5, a novel myxoma virus gene member of poxvirus host range superfamily, results in dramatic attenuation of myxomatosis in infected European rabbits. *J. Virol.* **70**, 4394–4410.
- Mossman, K., Nation, P., Macen, J., Garbutt, M., Lucas, A., and McFadden, G. (1996b). Myxoma virus M-T7, a secreted homolog of the interferon-gamma receptor, is a critical virulence factor for the development of myxomatosis in European rabbits. *Virology* **215**, 17–30.
- Mossman, K., Ostergaard, H., Upton, C., and McFadden, G. (1995). Myxoma virus and Shope fibroma virus encode dual-specificity tyrosine/serine phosphatases which are essential for virus viability. *Virology* **206**, 572–582.
- Niles, E. G., Condit, R. C., Caro, P., Davidson, K., Matusick, L., and Seto, J. (1986). Nucleotide sequence and genetic map of the 16-kb vaccinia virus HindIII D fragment. *Virology* **153**, 96–112.
- Niles, E. G., Lee-Chen, G. J., Shuman, S., Moss, B., and Broyles, S. S. (1989). Vaccinia virus gene D12L encodes the small subunit of the viral mRNA capping enzyme. *Virology* **172**, 513–522.
- Niles, E. G., and Seto, J. (1988). Vaccinia virus gene D8 encodes a virion transmembrane protein. *J. Virol.* **62**, 3772–3778.
- Ogawa, R., Calvert, J. G., Yanagida, N., and Nazerian, K. (1993). Insertional inactivation of a fowlpox virus homologue of the vaccinia virus F12L gene inhibits the release of enveloped virions. *J. Gen. Virol.* **74**, 55–64.
- Oguiura, N., Spehner, D., and Drillien, R. (1993). Detection of a protein encoded by the vaccinia virus C7L open reading frame and study of its effect on virus multiplication in different cell lines. *J. Gen. Virol.* **74**, 1409–1413.
- Opgenorth, A., Nation, N., Graham, K., and McFadden, G. (1993). Transforming growth factor alpha, Shope fibroma growth factor, and vaccinia growth factor can replace myxoma growth factor in the induction of myxomatosis in rabbits. *Virology* **192**, 701–709.
- Palaniyar, N., Fisher, C., Parks, R., and Evans, D. H. (1996). SFV topoisomerase: Sequence specificity in a genetically mapped interval. *Virology* **221**, 351–354.
- Palaniyar, N., Gerasimopoulos, E., and Evans, D. H. (1999). Shope fibroma virus DNA topoisomerase catalyses holliday junction resolution and hairpin formation in vitro. *J. Mol. Biol.* **287**, 9–20.
- Parkinson, J. E., and Smith, G. L. (1994). Vaccinia virus gene A36R encodes a M(r) 43–50 K protein on the surface of extracellular enveloped virus. *Virology* **204**, 376–390.
- Parks, R. J., and Evans, D. H. (1991a). Effect of marker distance and orientation on recombinant formation in poxvirus-infected cells. *J. Virol.* **65**, 1263–1272.
- Parks, R. J., and Evans, D. H. (1991b). Enhanced recombination associated with the presence of insertion and deletion mutations in poxvirus-infected cells. *Virology* **184**, 299–309.
- Parks, R. J., Lichty, B. D., Karakis, C., and Evans, D. H. (1994). Characterization of the Shope fibroma virus DNA ligase gene. *Virology* **202**, 642–650.
- Parks, R. J., Winchcombe-Forhan, C., DeLange, A. M., Xing, X., and Evans, D. H. (1998). DNA ligase gene disruptions can depress viral growth and replication in poxvirus-infected cells. *Virus Res.* **56**, 135–147.
- Patel, A. H., Gaffney, D. F., Subak-Sharpe, J. H., and Stow, N. D. (1990). DNA sequence of the gene encoding a major secreted protein of vaccinia virus, strain Lister. *J. Gen. Virol.* **71**, 2013–2021.
- Perkus, M. E., Goebel, S. J., Davis, S. W., Johnson, G. P., Limbach, K., Norton, E. K., and Paoletti, E. (1990). Vaccinia virus host range genes. *Virology* **179**, 276–286.
- Petit, F., Bertagnoli, S., Gelfi, J., Fassy, F., Boucraut-Baralon, C., and Milon, A. (1996). Characterization of a myxoma virus-encoded serpin-like protein with activity against interleukin-1 beta-converting enzyme. *J. Virol.* **70**, 5860–5866.
- Quick, S. D., and Broyles, S. S. (1990). Vaccinia virus gene D7R encodes a 20000-dalton subunit of the viral DNA-dependent RNA polymerase. *Virology* **178**, 603–605.
- Rajagopal, I., Ahn, B. Y., Moss, B., and Mathews, C. K. (1995). Roles of vaccinia virus ribonucleotide reductase and glutaredoxin in DNA precursor biosynthesis. *J. Biol. Chem.* **270**, 27415–27418.
- Reinecke, K., Wolf, B., Michelson, A. M., Puget, K., Steffens, G. J., and Flohe, L. (1988). The amino-acid sequence of rabbit Cu-Zn superoxide dismutase. *Biol. Chem. Hoppe Seyler* **369**, 715–725.
- Rodriguez, D., Esteban, M., and Rodriguez, J. R. (1995). Vaccinia virus A17L gene product is essential for an early step in virion morphogenesis. *J. Virol.* **69**, 4640–4648.
- Rodriguez, J. F., and Esteban, M. (1987). Mapping and nucleotide sequence of the vaccinia virus gene that encodes a 14-kilodalton fusion protein. *J. Virol.* **61**, 3550–3554.
- Rodriguez, J. F., and Smith, G. L. (1990). IPTG-dependent vaccinia virus: Identification of a virus protein enabling virion envelopment by Golgi membrane and egress. *Nucleic Acids Res.* **18**, 5347–5351.
- Rodriguez, J. R., Risco, C., Carrascosa, J. L., Esteban, M., and Rodriguez, D. (1998). Vaccinia virus 15-kilodalton (A14L) protein is essential for assembly and attachment of viral crescents to viroosomes. *J. Virol.* **72**, 1287–1296.
- Roper, R. L., Payne, L. G., and Moss, B. (1996). Extracellular vaccinia virus envelope glycoprotein encoded by the A33R gene. *J. Virol.* **70**, 3753–3762.
- Rosel, J., and Moss, B. (1985). Transcriptional and translational mapping and nucleotide sequence analysis of a vaccinia virus gene encoding the precursor of the major core polypeptide 4b. *J. Virol.* **56**, 830–838.
- Roseman, N. A., Evans, R. K., Mayer, E. L., Rossi, M. A., and Slabaugh, M. B. (1996). Purification and characterization of the vaccinia virus deoxyuridine triphosphatase expressed in *Escherichia coli*. *J. Biol. Chem.* **271**, 23506–23511.

- Roseman, N. A., and Slabaugh, M. B. (1990). The vaccinia virus *HindIII* F fragment: Nucleotide sequence of the left 6.2 kb. *Virology* **178**, 410–418.
- Rottger, S., Frischknecht, F., Reckmann, I., Smith, G. L., and Way, M. (1999). Interactions between vaccinia virus IEV membrane proteins and their roles in IEV assembly and actin tail formation. *J. Virol.* **73**, 2863–2875.
- Rypniewski, W. R., Mangani, S., Bruni, B., Orioli, P. L., Casati, M., and Wilson, K. S. (1995). Crystal structure of reduced bovine erythrocyte superoxide dismutase at 1.9 Å resolution. *J. Mol. Biol.* **251**, 282–296.
- Safronov, P. F., Petrov, N. A., Riazankina, O. I., Totmenin, A. V., Shchelkunov, S. N., and Sandakhchiev, L. S. (1996). [Genes of a circle of hosts for the cowpox virus]. *Dokl. Akad. Nauk.* **349**, 829–833.
- Sanderson, C. M., Parkinson, J. E., Hollinshead, M., and Smith, G. L. (1996). Overexpression of the vaccinia virus A38L integral membrane protein promotes Ca²⁺ influx into infected cells. *J. Virol.* **70**, 905–914.
- Sanz, P., and Moss, B. (1999). Identification of a transcription factor, encoded by two vaccinia virus early genes, that regulates the intermediate stage of viral gene expression. *Proc. Natl. Acad. Sci. USA* **96**, 2692–2697.
- Schnierle, B. S., Gershon, P. D., and Moss, B. (1992). Cap-specific mRNA (nucleoside-O²'-)-methyltransferase and poly(A) polymerase stimulatory activities of vaccinia virus are mediated by a single protein. *Proc. Natl. Acad. Sci. USA* **89**, 2897–2901.
- Schuler, G. D., Altschul, S. F., and Lipman, D. J. (1991). A workbench for multiple alignment construction and analysis. *Proteins* **9**, 180–190.
- Scott, C. B., Holdbrook, R., and Sell, S. (1981). Cell-mediated immune response to Shope fibroma virus-induced tumors in adult rabbits. *J. Natl. Cancer Inst.* **66**, 681–689.
- Sell, S., and Scott, C. B. (1981). An immunohistologic study of Shope fibroma virus in rabbits: Tumor rejection by cellular reaction in adults and progressive systemic reticuloendothelial infection in neonates. *J. Natl. Cancer Inst.* **66**, 363–373.
- Senkevich, T. G., Bugert, J. J., Sisler, J. R., Koonin, E. V., Darai, G., and Moss, B. (1996). Genome sequence of a human tumorigenic poxvirus: Prediction of specific host response-evasion genes. *Science* **273**, 813–816.
- Senkevich, T. G., Koonin, E. V., Bugert, J. J., Darai, G., and Moss, B. (1997). The genome of molluscum contagiosum virus: Analysis and comparison with other poxviruses. *Virology* **233**, 19–42.
- Shchelkunov, S. N., Massung, R. F., and Esposito, J. J. (1995). Comparison of the genome DNA sequences of Bangladesh-1975 and India-1967 variola viruses. *Virus Res* **36**, 107–118.
- Shchelkunov, S. N., and Totmenin, A. V. (1995). Two types of deletions in orthopoxvirus genomes. *Virus Genes* **9**, 231–245.
- Shisler, J. L., Senkevich, T. G., Berry, M. J., and Moss, B. (1998). Ultraviolet-induced cell death blocked by a selenoprotein from a human dermatotropic poxvirus [see comments]. *Science* **279**, 102–105.
- Shope, R. E. (1932a). A filterable virus causing a tumor-like condition in rabbits and its relationship to virus myxomatosis. *J. Exp. Med.* **56**, 803–822.
- Shope, R. E. (1932b). A transmissible tumor-like condition of rabbits. *J. Exp. Med.* **56**, 793–802.
- Shors, T., Keck, J. G., and Moss, B. (1999). Down regulation of gene expression by the vaccinia virus D10 protein. *J. Virol.* **73**, 791–796.
- Shuman, S. (1992). Vaccinia virus RNA helicase: An essential enzyme related to the DE-H family of RNA-dependent NTPases. *Proc. Natl. Acad. Sci. USA* **89**, 10935–10939.
- Slabaugh, M., Roseman, N., Davis, R., and Mathews, C. (1988). Vaccinia virus-encoded ribonucleotide reductase: Sequence conservation of the gene for the small subunit and its amplification in hydroxyurea-resistant mutants. *J. Virol.* **62**, 519–527.
- Smith, C. A., Davis, T., Anderson, D., Solam, L., Beckmann, M. P., Jerzy, R., Dower, S. K., Cosman, D., and Goodwin, R. G. (1990). A receptor for tumor necrosis factor defines an unusual family of cellular and viral proteins. *Science* **248**, 1019–1023.
- Smith, C. A., Davis, T., Wignall, J. M., Din, W. S., Farrah, T., Upton, C., McFadden, G., and Goodwin, R. G. (1991a). T2 open reading frame from the Shope fibroma virus encodes a soluble form of the TNF receptor. *Biochem. Biophys. Res. Commun.* **176**, 335–342.
- Smith, G. L., and Chan, Y. S. (1991). Two vaccinia virus proteins structurally related to the interleukin-1 receptor and the immunoglobulin superfamily. *J. Gen. Virol.* **72**, 511–518.
- Smith, G. L., Chan, Y. S., and Howard, S. T. (1991b). Nucleotide sequence of 42 kbp of vaccinia virus strain WR from near the right inverted terminal repeat. *J. Gen. Virol.* **72**, 1349–1376.
- Sodeik, B., Griffiths, G., Ericsson, M., Moss, B., and Doms, R. W. (1994). Assembly of vaccinia virus: Effects of rifampin on the intracellular distribution of viral protein p65. *J. Virol.* **68**, 1103–1114.
- Steinman, H. M., Naik, V. R., Abernethy, J. L., and Hill, R. L. (1974). Bovine erythrocyte superoxide dismutase. Complete amino acid sequence. *J. Biol. Chem.* **249**, 7326–7338.
- Strayer, D. S., Jerng, H. H., and O'Connor, K. (1991). Sequence and analysis of a portion of the genomes of Shope fibroma virus and malignant rabbit fibroma virus that is important for viral replication in lymphocytes. *Virology* **185**, 585–595.
- Strayer, D. S., Skaletsky, E., Cabirac, G. F., Sharp, P. A., Corbeil, L. B., Sell, S., and Leibowitz, J. L. (1983). Malignant rabbit fibroma virus causes secondary immunosuppression in rabbits. *J. Immunol.* **130**, 399–404.
- Stuart, D. T., Upton, C., Higman, M. A., Niles, E. G., and McFadden, G. (1993). A poxvirus-encoded uracil DNA glycosylase is essential for virus viability. *J. Virol.* **67**, 2503–2512.
- Takahashi, T., Oie, M., and Ichihashi, Y. (1994). N-terminal amino acid sequences of vaccinia virus structural proteins. *Virology* **202**, 844–852.
- Tartaglia, J., and Paoletti, E. (1985). Physical mapping and DNA sequence analysis of the rifampicin resistance locus in vaccinia virus. *Virology* **147**, 394–404.
- Tengelsen, L. A., Slabaugh, M. B., Bibler, J. K., and Hruby, D. E. (1988). Nucleotide sequence and molecular genetic analysis of the large subunit of ribonucleotide reductase encoded by vaccinia virus. *Virology* **164**, 121–131.
- Thompson, J. D., Higgins, D. G., and Gibson, T. J. (1994). CLUSTAL W: Improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* **22**, 4673–4680.
- Tonissen, K. F., and Krieg, P. A. (1993). Two neural-cell adhesion molecule (NCAM)-encoding genes in *Xenopus laevis* are expressed during development and in adult tissues. *Gene* **127**, 243–247.
- Traktman, P., Sridhar, P., Condit, R. C., and Roberts, B. E. (1984). Transcriptional mapping of the DNA polymerase gene of vaccinia virus. *J. Virol.* **49**, 125–131.
- Ulaeto, D., Grosenbach, D., and Hruby, D. E. (1996). The vaccinia virus 4c and A-type inclusion proteins are specific markers for the intracellular mature virus particle. *J. Virol.* **70**, 3372–3377.
- Upton, C., DeLange, A. M., and McFadden, G. (1987). Tumorigenic poxviruses: Genomic organization and DNA sequence of the telomeric region of the Shope fibroma virus genome. *Virology* **160**, 20–30.
- Upton, C., Macen, J. L., Maranchuk, R. A., DeLange, A. M., and McFadden, G. (1988). Tumorigenic poxviruses: Fine analysis of the recombination junctions in malignant rabbit fibroma virus, a recombinant between Shope fibroma virus and myxoma virus. *Virology* **166**, 229–239.
- Upton, C., Macen, J. L., Wishart, D. S., and McFadden, G. (1990a). Myxoma virus and malignant rabbit fibroma virus encode a serpin-like protein important for virus virulence. *Virology* **179**, 618–631.
- Upton, C., and McFadden, G. (1986). Identification and nucleotide sequence of the thymidine kinase gene of Shope fibroma virus. *J. Virol.* **60**, 920–927.
- Upton, C., Mossman, K., and McFadden, G. (1992). Encoding of a homolog of the IFN-γ receptor by myxoma virus. *Science* **258**, 1369–1372.

- Upton, C., Opgenorth, A., Traktman, P., and McFadden, G. (1990b). Identification and DNA sequence of the Shope fibroma virus DNA topoisomerase gene. *Virology* **176**, 439–447.
- Upton, C., Schiff, L., Rice, S. A., Dowdeswell, T., Yang, X., and McFadden, G. (1994). A poxvirus protein with a RING finger motif binds zinc and localizes in virus factories. *J. Virol.* **68**, 4186–4195.
- Upton, C., Stuart, D., and McFadden, G. (1991). Identification and DNA sequence of the large subunit of the capping enzyme from Shope fibroma virus. *Virology* **183**, 773–777.
- Upton, C., Stuart, D. T., and McFadden, G. (1993). Identification of a poxvirus gene encoding a uracil DNA glycosylase. *Proc. Natl. Acad. Sci. USA* **90**, 4518–4522.
- Van Meir, E., and Wittek, R. (1988). Fine structure of the vaccinia virus gene encoding the precursor of the major core protein 4 a. *Arch. Virol.* **102**, 19–27.
- Vanslyke, J. K., Whitehead, S. S., Wilson, E. M., and Hruby, D. E. (1991). The multistep proteolytic maturation pathway utilized by vaccinia virus P4a protein: A degenerate conserved cleavage motif within core proteins. *Virology* **183**, 467–478.
- Vazquez, M. I., Rivas, G., Cregut, D., Serrano, L., and Esteban, M. (1998). The vaccinia virus 14-kilodalton (A27L) fusion protein forms a triple coiled-coil structure and interacts with the 21-kilodalton (A17L) virus membrane protein through a C-terminal alpha-helix. *J. Virol.* **72**, 10126–10137.
- Vos, J. C., Sarker, M., and Stunnenberg, H. G. (1991). Vaccinia virus capping enzyme is a transcription initiation factor. *EMBO J.* **10**, 2553–2558.
- Wang, S. P., and Shuman, S. (1996). A temperature-sensitive mutation of the vaccinia virus E11 gene encoding a 15-kDa virion component. *Virology* **216**, 252–257.
- Weinrich, S. L., and Hruby, D. E. (1986). A tandemly-oriented late gene cluster within the vaccinia virus genome. *Nucleic Acids Res.* **14**, 3003–3016.
- Whitehead, S. S., and Hruby, D. E. (1994). A transcriptionally controlled trans-processing assay: Putative identification of a vaccinia virus-encoded proteinase which cleaves precursor protein P25K. *J. Virol.* **68**, 7603–7608.
- Wills, A., Delange, A. M., Gregson, C., Macaulay, C., and McFadden, G. (1983). Physical characterization and molecular cloning of the Shope fibroma virus DNA genome. *Virology* **130**, 403–414.
- Wright, C. F., Keck, J. G., Tsai, M. M., and Moss, B. (1991). A transcription factor for expression of vaccinia virus late genes is encoded by an intermediate gene. *J. Virol.* **65**, 3715–3720.
- Xiang, Y., Simpson, D. A., Spiegel, J., Zhou, A., Silverman, R. H., and Condit, R. C. (1998). The vaccinia virus A18R DNA helicase is a postreplicative negative transcription elongation factor. *J. Virol.* **72**, 7012–7023.
- Yang, W. P., and Bauer, W. R. (1988). Purification and characterization of vaccinia virus structural protein VP8. *Virology* **167**, 578–584.
- Yasui, A., Eker, A. P., Yasuhira, S., Yajima, H., Kobayashi, T., Takao, M., and Oikawa, A. (1994). A new class of DNA photolyases present in various organisms including aplacental mammals. *EMBO J.* **13**, 6143–6151.
- Yuen, L., and Moss, B. (1987). Oligonucleotide sequence signaling transcriptional termination of vaccinia virus early genes. *Proc. Natl. Acad. Sci. USA* **84**, 6417–6421.
- Zajac, P., Spehner, D., and Drillien, R. (1995). The vaccinia virus J5L open reading frame encodes a polypeptide expressed late during infection and required for viral multiplication. *Virus Res* **37**, 163–173.