

Groups and Rings - SF2729

Skeleton

Jim Holmström - 890503-7571

April 24, 2012

Exercise 1. Let R be a commutative ring with unity of prime characteristic p . Show that the map $\phi_p : R \rightarrow R$ given by $\phi_p(a) = a^p$ is a homomorphism.

Solution. In a commutative ring

$$(a + b)^n = \sum \binom{n}{k} a^k b^{n-k} \quad (1)$$

holds. Where

$$\binom{n}{k} = \frac{n!}{k!(n-k)!} \quad (2)$$

and we can see that we have if n is prime.

$$p \mid \binom{p}{i} \quad \forall i \in [1, p-1] \quad (3)$$

and hence the terms

$$\binom{p}{i} a^i b^{p-i} = 0 \quad (4)$$

in a commutative ring with characteristic p . This gives us the "freshman's dream"

$$(a + b)^p = a^p + b^p \quad (5)$$

With these facts its easy to show that ϕ is a homomorphism.

$$\phi_p(a + b) = (a + b)^p = a^p + b^p = \phi_p(a) + \phi_p(b) \quad (6)$$

and trivially since R is commutative

$$\phi_p(ab) = (ab)^p = a^p b^p = \phi_p(a) \phi_p(b) \quad (7)$$

and thus ϕ_p is a homomorphism. \square

Exercise 2. Prove that if F is a field, every proper nontrivial prime ideal of $F[x]$ is maximal.

Solution.