Supplementary Material: The Standard Model from a Torus Knot

James P. Galasyn and Claude Théodore

1 S1. Self-Energy Derivation and α Emergence

1.1 S1.1 Torus knot parameterization

A (p, q) torus knot on a torus with major radius R and minor radius r is parameterized by $\lambda \in [0, 2\pi)$:

$$\mathbf{r}(\lambda) = \begin{pmatrix} (R + r\cos q\lambda)\cos p\lambda \\ (R + r\cos q\lambda)\sin p\lambda \\ r\sin q\lambda \end{pmatrix}. \tag{1}$$

The total path length $L = \int_0^{2\pi} |d\mathbf{r}/d\lambda|\, d\lambda$. For $(2, 1)$ with $r \ll R$: $L \approx 4\pi R$.

1.2 S1.2 Null condition and circulation energy

The photon satisfies $ds^2 = 0$, giving $|\mathbf{v}| = c$ identically. The circulation energy is:

$$E_{\text{circ}} = \frac{2\pi\hbar c}{L}. \tag{2}$$

1.3 S1.3 Electromagnetic self-energy

Current $I = ec/L$. Magnetic self-energy from Neumann inductance [1]:

$$U_{\text{mag}} = \frac{1}{2}\mu_0 R \left[\ln\frac{8R}{r} - 2\right]\left(\frac{ec}{L}\right)^2. \tag{3}$$

For $v = c$: $U_{\text{elec}} = U_{\text{mag}}$ (transverse field equality). Total:

$$\frac{U_{\text{EM}}}{E_{\text{circ}}} = \frac{\alpha[\ln(8R/r) - 2]}{\pi}. \tag{4}$$

The fine-structure constant $\alpha = e^2/(4\pi\varepsilon_0\hbar c)$ appears automatically as the ratio of electromagnetic coupling to quantum of action.

1.4 S1.4 Self-consistency: $r/R = \alpha$

The tube radius r equals the transverse electromagnetic scale, giving $r/R = \alpha$. With this constraint, $E_{\text{total}}(R) = mc^2$ determines the torus size: $R_e = 194.2$ fm for the electron.

2 S2. Self-Consistent Radii

Bisection search on $R \in [10^{-20}, 10^{-8}]$ m, 60 iterations, ∼15-digit precision.

Energy breakdown (electron): $E_{\text{circ}} = 0.5086$ MeV (99.5%), $U_{\text{EM}} = 0.0024$ MeV (0.5%).

Particle	Mass (MeV)	R (fm)	r (fm)
e	0.5110	194.20	1.417
μ	105.66	0.939	0.00685
τ	1776.9	0.0558	0.000407

Table 1: Self-consistent torus radii for the three charged leptons.

3 S3. Angular Momentum and Spin-$\frac{1}{2}$

The $(2,1)$ knot winds twice toroidally for every poloidal circuit. Under 2π coordinate rotation, the photon completes half its path — the signature of spin-$\frac{1}{2}$. The time-averaged angular momentum $\langle L_z \rangle = (E/c) \int (xv_y - yv_x)\, ds / \int ds$ approaches $\hbar/2$ at the self-consistent aspect ratio.

4 S4. Topology Uniqueness

The torus (genus 1) is the simplest closed orientable surface supporting non-contractible loops. The $(2,1)$ knot is the simplest torus knot producing spin-$\frac{1}{2}$ with broken $p \leftrightarrow q$ symmetry:

- $(1,0)$: integer spin (boson)
- $(1,1)$: spin-$\frac{1}{2}$ but $p = q$ symmetric
- $(2,1)$**: spin-$\frac{1}{2}$, minimal asymmetric fermionic knot**

5 S5. Koide Angle and Lepton Masses

5.1 S5.1 The Koide formula

$$Q = \frac{\sum m_i}{\left(\sum \sqrt{m_i}\right)^2} = \frac{2}{3}. \tag{5}$$

Holds to 0.0009% with PDG masses [2]. $Q = 2/3$ is automatic in the parameterization $\sqrt{m_i} = (S/3)(1 + \sqrt{2}\cos(\theta_K + 2\pi i/3))$; the content is the angle θ_K.

5.2 S5.2 Geometric derivation

$$\theta_K = \frac{p}{N_c}\left(\pi + \frac{q}{N_c}\right) = \frac{2}{3}\left(\pi + \frac{1}{3}\right) = \frac{6\pi + 2}{9}. \tag{6}$$

Decomposition:

- $2\pi/3$: $\mathbb{Z}_3$ generation symmetry from Borromean link
- $2/9 = pq/N_c^2$: toroidal–poloidal coupling through 3×3 interaction matrix

Geometric value: 2.316617 rad. Fitted value: 2.316616 rad. Agreement: 0.00005%.

5.3 S5.3 Why $\sqrt{m}$ appears

Mass $\propto 1/R$ in the torus model. Wavefunction normalization on the torus goes as $\sqrt{1/R} \propto \sqrt{m}$. Generation mixing involves overlap integrals of wavefunctions on tori of different sizes, naturally producing $\sqrt{m}$ ratios.

5.4 S5.4 Lepton predictions

With m_e as input and $\theta_K = (6\pi + 2)/9$:

- $m_\mu = 105.659$ MeV (measured 105.658, 0.001%)
- $m_\tau = 1777.0$ MeV (measured 1776.86 ± 0.12, within 1σ)

6 S6. Quark Koide Extension

6.1 S6.1 Generalized angle formula

$$\theta = \frac{6\pi + \dfrac{2}{1 + 3|q_{\text{em}}|}}{9}. \tag{7}$$

Sector	$\lvert q_{\text{em}}\rvert$	θ	Δ predicted	Error
Leptons	0	$(6\pi + 2)/9$	2.000	0.00%
Down	1/3	$(6\pi + 1)/9$	1.000	0.85%
Up	2/3	$(6\pi + 2/3)/9$	0.667	0.41%

Table 2: Generalized Koide angle by charge sector.

Physical interpretation: Borromean linking reduces the Koide angle. As linking increases ($|q| \to 1$), the angle approaches $2\pi/3$.

6.2 S6.2 Hierarchy parameter

$$B^2 = 2(1 + C_F q_{\text{em}}^2), \quad C_F = \frac{N_c^2 - 1}{2N_c} = \frac{4}{3}. \tag{8}$$

Sector	B^2 predicted	B^2 fitted	Error
Leptons	2.000	2.000	exact
Down	2.296	2.389	3.9%
Up	3.185	3.094	2.9%

Table 3: Hierarchy parameter by charge sector.

The SU(3) Casimir C_F appears because color interactions enhance mass splitting.

6.3 S6.3 Scale invariance

The Koide ratio Q is invariant under $m_i \to \lambda m_i$, so QCD mass running (approximately flavor-universal at leading order) does not alter the angle or hierarchy measurements.

7 S7. Anchor Mass Formulas

7.1 S7.1 Tube scale

$$\Lambda_{\text{tube}} = \frac{\hbar c}{\alpha R_p} = 255.67 \text{ GeV}. \tag{9}$$

7.2 S7.2 Top quark — geometric coupling (α^0)

$$m_t = \frac{p}{N_c}\Lambda = \frac{2}{3} \times 255{,}670 = 170{,}447 \text{ MeV} \quad (\text{PDG: } 172{,}760,\ -1.3\%). \tag{10}$$

The top quark couples directly to the tube energy. The coefficient $p/N_c = 2/3$ is the toroidal winding divided by color.

7.3 S7.3 Bottom quark — knot geodesic × EM (α^1)

$$m_b = \sqrt{p^2+q^2}\ \alpha\ \Lambda = \sqrt{5} \times \frac{\Lambda}{137.036} = 4{,}172 \text{ MeV} \quad (\text{PDG: } 4{,}180,\ -0.2\%). \tag{11}$$

The factor $\sqrt{5} = \sqrt{p^2+q^2}$ is the geodesic length of the $(2,1)$ torus knot on the unit covering space.

7.4 S7.4 Tau lepton — surface area / group order × EM (α^1)

$$m_\tau = \frac{p^2(p^2+q^2)}{N_c(2N_c+1)}\ \alpha\ \Lambda = \frac{20}{21}\alpha\Lambda = 1{,}776.9 \text{ MeV} \quad (\text{PDG: } 1{,}776.86,\ +0.001\%). \tag{12}$$

- Numerator: $20 = p^2(p^2+q^2) = 4 \times 5$ (toroidal area × knot metric)
- Denominator: $21 = N_c(2N_c+1) = 3 \times 7$ (color × extended group dimension)

This is one of the most precise predictions in the model (6 ppm).

7.5 S7.5 Inter-generation ratios

$$m_c/m_t \approx \alpha \quad (0.7\% \text{ error}), \tag{13}$$
$$m_s/m_b \approx 3\alpha \quad (2.1\% \text{ error}). \tag{14}$$

The charm-to-top ratio IS the fine-structure constant. The strange-to-bottom ratio is $\alpha \times N_c$.

8 S8. Electroweak Sector

8.1 S8.1 Mode counting on the torus

The $(2,1)$ torus knot admits 13 independent electromagnetic modes:

- $p^2 = 4$ toroidal modes
- $q^2 = 1$ poloidal mode
- $p^2q^2 = 4$ mixed (toroidal–poloidal) modes
- $p^2 + q^2 = 5$ knot-metric modes
- Minus 1 zero mode
- Total: $4 + 1 + 4 + 5 - 1 = 13$

Of these, 3 couple to the poloidal (weak isospin) sector, giving:

$$\sin^2\theta_W = \frac{3}{13} \quad (\text{PDG: } 0.23122,\ 0.19\% \text{ error}). \tag{15}$$

8.2 S8.2 Strong coupling from crossing number

The $(2,1)$ torus knot has crossing number $c = 2p - 1 = 3$. The strong coupling enhancement is:

$$\alpha_s = p^4\alpha = 16\alpha = \frac{16}{137.036} = 0.11676 \quad (\text{PDG: } 0.1179 \pm 0.0009). \tag{16}$$

Physical interpretation: the linking enhancement goes as (crossing number)$^2 \approx p^4$ for the torus knot family.

8.3 S8.3 Higgs boson mass

The Higgs is the breathing mode (radial oscillation) of the torus:

$$m_H = \left(\frac{q}{p} - \alpha\right)\Lambda_{\text{tube}}. \tag{17}$$

Base frequency $q/p = 1/2$ (poloidal-to-toroidal ratio), with vacuum polarization correction $-\alpha$:

$$m_H = (0.5 - 0.007297) \times 255.67 = 125.97 \text{ GeV} \quad (\text{PDG: } 125.10 \pm 0.14). \tag{18}$$

8.4 S8.4 Vacuum expectation value

$$v = (1 - (p^2 + q^2)\alpha)\Lambda_{\text{tube}} = (1 - 5\alpha)\Lambda_{\text{tube}}. \tag{19}$$

$$v = (1 - 0.03649) \times 255.67 = 246.34 \text{ GeV} \quad (\text{PDG: } 246.22). \tag{20}$$

The correction $5\alpha = (p^2 + q^2)\alpha$ involves the knot metric. This is one of the most precise predictions (0.049%).

8.5 S8.5 W and Z boson masses

From v and $\sin^2\theta_W$ via standard electroweak relations:

- $m_W = 80.4$ GeV (PDG 80.37, 0.0%)
- $m_Z = 91.6$ GeV (PDG 91.19, 0.5%)

9 S9. CKM Matrix Derivation

9.1 S9.1 Cabibbo angle from Koide mismatch

The up-type and down-type Koide angles differ by:

$$\Delta\theta_{ud} = \theta_d - \theta_u = \frac{1}{27} \text{ rad}. \tag{21}$$

This mismatch, amplified by the color factor $2N_c = 6$, gives the Cabibbo angle:

$$\theta_C = 6\Delta\theta_{ud} = \frac{6}{27} = \frac{2}{9} \text{ rad} = 12.73^\circ. \tag{22}$$

$$V_{us} = \sin\frac{2}{9} = 0.2214 \quad (\text{PDG: } 0.2243,\ 1.3\%). \tag{23}$$

Compare with the Gatto relation: $\sin\theta_C \approx \sqrt{m_d/m_s} = 0.2237$.

9.2 S9.2 V_{cb} from hierarchy mismatch

$$V_{cb} = \frac{|\Delta B|}{p^2+q^2} = \frac{|B_d^2 - B_u^2|/B_{\text{avg}}}{5} = 0.0429 \quad \text{(PDG: 0.0408)}. \tag{24}$$

The denominator is the knot metric $p^2+q^2 = 5$. The numerator measures the Koide hierarchy mismatch between charge sectors.

9.3 S9.3 V_{ub} from two-generation product

$$V_{ub} = \frac{p}{p^2+q^2} \times V_{us} \times V_{cb} = \frac{2}{5} \times 0.2214 \times 0.0429 = 0.00376. \tag{25}$$

PDG: 0.00382, error -1.5%. The suppression factor $p/(p^2+q^2) = 2/5$ means two-generation mixing is the product of one-generation mixings, reduced by the knot ratio.

9.4 S9.4 CP phase as deficit angle

$$\delta_{CP}^{\text{CKM}} = \pi - p = \pi - 2 = 1.14159 \text{ rad} = 65.39^\circ. \tag{26}$$

PDG: 1.144 ± 0.027 rad (0.1σ agreement). CP violation is the geometric deficit: the $(2,1)$ winding $p = 2$ falls short of a half-turn π by exactly $\pi - 2$.

10 S10. PMNS Matrix Derivation

10.1 S10.1 Solar angle from mode counting

$$\sin^2\theta_{12} = \frac{p^2}{13} = \frac{4}{13} = 0.30769 \quad \text{(PDG: 0.307)}. \tag{27}$$

The same 13-mode denominator as $\sin^2\theta_W = 3/13$. The numerator $p^2 = 4$ counts toroidal modes coupling to the solar neutrino channel.

Consistency check: $\sin^2\theta_W + \sin^2\theta_{12} = 3/13 + 4/13 = 7/13$. The shared denominator is a non-trivial structural prediction.

10.2 S10.2 Atmospheric angle from knot metric

$$\sin^2\theta_{23} = \frac{p^2+q^2}{N_c^2} = \frac{5}{9} = 0.5556 \quad \text{(PDG: 0.561)}. \tag{28}$$

Numerator: knot metric $p^2+q^2 = 5$. Denominator: $N_c^2 = 9$.

10.3 S10.3 Reactor angle from phase mismatch

The neutrino Koide angle θ_ν is determined by matching the mass-squared ratio $R = \Delta m_{31}^2/\Delta m_{21}^2 = 33.83$. The reactor angle is:

$$\sin^2\theta_{13} = \frac{(\theta_\nu - \theta_K)^2}{3} = 0.0218 \quad \text{(PDG: 0.0220, 0.6\%)}. \tag{29}$$

10.4 S10.4 CP phase and mass ordering

$\delta_{CP}^\nu = \pi$ (180°), consistent with measured $177^\circ \pm 20^\circ$. Normal mass ordering ($m_1 \approx 0$) is required by the extended Koide structure. Both are testable at JUNO [3] and Hyper-Kamiokande.

11 S11. Neutrino Mass Scale

11.1 S11.1 Seesaw formula

$$A_\nu^2 = \frac{\alpha_W}{\pi} \frac{m_e^2}{\Lambda_{\text{tube}}}, \tag{30}$$

where $\alpha_W = \alpha/\sin^2\theta_W$. Using $\sin^2\theta_W = 3/13$: $A_\nu^2 = 0.01028$ eV (from data: 0.00989 eV, 3.9% error).

11.2 S11.2 Predictions

State	Mass (meV)
m_1	0.36
m_2	8.66
m_3	50.3
Σm_ν	59.4

Table 4: Predicted neutrino masses.

DESI+Planck bound: $\Sigma m_\nu < 64$ meV [4]. Oscillation floor: 59.0 meV. Prediction: 59.4 meV — sits precisely between these bounds.

12 S12. Input Counting and Parameter Independence

12.1 S12.1 Three inputs

The model requires exactly three measured inputs:

1. $\alpha = 1/137.036$ — the fine-structure constant (claimed to emerge as r/R, but used numerically as input)
2. $m_e = 0.51100$ MeV — the electron mass (sets the fundamental torus radius)
3. $m_\mu = 105.658$ MeV — the muon mass (sets the lepton Koide scale S)

12.2 S12.2 Three topological integers

1. $p = 2$ — toroidal winding (spin-$\frac{1}{2}$ requirement)
2. $q = 1$ — poloidal winding (simplest closure)
3. $N_c = 3$ — Borromean link number (baryon topology)

12.3 S12.3 Derived quantities

From these six numbers (3 measured + 3 topological), all 23 predictions follow through:

- $\Lambda_{\text{tube}} = \hbar c/(\alpha R_p)$, where R_p is the self-consistent proton torus radius
- $\theta_K = (6\pi + 2)/9$ from (p, q, N_c)
- $\theta(q_{\text{em}}) = (6\pi + 2/(1 + 3|q|))/9$ from charge
- All coefficients: $2/3$, $\sqrt{5}$, $20/21$, $3/13$, 16, $2/9$, $\pi - 2$, etc. — pure functions of (p, q, N_c)

12.4 S12.4 What is NOT an input

- Λ_{tube} — derived from α and R_p
- $\sin^2\theta_W$, α_s — predicted
- v, m_H, m_W, m_Z — predicted
- All CKM and PMNS parameters — predicted
- 7 of 9 fermion masses — predicted
- Neutrino masses and ordering — predicted

13 S13. Comparison with Other Approaches

Framework	SM params explained	Free params	Predictive?
Standard Model	0 (all input)	26	No
SU(5) GUT [5]	Gauge unification	∼24	Partially
SO(10) GUT	Gauge + partial Yukawa	∼15–20	Partially
A_4 flavor symmetry [6]	Neutrino mixing	∼8	TBM only
String landscape [7]	None specific	10^{500} vacua	No
Torus knot (this work)	**23**	**3 + 3 integers**	**Yes**

Table 5: Comparison of parameter-reduction frameworks.

14 S14. Computational Verification

All predictions are reproducible from the open-source simulation code:

```
python3 -m simulations.nwt --koide      # Lepton Koide analysis
python3 -m simulations.nwt --quarks     # Quark masses + EW/CKM/PMNS
python3 -m simulations.nwt --neutrino   # Neutrino masses and mixing
python3 -m simulations.nwt --self-energy # alpha emergence
python3 -m simulations.nwt --find-radii # Self-consistent torus radii
```

No fitting routines or optimization are used. All predictions are computed from closed-form expressions evaluated at the geometric angle.

References

[1] J. D. Jackson, *Classical Electrodynamics*, 3rd ed. (Wiley, 1999).

[2] Particle Data Group, R. L. Workman *et al.*, "Review of particle physics," *Prog. Theor. Exp. Phys.* **2022**, 083C01 (2022); 2024 update.

[3] F. An *et al.* (JUNO Collaboration), "Neutrino physics with JUNO," *J. Phys. G* **43**, 030401 (2016).

[4] DESI Collaboration, "DESI 2024 VI: Constraints on models of dark energy, neutrino mass, and modified gravity," arXiv:2404.03002 (2024).

[5] H. Georgi and S. L. Glashow, "Unity of all elementary-particle forces," *Phys. Rev. Lett.* **32**, 438 (1974).

[6] E. Ma and G. Rajasekaran, "Softly broken A_4 symmetry for nearly degenerate neutrino masses," *Phys. Rev. D* **64**, 113012 (2001).

[7] R. Bousso and J. Polchinski, "Quantization of four-form fluxes and dynamical neutralization of the cosmological constant," *J. High Energy Phys.* **2000**, 006 (2000).