


Modern Birkhäuser Classics


Logic for Computer Scientists
Uwe Schöningh

Prolog Programming for Artificial Intelligence
Horst R. Hell



FLORIDA
INTERNATIONAL
UNIVERSITY


First Order Logic (V)

Dr. Antonio L. Bajuelos


School of Computing &
Information Sciences

Note: The most of the information of these slides was extracted and adapted from Schöningh's book, "Logic for Computer Scientists". They are provided for COT-3541 students only. Not to be published or publicly distributed without permission by the publisher.

Resolution principle for FOL.



■ Using the **unification principle**, we are now in a situation to formulate the **resolution principle** for **FOL**.

DEFINITION (resolution in FOL):

Let C_1, C_2 and R be clauses (in predicate logic). Then R is called a *resolvent* of C_1, C_2 if the following holds.


1. There exist certain substitutions s_1 and s_2 which are variable renamings so that C_1s_1 and C_2s_2 do not contain the same variable.
2. There is a set of literals $L_1, \dots, L_m \in C_1s_1$ ($m \geq 1$) and $L'_1, \dots, L'_n \in C_2s_2$ ($n \geq 1$), such that $\mathbf{L} = \{\overline{L_1}, \overline{L_2}, \dots, \overline{L_m}, L'_1, L'_2, \dots, L'_n\}$ is unifiable. Let sub be a most general unifier for \mathbf{L} .
3. R has the form

$$R = ((C_1s_1 - \{L_1, \dots, L_m\}) \cup (C_2s_2 - \{L'_1, \dots, L'_n\}))sub .$$

2

Resolution principle for FOL.

- Example of the **resolvent**:



Substitutions???

3

Resolution principle for FOL.

Resolution Theorem (of FOL)

- Let F be a closed formula in **Skolem** form with its matrix F^* . Then, F is **unsatisfiable** iff $\square \in \text{Res}^*(F^*)$.

(see proof in the Textbook, Chapter 2, section 2.5)

4

Resolution principle for FOL.



Example:

- Consider the **two clauses**:
 $C1 = \{P(f(x), g(y)), Q(x, y)\}$
 $C2 = \{\neg P(f(f(a)), g(z)), Q(f(a), g(z))\}$
- These **two clauses** contain the following literals:
 $L1 = \{P(f(x), g(y))\}$
 $L2 = \{\neg P(f(f(a)), g(z))\}$
- It is easy to check that the literals in L1 and L2 are **unifiable** with **MGU** = $\{[x/f(a)], [y/z]\}$
- so the clauses C1 and C2 are clashing, and their **resolvent** is:
 $Res(C1, C2) = \{Q(f(a), z), Q(f(a), g(z))\}$

5

Resolution principle for FOL.



Example #1:

- In this example, we will prove that the set of clauses (1)-(4) is **unsatisfiable**.
 1. $\neg p(x, y) \vee p(y, x)$
 2. $\neg p(x, y) \vee \neg p(y, z) \vee p(x, z)$
 3. $p(x, f(x))$
 4. $\neg p(x, x)$
- **Resolution...**

3'. $p(x', f(x'))$	3 Rename x to x'
5. $p(f(x), x)$	1, 3' $[x'/x], [y/f(x)]$
3''. $p(x'', f(x''))$	3 Rename x to x''
6. $\neg p(f(x), z) \vee p(x, z)$	3'', 2 $[y/f(x)], [x''/x]$
5'''. $p(f(x'''), x''')$	5 Rename x to x'''
7. $p(x, x)$	6, 5 $[z/x], [x'''/x]$
4'''. $\neg p(x''', x''')$	4 Rename x to x'''
8. \square	7, 4''' $[x'''/x]$

6

Resolution principle for FOL.



Example #2:

- We will show, using the general resolution procedure, that the following set of clauses is **unsatisfiable**:
 1. $\neg p(x) \vee q(x) \vee r(x, f(x))$
 2. $\neg p(x) \vee q(x) \vee s(f(x))$
 3. $t(a)$
 4. $p(a)$
 5. $\neg r(a, y) \vee t(y)$
 6. $\neg t(x) \vee \neg q(x)$
 7. $\neg t(x) \vee \neg s(x)$
- **Resolution...**
 8. $\neg q(a)$ 3,6 [x/a]
 9. $\neg p(a) \vee s(f(a))$ 2,8 [x/a]
 10. $\neg p(a) \vee r(a, f(a))$ 1,8 [x/a]
 11. $s(f(a))$ 4,9
 12. $r(a, f(a))$ 4,10
 13. $t(f(a))$ 5,12 [y/f(a)]
 14. $\neg t(f(a))$ 11,7 [x/f(a)]
 15. \square 13,14

7

Resolution principle for FOL.



Example #3:

- Use resolution to prove that a **relation** $R \subseteq A \times A$ is **reflexive** if it is **transitive** and **symmetric**.

Solution:

- Translate in terms of logic.
- Define $R(x, y)$ to be true if x is related to y .
- Since A is the **domain** we have
$$\forall x \exists y R(x, y).$$
- The relation is **transitive**, that is,
$$\forall x \forall y \forall z ((R(x, y) \wedge R(y, z)) \rightarrow R(x, z)).$$
- The relation is **symmetric**, that is
$$\forall x \forall y (R(x, y) \rightarrow R(y, x)).$$
- We want to conclude from these premises that the relation is **reflexive**, that is,
$$\forall x R(x, x).$$
- To prove the theorem by **refutation**, negate the conclusion
$$\exists x \neg R(x, x).$$

8

Resolution principle for FOL.

Example #3 (cont...):

Original clauses are:

The **domain** we have

$$\forall x \exists y R(x, y)$$

Transitive:

$$\forall x \forall y \forall z ((R(x, y) \wedge R(y, z)) \rightarrow R(x, z))$$

Symmetric:

$$\forall x \forall y (R(x, y) \rightarrow R(y, x))$$

Reflexive:

$$\forall x R(x, x)$$

To prove the theorem by refutation, negate the conclusion

$$\exists x \neg R(x, x).$$

After Skolemization:

The **domain** we have

$$R(x, f(x))$$

Transitive:

$$\neg R(x, y) \vee \neg R(y, z) \vee R(x, z)$$

Symmetric:

$$\neg R(x, y) \vee R(y, x)$$

Reflexive:

To prove the theorem by refutation, negate the conclusion

$\neg R(a, a).$



9

Resolution principle for FOL.

Example #3 (cont...):

■ Clauses:

1. $\neg R(a, a)$ // Conclusion negation
2. $R(x, f(x))$
3. $\neg R(x, y) \vee \neg R(y, z) \vee R(x, z)$
4. $\neg R(x, y) \vee R(y, x)$

■ Resolution:

5. $\neg R(a, y) \vee \neg R(y, a) \vee R(a, a)$ 3, with $[x/a]$ $[z/a]$
6. $\neg R(a, y) \vee \neg R(y, a)$ resolve 1 and 5
7. $R(a, f(a))$ 2 with $[x/a]$
8. $\neg R(a, f(a)) \vee \neg R(f(a), a)$ 6 with $[y/f(a)]$
9. $\neg R(f(a), a)$ resolve 7 and 8
10. $\neg R(a, f(a)) \vee R(f(a), a)$ 4 with $[x/a]$ $[y/a]$
11. $\neg R(a, f(a))$ resolve 9 and 10
12. $R(a, f(a))$ 2 with $[x/a]$
13. \square 11, 12

empty clause – original claim is valid



10

Resolution principle for FOL.



Example #4:

Prove that everybody has a grandparent, provided everybody has a parent.

- Let $P(x,y)$ represent x is a parent of y . The **premise** can now be stated as

$$\forall x \exists y P(y,x).$$

- From this we must be able to **conclude** that there exists a parent of a parent, which can be expressed as

$$\forall x \exists y \exists z (P(z,y) \wedge P(y,x)).$$

- We must thus prove that

$$\forall x \exists y P(y,x) \models \forall x \exists y \exists z (P(z,y) \wedge P(y,x))$$

11

Resolution principle for FOL.



Example #4 (cont...):

Prove that everybody has a grandparent is a logical consequence that everybody has a parent.

$P(x,y)$: x is a parent of y

$$\forall x \exists y P(y,x) \models \forall x \exists y \exists z (P(z,y) \wedge P(y,x))$$

- We add the **negation of the conclusion** to the set of premises, which yields:

$$\forall x \exists y P(y,x) \wedge \exists x \forall y \forall z (\neg P(z,y) \vee \neg P(y,x))$$

- Eliminate the existential quantifiers (**skolemization** process):

$$\forall x \exists y P(y,x) \wedge \exists w \forall v \forall z (\neg P(z,v) \vee \neg P(v,w))$$

$$\forall x \exists y \exists w \forall v \forall z (P(y,x) \wedge (\neg P(z,v) \vee \neg P(v,w)))$$

$$\forall x \forall v \forall z (P(f(x),x) \wedge (\neg P(z,v) \vee \neg P(v,g(x))))$$

- After dropping the universal quantifiers, this yields:

$$S = \{ \{P(f(x),x)\}, \{\neg P(z,v), \neg P(v,g(x))\} \}$$

12

Resolution principle for FOL.



Example #4 (cont...):

Prove that everybody has a grandparent, provided everybody has a parent.

$$\forall x \exists y P(y, x) \models \forall x \exists y \exists z (P(z, y) \wedge P(y, x))$$



$\{ \{P(f(x), x)\}, \{ \neg P(z, v), \neg P(v, g(x)) \} \}$ is **unsatisfiable**

■ Resolution...

- | | |
|---|----------------------------------|
| 1. $P(f(x), x)$ | Given |
| 2. $\neg P(z, v) \vee \neg P(v, g(x))$ | Given |
| 3. $P(f(g(a)), g(a))$ | 1 with $[x/g(a)]$ |
| 4. $\neg P(z, f(g(a))) \vee \neg P(f(g(a)), g(a))$ | 2 with $[x/a]$ and $[v/f(g(a))]$ |
| 5. $\neg P(z, f(g(a)))$ | Resolve 3 and 4 |
| 6. $P(f(f(g(a))), f(g(a)))$ | 1 with $[x/f(g(a))]$ |
| 7. $\neg P(f(f(g(a))), f(g(a)))$ | 5 with $[z/f(f(g(a)))]$ |
| 8. \square empty clause (contradiction) from 6 and 7 | |

This means that the original argument is valid.

13