

Modern Birkhäuser Classics

Logic for Computer Scientists
Uwe Schöningh

Prolog Programming for Artificial Intelligence
Northern Institute

Peter Brönnimann




FLORIDA
INTERNATIONAL
UNIVERSITY


Propositional Logic CNF Examples

Dr. Antonio L. Bajuelos


School of Computing &
Information Sciences

Note: The most of the information of these slides was extracted and adapted from Schöningh's book, "Logic for Computer Scientists". They are provided for COT-3541 students only. Not to be published or publicly distributed without permission by the publisher.





Procedure for converting to CNF

- **To eliminate \leftrightarrow**
 - $(a \leftrightarrow b) \equiv (a \rightarrow b) \wedge (b \rightarrow a)$
- **To eliminate \rightarrow**
 - $(a \rightarrow b) \equiv \neg a \vee b$
- **Double negation \neg**
 - $\neg(\neg a) \equiv a$
- **De Morgan**
 - $\neg(a \wedge b) \equiv (\neg a \vee \neg b)$
 - $\neg(a \vee b) \equiv (\neg a \wedge \neg b)$
- **Distributivity of \wedge over \vee**
 - $(a \wedge (b \vee c)) \equiv ((a \wedge b) \vee (a \wedge c))$
- **Distributivity of \vee over \wedge**
 - $(a \vee (b \wedge c)) \equiv ((a \vee b) \wedge (a \vee c))$

2

Example #1

Reduce to Conjunctive Normal Form (CNF) the formula

$$\neg(\neg p \vee q) \vee (r \rightarrow \neg s)$$

1. $\neg(\neg p \vee q) \vee (\neg r \vee \neg s)$
2. $(\neg\neg p \wedge \neg q) \vee (\neg r \vee \neg s)$
3. $(p \wedge \neg q) \vee (\neg r \vee \neg s)$
4. $(p \vee \neg r \vee \neg s) \wedge (\neg q \vee \neg r \vee \neg s)$



3

Example #2

Reduce to Conjunctive Normal Form (CNF) the formula

$$(\neg p \rightarrow q) \rightarrow (q \rightarrow \neg r)$$

1. $\neg(\neg p \rightarrow q) \vee (q \rightarrow \neg r)$
2. $\neg(p \vee q) \vee (\neg q \vee \neg r)$
3. $(\neg p \wedge \neg q) \vee (\neg q \vee \neg r)$
4. $(\neg p \vee \neg q \vee \neg r) \wedge (\neg q \vee \neg r)$



4

Example #3

Reduce to Conjunctive Normal Form (CNF) the formula

$$\neg((\neg p \rightarrow \neg q) \wedge \neg r)$$

$$\begin{aligned}\neg((\neg p \rightarrow \neg q) \wedge \neg r) &\equiv \neg((\neg\neg p \vee \neg q) \wedge \neg r) \\ &\equiv \neg((p \vee \neg q) \wedge \neg r) \\ &\equiv \neg(p \vee \neg q) \vee \neg\neg r \\ &\equiv \neg(p \vee \neg q) \vee r \\ &\equiv (\neg p \wedge \neg\neg q) \vee r \\ &\equiv (\neg p \wedge q) \vee r \\ &\equiv (\neg p \vee r) \wedge (q \vee r)\end{aligned}$$

5

Example #4

Reduce to Conjunctive Normal Form (CNF) the formula

$$(p \rightarrow q) \rightarrow (\neg r \wedge q)$$

$$\begin{aligned}(p \rightarrow q) \rightarrow (\neg r \wedge q) &\equiv \neg(p \rightarrow q) \vee (\neg r \wedge q) \\ &\equiv \neg(\neg p \vee q) \vee (\neg r \wedge q) \\ &\equiv (\neg\neg p \wedge \neg q) \vee (\neg r \wedge q) \\ &\equiv (p \wedge \neg q) \vee (\neg r \wedge q) \\ &\equiv (p \vee (\neg r \wedge q)) \wedge (\neg q \vee (\neg r \wedge q)) \\ &\equiv (p \vee (\neg r \wedge q)) \wedge (\neg q \vee \neg r) \wedge (\neg q \vee q) \\ &\equiv (p \vee (\neg r \wedge q)) \wedge (\neg q \vee \neg r) \wedge \mathbf{T} \\ &\equiv (p \vee (\neg r \wedge q)) \wedge (\neg q \vee \neg r) \\ &\equiv (p \vee \neg r) \wedge (p \vee q) \wedge (\neg q \vee \neg r)\end{aligned}$$

6

Example #5

Reduce to Conjunctive Normal Form (CNF) the formula

$$(A \rightarrow B \vee C) \rightarrow (A \wedge D)$$

$$\begin{aligned} &\equiv \neg(A \rightarrow B \vee C) \vee (A \wedge D) \\ &\equiv (A \wedge \neg(B \vee C)) \vee (A \wedge D) \\ &\equiv (A \wedge \neg B \wedge \neg C) \vee (A \wedge D) \\ &\equiv ((A \wedge \neg B \wedge \neg C) \vee A) \wedge ((A \wedge \neg B \wedge \neg C) \vee D) \\ &\equiv A \wedge ((A \wedge \neg B \wedge \neg C) \vee D) \\ &\equiv A \wedge (A \vee D) \wedge (\neg B \vee D) \wedge (\neg C \vee D) \\ &\equiv A \wedge (\neg B \vee D) \wedge (\neg C \vee D) \end{aligned}$$



7

Example #6

Reduce to Conjunctive Normal Form (CNF) the formula

$$((p \wedge q) \vee (r \wedge s)) \vee (\neg q \wedge (p \vee t))$$

$$\begin{aligned} &\equiv (((p \wedge q) \vee r) \wedge ((p \wedge q) \vee s)) \vee (\neg q \wedge (p \vee t)) \\ &\equiv ((p \vee r) \wedge (q \vee r) \wedge (p \vee s) \wedge (q \vee s)) \vee (\neg q \wedge (p \vee t)) \\ &\equiv ((p \vee r) \vee (\neg q \wedge (p \vee t))) \wedge \\ &\quad ((q \vee r) \vee (\neg q \wedge (p \vee t))) \wedge \\ &\quad ((p \vee s) \vee (\neg q \wedge (p \vee t))) \wedge \\ &\quad ((q \vee s) \vee (\neg q \wedge (p \vee t))) \\ &\equiv (p \vee r \vee \neg q) \wedge (p \vee r \vee p \vee t) \wedge \\ &\quad (q \vee r \vee \neg q) \wedge (q \vee r \vee p \vee t) \wedge \\ &\quad (p \vee s \vee \neg q) \wedge (p \vee s \vee p \vee t) \wedge \\ &\quad (q \vee s \vee \neg q) \wedge (q \vee s \vee p \vee t) \end{aligned}$$



8