

Z-scores: Takeaways

by Dataquest Labs, Inc. - All rights reserved © 2020

Syntax

- Writing a function that converts a value to a z-score:

```
def z_score(value, array, bessell = 0):  
    mean = sum(array) / len(array)  
  
    from numpy import std  
    st_dev = std(array, ddof = bessell)  
  
    distance = value - mean  
    z = distance / st_dev  
  
    return z
```

- Standardizing a `Series`:

```
standardized_distro = Series.apply(  
    lambda x: (x - Series.mean()) /  
    Series.std()  
)
```

- Transforming a standardized distribution to a different distribution, with a predefined mean and standard deviation:

```
mean = some_mean  
st_dev = some_standard_deviation  
  
standardized_distro = Series.apply(  
    lambda z: z * st_dev + mean  
)
```

Concepts

- A **z-score** is a number that describes the location of a value within a distribution. Non-zero z-scores (+1, -1.5, +2, -2, etc.) consist of two parts:
 - A *sign*, which indicates whether the value is above or below the mean.
 - A *value*, which indicates the number of standard deviations that a value is away from the mean.
- The z-score of the mean is 0.
- To compute the z-score z for a value x coming from a population with mean μ and standard deviation σ , we can use this formula:

$$Z = \frac{x - \mu}{\sigma}$$

- To compute the z-score z for a value x coming from a sample with mean \bar{x} and standard deviation s , we can use this formula:

$$Z = \frac{x - \bar{x}}{s}$$

- We can **standardize** any distribution by transforming all its values to z-scores. The resulting distribution will have a mean of 0 and a standard deviation of 1. Standardized distributions are often called **standard distributions**.
- Standardization is useful for **comparing values** coming from distributions with different means and standard deviations.
- We can transform any population of z-scores with mean $\mu_z = 0$ and $\sigma_z = 1$ to a distribution with any mean μ and any standard deviation σ by converting each z-score z to a value x using this formula:

$$X = Z\sigma + \mu$$

- We can transform any sample of z-scores with mean $\bar{x}_z = 0$ and $s_z = 1$ to a distribution with any mean \bar{x} and any standard deviation s by converting each z-score z to a value x using this formula:

$$X = ZS + \bar{X}$$

Resources

- [The `z-score\(\)` function from `scipy.stats.mstats`](#) — useful for standardizing distributions.

- [The Wikipedia entry on z-scores.](#)



Takeaways by Dataquest Labs, Inc. - All rights reserved © 2020