Galaxy dynamics from Newtonian gravity and a non-baryonic mass set by the baryons

Can MOND's regularities follow from a mass that responds to the potential and enters the Poisson equation only through its source term?

James R. Marsen

v2.0.0, 2026-10-28. DOI 10.5281/zenodo.23748074. Supersedes GPRF Paper I v1.0.14 (Zenodo 10.5281/zenodo.20450130). CC0.

Abstract

Galaxies obey regularities that MOND predicted from the baryons alone: the baryonic Tully-Fisher relation (BTFR) and the radial acceleration relation (RAR). Weak lensing extends the RAR to accelerations two decades lower (Brouwer et al. 2021) and finds the circular speeds of isolated galaxies flat to 1 Mpc (Mistele et al. 2024b). We ask whether these regularities can follow, with gravity unmodified, from a non-baryonic mass whose density is set by the present baryons. We present a constitutive relation for such a mass, adapted from the cored isothermal sphere. Its velocity scale is fixed by the BTFR and its response saturates at the MOND acceleration, so MOND's two limits are inputs and the transition between them is an output, with one fitted constant, the central density. Each halo ends at an ambient density set by a posited smooth background. The mass gravitates and lenses as ordinary mass; a full dynamical treatment is deferred. The paper is conditional: if non-baryonic mass is distributed this way, then

(i) on the SPARC rotation curves a single central density reproduces the mean RAR, including the transition, to within 0.1 dex in every acceleration bin, inside the RAR's observed scatter ($\sim$0.13 dex; Lelli et al. 2017);

(ii) the fit improves when the central density depends on galaxy properties, and with one free parameter per galaxy it matches the empirical relation;

(iii) the lensing signal of isolated galaxies turns down within a few Mpc (closer in for galaxies with neighbours), at a radius growing as the fourth root of their baryonic mass, because the response ends at an ambient density set by a smooth background field; the downturn persists in voids, where MOND's external-field decline recedes;

(iv) there is no phantom density: the extra mass is real and positive everywhere, so where MOND's phantom density turns negative, in systems embedded in an external field, the two differ: wide binaries of low central density are Newtonian;

(v) dwarf galaxies do not depend on their environment.

It falls short in four ways, each of which weakens it:

- with a single central density its scatter about the RAR is larger than observed, by well over the relation's measured intrinsic scatter;
- it over-predicts the lowest accelerations, by up to 0.08 dex, and the outer speeds of low-surface-brightness galaxies, by a few per cent;
- the central density is fitted, not derived; and
- a first pre-registered test on dwarf dispersions misses its target.

We set out the tests that can decide.

Summary of results

	Result	Number	Section
Reproduced	Mean RAR, including the transition, one global ρ_c	within 0.1 dex per bin; rms 0.155 vs 0.133 empirical	3.1
	Individual curves, one parameter per galaxy	rms 0.081 vs 0.079 (RAR with M/L free)	3.2
	Milky Way decline (Gaia DR3)	rms 20 km/s; within 3 km/s of MOND	3.2, A
	Isolated-galaxy lensing, flat to 1 Mpc	by construction (BTFR tail)	C
	Core surface density	reproduced on average, not predicted	3.3
Conceded	Scatter exceeds the measured intrinsic scatter	$\sim$0.08 dex in quadrature vs 0.034	3.1
	Lowest accelerations over-predicted	up to 0.08 dex	3.1
	Outer speeds of low-surface-brightness galaxies over-predicted	3–8%	4.1
	Crater II, NGC 1052-DF2 and DF4 not reproduced at equilibrium	–	4.2, 4.3
	Pre-registered dwarf protocol fails its criterion	9 of 16	B.2
	ρ_c fitted, not derived	–	B.1
Predicted	Wide binaries, Gaia DR4	$\gamma = 1.000$ for low central density; killed by a confirmed $\gamma \geq 1.1$	6.1
	Lensing downturn, isolated galaxies	$r_t \propto M_b^{1/4}$, about 3–6 Mpc; killed by flatness to 5 Mpc (the closure value) or a bin ratio near 3.9	6.3, 6.5
	Environment	slope 0; killed by -0.08 ± 0.03	6.2
	Dwarfs, embedded against isolated	alike; killed by an offset > 0.15 dex	6.5
Open	Mpc budget (mean-field indication that it closes); dynamics and conservation; galaxy clusters	–	C, 7

1. Introduction

The question is whether real mass can do what MOND does. Milgrom's law (1983) and its field formulations (Bekenstein & Milgrom 1984; Milgrom 2010) reproduce the regularities of galaxy dynamics – the baryonic Tully-Fisher relation, the radial acceleration relation (McGaugh, Lelli & Schombert 2016), flat rotation curves, the halo surface-density scale (Milgrom 2009; Donato et al. 2009) – from the baryons alone, with one acceleration constant a_0. The case against dark matter made from these facts (Famaey & McGaugh 2012; McGaugh 2023) has never been that mass cannot exist; it is that an independent mass component cannot know about the baryons closely enough to produce them without fine-tuning. That objection targets one kind of real mass: an independent, history-laden particle population. A real mass whose distribution is a function of the present baryon field is a different hypothesis, and the objections become constraints on its relation. This note states such a relation and tests it. It offers potential explanations, not resolutions; where the relation falls short we say so, in our own terms.

Where this sits. Forty years of rotation curves have left two positions standing. Modified gravity reproduces the regularities of galaxies with one acceleration scale but needs a relativistic completion and falls short in galaxy clusters; particle dark matter has the cosmology but gets the galaxies only after fitting halos to them. Each side's success is the other's unsolved problem. Despite decades of searches, no particle dark-matter candidate has been confirmed non-gravitationally (LZ Collaboration 2025; XENONnT Collaboration 2025; PandaX-4T Collaboration 2025). The evidence for extra mass remains purely gravitational. That leaves the properties of the non-baryonic mass open, which is the space this note works in. Any particle candidate must also account for MOND's phenomenology; the relation studied here approaches that requirement from the other side. It is an attempt to put MOND's regularities where particle dark matter needs them to be – in a real, gravitating mass distribution – without the halo freedom. It uses different priors, not more priors: an isothermal form, a per-body velocity parameter fixed by the BTFR, and an ambient-density truncation, with one fitted constant, the central

density ρ_c. It takes MOND's two limits as inputs, obtains the transition between them from the self-consistent solve, and asks what follows when the mass that produces them is treated as real: a Newtonian limit for small bound systems, a truncation at the ambient density, and a mass that is truncated by tides, re-formed when bodies merge, and lensed. If a mass like this exists, three programmes approach it from different sides: superfluid and baryon-interacting dark matter from the particle side, the isothermal cores of self-interacting dark matter from the simulation side, and MOND phenomenology from the data. The relation can be read as a static limit compatible with all three. What it does not yet have is dynamics, and the places where that shows – momentum, conservation in mergers, the surface-density dependence of the outer curves – are where a completion is needed, and where this note says so.

Three things are bought by making the mass real. Lensing and dynamics follow from the same mass. The non-baryonic (nb) mass lenses as ordinary mass: it is non-relativistic ($\sigma/c \sim 10^{-3.5}$, so its pressure, however read, is negligible) and has no anisotropic stress, so in the weak field it enters both metric potentials equally. What a relativistic completion must still supply is conservation of the total stress-energy when the nb mass changes, as a body's baryons are merged or stripped (Section 7); static and slowly evolving systems are unaffected. The flat weak-lensing profiles of isolated galaxies to 1 Mpc (Mistele et al. 2024b) then follow by construction, since the BTFR fixes the isothermal tail, and are a consistency check rather than a test. There is no suppressing external-field effect: the internal dynamics of a bound system depend on their surroundings only through a small enhancing term, where MOND's depend on the external field at first order. And the relation is present-tense: identical baryons today give identical nb mass, whatever happened before, so nothing depends on how the baryons came to be where they are.

The nb mass here is real and gravitating but is not cold dark matter. It has no independent dynamics – it is a response to the present baryon field, not a population of particles with its own orbits – and no formation history, so none of CDM's halo phenomenology follows from it: no merger-built structure, no dependence of present structure on its past, no dark sub-haloes without galaxies, no stripping. It shares with CDM only that it gravitates and lenses as mass, which MOND lacks; it shares with MOND that the baryons fix everything, which CDM lacks. How a mass without its own dynamics is read (Section 2.2) and sourced (Section 2.3) is stated as assumptions. The note makes no cosmological claim; CDM's results on the microwave background and large-scale structure are outside its scope, as is the microphysics of the nb mass and of the background it exchanges with.

Prior art. Real-mass models built to reproduce MOND's phenomenology exist. Superfluid dark matter (Berezhiani & Khoury 2015) obtains a MOND-like force from phonons in a condensed medium; dipolar dark matter (Blanchet 2007; Blanchet & Le Tiec 2008) from gravitational polarisation; and Famaey, Khoury & Penco (2018) derive, as the equilibrium of dark-matter–baryon energy exchange, a near-isothermal dark-matter fluid whose velocity dispersion obeys $v^4 \approx a_0 G M_b(< r)$ – the dynamical counterpart of our velocity rule, with a local rather than a per-body scale and with mass and momentum conserved by construction; it requires the product of the cross section and the energy exchanged per collision to scale inversely with the dark-matter number density $n_{\rm DM}$, i.e. a cross section $\propto 1/(n_{\rm DM} v^2)$ if that energy scales as v^2, and our relation can be read as the static, per-body limit of such a fluid; Bruneton et al. (2009) asked whether dark matter could play MOND's phantom role; Khoury (2015) combined MOND-like gravity with a dark-matter fluid; and relativistic theories recover MOND in the weak field, through an emergent-gravity vector field (Hossenfelder 2017) or a scalar–vector extension of general relativity (Skordis & Zlosnik 2021). Two of these have been confronted with the isolated-galaxy lensing data at small accelerations: superfluid dark matter is in tension with it (Mistele, McGaugh & Hossenfelder 2023a), and the data show no clear indication of the deviations AeST predicts (Mistele, McGaugh & Hossenfelder 2023b) – a test the present relation passes by construction and only because its tail is the BTFR's; the nu-HDM picture (Angus 2009) adds a real hot component to MOND for galaxy clusters and cosmology. The present relation differs from all of these in having no new force and no particle dynamics: the mass is a response to the potential, written in the mathematical form of the cored isothermal sphere, truncated as in the lowered isothermal models of King (1966), the same form that self-interacting dark matter analyses fit as an equilibrium core (Kaplinghat, Tulin & Yu 2016): Ren et al. (2019) fit SPARC with $\rho_0 \exp[(\Phi_{\rm tot}(0) - \Phi_{\rm tot})/\sigma^2]$, formally eq. (1) without the saturation, matched to an outer NFW halo through the concentration–mass relation. We instead fix σ by the BTFR, saturate the baryon term at a_0, truncate at the ambient density rather than match a halo, and fit one parameter; the empirical overlap is close by construction, and the claims differ where those choices bite (Newtonian wide binaries, the Mpc truncation). The Burkert profile approximates the same form (Burkert 1995; Salucci & Burkert 2000). What those fits treat as the thermal state of a collisional particle fluid, this note treats as a constitutive relation (Section 2.2); the empirical content overlaps, the reading does not.

The background is where the relation departs from all of these. It is posited, smooth, and enters the relation through one number: the ambient density at which each halo's isothermal tail ends, a_0^2/Gc^2 as the global mean (Section 2.1). It carries no force. Because the relation is written as the excess over the background, a region below the ambient density acts as negative source without any negative mass; the nb density itself is positive everywhere, which is what separates the relation from MOND's phantom density in embedded systems (Section 6.2). In AeST and the superfluid picture the background medium produces the MOND-like force; here it only sets the outer boundary of each halo – at the mean density itself, several Mpc out, not at a virial overdensity – and takes up the balance of mass and momentum when a body's nb mass changes (Section 7). From that one role follow the predictions that separate the relation from MOND (Section 6.3): a truncation radius that follows the local density rather than an external field, a downturn that persists in voids, no suppression of internal dynamics, and inverse-square attraction between bodies beyond their truncation radii. Two things remain open: the value of the ambient density is posited, and the lensing data so far bound it from above only (Appendix C); and the exchange with the background has no transport equation (Section 7). McGaugh (2026) sets as the test of any dark-matter candidate that it reproduce the MOND phenomenology without being fitted to it; the relation takes the two MOND limits as inputs and is tested on the transition and on where it departs from MOND (Sections 6.2–6.3).

Relation to MOND. The relation keeps the Poisson equation and puts the only new element in the source: $\nabla^2\Phi = 4\pi G\,(\rho_b + \rho_{\rm nb}[\Phi])$. MOND modifies the operator (AQUAL) or, equivalently, prescribes a phantom source fixed by $\nu(g_N/a_0)$ (QUMOND) that is not mass and can be negative. The present model takes MOND's phenomenology in its QUMOND form and replaces the phantom prescription by a real, positive density obeying a stated constitutive relation. The structure is inherited; the closure is posited. The two coincide where $\rho_{\rm nb} \sim \rho_{\rm ph}$ (galaxy centres) and differ at potential saddles, in environment, and at wide binaries; where they differ, the data decide. It is not a reformulation of MOND.

Scope. This note supersedes the author's v1.0.14 preprint (Appendix F), posts before the Gaia DR4 wide-binary results, and is deliberately narrow. In scope: the rotation curves and acceleration relation of disc galaxies, dwarf dispersions, wide binaries, and the weak-lensing profiles of isolated galaxies. Out of scope: cosmology and galaxy clusters. The cosmic background enters only through the baseline of the potential and the ambient term (Section 2.1), whose global mean we take at the closure value $a_0^2/Gc^2 \approx 0.8\,\rho_m$ (Section 2.1), a numerical relation we use but do not derive; whether the nb mass is the cosmological dark matter is not addressed. Galaxy clusters are deferred: the source rule of Section 2.3 counts a cluster's hot gas as a body, which is where a treatment would start (Section 7). The first problem beyond galaxies is the Mpc-scale budget of the nb mass (Appendix C).

2. The relation

2.1 Statement

The bump picture. The relation responds to the potential measured from a baseline, not from zero. In an empty universe a body's potential could be set to zero at infinity; the universe is not empty. Summed over the matter within the causal horizon, the potential at any point is of order $\Phi_{\rm bg} \sim G\bar{\rho}L^2$, and Sciama's condition (Sciama 1953; Brans & Dicke 1961) sets it at $\Phi_{\rm bg} \sim c^2$, which fixes the horizon length as $L \sim c/\sqrt{G\bar{\rho}}$; the sum is finite only because it is cut there. The background is uniform on galactic scales, so it exerts no force, but it sets the level from which each body's potential rises: $\varphi_{\rm loc} = \varphi_{\rm tot} - \Phi_{\rm bg} > 0$ inside any bound structure. This is the bump picture, and it is why the relation can respond to the value of the potential and not only to its gradient. The same condition gives the background a gradient scale $\Phi_{\rm bg}/L \sim c\sqrt{G\bar{\rho}}$; setting it equal to a_0 gives the closure value of the ambient density below, up to a factor of order unity and the choice of mean density (all matter, or the nb mass alone, differing by ∼8% in a_0); both are subject to further investigation (Appendix G).

Let $\varphi_b(r) \geq 0$ be the potential of the body's own baryons in the bump convention (positive, rising inward, zero at infinity). It is the baryonic part of the local bump $\varphi_{\rm loc} = \varphi_{\rm tot}$ - $\Phi_{\rm bg}$; the nb part is $\varphi_{\rm nb}$ below. A constant external potential belongs to $\Phi_{\rm bg}$ and does not enter the relation; an external gradient enters only through the truncation radius (tides) and the nesting rule of Section 2.3. This is the definition under which the relation has no suppressing external-field effect; it is an assumption of the source variable, not a derived property, and Section 4.4 and 6.2 test it. Let $\varphi_{\rm nb}$ be the potential of the nb mass itself, in the same convention; both potentials enter the relation as their drop from the centre. The nb density is

$$\rho_{\rm nb}(r) = \rho_c \exp\left[S\big(x_b(r)\big) - S(x_{b0}) - \frac{\delta\varphi_{\rm nb}(r)}{\sigma^2}\right] - \rho_{\rm amb}, \qquad x_b = \frac{\varphi_b}{\sigma^2}, \tag{1}$$

with $S(x) = 2\tanh(x/2)$, which equals x for $x \ll 2$ and saturates at 2; x_{b0} is the central value; $\delta\varphi_{\rm nb}(r) = \varphi_{\rm nb}(0) - \varphi_{\rm nb}(r)$ is the nb mass's own potential drop from the centre (the baryon term's drop is $\sigma^2\,[x_{b0} - x_b(r)]$); and

$$\sigma^2 = \tfrac{1}{2}\sqrt{G\,a_0\,M_b}, \tag{2}$$

is the velocity parameter, fixed for each galaxy by its baryonic mass through the baryonic Tully-Fisher relation with $a_0 = 1.2 \times 10^{-10}\,{\rm m\,s^{-2}}$. We call eqs. (1)–(2) the saturated isothermal relation: saturated because the baryon response S is capped, isothermal because the nb mass takes the cored isothermal form with velocity parameter σ. It is *the relation* below, and *this work* in the figures. ρ_c is the central nb density, one number per galaxy, and $\rho_{\rm amb}$ is the ambient nb density of the surroundings; eq. (1) is the EXCESS over ambient, which is what sources the body's own potential and its lensing signal, and it vanishes where the exponential tail falls to $\rho_{\rm amb}$. (Every solve in this note includes the subtraction; earlier statements of the relation without it were shorthand.) The total potential obeys the unmodified Poisson equation with source $\rho_b + \rho_{\rm nb}$, so (1) is solved self-consistently (the $\varphi_{\rm nb}$ term is the resummed self-gravity of the response; nothing is added twice). Far from the baryons the solution is the cored isothermal tail, $\rho_{\rm nb} \sim 1/r^2$ and a flat circular speed $v^2 = 2\sigma^2$, out to the radius $r_t = \sigma/\sqrt{2\pi G\rho_{\rm amb}}$ at which the tail density $\sigma^2/(2\pi G r^2)$ meets the ambient density $\rho_{\rm amb}$ (taken as $35.5\ M_\odot\,{\rm kpc^{-3}} = a_0^2/Gc^2$, about 0.8 of the cosmic mean matter density, for isolated galaxies; this is the numerical coincidence $a_0 \sim cH_0$ noted by Milgrom (1983), written as a density, which we use but do not derive (Appendix G); the closure value is posited as the global mean of a smooth background field, and the local ambient density, around galaxies or in voids, may lie below it, Section 7); the core radius is $r_c = 3\sigma/\sqrt{4\pi G\rho_c}$.

What is assumed and what is derived is set out in Table 1. a_0 enters once, through σ, and from there twice: as the low-acceleration normalisation ($v^2 = 2\sigma^2$ gives $v^4 = Ga_0M_b$, the BTFR) and as the saturation point, since for a point mass $x_b = 2\sqrt{g_b/a_0}$ and the cap $S = 2$ is reached where $g_b = a_0$.

Table 1. Inputs and outputs of the relation.

INPUT	value	how fixed
a_0	$1.2 \times 10^{-10}\,{\rm m\,s^{-2}}$	BTFR normalisation (the only acceleration scale)
$\sigma^2 = \sqrt{Ga_0M_b}/2$	per galaxy	BTFR as input; a_0 enters here
saturation of S at 2	fixed	the point-mass identity $x_b = 2$ at $g_b = a_0$ (the Newtonian limit, an input); the acceleration relation prefers 1.7-2 (0.07 dex per unit); required given the BTFR σ (rotation-curve χ^2 halves; with σ free the unsaturated relation fits as well)
form of S	tanh	up to 0.04 dex at fixed cap (0.01–0.02 for erf and exponential forms, 0.04 for a hard cap or a rational form); tilts the transition, per-point rms unchanged (0.155–0.160); fixed before any fit
$\rho_{\rm amb}$	$35.5\ M_\odot\,{\rm kpc^{-3}}$	$= a_0^2/Gc^2 \sim 0.8\,\rho_m$ (posited global mean; local values may be lower); the subtracted ambient term in eq. (1); sets the truncation radius
stellar M/L	0.5 (disc), 0.7 (bulge)	SPARC convention; fixed
ρ_c	one per galaxy (full-sample median $10^{7.42}$, spread 0.68 dex); $10^{7.45}$ adopted for the no-parameter results (the median of an earlier 120-galaxy fit)	fitted; the sample median for the global value

INPUT	value	how fixed
OUTPUT		
the BTFR	exact	by construction (input)
the low-acceleration limit $g \to \sqrt{g_b a_0}$	exact (input)	isothermal tail with the BTFR sigma
the Newtonian limit	exact (input)	saturation at $g_b = a_0$
the transition shape	follows, with one fitted global density	self-consistent solve; no interpolating function fitted (Section 3.1)
core surface density $\sim a_0/2\pi G$	reproduced on average	from the fitted ρ_c (Section 3.3)
wide binaries Newtonian	conditional	provided the binary's own central density is below $\sim 10^{10}\, M_\odot\,\mathrm{kpc}^{-3}$ (Section 6.1)

2.2 Reading

The relation adapts the mathematical form of the cored isothermal sphere into a constitutive relation: the isothermal response to the potential, with the baryon term saturating where $g_b = a_0$ and the ambient density subtracted. σ is a parameter of the relation, not the temperature of a dynamical medium. The relation fixes where the nb mass is at each instant, not how it gets there. We take it to be re-established around each body almost instantaneously (on timescales short compared with dynamical times, not faster than light: the relation is elliptic, as the Newtonian Poisson equation is, and a relativistic completion must supply the propagation), in the configuration the relation prescribes for the present baryons; whether the nb mass is carried with its body or re-formed in place the relation does not specify. A lag or a wake would appear only under accelerations fast compared with its response, where a dynamical completion is needed. This dynamical assumption is separate from the relation; the consequences that rest on it (no dynamical friction, Section 6.4; the momentum and mass bookkeeping, Section 7) are labelled as such. Stability limits for self-gravitating gas or stellar systems therefore do not apply; the relevant question is well-posedness, and beyond a central contrast of ~14 the bounded equation admits several solutions for the same outer boundary. Integrating from the centre at given ρ_c is unique, so the multiplicity is absorbed by ρ_c: this is a candidate reason the relation does not fix ρ_c.

2.3 Source definition (assumptions)

A body is a baryonic system taken as a whole: a star, a star cluster, a galaxy, a binary star, a gas cloud, or a galaxy cluster's hot gas. Every system sources, and the nb mass it carries is set by its ρ_c: low for diffuse systems, whose cores are then comparable to their own size, so that a diffuse system carries some nb mass but may not be bound by it; whether a system ends up bound is an outcome, not a rule. Individual particles must carry no excess (ρ_c at the ambient value), which the relation does not yet derive (Section 7). The amplitude of a body's response is set by its total bound baryons, gas included (in SPARC, with stars alone the ratio of observed to predicted outer speed is 1.45 for galaxies with gas fraction above 0.5 and 1.74 above 0.8, against 1.01 with gas included; N = 43 and 11); the location is the body's. Each nb population carries the velocity parameter of the body that sources it; a host's nb mass inside a nested body is boosted by $\exp(\varphi_{\mathrm{body}}/\sigma^2_{\mathrm{host}})$, capped at e^2, which reduces to pass-through below the LMC scale (boost 1.000 for Palomar 5, 1.008 for Fornax, 1.36 for the LMC, with host $\sigma \approx 125$ km/s from the Milky Way's BTFR). These are assumptions. Which collections count as bodies is chosen by the analyst in each application (the galaxy for rotation curves, the pair for wide binaries, the satellite within its host for dwarfs); the nesting rule fixes how nested bodies combine but does not derive the decomposition (Section 7). A diffuse gas halo has no central concentration, which is consistent with the merging-cluster population in which the lensing peaks sit with the galaxies and not with the gas (Harvey et al. 2015); they enter the dwarf protocol of Appendix B through the host's ambient term and the satellite truncation of Appendix A; their direct tests are named in Section 7. Appendix F of v1.0.14, which argued the uniqueness of the linear law, is superseded; no replacement uniqueness claim is made.

3. What the relation reproduces

Sample and method. We use the SPARC rotation curves (Lelli, McGaugh & Schombert 2016a). We adopt the selection of McGaugh et al. (2016): quality flag ≤ 2, inclination ≥ 30 deg, and points with velocity errors below 10%, which leaves 2696 points in 147 of the 153 galaxies that pass the first two cuts (the six without any point below 10%: UGC 05005, UGC 07559, UGC 07577, UGC 07866, UGC 09992, UGCA 444) (McGaugh et al. report 2693). The relation's velocity parameter comes from the baryonic mass (eq. 2), so no flat velocity is needed; the 27 galaxies whose curves do not reach a flat part (Lelli et al.'s $V_f = 0$) are kept and reported separately, since they populate the lowest accelerations. (Earlier versions required $V_f > 0$, which removed them and flattered the model there.) Stellar mass-to-light ratios 0.5 (disc) and 0.7 (bulge), gas mass 1.33 $\times$ HI for helium (the SPARC gas velocities already include this factor). Each galaxy is solved spherically in the observed baryon profile. The baryon potential is built from the SPARC baryonic rotation curve as $\varphi_b(r) = \int_r^{r_{\rm last}} v_b^2\, dr'/r' + v_b^2(r_{\rm last})$, i.e. a point-mass continuation beyond the last measured radius, with negative $v_{\rm gas}^2$ set to zero; x_{b0} is therefore sensitive to gas beyond the last point in gas-rich galaxies, a sensitivity not yet quantified; the fit statistic is $\chi^2 = \sum w\,(v_{\rm model} - v_{\rm obs})^2$ with $w = (0.1\, v_{\rm obs})^{-2}$ (not divided by N; ~24 points per galaxy, so $\chi^2 \sim 14$ is $\chi^2/N \sim 0.6$).

3.1 A cored isothermal response with BTFR normalisation reproduces the acceleration relation; the transition shape follows from the self-consistent solve

Fix $\rho_c = 10^{7.45}\ M_\odot\,{\rm kpc}^{-3}$ (the sample median) for every galaxy, so the model has no per-galaxy parameter, and solve all of them. Figure 1 shows the emergent ratio $\nu_{\rm eff} = g_{\rm model}\ /\ g_{\rm bar}$ at the data radii, binned in $\log(g_{\rm bar}/a_0)$, against the binned data. On the full sample the model follows the binned data to within 0.04 dex for $\log(g_{\rm bar}/a_0) \geq -1.25$, but lies above the data by 0.05, 0.08 and 0.06 dex in the three bins from -2.0 to -1.25, where the fitted function of McGaugh et al. (2016) stays within 0.033 dex; the galaxies without a flat part, kept now, are the worst (median 0.07 dex above). Overall the per-point rms is 0.155 dex for the model at this single density against 0.133 for the fitted function. Setting the density instead by the empirical relation of Appendix B, $\rho_c(M_b, \Sigma_h, f_{\rm gas})$ with four global constants and no per-galaxy freedom (here $\Sigma_h = M_b/2\pi R_h^2$, in $M_\odot\,{\rm pc}^{-2}$ wherever relation (3) is evaluated, with $R_h = 1.678 R_d$ the projected half-mass radius of the SPARC exponential disc, i.e. a mean baryonic surface density inside the half-mass radius, not the SPARC $\Sigma_{\rm eff}$), brings the transition bins to within 0.01 dex of the data, halves the radial trend of the residuals, and gives an rms of 0.140 – but leaves the low-acceleration offsets (0.04–0.08 dex) and the surface-density trend of the outer speeds (Section 4.1) in place: that residual is not in ρ_c. The model's own scatter about the acceleration relation (0.07–0.11 dex at low accelerations, from the variety of baryon profiles) is not to be compared with the observed scatter directly: injecting the catalogue distance and inclination errors, 0.1 dex in the mass-to-light ratio and the velocity errors into the model curves gives 0.11–0.16 dex per bin, against 0.10–0.23 in the data; from $\log(g_{\rm bar}/a_0) = -1.75$ to -0.25 the model with errors already scatters as much as the data or more (0.13–0.16 against 0.12–0.16), leaving no room for intrinsic scatter there. That is a tension, not a consistency check. The residuals of the model also trend with radius (Spearman -0.17 per point, against -0.04 for the fitted function) and slightly more strongly with surface density and gas fraction. Figure 1 shows the full sample.

The two limits are exact by construction; the transition shape between them follows from the self-consistent solve: no interpolating function is fitted, the saturation is fixed at $g_b = a_0$ before any fit, its functional form shifts the binned relation by at most 0.04 dex (below), and the only fitted quantity is one global density. The count is therefore a_0 plus one fitted normalisation, against $g_\dagger$ plus a chosen function for the empirical relation. Its sensitivity: changing the FORM of S at the same saturation value moves the binned relation by up to 0.04 dex (0.01–0.02 for erf and exponential forms, 0.04 for a hard cap or a rational function), tilting the transition without changing the per-point rms (0.155–0.160) or the scatter comparison below; the hyperbolic tangent is fixed before any fit; changing the VALUE of the saturation moves it by ~0.07 dex per unit around 2 (Figure 1b), and the empirical relation itself prefers a value between 1.7 and 2, i.e.~the identity $g_b = a_0$. The single density moves the relation nearly uniformly (~0.10 dex per 0.35 dex in ρ_c): ρ_c is a normalisation, the shape is the model's.

Three honest limits of this result.

(i) The normalisation is one global number taken from the data (the sample median); an independent choice, ρ_c from the Donato et al. core surface density, gives $10^{7.2}$ and lands within 0.1 dex of the data, uniformly low.

(ii) MOND predicts zero intrinsic scatter about the relation; this model at universal ρ_c predicts 0.07–0.11 dex at low accelerations, because galaxies with different baryon profiles reach different accelerations at the




Figure 1: Figure 1. (a) The radial acceleration relation on the full sample (147 galaxies, 2696 points; selection of McGaugh et al. 2016): binned data (squares), the empirical function (dashed), and the model at the universal density $10^{7.45}$ (red), with ρ_c from relation (3) (blue), and with ρ_c fitted per galaxy (green). (b) The median residual $\log(g_{\rm obs}/g_{\rm model})$ in octiles of baryonic surface density Σ_h: the model at universal density runs from -0.10 dex at the lowest surface densities to $+0.05$ at the highest, relation (3) halves that, and the empirical function itself is not flat (-0.04 to $+0.03$). The saturation-value and form sensitivity formerly shown here is unchanged (0.07 dex per unit; form up to 0.04 dex, Section 3).

same $g_{\rm bar}$. The tightness of the relation is itself one of the regularities to be explained, so matching its mean is not enough. The intrinsic scatter has been measured: 0.034 ± 0.001 (stat) ± 0.001 (sys) dex on SPARC, assuming the SPARC error model is reliable (Desmond 2023; Li et al. 2018 account for most of the observed 0.057 dex with errors); both this estimate and our error-injection test rest on that model, and uncertainties overstated in it would leave more room for intrinsic scatter. That estimate marginalises over mass-to-light ratio, distance and inclination, and our 0.07–0.11 dex is at fixed values, so the two are not directly comparable; but at the same baryon models the model's per-point rms exceeds the acceleration relation's by 0.08 dex in quadrature (0.155 against 0.133). A universal ρ_c is therefore in tension with the data; a marginalised fit of the model, which we have not done, would decide. A density that varies with galaxy properties reduces the excess but does not remove it (relation (3): 0.140, 0.044 dex in quadrature, still above the measured 0.034); both quadrature figures subtract rms values of different models on the same points and are indicative only. Each galaxy is fit as well as by the empirical relation with its mass-to-light ratio free (Table 2); what is not yet explained is why the central density varies with galaxy properties just enough to keep the relation tight.

(iii) At the single value the model under-performs the empirical relation in gas-dominated and slowly rising galaxies, where the fitted densities are lower (median $10^{7.19}$ for the galaxies without a flat part); that is where the low-acceleration offset comes from.

3.2 Individual rotation curves, and the comparison with the acceleration relation

On the full sample the median per-galaxy χ^2 is 25.5 at the universal density and 6.0 with ρ_c free (Figure 2). The relation with its BTFR σ does not describe individual curves to the data's precision; freeing σ per galaxy roughly halves the median χ^2 again (Appendix A.1).

Table 2 places the model and the empirical acceleration relation on the same footing. On the full sample, per point, the model at a single ρ_c (0.155 dex) is worse than the relation at fixed M/L (0.133); with one free parameter each the two are equal within the noise (0.081 for the model with ρ_c free, 0.079 for the relation with the disc M/L free), and the relation with M/L, distance and inclination free under the SPARC priors reaches 0.067. The relation does slightly better in the galaxies without a flat part (0.072 vs 0.082). The fitted densities have median $10^{7.42}$ and spread 0.68 dex. (An earlier 120-galaxy comparison favoured the model, 0.108 vs 0.125; it does not




Figure 2: Figure 2. Per-galaxy χ^2 (10% velocity weights) on the full sample at the universal density (median 25.5) and with ρ_c fitted (median 6.0). The earlier three-way comparison (no saturation, saturated, σ free) is given in Appendix A.1.




Figure 3: Figure 3. Three rotation curves: observed (points), baryons (dotted), the model at the single density $10^{7.45}$ (blue) and at the fitted density (red). DDO 154 is gas-dominated; NGC 2403 intermediate; NGC 2841 a massive bulge galaxy whose outer curve the model under-predicts by ~25% at either density – the kind of residual that the free-sigma fit of Figure 2 absorbs and the BTFR σ does not.

survive the matched sample.) We draw no conclusion beyond "comparable". For scale, self-interacting dark matter fits to SPARC carry two halo parameters per galaxy in addition to the mass-to-light ratio (Kaplinghat, Ren & Yu 2020); this model carries one, with the mass-to-light ratio fixed.

Table 2. Per-point rms of $\log(g_{\rm obs}/g_{\rm model})$, 2696 points, 147 galaxies (selection of McGaugh et al. 2016).

model	free per galaxy	rms (dex)
acceleration relation, M/L fixed	none	0.133
this model, $\rho_c = 10^{7.45}$	none	0.155
acceleration relation, M/L_disc free	1	0.079
this model, ρ_c free	1	0.081
acceleration relation, M/L, D, i free (SPARC priors)	3	0.067
(Li et al. 2018, MCMC, per-point errors)	3	0.057 – not like-for-like

The Milky Way is a case the sample does not contain: a compact, massive disc whose Sun sits at about one MOND radius, and whose Gaia DR3 circular-velocity curve declines from 233 km/s at the Sun to ~175 at 27 kpc (Ou et al. 2024). The relation is consistent with that decline, from their baryon model at the universal ρ_c (rms 20 km/s over 8–27 kpc, slope −1.9 against the observed −2.5 km/s/kpc), within 3 km/s of MOND on the same baryons at every radius; with ρ_c fitted ($10^{7.88}$, inside the sample spread) the rms is 9 km/s (slope −3.0 against the observed −2.5). The decline is the transition toward the BTFR speed of 177 km/s, not a Keplerian fall: at the Sun the baryons alone give 173 km/s and the nb mass the rest. Appendix A gives the numbers and the outer prediction.

3.3 The core surface density

MOND obtains the halo core surface density from a_0 (Milgrom 2009). In galaxies whose mean acceleration is at or above a_0 it is $\gamma\Sigma_M$, with $\Sigma_M = a_0/2\pi G \approx 138\,M_\odot\,{\rm pc}^{-2}$ and γ of order unity, fixed by the interpolating function. At low surface density it falls as $\sqrt{\Sigma_b}$. Donato et al. (2009) measure $\log\,\mu_0 = 2.15 \pm 0.2$. The fitted ρ_c of this relation implies a median $\log\,\mu_0 = 2.24 \pm 0.39$, with $\mu_0 = \rho_c r_c$: reproduced on average, not predicted. Figure 4 compares the two galaxy by galaxy. MOND's value is Milgrom's estimate (his eq. 11), which he calls rough. We evaluate it with the interpolating function of the acceleration relation, which gives $\gamma = 2.09$, and with Σ_b the mean baryonic density inside the half-mass radius times that radius ($\Sigma_b = 0.75\,\Sigma_h$). Against baryonic mass the SPARC Burkert fits of Li et al. (2020) rise with slope +0.21. This relation's values rise with +0.24. MOND's rise with +0.15 (+0.13 to +0.16 with Σ_b halved or doubled). Fitted on mass and surface density together, the Burkert values go as $M_b^{0.14\pm0.04}\Sigma_h^{0.20\pm0.07}$ and this relation's as $M_b^{0.16\pm0.05}\Sigma_h^{0.22\pm0.07}$. MOND's depend on Σ_b alone, as $\Sigma_h^{0.40}$, with no mass term. Galaxy by galaxy the rms offset from the Burkert values is 0.26 dex for this relation and 0.30 for MOND (109 galaxies in both). Two cautions keep this from being a discriminator. First, this relation's values come from one density fitted per galaxy to the same rotation curves the Burkert fits use, so the agreement is partly shared input; MOND's values carry no per-galaxy freedom. Second, the mass trend depends on the mass-to-light ratio. At a disc M/L of 0.7, or a mass-dependent one, ρ_c falls with mass (slope −0.26 to −0.30) and the implied μ_0 slope drops to ~0.12; MOND's slope at that M/L was not recomputed. We read the comparison as Table 2 reads the rotation curves: a comparable description with a different parameter count. What the relation lacks is in the amplitude of the tail, which it sets by M_b alone (Section 4.1).

4. Where the model is incomplete

4.1 Surface-density dependence at fixed mass

The dependence on surface density at fixed baryonic mass, which MOND predicts, is present in the data. In the relation it is missing from the tail, whose amplitude σ is set by M_b alone. (The fitted ρ_c does depend on surface density – Section 3.3 and relation (3) – but ρ_c sets the core, not the tail.) It shows directly in the outer curves: at the single density the ratio of model to observed speed at the last measured point has median 1.02 over 147 galaxies, with no trend in baryonic mass but a clear one in surface density (Spearman −0.30, $p = 2\times10^{-4}$), from 1.085 in the lowest-surface-density quartile to 0.98 in the highest. The excess does not correlate with inclination,




Figure 4: Figure 4. Halo core surface density μ_0 against baryonic mass, 120 SPARC galaxies at the SPARC mass-to-light ratio. Squares: Burkert fits to the data (Li et al. 2020); dotted line, their trend. Red points: this work, $\mu_0 = \rho_c r_c$ with ρ_c fitted per galaxy and the King radius $r_c = 3\sigma/\sqrt{4\pi G\rho_c}$; this is a King core, not a Burkert refit, so the comparison with Burkert $\rho_0 r_0$ is approximate. Open triangles: MOND's value for each galaxy from its baryons (Milgrom 2009, eq. 11), with the interpolating function of the acceleration relation ($\gamma = 2.09$) and $\Sigma_b = 0.75\,\Sigma_h$. Solid lines: least-squares trends. Dashed: $\Sigma_M = a_0/2\pi G$. Band: Donato et al. (2009), 2.15 ± 0.2. The red points use one fitted density per galaxy; the triangles use none (Section 3.3).

inclination error or distance error, but it does shrink with data quality: in the 69 galaxies with quality flag 1, inclination $\geq 45^\circ$ and inclination error $\leq 5^\circ$ the trend is not significant (Spearman -0.15, $p = 0.22$) and the lowest-quartile excess is 4%, against -3% in the highest. Correcting the outer speeds for pressure support in the gas (asymmetric drift; gas dispersion 6–12 km/s, as measured in dwarfs and outer discs by Ianjamasimanana et al. 2015 and Iorio et al. 2017; exponential outer gas disk) removes about a third of the lowest-quartile excess at the universal density (8.5% to 5.5% at 8 km/s) and little of it under relation (3); with the drift term and $V_{\rm max}$ controlled, the surface-density term survives at 2.4σ. We read it as a residual of 3–8% with MOND's sign, partly data quality and partly pressure support; it tracks gas fraction, and it is not removed by any choice of ρ_c (Section 3.1), so it sits in the amplitude of the tail, which the relation fixes by M_b alone. The stellar mass-to-light ratio does not absorb it: at a disc M/L of 0.4, 0.5 and 0.7 the lowest-quartile excess is 5, 8 and 13% and the highest-quartile ratio 0.94, 0.98 and 1.09, so a higher M/L narrows the differential only by over-predicting every galaxy, and the low-acceleration offsets grow with M/L; 0.5, the population-synthesis value (Schombert, McGaugh & Lelli 2019), gives the best rms of the three. Freeing the velocity parameter per galaxy shows where the residual lives: the BTFR value is right on average (median offset -0.006 dex in σ^2), and the fitted offset tracks gas fraction (-0.49 ± 0.15 per unit $f_{\rm gas}$), not surface density, once both are included. Setting σ by the baryonic mass enclosed within the last measured point, instead of the total, removes the gas-fraction trend and most of the low-surface-density excess, but a velocity parameter that depends on radius is a different relation (Section 7) and is not attempted here. The isothermal form also overshoots its asymptote near three core radii, and the residuals at the last point show that hump (peak at $R \approx 2.8\,r_0$, 3σ), independent of the surface-density trend. Over the galaxies with a flat part the relation reaches V_f as the data do (median model 1.002, observed 1.006 of V_f at the last point), so the shortfall seen in individual massive discs (Figure 3) is not a sample property.

4.2 Crater II

McGaugh (2016) predicted $\sigma = 2.1$ (+0.9/-0.6) km/s from the external-field effect before Caldwell et al. (2017) measured 2.7 ± 0.3. At $\rho_c = 10^{7.45}$ the relation gives 5.2 km/s; it reaches 2.7 only with $\rho_c \sim 10^{6.1}$, 1.3 dex below the median. The surface-density trend alone (ρ_c against M_b and Σ_h) supplies a third of that, predicting $10^{7.0}$; the full relation (3), in which the gas-fraction term raises ρ_c for a gas-free object, predicts $10^{8.7}$, i.e.~it moves the wrong way; the dispersion there is 5.1-5.4 km/s because above $\sim 10^{7.5}$ the projected dispersion saturates near the relation's velocity parameter (5.7 km/s for Crater II). Crater II's tidal disruption has been reported (Vivas et al. 2025, preprint). Quantified under the relation, with its own Milky Way curve and the pericentre of Ji et al. (2021), 21.7 kpc, truncation at the pericentric Jacobi radius brings the universal-ρ_c value to 3.1–2.8 km/s, within the bar of Caldwell et al. and, in the stronger case, of Ji et al.'s 2.34 km/s; Antlia II, observed to be disrupting, passes in the same cases (Appendix B). Because that radius (0.7 kpc) lies inside today's half-light radius, these are bounds, not a model; a stripping calculation is owed. The mechanism is the tidal one CDM uses (Sanders, Evans & Dehnen 2018; Borukhovetskaya et al. 2022). Status: conceded at equilibrium; the tidal reading is listed as open until a stripping calculation exists (Section 7).

4.3 NGC 1052-DF2 and DF4

Both have stellar dispersions consistent with their stars alone: 8.4 ± 2.1 km/s (Danieli et al. 2019) and 8.0 (+2.3/-1.9) (Shen et al. 2023), at 20-22 Mpc (Shen et al. 2021), with $M_* \sim 2 \times 10^8$ (van Dokkum et al. 2018) and ~1.5e8 $M_\odot$ (Shen et al. 2023 adopt $M_* = 7.5 \pm 2 \times 10^7$ inside the 3-D half-light radius at M/L = 2, from van Dokkum et al. 2019; their dynamical mass there is $8\,(+6/-4) \times 10^7$ against a stars-only expectation of 7 $\pm$ 1 km/s). The relation at the sample density gives 17-22 km/s; it needs ρ_c about 1.5 and 1 dex below the SPARC galaxies of the same mass and surface density, which is not an extrapolation of relation (3) but an outlier to it. The relation has no lever here. They are anomalous for isolated MOND as well (~20 km/s); MOND recovers 13.4 (+4.8/-3.7) through the external field of NGC 1052 at a separation not known in three dimensions (Famaey, McGaugh & Milgrom 2018; Haghi et al. 2019). We list them as unexplained because the relation has no comparable lever, not because MOND predicts them.

4.4 Environment

The relation predicts that the shape of a rotation curve does not depend on the external field of neighbouring galaxies: the ambient density at a field galaxy is $<\sim 630\ M_\odot$ kpc^-3 (Appendix C) against a core density $\sim 10^{7.5}$, and the nesting boost is ~1. On SPARC the low-acceleration residual from the acceleration relation varies with

the environmental field of Chae et al. (2021) as -0.12 ± 0.16 dex per dex at fixed mass and surface density; the external-field effect predicts -0.08 for this sample, the relation 0; the sample does not separate them. Applying their own statistic (a fitted external-field parameter per galaxy) to the relation's curves returns a positive median too, so their median detection does not discriminate the relation; the dependence of that parameter on the field is weak and not of the external-field sign on the galaxies with field values; their cosmic-web regional contrast is untested here. Appendix D. This is the note's open weak point.

Three of these (4.2-4.4) are one problem stated three times: the relation's environmental effect is enhancing and small, so wherever MOND explains a low dispersion by suppression from an external field, this relation must explain it with a low ρ_c – and it does so for And XIX, not for Crater II (the surface-density trend alone goes a third of the way; the full relation does not; a tidal reading is open, Section 7), and not for DF2 and DF4.

5. What real mass buys

The three purchases of the introduction, with their evidence in Appendix C: lensing and dynamics from the same mass (it lenses as ordinary mass), compatible with the flat isolated-galaxy profiles of Mistele et al. (2024b) to 1 Mpc, with a truncation predicted at 0.8–3.4 Mpc for an isolated L^* galaxy, and a Mpc-scale budget that is open, with a mean-field indication that it can close (Appendix C); no suppressing external-field effect (the environmental coupling is small and enhancing); and a present-tense relation, so that identical baryons today give identical nb mass whatever came before.

The ambient term does four things at once:

- it truncates each halo where the tail meets the local nb density, so the cut tracks environment through density rather than through an external field;
- it keeps the downturn in voids, at larger radii;
- it carries no external-field suppression of internal dynamics;
- and it makes the force between bodies farther apart than their truncation radii inverse-square, as the pairwise kinetic Sunyaev-Zel'dovich measurement finds (Gallardo et al. 2026; Appendix E).

These, not the galaxy-scale fits, are where the relation and MOND part.

Positivity. In MOND's field formulations the phantom density turns negative in a region around the point where an embedded system's internal field cancels the external one (Milgrom 1986). For MOND this is harmless, since the phantom density is not mass. For a theory in which the missing mass is real it is a constraint: real mass cannot be negative, so such a theory cannot follow MOND there. This relation has no phantom density: the extra mass is real nb mass, positive everywhere, and the departure is a prediction: Newtonian dynamics for wide binaries in the Galactic field, provided their own central density stays low (Section 6.1). The argument does not show that MOND fails; it shows where a real-mass account must differ from it.

6. Predictions and falsifiers

6.1 Wide binaries

Status: live; the Newtonian value is shared with Newton and ΛCDM, so this test discriminates against MOND only. Let γ be the ratio of the observed to the Newtonian relative acceleration of a wide pair at given separation. Because the response density is bounded above by $e^2\rho_c$ – below a few solar masses per cubic parsec for any fitted ρ_c – a wide binary encloses a negligible nb mass: for a 2 $M_\odot$ pair at 7-28 kAU, γ - 1 = 2e-5 to 2e-3, under either choice of velocity parameter and in the halo as in the disc. To reach $\gamma = 1.1$ at 28 kAU would require 19 $M_\odot\,\mathrm{pc}^{-3}$ inside the orbit, two decades above any density on file. The relation predicts $\gamma = 1.000$ ($+< 0.003$) provided the binary's own central density is below $\sim 10^{10}\,M_\odot\,\mathrm{kpc}^{-3}$ ($\gamma - 1 < 0.1$ at 28 kAU); the globular-cluster bound ($\lesssim 10^{7.2}$, Appendix A) clears this by three decades, and relation (3), calibrated on discs, does not extend to stellar systems. Conversely, Newtonian data at 1% precision would bound it below $10^{9.2}$. We state in advance what is being tested: eq. (1) with the central density of stellar systems bounded by the globular-cluster value. A confirmed $\gamma \geq 1.1$ falsifies that model; raising the binary's density afterwards to $\sim 10^{10}$ would be a new hypothesis, not a rescue of this one. A variant in which the host's nb mass is depleted around the

pair, keeping the total non-negative, gave $\gamma \approx 1.1$–1.25 (Appendix F); it is a different relation from eq. (1), and DR4 separates 1.00, ∼1.2 and 1.3. MOND with the Galactic external field predicts $\gamma = 1.28$-1.36 at separations of 1-4 MOND radii, i.e.∼7-28 kAU for a 1 $M_\odot$ system and 10-40 kAU for 2 $M_\odot$ (two-body QUMOND formalism: Pflamm-Altenburg 2025); the relation's γ - 1 < 0.003 holds across that range. The present data are disputed. Cookson et al. (2026), Banik et al. (2024) and Pittordis et al. (2025) find Newtonian behaviour at all separations (Newton up to 1500x more likely than MOND in the cleanest sample of Cookson et al.). Chae (2023, 2024a,b, 2025) reports a boost at $g_N \lesssim 10^{-9}\,\mathrm{m\,s^{-2}}$ ($\gamma_g \approx 1.3$–1.5), with Newton rejected at 5–10σ in his samples, and Chae et al. (2026) find $\gamma = 1.60^{+0.24}_{-0.14}$ in 36 systems with accurate 3-D velocities; Hernandez et al. (2024) also report a boost; and Chae & Yoon (2026) argue that the data-quality cuts used by Banik et al. and Cookson et al. bias the result toward Newton. The relation's $\gamma = 1.000$ is therefore already disfavoured by one side's analyses and consistent with the other's. The falsifier we commit to is a confirmed $\gamma \geq 1.1$ from Gaia DR4 under a selection both sides accept: an error cut on the projected velocity in $\mathrm{m\,s^{-1}}$ rather than on the scaled velocity, a multiplicity fraction calibrated in the Newtonian regime of the same sample, hundreds of pairs beyond one MOND radius, and a stated γ convention. The MOND reference values above are a two-body QUMOND calculation at fixed separation, benchmarked against the independent two-body solutions of Pflamm-Altenburg (2025) to within 0.03 at 7-28 kAU; his finding that the one-body (test-particle) approximation overstates the boost (up to 1.44 against 1.2-1.3) applies to analyses that use it, not to these values.




Figure 5: Figure 5. Wide binaries: the relation predicts $\gamma = 1.000$ (+< 0.003) at all separations for binary central densities below $\sim 10^{10}\,M_\odot\,\mathrm{kpc}^{-3}$ (red); MOND with the Galactic external field, two-body QUMOND for a 1 $M_\odot$ system, gives 1.28-1.36 (blue; separations scale with mass$^{1/2}$). The green band marks the Newtonian results of Cookson et al. (2026), Banik et al. (2024) and Pittordis et al. (2025); the magenta point is Chae (2023). The status of these data is disputed (Section 6.1).

6.2 Environment

Status: live; paired with 6.1, opposite MOND expectations. Observable: the mean residual $\log(g_{\rm obs}/g_{\rm RAR})$ over points with $g_{\rm bar} < a_0/3$, per galaxy, against the maximum-clustering environmental field log e_N of Chae et al. (2021), at fixed log M_b and log Σ_h. The relation predicts slope 0; the external-field effect predicts -0.08 dex per dex for the SPARC sample. A slope of -0.08 ± 0.03 or tighter on a larger sample falsifies the relation's

environmental prediction and, with it, the source rule of Section 2.3.

6.3 The ambient truncation

Status: live (Appendix C, Figure 7). The stacked lensing circular velocity of isolated galaxies turns down at $r_t = \sigma/\sqrt{2\pi G\rho_{\rm amb}} \propto M_b^{1/4}\rho_{\rm amb}^{-1/2}$, with, for isolated galaxies, one $\rho_{\rm amb}$ for every mass bin, the most massive bin last, and $v_c \propto r^{-1/2}$ beyond. The mass ordering is not by itself a discriminator: an external-field decline in MOND sets in where the internal field falls to the external one, at a radius $\propto M_b^{1/2}$ at fixed field, more steeply mass-ordered than $M_b^{1/4}$; massive galaxies also sit in denser surroundings, so $\rho_{\rm amb}$ correlates with M_b; and the two-halo term is mass-ordered too. What distinguishes the two is the shape (a cut at an ambient density against a roll-over set by the external field) and the dependence on environment at fixed mass; the separation must be shown with mock stacks and an orientation-averaged external-field profile, neither of which is done here. The value of $\rho_{\rm amb}$ is the one free parameter of the forecast. The data within 1 Mpc bound it from above only (< 630–$710\,M_\odot\,{\rm kpc}^{-3}$ at 95%, depending on the radial range fitted, and 160–355 with the amplitude fixed by the relation; Appendix C); the points beyond 1 Mpc mildly prefer a truncation, but the ambient density they would imply ($\sim$350, nine times the cosmic mean around galaxies selected as isolated) would itself lens and is not credible, so we do not quote it as a fit. The closure identity $a_0^2/Gc^2 = 35.5$ ($r_t \approx 3.4$ Mpc for L^*) is the one physically motivated value, and the forecast below is stated for it and for the bound. A forecast with V_f free, the covariance of the deprojected points, the extrapolation systematics and a two-halo nuisance term is owed and not attempted here; qualitatively, a truncation at the bound (0.8 Mpc for L^*) is already within reach of present data if the two-halo term can be controlled, and the closure value (3.4 Mpc) is beyond isolated-lens selection; the closure value needs 5 Mpc, beyond where isolated-lens selection holds against the two-halo term. Outcomes: a decline with the predicted shape and a common $\rho_{\rm amb}$ across mass bins detects the ambient term and measures it; flatness to 2 Mpc at better precision pushes $\rho_{\rm amb}$ below $\sim$60 and leaves the closure value untested; a decline that follows environment at fixed mass rather than a common ambient density favours the external-field reading. Lensing flatness alone is therefore not a falsifier of the relation; flatness to 5 Mpc at the forecast precision falsifies the closure value of the ambient density for isolated galaxies (Section 6.5), not eq. (1).

6.4 Further predictions

Three predictions are detailed in Appendix E:

- the starless cloud Cloud-9 is bound only if its own nb mass is large enough ($\log\rho_c \gtrsim 6$, giving 1–$7\times10^7\,M_\odot$ inside 1.4 kpc, of the order its line width implies);
- early- and late-type lensing relations identical at fixed baryonic mass under the dynamical assumption of Section 2.2, whereas a conserved completion predicts an early-type excess growing with merger history (about 0.04–0.075 dex in $g_{\rm obs}$ per merger); the data show a $\geq 6\sigma$ difference in Brouwer et al. (2021) and none significant after the stellar-mass, hot-gas and isolation corrections of Mistele et al. (2024a,b), who recover Brouwer's difference with Brouwer's choices;
- the pairwise force law at 30-230 Mpc.

No dynamical friction from the nb mass, under the dynamical assumption of Section 2.2 (not a consequence of the relation): its response is fast compared with the bar's rotation, so a bar rotating through it leaves no lagging wake and loses no angular momentum to it. Fast bars ($\mathcal{R} = R_{\rm CR}/R_{\rm bar} \approx 1$) are then expected, as in MOND; in ΛCDM simulations halo friction slows bars (Debattista & Sellwood 2000), a tension reported at $> 9\sigma$ for EAGLE and IllustrisTNG by Roshan et al. (2021) and reduced in the more baryon-dominated Auriga discs by Fragkoudi et al. (2021). Shared with MOND; a discriminator against massive dynamical haloes only.

6.5 Tests in reach

The predictions above can be tested with data that exist or are scheduled; we state each with its main systematic so that each is decidable.

(i) Lensing beyond 1 Mpc. Stacked galaxy-galaxy lensing to $\sim$5 Mpc, binned by baryonic mass and by local density measured from spectroscopic redshifts (DESI overlapping KiDS or HSC now; Euclid later). The relation predicts a downturn whose radius scales as $M_b^{1/4}\rho_{\rm amb}^{-1/2}$: about 3–6 Mpc across the four mass bins of

Mistele et al. (2024b) at the closure density (Appendix C), with the excess surface density at 5 Mpc falling to 0.49–0.91 of the untruncated profile (for baryonic masses 0.1 dex above nominal, as the lensing amplitude prefers); at an ambient density of 160 the downturn moves to 1.4–2.7 Mpc. The radius ratio between the heaviest and lightest bins is about 2.0, against about 3.9 for a decline set by MOND's external field at fixed field strength, and the relation's radius follows the ambient density rather than the field. Because the ambient density is set by a smooth background field rather than by the surrounding galaxies, the downturn persists in voids, at larger radii where the background lies below its global mean, while MOND's external-field decline recedes. Main systematic: the two-halo term, which must be forward-modelled on mocks with the same selection. Falsified by: flatness to 5 Mpc at the closure precision (which falsifies the closure value for isolated galaxies, not eq. (1)), or a radius ratio near 3.9.

(ii) Wide binaries (Gaia DR4). $\gamma = 1.00$, provided the binary's own central density is below $\sim 10^{10}\,M_\odot\,\mathrm{kpc}^{-3}$ (Section 6.1). Main systematic: quality cuts and hidden companions. Falsified by: a confirmed $\gamma \geq 1.1$ (of eq. (1) with the globular-cluster density bound, declared in Section 6.1).

(iii) Dwarfs, predictions first. For dwarfs without measured dispersions, the relation's and MOND's (with the external field) predictions are to be published before the data, under the protocol of Appendix B (its first pass is reported there). The relation predicts embedded and isolated dwarfs alike; MOND predicts the embedded ones lower. Main systematic: tides and non-equilibrium. Falsified by: embedded dwarfs below isolated ones by more than 0.15 dex on average.

(iv) Bars. With no friction from the nb mass under the dynamical assumption (Section 2.2), bars stay fast (Section 6.4). Main systematic: pattern-speed measurement (Tremaine-Weinberg). Falsified by slow bars as the norm – which would falsify the dynamical assumption, not the relation.

7. Open list

- ρ_c: what sets it (solution selection; relation (3) is empirical).
- A closure for ρ_c remains open. MOND's core surface density (Milgrom 2009) fails as one on the dwarfs (Appendix B.1).
- The extended-gas source: how much nb mass a cloud carries is set by its ρ_c (Section 2.3), which is not understood; tested by Abell 520 (a gas-centred lensing core, if confirmed on undisputed data, contradicts it) and Cloud-9.
- The nesting rule: the LMC should carry a 1.36x excess of Milky Way nb mass inside it. The body decomposition: which collections of baryons count as bodies is chosen in each application, not derived (Section 2.3).
- Galaxy clusters: counting the hot gas as part of the cluster body (Section 2.3), the relation reaches MOND parity inside 2 Mpc for a Coma-like cluster ($7 \times 10^{14}\,M_\odot$ of nb against $\sim 8.5 \times 10^{14}$ needed) and keeps MOND's inner-Mpc deficit; the nb released by member galaxies is about three-quarters of the cluster's own halo out to its truncation radius, so a conserved completion closes the bookkeeping and predicts dense outskirts ($\sim 30\,\rho_m$ at 5 Mpc), testable with stacked cluster lensing.
- The Mpc-scale nb budget (a mean-field estimate with galaxy clustering closes it; the calculation over the distribution of local densities is owed; Appendix C); the lensing amplitude; pairwise kSZ; a $\Delta\Sigma$-space fit with the satellite-body term included.
- The microphysics of the nb mass and of the background it exchanges with.
- The self-term identity $\delta\varphi_{\rm nb}/\sigma^2 \sim 2\ln(r_t/r_c) \sim 12$.
- The response time of the nb mass and of the background it exchanges with. Both are assumed fast compared with dynamical times; this is an assumption, not a consequence of the relation. The background need only keep pace with the rate at which a body's baryonic mass changes, which asks for redistribution speeds of at least $\sim 10^3$ km/s (a few Mpc in a few Gyr), and causality bounds it by c. Within that window a finite response time would show only where the demand changes fast – mergers, gas stripping, first infall – and would introduce a dependence on history; history-blindness (Appendix C) then holds at fixed baryons and

fixed local background. A measured lag or wake of the nb mass in a merger would falsify the assumption. One candidate reading is two-phase: a fast, smooth background exchanging with cold isothermal halos; the carrier is not specified.

- The relation has no action. Momentum is not conserved by the relation as written. If the nb mass is re-formed in place rather than carried, its configuration carries no momentum, and the pull of one body on another's nb mass has a recipient only if the reaction is taken up by the background with which the nb mass exchanges; without one, unequal pairs would drift: negligibly for the Milky Way and M31 (~1 km/s over 10 Gyr), by ~20 km/s over 10 Gyr for the Milky Way and the LMC (0.36 of the mutual force) – the problem of Milgrom's original formulation (Felten 1984; Bekenstein & Milgrom 1984). The obvious action, a perfect-fluid pressure $P = \sigma^2 \rho_{\rm nb}$ coupled to the baryon potential through the saturation, conserves momentum but makes the baryons feel only $\rho_{\rm nb} S'(x_b)$, and fails the Milky Way curve (rms 50 against 20 km/s). More generally, with canonical auxiliary fields and local couplings (the class of Milgrom's 2010 quasi-linear formulation, which absorbs the nonlocal baryon potential without a reaction force) an action exists only if the response depends on a single combination of the baryon and nb potentials; the tail fixes that combination to the total potential, and saturating on the total potential removes the tail. Saturation on the baryons together with an isothermal self-tail is therefore not variational in that class; other classes are untested.
- The LMC: the relation gives it $\sim 4 \times 10^{10}\, M_\odot$ (its tail cut by the Milky Way's density at 50 kpc, with the nesting boost), four to five times less than the 1–2 $\times 10^{11}$ inferred from the Milky Way's reflex motion and from stream fits (Chandra et al. 2024; Erkal et al. 2019; Vasiliev et al. 2021); a conserved completion that keeps the first-infall halo may close the gap, a calculation owed.
- Mass is not conserved by the relation as written: a body's total nb mass, $(4/3)\sigma^2 r_t/G$, scales as $M_b^{3/4}$, so a merger remnant carries less than its progenitors (16% less for equal masses, 9% for 10:1). We take the difference to be returned to the environment, raising the local ambient density; conservation then holds globally but needs transport. If the nb mass is re-formed in place, the same exchange is needed locally and continuously, not only in mergers: local conservation then requires a source term, an exchange with a background component (identified here only with the ambient density), and a relativistic completion must conserve the total stress-energy, so that background carries the balance. A collisional fluid of the Famaey–Khoury–Penco (2018) type conserves mass and momentum, but its velocity scale is local ($v^4 \approx a_0 G M_b(<r)$), not per-body. Adopting it would change the relation, not complete it: σ would vary with radius inside each body and could depend on the baryons' surface density, which the present relation lacks (Section 4.1). Whether that change supplies the missing ingredient or creates new problems is open; we take it as the motivation for a local-σ variant, not as a completion of this one.
- Where nb mass goes when a body loses it, in a merger or to tides, depends on its transport, which the static relation does not specify; whether it spreads with the orbits of the bodies that lost it or flows toward the most concentrated masses decides how cluster cores are filled, and is left open.
- Owed test: Chae et al.'s regional contrast under this relation (needs their region assignments).

Appendix A. Individual curves, the Tully-Fisher relation, features, and globular clusters

A.1 The Tully-Fisher relation and curve features

The BTFR is an input (eq. 2), not a prediction; we say so. What is checked is that the per-galaxy spread of ρ_c does not reappear as BTFR scatter: the residual of the model's speed at the last measured point at fixed M_b is 0.044 dex against 0.059 for the observed speed at the same point (0.029 at universal ρ_c); this is not the intrinsic scatter of V_f as defined by Lelli, McGaugh & Schombert (2016b), which is smaller (6%, ~0.025 dex; Lelli et al. 2019), and the slope is imposed by eq. (2). Features in the baryon potential are carried into the response: detrended curve features in the model correlate with those observed at r = 0.33 over the sample (0.24 for the acceleration relation, 0.27 for the baryons alone); but the response is to the potential, which smooths the sharpest density features, and on IC 2574 the model carries only ~40% of the observed dip at 4 kpc where the acceleration relation carries it fully. We do not claim Renzo's rule (Sancisi 2004).

Earlier 120-galaxy comparison (Section 3.2). On the earlier 120-galaxy comparison, with ρ_c free per galaxy the median χ^2 falls from 28.0 (no saturation) to 13.8 (the acceleration relation with the disc mass-to-light ratio free: 16.4; with distance and inclination also free: 12.5); with σ also free it reaches 6.1 (5.6 for the unsaturated relation), so the BTFR σ does not describe individual curves to the data's precision.

The Milky Way (Section 3.2). Baryons: Ou et al. (2024) Table 2 (Hernquist bulge 1.55×10^{10} $M_\odot$, $a = 0.70$ kpc; exponential stellar disc 3.65×10^{10}, $R_d = 2.35$; HI 8.2×10^9, $R_d = 18.2$; H_2 1.3×10^9, $R_d = 2.57$; total 6.15×10^{10}), disc terms by the thin-disc formula, nb response spherical as elsewhere; $\sigma = 125$ km/s, $v_{\rm flat} = 177$. Data: their Table 1 (37 bins, 6.3–27.3 kpc), errors with a 5 km/s floor declared before the run. Newton alone: rms 87 km/s, slope −4.1. MOND (RAR ν, same baryons): rms 17, slope −1.7, 218 km/s at the Sun. Law at $10^{7.45}$: rms 20, slope −1.9, 215 at the Sun. Law at the fitted $10^{7.88}$: rms 9, slope −3.0, 238 at the Sun. Both MOND and the relation are ~15 km/s low at the Sun under this baryon model; McGaugh (2019) fixes the Milky Way's disc from the terminal-velocity features instead, with a heavier disc, and obtained the declining outer curve as a prediction before the Gaia DR3 measurements. We kept the Ou et al. model to keep the comparison at fixed baryons. Beyond the data the relation's curve passes 5–10% below the BTFR speed at 40–100 kpc (169 km/s at 100 kpc at the universal density) before converging – the isothermal sphere's approach to $\sqrt{2}\sigma$ – where MOND converges from above (182 at 100 kpc). The steeper, near-Keplerian decline reported beyond 20 kpc by Jiao et al. (2023) falls where Ou et al. put systematics above 15% from the neglected asymmetric drift; whether the outer curve is a transition decline or a Keplerian one is what a cleaner outer-tracer curve would settle, and the two models predict the same answer. Not a discriminator between them.

A.2 Dwarf spheroidals and globular clusters

With $\rho_c = 10^{7.45}$ the relation gives And XIX a dispersion of 8.3 km/s against the observed 7.8 (+1.7/-1.5) (Collins et al. 2020), where our own estimates of isolated MOND (5.95, from $\sigma^4 = 4$ G a_0 M/81) and of the quasi-Newtonian external-field value (2.1) bracket it. Truncation by the host's nb mass changes satellite dispersions by 3-15% only. Globular-cluster cores are clean: the saturation holds the nb fraction inside the half-light radius to ~1e-4 for 47 Tuc and omega Cen and ~2e-3 for NGC 2419; Palomar 5 excludes a dSph-level ρ_c (10^8) for globular clusters at $\sim 3\sigma$ and the SPARC median at $\sim 1.4\sigma$, a soft bound $\rho_c({\rm GC}) \lesssim 10^{7.2}$ – so ρ_c already depends on the type of body.

Appendix B. What sets ρ_c, and the dwarf-dispersion protocol

B.1 What sets ρ_c

Across SPARC the fitted ρ_c spans 0.65 dex after mass and surface density are removed. Its marginal trend with mass is flat (slope -0.02 over 3.4 dex in M_b; the -0.365 in relation (3) is the partial slope at fixed surface density and gas fraction), with the sign of any trend set by the stellar mass-to-light ratio (max-disc gives the Kormendy & Freeman direction); it rises with baryonic surface density at fixed mass (+0.37 dex per dex, 95% [+0.14, +0.62], with mass as the only other variable; +0.27 once gas fraction is also included) and falls with gas fraction (-0.93 dex per unit, [-1.71, -0.14]). The published relation is

$$\log \rho_c = 11.06 - 0.365 \log M_b + 0.268 \log \Sigma_h - 0.93\, f_{\rm gas} \tag{3}$$

used in Appendix B with no per-object freedom. ρ_c is the one unexplained number of the model, and it knows about the baryons in the way any such density must; we take that as the present-tense form of the fine-tuning problem, restated, not solved. Two closures were tried and failed: ρ_c from a_0 alone ($\rho_c = a_0^2/9\pi G\sigma^2$, which fixes the core surface density at $a_0/2\pi G$ exactly) gets the median right and the mass dependence wrong; the gas-fraction dependence runs the wrong way for the objects that need a low density (Section 4). We read the correlations of ρ_c with M_b, Σ_h and $f_{\rm gas}$ as the data on a rule the relation does not yet contain; the same mass and surface-density dependences appear in the core surface density (Section 3.3). A closure through MOND's core surface density (Milgrom 2009), with $\rho_c r_c$ set equal to it galaxy by galaxy, was pre-registered and tested. It fails on the dwarfs of Appendix B.2 (6 of 16 rows within the bar, rms 0.18 dex), while the SPARC curves tolerate it (per-point rms 0.149, against 0.140 for relation (3)). On the dwarfs it reproduces MOND's isolated dispersions to 0.006 dex: in the low-surface-density limit r_h cancels and $\sigma^4_{\rm los} \propto G a_0 M_b$.

B.2 Dwarf dispersions, pre-registered

Protocol: ρ_c from relation (3) with $f_{\rm gas} = 0$ (an extrapolation beyond the gas-bearing discs on which it was calibrated; dwarf spheroidals lie on the same acceleration relation as discs, Lelli et al. 2017); σ from eq. (2) with $M_b = 1.6$ L_V; Plummer stars; the host's nb mass as ambient; projected dispersion within the half-light radius; no per-object freedom. First pass on 15 measured objects, 16 rows (eight classical Milky Way satellites with Walker et al. 2009 inputs, plus Crater II; And II, XIX, XXI; Cetus with two published dispersions, Tucana and And XVIII as isolated systems; And XVIII's dispersion is disputed): 9 of 16 rows within the per-object bar: the pre-registered two-thirds criterion is FAILED on this pass, and the misses are structural rather than random (the two with the highest dispersion for their luminosity under-predicted, the massive extended and the diffuse pairs over-predicted); rms 0.15 dex, mean +0.03; embedded (+0.03) and isolated (+0.01) residuals agree, with three isolated objects. The model over-predicts the massive extended pair (Fornax, And II) and the diffuse pair (Crater II, And XXI) and under-predicts the two with the highest dispersion for their luminosity (Draco, Ursa Minor): ρ_c depends on something beyond (M_b, Σ_h, $f_{\rm gas}$). Relation (3) is calibrated at $\log M_b \gtrsim 8$, on curves that at low mass end near 1.5–2 r_c; the dwarfs lie 0.7–2.6 dex below that range, so this tests an extrapolation of an empirical fit, not eq. (2). A uniform undercount of baryons does not help: the predicted dispersion scales as $M_b^{0.22}$, and the mean residual is already positive. CDM is not scored here: it gives no dispersion from the baryons alone and needs a halo mass for each object. MOND's own a-priori predictions for the Andromeda set are in McGaugh & Milgrom (2013a,b). Antlia II (Ji et al. 2021: $\sigma = 5.98$ km/s, r_h 2.5 kpc, pericentre 38.6 kpc) is the control for Crater II's tidal reading: 7.6–7.9 km/s at equilibrium, 5.8–6.1 with the pericentric truncation. The same protocol will be applied to dwarfs without measured dispersions, to be named after a literature check, before their dispersions are measured; if the embedded objects come out below the protocol's predictions by more than 0.15 dex on average while the isolated ones do not, the model is dead regardless of SPARC.




Figure 6: Figure 6. The dwarf protocol, first pass: predicted dispersion from relation (3) with no per-object freedom against the observed value, for Milky Way satellites, M31 satellites and isolated dwarfs; the band is ± 0.1 dex. Nine of sixteen rows fall inside the per-object bar, short of the pre-registered two-thirds; the residuals show no dependence on the external field (three isolated objects).

Appendix C. Lensing: compatibility, truncation, satellites and the mass budget

The three purchases stated in the introduction rest on the following. Lensing: the nb mass lenses as ordinary non-relativistic mass (Section 1: its pressure and anisotropic stress are negligible), so the weak-lensing profile of an isolated galaxy is the dynamical one. Down to the truncation radius the tail is a singular isothermal sphere with $v^4 = Ga_0 M_b$, so lensing, dynamics and the MOND prediction coincide there; the truncation sits at $r_t = \sigma/\sqrt{2\pi G\rho_{\rm amb}}$: for the four stellar-mass bins of Mistele et al. (2024b) ($\log M_*/M_\odot$ = 9.0–10.5, 10.5–10.8, 10.8–11.1, 11.1–11.5; V_f = 141, 191, 214, 259 km/s) this is 3.2–5.9 Mpc at the closure value 35.5 and 0.8–1.4 Mpc at the 95% bound 630 (for an L^* galaxy, $M_b = 3\times10^{10}$, 3.4 and 0.8 Mpc). r_t is the matching radius of the isothermal tail; in the full solve with the subtraction the enclosed excess peaks about 20% beyond it and the 10% decline in v_c sets in about 30% inside it; r_t is used for scaling and the solve for the forecast (Figure 7). On the baryonic-acceleration axis of the lensing relation this is $g_{\rm bar}(r_t) = GM_b/r_t^2 \approx 4\times10^{-16}\,{\rm m\,s^{-2}}$ for an L^* galaxy at the closure value (6.5×10^{-15} at the bound), at or just beyond the last bins of the stacks (Brouwer et al. 2021; Mistele et al. 2024b); the observed acceleration there is $g_{\rm obs}(r_t) = 2\sigma\sqrt{2\pi G\rho_{\rm amb}} = 2\sqrt{2\pi}\,(\sigma/c)\,a_0 \approx 2\times10^{-13}\,{\rm m\,s^{-2}}$ at the closure value. MOND expects a decline at large radius too, from the external-field effect; Mistele et al. (2024b, Discussion) write that it "might become possible to perceive" with better data, that it "may be undetectable in the present data", and that beyond 1 Mpc "some mass bins possibly decline, but there is no clear departure from flatness". The two expectations differ in kind – an external-field decline is ordered by environment, the ambient truncation by $M_b^{1/4}$ at fixed $\rho_{\rm amb}$ (Section 6.3) – but a quantitative comparison needs the orientation-averaged stacked profile in a non-uniform field, which we have not computed. A fit of the truncated profile to their deprojected circular velocities (four mass bins, 41 points beyond 100 kpc, statistical and systematic errors in quadrature, V_f free per bin) gives $\rho_{\rm amb} < 630$ at 95% ($\chi^2 = 21.5$ for 36 degrees of freedom). With the amplitude fixed by the relation rather than fitted ($V_f^4 = Ga_0M_b$, also MOND's asymptote), the lensing flat speeds lie 5–13% above the prediction at the stacked baryonic masses ($\chi^2 = 74$ for 41 points); a 0.1 dex increase in M_b, within the stellar-mass and gas systematics, gives $\chi^2 = 33$, and the ambient density is then bounded by 160–355 (95%). The amplitude offset is shared with MOND and is not a discriminator. Among the fits with V_f free, a fit in the excess surface density gives < 630 without a satellite term and < 1000 with one. Restricted to 100 kpc–1 Mpc the data bound $\rho_{\rm amb}$ from above only (< 710 at 95%): they do not detect a truncation. The mild preference for one over an untruncated tail ($\Delta\chi^2 \approx 6$) comes from the points beyond 1 Mpc, where isolation, the two-halo term and the deprojection's assumed continuation are least secure (the systematic column grows to 40–150 km/s there), so it is not a detection. The errors are treated as uncorrelated, and the systematic column (the uncertainty of their extrapolation beyond the last bin) is coherent across radius, so these numbers are indicative and the χ^2 is too low; a forward model of $\Delta\Sigma$ with the KiDS covariance and a two-halo term is owed. Under the source rule of Section 2.3 the satellites of an isolated galaxy are bodies with their own nb mass, each truncated by the host's tail; for a plausible ten satellites above $10^8\ M_\odot$ within 1 Mpc they add ~20–45% to the host's enclosed nb mass at 1 Mpc (more for low-mass hosts) and little inside 0.5 Mpc, so the stacked profile beyond ~0.5 Mpc is not the host's halo alone; the present data neither require nor exclude that term ($\Delta\chi^2 \approx 0.3$). Beyond r_t the relation predicts a truncation of the excess mass whose radius scales as $M_b^{1/4}$ and $\rho_{\rm amb}^{-1/2}$, turning $\Delta\Sigma$ from $1/R$ toward $1/R^2$ between 1 and 5 Mpc (at the closure value, 0.95 of the untruncated value at 3 Mpc and 0.71 at 5), mass-ordered like the correlated-structure term, so separating the two needs mock stacks; that is a forecast for mass-binned stacks, not a present test. Early- and late-type lensing: Section 6.4. Mass budget. The closure value is taken as the global mean of all nb mass (Section 2.1), so the halos are drawn from the background and the budget closes by construction; the question is whether a non-negative background can supply them. Summed over the galaxy stellar mass function, isolated halos would carry a mean excess of $\sim$110 $M_\odot\,{\rm kpc}^{-3}$, 3.1 times the closure value: they cannot all exist. They need not: r_t (3–6 Mpc) exceeds the typical separation of galaxies (a few Mpc), so a tail is cut by its neighbours' before it reaches the background, and isolated-halo sums overcount the same mass. A mean-field estimate with the observed galaxy clustering (Zehavi et al. 2011), in which each tail ends where it meets the mean nb density around its galaxy, brings the halo excess to 0.4 of the closure value and leaves a background of 0.5–0.7 of it between galaxies, while isolated galaxies keep truncation radii of 2.7–5.2 Mpc. The estimate is biased low: it places every galaxy at the mean density around it, and for dwarfs that mean is set by the few that sit inside a host; if galaxies below $M_* \sim 10^{9.5}\,M_\odot$ are instead left untruncated, the background reaches zero. The budget is therefore open, with a mean-field indication that it can close; a calculation over the distribution of local densities, satellites included, is owed (Section 7). No suppressing external-field effect: the internal dynamics of a bound system depend on their surroundings only through the small enhancing ambient

and nesting terms (Section 2.3), because the response is to the body's own potential; MOND's external-field effect acts at first order, and the difference is the discriminator of Section 4.4 and 6.2. History-blindness: identical present baryons in the same local background give identical nb mass. This predicts no dependence of present structure on its past and no dark sub-haloes without galaxies; it is untested here and is listed as a prediction. The baryons come first: a cloud that becomes self-gravitating acquires its nb mass on its own dynamical time, not after a halo has formed.




Figure 7: Figure 7. The ambient truncation against the stacked weak-lensing circular velocities of Mistele et al. (2024b), Table 1, four stellar-mass bins, on linear axes. Points: their deprojected v_c with statistical and systematic errors in quadrature, normalised by the bin's flat speed V_f fitted at $\rho_{amb} = 35.5$ (which favours the closure curve visually). Curves: the isothermal tail with the ambient term subtracted and cut at $r_t = \sigma/\sqrt{2\pi G\rho_{amb}}$, for $\rho_{amb} = 35.5$ (closure), 355 (the value the points beyond 1 Mpc would prefer, shown for reference and not adopted) and 630 (95% bound) $M_\odot\,\mathrm{kpc}^{-3}$. The decline begins before r_t because the subtracted ambient term already lowers the enclosed excess; beyond r_t the excess mass is constant and v_c falls as $r^{-1/2}$. The most massive bin truncates last ($r_t \propto M_b^{1/4}$) and is the flattest.

Appendix D. The environment test in full

Chae et al. (2020, 2021) fit each SPARC curve with an external-field parameter e-tilde, marginalising mass-to-light ratios, distance and inclination, on a sample selected to be unbiased with respect to outer decline, and detect e-tilde > 0 on the sample median (4.4σ; 97 of 143 positive) with a contrast between underdense and overdense regions of the cosmic web; they also report that the galaxy-by-galaxy correlation of e-tilde with the environmental field is not recoverable at current errors (their sec 4.2). Our test is weaker than theirs: the low-acceleration residual from the acceleration relation, at fixed mass-to-light ratio, distance and inclination, over points with g_{bar}

$< a_0/3$ (a selection their Appendix B notes favours less-declining curves), regressed on their environmental field. It gives -0.12 ± 0.16 dex per dex at fixed mass and surface density. The external-field effect, evaluated galaxy by galaxy with Chae et al.'s own formula at their field values, predicts -0.08 for this sample; the relation predicts 0. The sample does not separate the two; the point estimate lies at the external-field value. The outer-slope statistic is null (-0.03 ± 0.21). Chae et al. (2020) report a detection with the external-field sign from per-galaxy fits; Paranjape & Sheth (2022, arXiv:2112.00026; abstract) argue its significance is compromised by rotation-curve uncertainties and the simulation-based field estimates, and that a qualitatively similar signal is expected in LCDM from clustering-property correlations. We then applied their statistic itself: their MOND-plus-external-field model fitted to each curve with mass-to-light ratios, distance and inclination marginalised under the SPARC priors, on their $x_{0,3}$ selection (102 galaxies). On the observed curves this reproduces their per-galaxy values (rank correlation 0.88) and a positive median, +0.016 (their tabulated fits: +0.025, 61% positive). Fitted to the relation's own curves, which contain no external field, the same model returns a positive median too: +0.012 (57% positive) at the single density $10^{7.45}$, and +0.022 (60%) at the fitted densities. The cored-isothermal outer shape is read by their model as an external field of about half the size they detect; the median-e-tilde detection therefore does not by itself discriminate suppression from this relation. What would is the dependence of e-tilde on the environmental field; on this subsample it is weak and not of the external-field sign (rank correlation -0.26 for 60 galaxies with field values; -0.10 for their tabulated values), consistent with their own statement that the galaxy-by-galaxy correlation is not recoverable at current errors. Splitting the 60 galaxies with field values into terciles, e-tilde is higher in the LOW-field tercile for the observed curves (+0.05 against 0.00, p = 0.03) and for their own fits (+0.09 against +0.03), and the relation's curves show the same contrast (p = 0.02); the observed-minus-model difference has no field dependence (p = 0.76). The regional contrast they report (underdense against overdense regions of the cosmic web) uses region assignments not available to us and is the remaining discriminator, untested here. This is the note's open weak point.

Appendix E. Cloud-9, lensing by morphology, pairwise forces

E.1 Cloud-9

The starless HI cloud near M94 (M_HI ~ 1e6 $M_\odot$, r ~ 1.4 kpc, W50 = 12 km/s, non-rotating, D ~ 4.4 Mpc, no stars above $10^{3.5}$ $M_\odot$; Benitez-Llambay et al. 2024, arXiv:2406.18643; Anand et al. 2025, arXiv:2508.20157;) has a random speed (~5 km/s) above its self-gravity speed (~1.8 km/s) and a Jeans length (~8 kpc) larger than itself, so its gas alone does not bind it. Under the source rule of Section 2.3 it nonetheless carries nb mass, set by its ρ_c: for $\log \rho_c \lesssim 5.5$ too little to bind it (it is then confined by its surroundings or dispersing), for $\log \rho_c \gtrsim 6$ enough ($1–7 \times 10^7$ $M_\odot$ inside 1.4 kpc, of the order of the dynamical mass $\sim 2.5 \times 10^7$ $M_\odot$ its line width implies). LCDM (RELHIC) places it in a ~5e9 $M_\odot$ halo; MOND gives a modest external-field boost. One object, three different answers.

E.2 Lensing by morphology

Early- and late-type lensing relations are identical at fixed baryonic mass under the dynamical assumption of Section 2.2, whereas a conserved completion predicts an early-type excess growing with merger history (about 0.04–0.075 dex in $g_{\rm obs}$ per merger); the data show a $\geq 6\sigma$ difference in Brouwer et al. (2021) and none significant after the stellar-mass, hot-gas and isolation corrections of Mistele et al. (2024a,b), who recover Brouwer's difference with Brouwer's choices (Appendix C and the merger estimate of Section 7). Pairwise forces at 30-230 Mpc.

E.3 Pairwise forces

Gallardo et al. (2026) find an inverse-square pairwise force law from the kinetic Sunyaev-Zel'dovich effect: the acceleration between halo pairs scales as r^{-n} with $n = 2.1 \pm 0.3$ at 30–230 Mpc (ACT with SDSS). Any real-mass theory passes this in shape by construction; the amplitude requires the Mpc-scale nb budget and is open. The external field invoked to recover inverse-square behaviour in MOND (a few per cent of a_0) is at the upper end of, or above, the values Chae et al. (2021) assign to SPARC galaxies (0.002–0.018 a_0) and the 1–2% estimate for the field of large-scale structure (McGaugh 2026).

Appendix F. Relation to v1.0.14

The v1.0.14 preprint derived its augmented Poisson equation with the wrong operator sign ($-k^2$; the bump substitution gives $+k^2$; erratum posted 2026-10), withdrew its Yukawa exterior and nu-function with it, and argued the uniqueness of a linear law (Appendix F) that the present nonlinear relation supersedes. Its account of the Bullet cluster (p.~25, Fig. 6), that lensing responds to the gradient of surface density, is withdrawn: convergence is surface density, and the Clowe et al. (2006) apertures show baryons at the plasma peaks equal to or above those at the galaxies. A post-v1.0.14 working wide-binary number, $\gamma = 1.13$-1.26 (2026-10-15), came from a floor-clipped phantom-density construction under an intermediate law and is superseded by the Newtonian prediction of Section 6.1. Its three-regime picture (Newtonian, MOND-like, and an exponential falloff beyond) and its claim that the Gaia DR3 decline of the Milky Way curve is a regime MOND cannot predict are withdrawn: the Yukawa cutoff at $k^{-1} \approx 14$ kpc that defined the outer regime went with the operator sign, and the MOND comparison (rising to ~204 km/s at 27 kpc) normalised MOND at 10 kpc rather than computing it from the baryons. Computed from the same baryons, MOND and the present relation both decline across that range (Section 3.2, Appendix A); the decline is the transition, and the relation's departure from MOND phenomenology lies at the ambient truncation, at 0.8–3.4 Mpc (Section 6.3). Same concept DOI; this is v2.0.0.

Appendix G. A numerical correspondence

We record a numerical correspondence without drawing a cosmological conclusion from it. The global mean of the ambient density is taken at $a_0^2/Gc^2 = 35.5\, M_\odot\,\mathrm{kpc}^{-3}$ (Section 2.1). The Planck physical densities, which do not depend on the Hubble constant, are $\Omega_c h^2 = 0.120$ for non-baryonic matter and $\Omega_b h^2 = 0.0224$ for baryons, i.e. 33.3 and 6.2 $M_\odot\,\mathrm{kpc}^{-3}$ today (Planck Collaboration 2020). The closure value is 1.06 times the first; added to the baryons it is 1.05 times the total matter density. Three cautions keep this a correspondence and not a result:

(i) a_0 is known to about 20% (systematic; McGaugh, Lelli & Schombert 2016), so the closure density is uncertain by about 40%; agreement at the 5% level is better than the inputs justify.

(ii) The closure carries an undetermined factor of order unity (a uniform sphere cut off at the causal length gives 2π rather than 1), so the match depends on a choice we have not derived.

(iii) Whether the nb mass of this note is the non-baryonic matter of cosmology is not addressed here (Section 1).

We note it because the relation's one cosmological input, set by an acceleration measured in galaxies, lands on the measured non-baryonic density, and because the same a_0 fixes where the baryon term saturates (Section 2.1).

AI usage

This work was carried out with extensive assistance from Anthropic's Claude models: literature search and verification of references, code for the solvers and fits, numerical runs, drafting and editing of text, and simulated referee reports used as stress tests. The author set the questions and the hypotheses, pre-registered the tests, decided every result's status and every claim's wording, and takes full responsibility for the content.

References

Anand, G. S., Benítez-Llambay, A., Beaton, R., et al. 2025, ApJL 993, L55 (arXiv:2508.20157)
Angus, G. W. 2009, MNRAS 394, 527 (arXiv:0805.4014)
Banik, I., et al. 2024, MNRAS 527, 4573 (arXiv:2311.03436)
Bekenstein, J., & Milgrom, M. 1984, ApJ 286, 7
Benítez-Llambay, A., Dutta, R., Fumagalli, M., et al. 2024, arXiv:2406.18643 (submitted to ApJ)
Berezhiani, L., & Khoury, J. 2015, PRD 92, 103510 (arXiv:1507.01019)
Blanchet, L. 2007, CQG 24, 3529 (arXiv:astro-ph/0605637)
Blanchet, L., & Le Tiec, A. 2008, PRD 78, 024031 (arXiv:0804.3518)
Borukhovetskaya, A., et al. 2022, MNRAS 512, 5247 (arXiv:2112.01540)
Brans, C., & Dicke, R. H. 1961, Phys. Rev. 124, 925
Brouwer, M. M., et al. 2021, A&A 650, A113 (arXiv:2106.11677)

Bruneton, J.-P., Liberati, S., Sindoni, L., & Famaey, B. 2009, JCAP 03, 021 (arXiv:0811.3143)
Burkert, A. 1995, ApJL 447, L25 (arXiv:astro-ph/9504041)
Caldwell, N., et al. 2017, ApJ 839, 20 (arXiv:1612.06398)
Chae, K.-H. 2023, ApJ, doi:10.3847/1538-4357/ace101 (arXiv:2305.04613)
Chae, K.-H. 2024a, arXiv:2309.10404
Chae, K.-H. 2024b, arXiv:2402.05720
Chae, K.-H. 2025, arXiv:2502.09373
Chae, K.-H., et al. 2020, ApJ 904, 51
Chae, K.-H., et al. 2021, ApJ 921, 104 (arXiv:2109.04745)
Chae, K.-H., Lee, B.-C., Hernandez, X., et al. 2026, arXiv:2601.21728
Chae, K.-H., & Yoon, Y. 2026, arXiv:2607.14450
Chandra, V., Naidu, R. P., Conroy, C., et al. 2024, ApJ, doi:10.3847/1538-4357/addab6 (arXiv:2406.01676)
Clowe, D., et al. 2006, ApJL 648, L109 (arXiv:astro-ph/0608407)
Collins, M. L. M., et al. 2020, MNRAS 491, 3496 (arXiv:1910.12879)
Cookson, S. A., Banik, I., El-Badry, K., et al. 2026, MNRAS, doi:10.1093/mnras/stag342 (arXiv:2602.24035)
Danieli, S., et al. 2019, ApJL 874, L12 (arXiv:1901.03711)
Debattista, V. P., & Sellwood, J. A. 2000, ApJ 543, 704 (arXiv:astro-ph/0006275)
Desmond, H. 2023, MNRAS 526, 3342 (arXiv:2303.11314)
Donato, F., et al. 2009, MNRAS 397, 1169 (arXiv:0904.4054)
Erkal, D., et al. 2019, MNRAS 487, 2685 (arXiv:1812.08192)
Famaey, B., Khoury, J., & Penco, R. 2018, JCAP 03, 038 (arXiv:1712.01316)
Famaey, B., & McGaugh, S. S. 2012, Living Rev. Relativ. 15, 10 (arXiv:1112.3960)
Famaey, B., McGaugh, S., & Milgrom, M. 2018, MNRAS 480, 473 (arXiv:1804.04167)
Felten, J. E. 1984, ApJ 286, 3
Fragkoudi, F., et al. 2021, A&A 650, L16 (arXiv:2011.13942)
Gallardo, P. A., et al. 2026, PRL 136, 151002 (doi:10.1103/rk8v-rcm3)
Haghi, H., et al. 2019, MNRAS 487, 2441 (arXiv:1906.03268)
Harvey, D., et al. 2015, Science 347, 1462 (arXiv:1503.07675)
Hernandez, X., Chae, K.-H., & Aguayo-Ortiz, A. 2024, arXiv:2312.03162
Hossenfelder, S. 2017, PRD 95, 124018 (arXiv:1703.01415)
Ianjamasimanana, R., et al. 2015, AJ 150, 47 (doi:10.1088/0004-6256/150/2/47)
Iorio, G., et al. 2017, MNRAS (arXiv:1611.03865)
Ji, A. P., et al. 2021, ApJ 921, 32 (arXiv:2106.12656)
Jiao, Y., et al. 2023, A&A 678, A208 (arXiv:2309.00048)
Kaplinghat, M., Ren, T., & Yu, H.-B. 2020, JCAP 06, 027 (arXiv:1911.00544)
Kaplinghat, M., Tulin, S., & Yu, H.-B. 2016, PRL 116, 041302 (arXiv:1508.03339)
Khoury, J. 2015, PRD 91, 024022 (arXiv:1409.0012)
King, I. R. 1966, AJ 71, 64
Kormendy, J., & Freeman, K. C. 2016, ApJ 817, 84 (arXiv:1411.2170)
Lelli, F., McGaugh, S. S., & Schombert, J. M. 2016a, AJ 152, 157 (arXiv:1606.09251)
Lelli, F., McGaugh, S. S., & Schombert, J. M. 2016b, ApJL 816, L14 (arXiv:1512.04543)
Lelli, F., et al. 2017, ApJ 836, 152 (arXiv:1610.08981)
Lelli, F., McGaugh, S. S., Schombert, J. M., Desmond, H., & Katz, H. 2019, MNRAS 484, 3267 (arXiv:1901.05966)
Li, P., et al. 2018, A&A 615, A3 (arXiv:1803.00022)
Li, P., Lelli, F., McGaugh, S., & Schombert, J. 2020, ApJS, doi:10.3847/1538-4365/ab700e (arXiv:2001.10538)
LZ Collaboration 2025, PRL 135, 011802 (arXiv:2410.17036)
Marsen, J. R. 2026, GPRF Paper I, v1.0.14, Zenodo, doi:10.5281/zenodo.20943223
McGaugh, S. S. 2016, ApJL 832, L8 (arXiv:1610.06189)
McGaugh, S. S. 2019, ApJ 885, 87 (arXiv:1909.11158)
McGaugh, S. 2023, 'Required dark matter properties', Triton Station (blog), 17 Aug 2023
McGaugh, S. 2026, 'The External Field Effect and Tests of MOND on Cosmological Scales', Triton Station (blog), 27 Aug 2026, doi:10.59350/57294-avw93
McGaugh, S. S., Lelli, F., & Schombert, J. M. 2016, PRL 117, 201101 (arXiv:1609.05917)
McGaugh, S. S. 2026, Why the search for particle dark matter is hopeless, part 1, Triton Station (blog), 26 October 2026, https://tritonstation.com/2026/10/26/why-the-search-for-particle-dark-matter-is-hopeless-part-1/
McGaugh, S. S., & Milgrom, M. 2013a, ApJ 766, 22 (arXiv:1301.0822)

McGaugh, S. S., & Milgrom, M. 2013b, ApJ 775, 139 (arXiv:1308.5894)
Milgrom, M. 1983, ApJ 270, 365
Milgrom, M. 1986, ApJ 306, 9
Milgrom, M. 2009, MNRAS 398, 1023 (arXiv:0909.5184)
Milgrom, M. 2010, MNRAS 403, 886 (arXiv:0911.5464)
Mistele, T., McGaugh, S., & Hossenfelder, S. 2023a, JCAP 09, 004 (arXiv:2303.08560)
Mistele, T., McGaugh, S., & Hossenfelder, S. 2023b, A&A 676, A100 (arXiv:2301.03499)
Mistele, T., et al. 2024a, JCAP 04, 020 (arXiv:2310.15248)
Mistele, T., et al. 2024b, ApJL 969, L3 (arXiv:2406.09685)
Ou, X., et al. 2024, MNRAS 528, 693 (arXiv:2303.12838)
PandaX-4T Collaboration 2025, PRL 134, 011805 (arXiv:2408.00664)
Paranjape, A., & Sheth, R. K. 2022, MNRAS 517, 130 (arXiv:2112.00026)
Pflamm-Altenburg, J. 2025, A&A 703, A68
Planck Collaboration 2020, A&A 641, A6 (arXiv:1807.06209)
Pittordis, C., et al. 2025, Open J. Astrophys., doi:10.33232/001c.142887 (arXiv:2504.07569)
Ren, T., Kwa, A., Kaplinghat, M., & Yu, H.-B. 2019, PRX 9, 031020 (arXiv:1808.05695)
Roshan, M., et al. 2021, MNRAS 508, 926 (arXiv:2106.10304)
Salucci, P., & Burkert, A. 2000, ApJL 537, L9 (arXiv:astro-ph/0004397)
Sancisi, R. 2004, IAU Symp. 220, 233 (arXiv:astro-ph/0311348)
Sanders, J. L., Evans, N. W., & Dehnen, W. 2018, MNRAS, doi:10.1093/mnras/sty1278 (arXiv:1802.09537)
Sciama, D. W. 1953, MNRAS 113, 34
Schombert, J., McGaugh, S., & Lelli, F. 2019, MNRAS 483, 1496 (arXiv:1811.10579)
Shen, Z., et al. 2021, ApJL 914, L12 (arXiv:2104.03319)
Shen, Z., et al. 2023, ApJ 957, 6 (arXiv:2309.08592)
Skordis, C., & Zlosnik, T. 2021, PRL 127, 161302 (arXiv:2007.00082)
van Dokkum, P., et al. 2018, Nature 555, 629 (arXiv:1803.10237)
van Dokkum, P., Danieli, S., Abraham, R., et al. 2019, ApJL, doi:10.3847/2041-8213/ab0d92 (arXiv:1901.05973)
Vasiliev, E., Belokurov, V., & Erkal, D. 2021, MNRAS, doi:10.1093/mnras/staa3673 (arXiv:2009.10726)
Vivas, A. K., Walker, A., Martínez-Vázquez, C., et al. 2025, arXiv:2512.05535
Walker, M. G., et al. 2009, ApJ 704, 1274; erratum 2010, ApJ 710, 886 (arXiv:0906.0341)
XENONnT Collaboration 2025, PRL 135, 221003 (doi:10.1103/msw4-t342)
Zehavi, I., et al. 2011, ApJ 736, 59 (arXiv:1005.2413)