

MAT327 - Introduction to Topology

JIMMY XU

Summer 2024

Contents

1	Day 1 : Introduction, Topological Spaces and Bases	2
1.1	Marking Scheme	2
1.2	Open sets in the Real Numbers	2
1.3	Topological Spaces	3
1.4	Bases	4
2	Day 2 : Bases and Sub-bases	7
2.1	Sub-Bases	8
3	Day 3 : Axiom of Choice, Zorn's Lemma and Ordering	9
3.1	Axiom of Choice	9
3.2	Orderings of Sets	9
3.3	Zorn's Lemma	10
3.4	Order Topologies	10
4	Day 4 : Finite Products, Product Topology and Projections	13
4.1	Finite Product and Product Topology	13
4.2	Projections	14

Ω1 Day 1 : Introduction, Topological Spaces and Bases

This class is taught by Daniel Wilches Calderon during the Summer 2024 semester. We will mostly be following Topology by Munkres, with some suggested readings on the Quercus page

Ω1.1 Marking Scheme

1. Final Exam - 30%
2. Term Test - 20%
3. Quizzes - 50% (Best 8 of 10)

Every week we will be given a problem set, and during tutorial one of those questions will be tested

Ω1.2 Open sets in the Real Numbers

Recall that $U \subseteq \mathbb{R}$ is open if

$$\text{for all } x \in U, \text{ there exists } \varepsilon > 0 \text{ such that } x \in (x - \varepsilon, x + \varepsilon) \subseteq U$$

Open sets have some properties. That the arbitrary union and finite intersection of open sets is still open

1. Arbitrary Union

Let $\{U_\alpha | \alpha \in \Lambda\}$ ¹ be an arbitrary collection of open sets, if $x \in \bigcup_{\alpha \in \Lambda} U_\alpha$, then $x \in U_\alpha$ for some $\alpha \in \Lambda$, and since U_α is open, we have that there exists some $\varepsilon > 0$ such that

$$x \in (x - \varepsilon, x + \varepsilon) \subseteq U_\alpha \subseteq \bigcup_{\alpha \in \Lambda} U_\alpha$$

This same result doesn't hold true for arbitrary intersection, but it does for finite.

2. Finite intersection

Let $\{U_k | k \leq n\}$ be a finite collection of open sets. if $x \in \bigcap_{k \leq n} U_k$, then $x \in U_k$ for all $k \leq n$. Since each U_k is open, there are $\varepsilon_i > 0$ such that $x \in (x - \varepsilon_i, x + \varepsilon_i) \subseteq U_i$ for $i = 1, 2, \dots, n$. Then pick $\varepsilon = \min\{\varepsilon_1, \dots, \varepsilon_n\}$, then it follows that

$$x \in (x - \varepsilon, x + \varepsilon) \subseteq (x - \varepsilon_k, x + \varepsilon_k) \subseteq U_k$$

for all $k \leq n$, so $(x - \varepsilon, x + \varepsilon) \subseteq \bigcap_{k \leq n} U_k$.

¹We use Λ to denote an arbitrary indexing set

Ω1.3 Topological Spaces

Definition 1.1 (Topological Space). A topology, τ , on X is a collection of subsets of X that satisfy

1. $X, \emptyset \in \tau$
2. τ is closed under arbitrary Union
3. τ is closed under finite intersection

The double (X, τ) is called a **Topological Space**. Furthermore, the elements of τ are called open sets, and the elements of X are called points.

We first notice that showing the intersection of 2 sets of τ is still in τ is equivalent to showing the finite intersection of sets of τ will still be in τ . This is true by an inductive argument.

The following are some examples of topologies on a set with 2 elements, the prof briefly went over the topologies on a set with 3 elements, but basically left it as an exercise.

Example 1.2 (Topologies on 2 Point Sets)

Let $X = \{x, y\}$, we have the following are possible topologies on X

1. $\{X, \emptyset\}$

This is often called the indiscrete/trivial topology

2. $\{X, \emptyset, \{x\}, \{y\}\}$

This is often called the discrete topology

3. **Sierpinski Spaces**

The main characteristic of these spaces are that one point is open, while the other is not. More explicitly, the following are the Sierpinski Spaces

- a) $\{X, \{x\}, \emptyset\}$
- b) $\{X, \{y\}, \emptyset\}$

You might've noticed that we labelled some of those examples as the **Discrete Topology** and the **Indiscrete/Trivial Topology**.

Definition 1.3 (Discrete and Indiscrete Topology). Given a set X , the discrete topology on X is equal to the Power Set of X , and the indiscrete topology is equal to $\{x, \emptyset\}$

Definition 1.4 (Cofinite Topology). Let X be a set, the **Cofinite Topology** on X is the collection

$$\tau = \{U \subseteq X \mid X \setminus U \text{ is finite}\} \cup \{\emptyset\}$$

Now to prove that this is a topology, we must prove 3 things. The entire set and the empty set exist within τ . τ is closed under arbitrary union and finite intersection. The first of these is almost immediate as $\emptyset \in \tau$ by definition, and $X \setminus X = \emptyset$, which is a finite set.

1. Arbitrary Union

Let $\{U_\alpha | \alpha \in \Lambda\}$ be open subsets of X , we have that $X \setminus U_\alpha$ is open. Then notice that

$$X \setminus \left(\bigcup_{\alpha \in \Lambda} U_\alpha \right) = \bigcap_{\alpha \in \Lambda} (X \setminus U_\alpha)$$

by DeMorgan's, which is an intersection of finite sets, so it must also be finite.

2. Finite Intersection

Let $\{U_k | k \leq n\}$ be open, we have that $X \setminus U_k$ are finite, so then we notice that

$$X \setminus \left(\bigcap_{k \leq n} U_k \right) = \bigcup_{k \leq n} (X \setminus U_k)$$

By DeMorgan's, which is a finite union of finite sets, which implies that it is finite, implying it must be in τ

Definition 1.5 (Cocountable Topology). The **Cocountable Topology** is defined synonymously to the cofinite topology. It is the collection

$$\tau = \{U \subseteq X | X \setminus U \text{ is countable} \} \cup \{\emptyset\}$$

Prove that this is a topology

Definition 1.6. Let τ, τ' be topologies on X , if $\tau \subseteq \tau'$, then we say

- τ is coarser than τ'
- τ' is finer than τ

As a more general statement, if $\tau \subseteq \tau'$ or $\tau' \subseteq \tau$, then we say that τ and τ' are comparable.

Ω1.4 Bases

These are very similar to the bases you're probably familiar with from linear algebra, where the linear combinations form the vector space, where as these bases are sets that "Unionize" to form the topological space.

Definition 1.7 (Bases). A bases for a topology on a set X is a collection of subsets of X , \mathcal{B} such that

- for all $x \in X$, there exists $B \in \mathcal{B}$ such that $x \in B$
- if $x \in B_1 \cap B_2$, then there exists $B_3 \in \mathcal{B}$ such that $x \in B_3 \subseteq B_1 \cap B_2$

Lemma 1.8 (Generated Topologies)

Let \mathcal{B} be a basis of X , then the collection of all subsets U of X such that for all $x \in U$ there exists $B \in \mathcal{B}$ such that $x \in B \subseteq U$ is a topology on X . This is called the **Generated Topology**

Proof. We now need to prove that this is indeed a topology. We first have that $\emptyset \in \tau$ for vacuous reasons, then we also have that $X \in \tau$ because by letting $x \in X$ be arbitrary, we have that, by definition, there exists a $B \in \mathcal{B}$ such that $x \in B \subseteq X$.

1. Union

Let $\{U_\alpha | \alpha \in \Lambda\}$ be some collection of open sets, let $x \in \bigcup_{\alpha \in \Lambda} U_\alpha$, then there is some $\alpha \in \Lambda$ such that $x \in U_\alpha$, then let $B \in \mathcal{B}$ be such that $x \in B \subseteq U_\alpha \subseteq \bigcup_{\alpha \in \Lambda} U_\alpha$

2. Intersection

Let $U_1, U_2 \in \tau$, let $x \in U_1 \cap U_2$, since $U_1, U_2 \in \tau$, there exists B_1, B_2 such that

- $x \in B_1 \subseteq U_1$
- $x \in B_2 \subseteq U_2$

so there must exist B_3 such that $x \in B_3 \subseteq B_1 \cap B_2$

□

Example 1.9 (Circles)

The bases of open disks in \mathbb{R}^2

$$\tau = \{\{x \in \mathbb{R}^2 | d(x, y) < \varepsilon\} | y \in \mathbb{R}^2, \varepsilon > 0\}$$

Example 1.10 (Rectangles)

$$\tau = \{(a_1, b_1) \times (a_2, b_2) | a_1, b_1, a_2, b_2 \in \mathbb{R}\}$$

Lemma 1.11

For subset \mathcal{B} of τ on X , The following are equivalent

- \mathcal{B} is a basis generating τ
- every non-empty element of τ is a union of elements of \mathcal{B}

Proof. Letting (X, τ) be a topological space, we begin with the forwards direction. Let $U \in \tau$ and let $x \in U$, since τ is generated by \mathcal{B} , there exists some $B_x \in \mathcal{B}$ such that $x \in B_x \subseteq U$, so it follows that

$$U = \bigcup_{x \in U} B_x^2$$

Conversely, let $x \in X$ be arbitrary. Since $X \in \tau$, $X = \bigcup_{\alpha \in \Lambda} U_\alpha$ with $B_\alpha \in \mathcal{B}$ it follows that $x \in B_\alpha$ for some $\alpha \in \Lambda$. Now let $B_1, B_2 \in \mathcal{B}$ with x existing in their intersection. We first have that $B_1 \cap B_2$ is open, so we may represent it as a union,

$$B_1 \cap B_2 = \bigcup_{\alpha \in \Lambda} B_\alpha$$

so $x \in B_\alpha$ for some $\alpha \in \Lambda$. so $x \in B_\alpha \subseteq B_1 \cap B_2$

□

²This equality wasn't immediately obvious to me why this is true, but you can prove it with double inclusion

Ω2 Day 2 : Bases and Sub-bases

Proposition 2.1

Let \mathcal{B} and \mathcal{B}' be bases for topologies τ and τ' respectively on X . Then the following are equivalent.

- τ' is finer than τ
- for all $x \in X$ and for all $B \in \mathcal{B}$. if $x \in B$, then there exists $B' \in \mathcal{B}'$ such that $x \in B' \subseteq B$

Proof. Let $x \in X$ and $B \in \mathcal{B} \subseteq \tau$ so B is open with respect to τ , since τ' is finer than τ , we have that $B \in \tau'$, and therefor there is some $B' \in \mathcal{B}'$ such that $x \in B' \subseteq B$.

Conversely let $U \in \tau$ be arbitrary. Since $U \in \tau$, we have that

$$U = \bigcup_{\alpha \in \Lambda} B_{\alpha}$$

for some collection $B_{\alpha} \in \mathcal{B}$. We further have that each $B_{\alpha} = \bigcup_{\gamma \in \Lambda_{\alpha}} B'_{\gamma}$ with $B'_{\gamma} \in \mathcal{B}'$. Then rewriting U yields

$$U = \bigcup_{\alpha \in \Lambda} \left(\bigcup_{\gamma \in \Lambda_{\alpha}} B'_{\gamma} \right)$$

Thus U is a union of elements from \mathcal{B}' , so $U \in \mathcal{B}'$ □

Recall from last lecture the topology generated by the bases of open disks and open rectangles of \mathbb{R}^2 , using this proposition we may show that both topologies are equivalent, as we may always find sufficiently small circles and rectangles that fit within other circles and rectangles.

Example 2.2 (The Line)

The underlying set is \mathbb{R} and τ is generated by the open intervals (x, y)

Example 2.3 (Sorgenfrey Line)

This is also commonly referred to as the **Lower Limit Topology**, the underlying set is \mathbb{R} and τ is generated by clopen intervals $[x, y)$

Example 2.4 (K-Topology)

The **K-Topology** has an underlying set of \mathbb{R} and τ is generated by

$$(x, y) \qquad (x, y) \setminus K$$

where $K = \{\frac{1}{n} | n \in \mathbb{N}\}$

An exercise left by the prof is to show that the basis generating the K-topology is indeed a basis.

Ω2.1 Sub-Bases

Definition 2.5 (Sub-Bases). A **Sub-Bases** on X is a collection, \mathcal{S} of subsets of X such that \mathcal{S} covers X

Lemma 2.6

The finite intersection of elements of a sub-bases, \mathcal{S} , for a topology on X is a bases for a topology calle dthe basis generated by \mathcal{S}

Proof. Let \mathcal{B} bet he collection of finite intersection of elements of \mathcal{S} , to prove \mathcal{B} is a basis we must check that it covers X , which is true because $\mathcal{S} \subseteq \mathcal{B}$ and \mathcal{S} covers X . and we may find a basis element inside the intersection of two elements. to do this we let $B_1, B_2 \in \mathcal{B}$ be arbitrary, let $x \in B_1 \cap B_2$, by definition, we have that B_1 and B_2 are a finite intersection of elements from \mathcal{S} , so $B_1 \cap B_2$ is also a finite inetrsection of elements of \mathcal{S} , so picking $B_3 = B_1 \cap B_2 \in \mathcal{B}$, it follows that

$$x \in B_3 = B_1 \cap B_2 \subseteq B_1 \cap B_2$$

Thus verifying that it is a bases

□

Ω3 Day 3 : Axiom of Choice, Zorn's Lemma and Ordering

Ω3.1 Axiom of Choice

Theorem 3.1 (Axiom of Choice)

if $\{A_\alpha | \alpha \in \Lambda\}$ is a collection of non-empty sets, then their product

$$\prod_{\alpha \in \Lambda} A_\alpha$$

is non-empty as well

Image1

Remark 3.2. The Axiom of Choice is used to pick from a **non-finite** product of non-empty sets.

Picking from a finite product is pretty straight forward, take \mathbb{R}^n as an example.

Ω3.2 Orderings of Sets

Definition 3.3 (Partial Order). A **Partial Order**, \leq , on a set P , is a binary relation that satisfies

1. (Reflexitivity) for all $p \in P$, $p \leq p$
2. (Anti-Symmetry) for all $p, q \in P$ if $p \leq q$ and $q \leq p$, then $p = q$
3. (Transitivity) for all $p, q, r \in P$ if $p \leq q$ and $q \leq r$, then $p \leq r$

For any $p, q \in P$ if either $p \leq q$ or $q \leq p$ is always true, then we say that \leq is total³

Let (P, \leq) be a partially ordered set,

1. $p \in P$ is maximal if there is no q such that $p < q$
2. $q \in P$ is minimal if there is no p such that $p < q$
3. $Q \subset P$ is bounded, if there exists an $r \in P$ such that $q \leq r$ for all $q \in Q$
4. $C \subseteq P$ is a **Chain** if the restriction of \leq to C is total

Further more we say that a totally ordered set P is **Well-Ordered** if every non-empty subset has a minimal element.

³or linear, but I will usually be using total

Theorem 3.4 (Well-Ordering Principle)

Every non-empty set is well-orderable

Proof. The proof of this requires the Axiom of Choice, the proof wasn't covered in class but you're interested [heres a link](#) \square

Proposition 3.5

In a well ordered set S , every element, which is not maximal, has an immediate successor

Remark 3.6. The intuition for this is that given a non maximal element, x , there exists nothing between x and it's successor. or more formally, for $x \in S$, there exists $y \in S$ such that $x < y$, and if $x < z$ it follows that $y \leq z$

Proof. let $x \in S$ be an arbitrary non-maximal element, then $\{z \in S | x < z\} = A$ is a non-empty set. By Well-Ordering, let $y = \min A$, then we have that $y > x$ and for all $z \in A$, $y \leq z$, by definition. \square

 Ω 3.3 Zorn's Lemma**Lemma 3.7** (Zorn's Lemma)

Every non-empty partially ordered set in which every chain is bounded has a maximal element.

The proof for this is rather complicated and technical, much too far out of the scope for this course, but if you're interested you can look it up [here](#). We're asked to prove a similiar result on the problem set, but we've been given the extra hypothesis of that every chain is finite. A rather surprising result is that in some axiomatic settings, Zorn's Lemma, the axiom of choice, and the well-ordering principle are all equivalent. You can again read about it [here](#).

 Ω 3.4 Order Topologies

Definition 3.8 (Order Topology). Given an ordered set $(X, <)$ with at least 2 elements, the order topology on X is the topology generated by the basis consisting of intervals

1. $(x, y) = \{z \in X | x < z < y\}$
2. $[x_{\min}, y) = \{z \in X | x_{\min} \leq z < y\}$
3. $(x, y_{\max}] = \{z \in X | x < z \leq y_{\max}\}$

We notice that the minimum and maximum might not exist. If either exists we add 2 and 3 to our basis accordingly

Before we delve into some examples, it is important to recall that a set X is countable if there exists a surjection

$$f : \mathbb{N} \rightarrow X$$

We require that it is surjective so that finite sets are also countable.⁴

Example 3.9 (The Line)

The underlying set is \mathbb{R} , with the usual order. The order topology is generated by the open intervals of \mathbb{R} , as we have that \mathbb{R} does not have a maximum

Example 3.10 (The Lexicographic Plane)

The underlying set is \mathbb{R}^2 and the order relation is given by

$$\langle a, b \rangle^a <_{lex} \langle c, d \rangle \text{ if either}$$

1. $a <_{\mathbb{R}} c$
2. $a = b$ and $b <_{\mathbb{R}} d$

^aWe use this notation to represent a tuple, to avoid confusion with intervals

Example 3.11 (The Natural Numbers)

The underlying set are the Natural Numbers. the order is as usual. We notice that \mathbb{N} isn't generated by the open intervals, because 0 doesn't exist within any open interval. But we have that the basis for the order topology of \mathbb{N} is indeed a basis, because we first may let $n \in \mathbb{N}$ be arbitrary, if $n = 0$ then $n \in [0, 1)$, otherwise we have that $n \in (n - 1, n + 2)$, which are both sets that are in the basis. We further have that the order and discrete topologies on \mathbb{N} are equal.

Example 3.12 (Two Copies of the Natural Numbers)

The underlying set is $\{0, 1\} \times \mathbb{N}$, and the order is the lexicographic order. We actually have that every point in this topology is open, apart from $\langle 1, 0 \rangle$

⁴It is convention that the empty set is also countable

Example 3.13 (ω_1 , The Smallest Uncountable Ordinal)

This is an uncountable, well ordered set such that $\forall \alpha \in \omega_1$, the set $\{\gamma \mid \gamma < \alpha\}$ is countable

Theorem 3.14

ω_1 exists

Proof. Let X be an uncountable set and let \leq be a well ordering on X . Then consider the lexicographic order in $\{0, 1\} \times X$. We have the following two facts

- $\{x \in \{0, 1\} \times X \mid \{y \leq_{lex} x\} \text{ is uncountable}\}$ is non empty.

This is true because $\langle 1, \min X \rangle$ exists within the set

- By Well-Ordering, there exists a minimum, x , with uncountably many predecessors, so the set of predecessors of x is uncountable.

□

Ω4 Day 4 : Finite Products, Product Topology and Projections

Ω4.1 Finite Product and Product Topology

First recall that the **Cartesian Product** between two sets, X, Y is defined to be

$$X \times Y = \{\langle x, y \rangle | x \in X, y \in Y\}$$

Then we further define the Product topology as follow,

Definition 4.1 (Product Topology). Let $(X, \tau), (Y, \mu)$ be topological spaces, the **Product Topology** on $X \times Y$ is the topology generated by the basis

$$\mathcal{B} = \{U \times V | U \in \tau \text{ and } V \in \mu\}$$

Proposition 4.2

\mathcal{B} is a basis for a topology

Proof. We first begin by proving that \mathcal{B} covers $X \times Y$, let $\langle x, y \rangle \in X \times Y$ be arbitrary, we notice that $X \in \tau$ and $Y \in \mu$, so we have that $\langle x, y \rangle \in X \times Y \in \mathcal{B}$.

Now let $B_1, B_2 \in \mathcal{B}$, by definition we have $B_1 = U_1 \times V_1$ and $B_2 = U_2 \times V_2$ for suitable sets. Let $\langle x, y \rangle \in (U_1 \times V_1) \cap (U_2 \times V_2)$. Then we notice that

$$\langle x, y \rangle \in (U_1 \cap U_2) \times (V_1 \cap V_2) \subseteq (U_1 \times V_1) \cap (U_2 \times V_2)$$

□

Theorem 4.3

If \mathcal{B} and \mathcal{S} are bases for X and Y , respectively. Then the set

$$\mathcal{D} = \{B \times C | B \in \mathcal{B}, C \in \mathcal{S}\}$$

is a basis for the product topology

Proof. We wish to show that it indeed is a basis, and that it does generate the product topology.

We begin by showing that it covers. Let $\langle x, y \rangle \in X \times Y$ be arbitrary, we then notice two things

1. Since \mathcal{B} is a basis for X , there exists $B \in \mathcal{B}$ such that $x \in B$
2. Since \mathcal{S} is a basis for Y , there exists $S \in \mathcal{S}$ such that $y \in S$

So we have $\langle x, y \rangle \in B \times S \in \mathcal{D}$

Further let $B_1 \times C_1, B_2 \times C_2 \in \mathcal{D}$, let $\langle x, y \rangle \in (B_1 \times C_1) \cap (B_2 \times C_2)$. Notice that since B_1 and B_2 belong to the same topological space, and that x exists in their intersection, we have the existence of some $B_3 \in \mathcal{B}$ such that $x \in B_3 \subseteq B_1 \cap B_2$. We may similarly conclude the existence of a C_3 . Thus we have

$$\langle x, y \rangle \in B_3 \times C_3 \subseteq (B_1 \times C_1) \cap (B_2 \times C_2)$$

Now that we have shown that \mathcal{D} is indeed a bases, we are left to show that it generates the product topology. Let $U \in X$ and $V \in Y$ be open. We may rewrite them as

$$U = \bigcup_{\alpha \in \Lambda} B_\alpha, B_\alpha \in \mathcal{B} \text{ and } V = \bigcup_{\gamma \in \Pi} C_\gamma, C_\gamma \in \mathcal{C}$$

Thus it follows That

$$\begin{aligned} U \times V &= \bigcup_{\alpha \in \Lambda} B_\alpha \times \bigcup_{\gamma \in \Pi} C_\gamma \\ &= \bigcup_{\alpha \in \Lambda, \gamma \in \Pi} B_\alpha \times C_\gamma \end{aligned}$$

□

Proposition 4.2 and Theorem 4.3 basically say that we can mix and match topologies and bases. An example of this is the plane

Example 4.4 (The Plane)

Given $\mathbb{R}^2 = \mathbb{R} \times \mathbb{R}$ with each copy of the real line endowed with the usual topology, we have that the topology of \mathbb{R}^2 is generated by open rectangles.

Ω4.2 Projections

Definition 4.5 (Projection). The functions $\pi_1 : X \times Y \twoheadrightarrow X$ and $\pi_2 : X \times Y \twoheadrightarrow Y$ are given by

$$\pi_1 \langle x, y \rangle = x \qquad \pi_2 \langle x, y \rangle = y$$

are called the projections of $X \times Y$ onto X and Y respectively.

⁵This arrow is to symbolize that the function is surjective

Proposition 4.6

Given topological spaces, (X, τ) and (Y, μ) , the collection

$$\mathcal{S} = \{\pi_1^{-1}(U), \pi_2^{-1}(V) | U \in \tau, V \in \mu\}$$

form a subbases for the product topology on $X \times Y$

Remark 4.7. The pre-image of U and V are actually elements of $X \times Y$, this kinda confused me in lecture.

Proof. We begin by showing covering, we first notice that the preimage of X is necessarily $X \times Y$, thus we have covering.

Now to show that \mathcal{S} does generate the product topology. Let $U \in \tau, V \in \mu$, we want to express $U \times V$ as a finite intersection of elements of \mathcal{S} . So we may notice that

$$U \times V = \pi_1^{-1}(U) \cap \pi_2^{-1}(V)$$

□