

# Modeling plant stable isotopes using the 3PG forest growth model in R

*Daniel M Griffith*

*April 12, 2018*

The r3PG R package implements the 3-PG (Physiological Processes Predicting Growth) model which was originally developed by Landsberg and Waring (1997). Specifically, this R implementation of the model is based directly on the Python version created by Wei et al. (2014) to incorporate predictions of stable isotope compositions. This vignette is focused specifically on using r3PG and predicting carbon and oxygen isotopic values. For a thorough introduction to 3PG and raw data for many of the examples presented here, check out the course materials available online.

As implemented in R, the 3PG model requires two types of input which are climate data and information about the site and tree species. This structure is a legacy from the earlier versions, and for compatibility “.cfg” files used in the Python version can also be used. However, r3PG also accepts native R objects and includes functions for reading and editing model configuration data within R.

Below, we provide a workflow for comparing the stable isotope predictions for a variety of tree species at four sites with different climates.

## Installation

Install r3PG from GitHub using devtools.

```
install.packages("devtools", repos = "http://cran.us.r-project.org")
library(devtools)
install_github(repo = "griffithdan/r3PG")
library(r3PG)
```

## Load example climate data

The “r3PG” R package has monthly climate normals available for four example sites.


```
data("clim.WesternOR")
data("clim.Argentina")
data("clim.NorthCarolina")
data("clim.BritishColumbia")

par(mfrow = c(1,2))
with(clim.WesternOR, plot(Rain ~ c(1:12), type = "l", ylab = NA, xlab = "",
                        xaxt = "n", col = "dodgerblue", main = "Western Oregon"))
axis(side = 1, at = c(1:12), labels = month.abb, las = 2)
axis(side = 2, col = "dodgerblue", col.axis = "dodgerblue")
mtext(side = 2, line = 3, "Rain (mm)", col = "dodgerblue")
par(new = T)
with(clim.WesternOR, plot(Tav ~ c(1:12), pch=16, axes=F, xlab=NA, ylab=NA,
                        type = "l", col = "darkorange"))
axis(side = 4, col = "darkorange", col.axis = "darkorange")
mtext(side = 4, line = 3, "Temperature (c)", col = "darkorange")
with(clim.NorthCarolina, plot(Rain ~ c(1:12), type = "l", ylab = NA,
```

```

xlab = "", xaxt = "n", col = "dodgerblue",
main = "North Carolina"))
axis(side = 1, at = c(1:12), labels = month.abb, las = 2)
axis(side = 2, col = "dodgerblue", col.axis = "dodgerblue")
mtext(side = 2, line = 3, "Rain (mm)", col = "dodgerblue")
par(new = T)
with(clim.NorthCarolina, plot(Tav ~ c(1:12), pch=16, axes=F,
xlab=NA, ylab=NA, type = "l",
col = "darkorange"))
axis(side = 4, col = "darkorange", col.axis = "darkorange")
mtext(side = 4, line = 3, "Temperature (c)", col = "darkorange")

```



Load example configuration data

Compare species


```

data("cfg.Pseudotsuga.menziesii")
data("cfg.Pinus.ponderosa")
pipo.run <- run_3pg(config = cfg.Pinus.ponderosa,
  climate = example.clim,
  output = FALSE,
  python_indexing = FALSE)
doug.run <- run_3pg(config = cfg.Pseudotsuga.menziesii,
  climate = example.clim,
  output = FALSE,
  python_indexing = FALSE)

```

```
example.clim$date <- apply(X = cbind(example.clim$Year, example.clim$Month), MARGIN = 1, FUN = function
example.clim$date <- strptime(x = example.clim$date, format = "%Y-%m-%d")

plot(tail(pipo.run$D13CTissue+1.5,100) ~ as.Date(tail(example.clim$date, 100)), type = "l", col = "dodgerblue",
lines(tail(doug.run$D13CTissue+1.5,100) ~ as.Date(tail(example.clim$date, 100)), type = "l", col = "darkorange")
```



## Compare species at sites

```
data("clim.Argentina")
data("clim.BritishColumbia")
data("clim.NorthCarolina")
data("clim.WesternOR")

cfg.Pinus.ponderosa$TimeRange
cfg.Pinus.ponderosa$TimeRange$EndAge <- 25
cfg.Pinus.ponderosa$TimeRange$EndYear <- 1975

pipo.Arg.run <- run_3pg(config = cfg.Pinus.ponderosa,
                        climate = clim.Argentina,
                        output = FALSE,
                        python_indexing = FALSE)
```


```

pipo.WOR.run <- run_3pg(config = cfg.Pinus.ponderosa,
  climate = clim.WesternOR,
  output = FALSE,
  python_indexing = FALSE)

pipo.Arg.run$D13CTissue <- pipo.Arg.run$D13CTissue + 1.5
pipo.WOR.run$D13CTissue <- pipo.WOR.run$D13CTissue + 1.5

plot(tail(pipo.Arg.run$D13CTissue, 12) ~ 1, xaxt = "n", type = "l", col = "dodgerblue", ylab = "d13C wood",
  lines(tail(pipo.WOR.run$D13CTissue, 12) ~ 1, type = "l", col = "darkorange", ylab = "d13C wood", xlab = "month",
  axis(side = 1, at = c(1:12), labels = month.abb))

```



```

data("cfg.Eucalyptus.globulus")
data("cfg.Picea.sitchensis")
data("cfg.Pinus.contorta")
data("cfg.Pinus.radiata")
data("cfg.Pinus.taeda")

```

## References

Landsberg, J.J., and R.H. Waring. “A Generalised Model of Forest Productivity Using Simplified Concepts of Radiation-Use Efficiency, Carbon Balance and Partitioning.” *Forest Ecology and Management* 95, no. 3 (August 1997): 209-28. [https://doi.org/10.1016/S0378-1127\(97\)00026-1](https://doi.org/10.1016/S0378-1127(97)00026-1).

Wei, Liang, John D. Marshall, Jianwei Zhang, Hang Zhou, and Robert F. Powers. “3-PG Simulations of Young Ponderosa Pine Plantations under Varied Management Intensity: Why Do They Grow so Differently?” *Forest Ecology and Management* 313 (February 2014): 69-82. <https://doi.org/10.1016/j.foreco.2013.10.035>.