Exploratory Data Analysis Lecture 10

Corina Besliu

Technical University of Moldova

October 3, 2021



Formulate the hypothesis

 $H_0: \beta_k = 0$ $H_A: \beta_k \neq 0$

Formulate the hypothesis

$$H_0: \beta_k = 0$$

$$H_A: \beta_k \neq 0$$

$$t_k = \frac{(\hat{\beta}_k - \beta_{H_0})}{SE(\hat{\beta}_k)} \qquad (k = 1, 2, \dots, K)$$

- **③** Find t-critical in the t-table for chosen significance level α (commonly $\alpha=0.05$) and respective degrees of freedom (df = N K 1)
- Reject H_0 if $-t_c < t_k > t_c$

Formulate the hypothesis

$$H_0: \beta_k = 0$$

 $H_A: \beta_k \neq 0$

$$t_k = \frac{(\hat{\beta}_k - \beta_{H_0})}{SE(\hat{\beta}_k)} \qquad (k = 1, 2, \dots, K)$$

- § Find t-critical in the t-table for chosen significance level α (commonly $\alpha=0.05$) and respective degrees of freedom (df = N K 1)
- Reject H_0 if $-t_c < t_k > t_c$

Formulate the hypothesis

$$H_0: \beta_k = 0$$

 $H_A: \beta_k \neq 0$

$$t_k = \frac{(\hat{\beta}_k - \beta_{H_0})}{SE(\hat{\beta}_k)} \qquad (k = 1, 2, \dots, K)$$

- § Find t-critical in the t-table for chosen significance level α (commonly $\alpha=0.05$) and respective degrees of freedom (df = N K 1)
- Reject H_0 if $-t_c < t_k > t_c$


Formulate the hypothesis

$$H_0: \beta_k = 0$$

 $H_A: \beta_k \neq 0$

$$t_k = \frac{(\hat{\beta}_k - \beta_{H_0})}{SE(\hat{\beta}_k)} \qquad (k = 1, 2, \dots, K)$$

- § Find t-critical in the t-table for chosen significance level α (commonly $\alpha=0.05$) and respective degrees of freedom (df = N K 1)



A p-value for a t_k is the probability of observing a t-score that size or larger (in absolute value) if H_0 were true $(0 \ge p - value \le 1)$.

In other words

p-value is the probability that you rejected a H_0 that was correct (assuming that the estimate is in the expected direction).

A p-value for a t_k is the probability of observing a t-score that size or larger (in absolute value) if H_0 were true $(0 \ge p - value \le 1)$.

In other words:

p-value is the probability that you rejected a H_0 that was correct (assuming that the estimate is in the expected direction).

A p-value for a t_k is the probability of observing a t-score that size or larger (in absolute value) if H_0 were true $(0 \ge p - value \le 1)$.


In other words:

p-value is the probability that you rejected a H_0 that was correct (assuming that the estimate is in the expected direction).

A p-value for a t_k is the probability of observing a t-score that size or larger (in absolute value) if H_0 were true $(0 \ge p - value \le 1)$.

In other words:

p-value is the probability that you rejected a H_0 that was correct (assuming that the estimate is in the expected direction).



To cofidently reject a null, you will want a <u>low p-value</u>

 $p-value < 0.05 \rightarrow reject H_0$

If $p-value > 0.05 \rightarrow \alpha > 0.05 \rightarrow P(\text{Type I Error}) > 0.05$ Not Great!

To cofidently reject a null, you will want a <u>low p-value</u> **Rule of Thumb:**

$$p-value < 0.05
ightarrow {
m reject} \ H_0$$

If
$$p-value > 0.05 \rightarrow \alpha > 0.05 \rightarrow \frac{P(\text{Type I Error}) > 0.05}{\text{Not Great!}}$$

To cofidently reject a null, you will want a <u>low p-value</u> **Rule of Thumb:**

$$p-value < 0.05 \rightarrow reject H_0$$

If $p-value > 0.05 \rightarrow \alpha > 0.05 \rightarrow \frac{P(\text{Type I Error}) > 0.05}{\text{Not Great!}}$

To cofidently reject a null, you will want a <u>low p-value</u> **Rule of Thumb:**

$$p-value < 0.05 \rightarrow \text{reject } H_0$$

If $p-value > 0.05 \rightarrow \alpha > 0.05 \rightarrow P(Type \ I \ Error) > 0.05$ Not Great!

To cofidently reject a null, you will want a <u>low p-value</u> **Rule of Thumb:**

$$p-value < 0.05 \rightarrow \text{reject } H_0$$

If
$$p-value > 0.05 \rightarrow \alpha > 0.05 \rightarrow \frac{P(\text{Type I Error}) > 0.05}{\text{Not Great!}}$$

To cofidently reject a null, you will want a <u>low p-value</u>

Rule of Thumb:

$$p-value < 0.05 \rightarrow \text{reject } H_0$$
 If $p-value > 0.05 \rightarrow \alpha > 0.05 \rightarrow \frac{\text{P(Type I Error)} > 0.05}{\text{Not Great!}}$

Regression Results

OLS Regression Results

Dep. Variable:	Profit	R-squared (uncentered):	0.984
Model:	OLS	Adj. R-squared (uncentered):	0.982
Method:	Least Squares	F-statistic:	839.8
Date:	Wed, 29 Sep 2021	Prob (F-statistic):	2.95e-49
Time:	11:27:08	Log-Likelihood:	-661.85
No. Observations:	60	AIC:	1332.
Df Residuals:	56	BIC:	1340.
Df Model:	4		
Covariance Type:	nonrobust		

	coef	std err	t	P> t	[0.025	0.975]
RD	0.7145	0.068	10.490	0.000	0.578	0.851
Admin	0.2546	0.043	5.927	0.000	0.169	0.341
Marketing	0.0962	0.025	3.915	0.000	0.047	0.145
Office	0.9425	0.663	1.421	0.161	-0.386	2.271
Omnibus:		6.4	29 Durbin	-Watson:		1.745
Prob(Omnibus):	0.0	40 Jarque	-Bera (JB):		9.643
Skew:		-0.1	96 Prob(J	B):		0.00805
Kurtosis:		4.9	24 Cond.	No.		91.1

Although R^2 and \bar{R}^2 measure the overall degree of fit of an equation, they don't provide a formal hypothesis test of that overall fit.

But we have the F-test!

The F-test is a joint hypothesis test.

 $H_0: \beta_1 = \beta_2 = ... = \beta_k = 0$ $H_A: H_0$ in not true

Although R^2 and \bar{R}^2 measure the overall degree of fit of an equation, they don't provide a formal hypothesis test of that overall fit.

But we have the F-test!

The F-test is a joint hypothesis test.

$$H_0: \beta_1 = \beta_2 = ... = \beta_k = 0$$

 $H_A: H_0$ in not true

Although R^2 and \bar{R}^2 measure the overall degree of fit of an equation, they don't provide a formal hypothesis test of that overall fit.

But we have the F-test!

The F-test is a joint hypothesis test.

$$H_0: \beta_1 = \beta_2 = ... = \beta_k = 0$$

 $H_A: H_0$ in not true

Although R^2 and \bar{R}^2 measure the overall degree of fit of an equation, they don't provide a formal hypothesis test of that overall fit.

But we have the F-test!

The F-test is a joint hypothesis test.

$$H_0:\beta_1=\beta_2=...=\beta_k=0$$

 $H_A: H_0$ in not true

$$H_0: \beta_1 = \beta_2 = ... = \beta_k = 0$$

 $H_A: H_0$ in not true

• Translate the null hypothesis into constraints on the equation. \rightarrow Constrained equation: $Y_i = \beta_0 + \epsilon_i$ Unconstrained equation: $Y_i = \beta_0 + \beta_1 + \beta_2 + ... + \beta_k + \epsilon_i$

$$H_0: \beta_1 = \beta_2 = ... = \beta_k = 0$$

 $H_A: H_0$ in not true

lacktriangle Translate the null hypothesis into constraints on the equation. ightarrow

Unconstrained equation: $Y_i = \beta_0 + \beta_1 + \beta_2 + ... + \beta_k + \epsilon_i$

$$H_0: \beta_1 = \beta_2 = ... = \beta_k = 0$$

 $H_A: H_0$ in not true

• Translate the null hypothesis into constraints on the equation. \rightarrow Constrained equation: $Y_i = \beta_0 + \epsilon_i$

Unconstrained equation: $Y_i = \beta_0 + \beta_1 + \beta_2 + ... + \beta_k + \epsilon_i$

$$H_0: \beta_1 = \beta_2 = ... = \beta_k = 0$$

 $H_A: H_0$ in not true

• Translate the null hypothesis into constraints on the equation. \rightarrow Constrained equation: $Y_i = \beta_0 + \epsilon_i$ Unconstrained equation: $Y_i = \beta_0 + \beta_1 + \beta_2 + ... + \beta_k + \epsilon_i$

$$H_0: \beta_1 = \beta_2 = ... = \beta_k = 0$$

 $H_A: H_0$ in not true

- **1** Translate the null hypothesis into constraints on the equation. \rightarrow Constrained equation: $Y_i = \beta_0 + \epsilon_i$ Unconstrained equation: $Y_i = \beta_0 + \beta_1 X_{1i} + \beta_2 X_{2i} + ... + \beta_k + \epsilon_i$
- Estimate the constrained equation and the unconstrained equation to compute:

$$F = \frac{(RSS_{M} - RSS)/M}{RSS/(N - K - 1)}$$

where

 ${\it M}$ - degrees of freedom in numerator (nb. of constraints ${\it N}-{\it K}-1$ - degrees of freedom in the denominator

 $\begin{array}{ll} \text{ Reject} & H_0 \text{ if } F > F_c \\ \text{ Do not reject} & H_0 \text{ if } F < F_c \end{array}$

$$H_0: \beta_1 = \beta_2 = ... = \beta_k = 0$$

 $H_A: H_0$ in not true

- **1** Translate the null hypothesis into constraints on the equation. \rightarrow Constrained equation: $Y_i = \beta_0 + \epsilon_i$ Unconstrained equation: $Y_i = \beta_0 + \beta_1 X_{1i} + \beta_2 X_{2i} + ... + \beta_k + \epsilon_i$
- Estimate the constrained equation and the unconstrained equation to compute:

$$F = \frac{(RSS_M - RSS)/M}{RSS/(N - K - 1)}$$

where,

 ${\it M}$ - degrees of freedom in numerator (nb. of constraints)

 ${\it N}-{\it K}-1$ - degrees of freedom in the denominator

$$H_0: \beta_1 = \beta_2 = ... = \beta_k = 0$$

 $H_A: H_0$ in not true

- **1** Translate the null hypothesis into constraints on the equation. \rightarrow Constrained equation: $Y_i = \beta_0 + \epsilon_i$ Unconstrained equation: $Y_i = \beta_0 + \beta_1 X_{1i} + \beta_2 X_{2i} + ... + \beta_k + \epsilon_i$
- Estimate the constrained equation and the unconstrained equation to compute:

$$F = \frac{(RSS_M - RSS)/M}{RSS/(N - K - 1)}$$

where,

 ${\it M}$ - degrees of freedom in numerator (nb. of constraints)

N-K-1 - degrees of freedom in the denominator

3 Reject H_0 if $F > F_c$ Do not reject H_0 if $F < F_c$

F in the Regression Results

OLS Regression Results

Dep. Variabl	e:	Prof	fit R-squa	ared (uncent	ered):		0.984
Model:		(OLS Adj. I	R-squared (u	ncentered):		0.982
Method:		Least Squar	res F-stat	tistic:			839.8
Date:	We	d, 29 Sep 20	21 Prob	(F-statistic):		2.95e-49
Time:		11:27:	:08 Log-L:	ikelihood:			-661.85
No. Observat	ions:		60 AIC:				1332.
Df Residuals	:		56 BIC:				1340.
Df Model:			4				
Covariance T	vpe:	nonrobu	ıst				
	coef	std err	t	P> t	[0.025	0.975]	
RD	0.7145	0.068	10.490	0.000	0.578	0.851	
Admin	0.2546	0.043	5.927	0.000	0.169	0.341	
Marketing	0.0962	0.025	3.915	0.000	0.047	0.145	
Office	0.9425	0.663	1.421	0.161	-0.386	2.271	
Omnibus:		6.4	======================================	======== n-Watson:		1.745	
Prob(Omnibus):			e-Bera (JB):		9.643	
Skew:	, -	-0.1		, ,		0.00805	
Kurtosis:			924 Cond.	,		91.1	

OLS Assumptions

Notation	Meaning
1. $E(u_i) = 0$	Average value of residuals is zero
$2.\text{Var}\left(u_i\right) = \sigma^2$	The variance of the residuals is constant
$3.\text{Cov }(u_i,x_i)=0$	There is no linear relationship between residues and x
$4. u_i \sim N(0, \sigma^2)$	Residuals have a normal distribution
$5.\mathrm{Cov}\;(u_i,u_j)=0$	Residuals don't depend on each other
6. Multicollinearity	Independent variables are not correlated with each other

- Choosing the correct independent variables
- ② Choosing the correct functional form
- Ohoosing the correct form of the stochastic error term

- Choosing the correct independent variables
- ② Choosing the correct functional form
- Ochoosing the correct form of the stochastic error term

- Choosing the correct independent variables
- Choosing the correct functional form
- Ohoosing the correct form of the stochastic error term

- Ohoosing the correct independent variables
- 2 Choosing the correct functional form
- Ohoosing the correct form of the stochastic error term

Choosing the Correct Independent Variables

There are two mistakes one can make when choosing the independent variables for the model:

- Omitting an important variable
- ② Including an irrelevant variable.

Choosing the Correct Independent Variables

There are two mistakes one can make when choosing the independent variables for the model:

- Omitting an important variable.
- 2 Including an irrelevant variable.

Choosing the Correct Independent Variables

There are two mistakes one can make when choosing the independent variables for the model:

- Omitting an important variable.
- Including an irrelevant variable.

Omitting an Important Variable

Omitted Variable Bias

True Model

$$Y_i\,=\,\beta_0+\beta_1X_{1i}+\beta_2X_{2i}+\varepsilon_i$$

$$Y_{i} = \beta_{0}^{*} + \beta_{1}^{*}X_{1i} + \epsilon_{i}^{*}$$

$$\downarrow$$

$$\epsilon_{i}^{*} = \epsilon_{i} + \beta_{2}X_{2i}$$

If
$$X_1$$
 and X_2 are correlated Cov $(X_{1i}, \epsilon^*) \neq 0 \rightarrow \hat{\beta}_1^* \neq \beta_1$
Biased!

Omitting an Important Variable

Omitted Variable Bias

True Model

$$Y_i\,=\,\beta_0+\beta_1X_{1i}+\beta_2X_{2i}+\varepsilon_i$$

$$Y_{i} = \beta_{0}^{*} + \beta_{1}^{*}X_{1i} + \epsilon_{i}^{*}$$

$$\downarrow$$

$$\epsilon_{i}^{*} = \epsilon_{i} + \beta_{2}X_{2i}$$

If
$$X_1$$
 and X_2 are correlated Cov $(X_{1i}, \epsilon^*) \neq 0 \rightarrow \frac{\hat{\beta}_1^* \neq \beta_1}{\text{Biased}!}$

Omitting an Important Variable

Omitted Variable Bias

True Model

$$Y_i\,=\,\beta_0+\beta_1X_{1i}+\beta_2X_{2i}+\varepsilon_i$$

$$Y_{i} = \beta_{0}^{*} + \beta_{1}^{*}X_{1i} + \epsilon_{i}^{*}$$

$$\downarrow$$

$$\epsilon_{i}^{*} = \epsilon_{i} + \beta_{2}X_{2i}$$

If
$$X_1$$
 and X_2 are correlated Cov $(X_{1i}, \epsilon^*) \neq 0 \rightarrow \frac{\hat{\beta}_1^* \neq \beta_1}{\text{Biased!}}$

Including an irrelevant variable won't bias $\hat{\beta}_{\pmb{k}}$

$$Y_i = \, \beta_0 + \beta_1 X_{1i} + \varepsilon_i$$

Specified Model

$$\begin{aligned} Y_i &= \beta_0 + \beta_1 X_{1i} + \beta_2 X_{2i} + \varepsilon_i^{**} \\ \downarrow \\ \varepsilon_i^{**} &= \varepsilon_i - \beta_2 X_{2i} \end{aligned}$$

$$\hat{eta}_1^*=eta_1$$
 but Var $(\hat{eta}_1)\uparrow$ thus $\underbrace{t_{X_1}\downarrow}_{ ext{Biased}}$

Also $R^2 \uparrow$, but $\bar{R}^2 \downarrow$

Including an irrelevant variable won't bias $\hat{\beta}_{\pmb{k}}$

True Model
$$Y_i = \beta_0 + \beta_1 X_{1i} + \epsilon_i$$

$$\begin{aligned} Y_i &= \beta_0 + \beta_1 X_{1i} + \beta_2 X_{2i} + \varepsilon_i^{**} \\ \downarrow \\ \varepsilon_i^{**} &= \varepsilon_i - \beta_2 X_{2i} \end{aligned}$$

$$\hat{eta}_1^* = eta_1$$
 but $\mathsf{Var}\;(\hat{eta}_1)\uparrow\mathsf{thus}\;\underbrace{t_{X_1}\downarrow}_{\mathsf{Biased}}$

Also
$$R^2 \uparrow$$
, but $\bar{R}^2 \downarrow$

Including an irrelevant variable won't bias \hat{eta}_k

True Model
$$Y_i = \beta_0 + \beta_1 X_{1i} + \epsilon_i$$

Specified Model

$$\begin{aligned} Y_i &= \beta_0 + \beta_1 X_{1i} + \beta_2 X_{2i} + \varepsilon_i^{**} \\ \downarrow \\ \varepsilon_i^{**} &= \varepsilon_i - \beta_2 X_{2i} \end{aligned}$$

$$\hat{eta}_1^* = eta_1$$
 but $\mathsf{Var}\;(\hat{eta}_1)\uparrow\mathsf{thus}\; \underbrace{t_{X_1}\downarrow}_{\mathsf{Biased}}$

Also $R^2 \uparrow$, but \bar{R}^2

Including an irrelevant variable won't bias $\hat{\beta}_{\pmb{k}}$

True Model
$$Y_i = \beta_0 + \beta_1 X_{1i} + \epsilon_i$$

Specified Model

$$\begin{aligned} Y_i &= \beta_0 + \beta_1 X_{1i} + \beta_2 X_{2i} + \varepsilon_i^{**} \\ \downarrow \\ \varepsilon_i^{**} &= \varepsilon_i - \beta_2 X_{2i} \end{aligned}$$

$$\hat{eta}_1^* = eta_1$$
 but $\operatorname{Var}\left(\hat{eta}_1
ight) \uparrow$ thus $\underbrace{t_{X_1}\downarrow}_{\mathsf{Biased}!}$

Also $R^2 \uparrow$, but $\bar{R}^2 \downarrow$

Including an irrelevant variable won't bias \hat{eta}_k

$$Y_i = \beta_0 + \beta_1 X_{1i} + \epsilon_i$$

$$\begin{aligned} Y_i &= \beta_0 + \beta_1 X_{1i} + \beta_2 X_{2i} + \epsilon_i^{**} \\ \downarrow \\ \epsilon_i^{**} &= \epsilon_i - \beta_2 X_{2i} \end{aligned}$$

$$\hat{\beta}_1^* = \beta_1$$
 but $\operatorname{Var}\left(\hat{\beta}_1\right) \uparrow$ thus $\underbrace{t_{X_1} \downarrow}_{\mathsf{Biased}!}$

Also
$$R^2 \uparrow$$
, but $\bar{R}^2 \downarrow$

- Theory: Is the variables place in the equation unambiguous and theoretically sound?
- 2 t-Test: Is the variables estimated coefficient significant in the expected direction?
- ② \bar{R}^2 : Does the overall fit of the equation (adjusted for degrees of freedom) improve when the variable is added to the equation?
- Bias: Do other variables coefficients change significantly when the variable is added to the equation?

- Theory: Is the variables place in the equation unambiguous and theoretically sound?
- 2 t-Test: Is the variables estimated coefficient significant in the expected direction?
- **3** \bar{R}^2 : Does the overall fit of the equation (adjusted for degrees of freedom) improve when the variable is added to the equation?
- Bias: Do other variables coefficients change significantly when the variable is added to the equation?

- Theory: Is the variables place in the equation unambiguous and theoretically sound?
- 2 t-Test: Is the variables estimated coefficient significant in the expected direction?
- **3** \bar{R}^2 : Does the overall fit of the equation (adjusted for degrees of freedom) improve when the variable is added to the equation?
- Sias: Do other variables coefficients change significantly when the variable is added to the equation?

- Theory: Is the variables place in the equation unambiguous and theoretically sound?
- 2 t-Test: Is the variables estimated coefficient significant in the expected direction?
- **3** \bar{R}^2 : Does the overall fit of the equation (adjusted for degrees of freedom) improve when the variable is added to the equation?
- Bias: Do other variables coefficients change significantly when the variable is added to the equation?

Should we include the intercept β_0 ?

- Beginner researchers may want to supress the intercept when there is no logical interpretation for β_0 .
- This would be a mistake. β_0 should be suppressed only on very rare occasions.
 - For example: $C_i = \beta_0 + \beta_1 Q_i + \epsilon_i$ where.
 - C_i are the fixed and variable costs of businesses in a branch
- Excluding β_0 in this context would preserve one degree of freedom and would supposedly make the estimate of β_1 more accurate.
- However such examples in real life are rare
- The risks associated with supressing β_0 most times outweight the benefits of doing so.

Should we include the intercept β_0 ?

- Beginner researchers may want to supress the intercept when there is no logical interpretation for β_0 .
- This would be a mistake. β_0 should be suppressed only on very rare occasions.
 - For example: $C_i = \beta_0 + \beta_1 Q_i + \epsilon_i$
 - C_i are the fixed and variable costs of businesses in a branch
- Excluding β_0 in this context would preserve one degree of freedom and would supposedly make the estimate of β_1 more accurate.
- However such examples in real life are rare
- The risks associated with supressing β_0 most times outweight the benefits of doing so.

Should we include the intercept β_0 ?

- Beginner researchers may want to supress the intercept when there is no logical interpretation for β_0 .
- This would be a mistake. β_0 should be suppressed only on very rare occasions.

```
For example: C_i = \beta_0 + \beta_1 Q_i + \epsilon_i where
```

- Excluding β_0 in this context would preserve one degree of freedom and would supposedly make the estimate of β_1 more accurate.
- However such examples in real life are rare
- The risks associated with supressing β_0 most times outweight the benefits of doing so.

Should we include the intercept β_0 ?

- Beginner researchers may want to supress the intercept when there is no logical interpretation for β_0 .
- This would be a mistake. β_0 should be suppressed only on very rare occasions.

For example:
$$C_i = \beta_0 + \beta_1 Q_i + \epsilon_i$$

where

- C_i are the fixed and variable costs of businesses in a branch
- Excluding β_0 in this context would preserve one degree of freedom and would supposedly make the estimate of β_1 more accurate.
- However such examples in real life are rare
- The risks associated with supressing β_0 most times outweight the benefits of doing so.

Should we include the intercept β_0 ?

- Beginner researchers may want to supress the intercept when there is no logical interpretation for β_0 .
- This would be a mistake. β_0 should be suppressed only on very rare occasions.

For example: $C_i = \beta_0 + \beta_1 Q_i + \epsilon_i$ where.

- Excluding β_0 in this context would preserve one degree of freedom and would supposedly make the estimate of β_1 more accurate.
- However such examples in real life are rare
- The risks associated with supressing β_0 most times outweight the benefits of doing so.

Should we include the intercept β_0 ?

- Beginner researchers may want to supress the intercept when there is no logical interpretation for β_0 .
- This would be a mistake. β_0 should be suppressed only on very rare occasions.

For example: $C_i = \beta_0 + \beta_1 Q_i + \epsilon_i$ where,

- Excluding β_0 in this context would preserve one degree of freedom and would supposedly make the estimate of β_1 more accurate.
- However such examples in real life are rare
- The risks associated with supressing β_0 most times outweight the benefits of doing so.

Should we include the intercept β_0 ?

- Beginner researchers may want to supress the intercept when there is no logical interpretation for β_0 .
- This would be a mistake. β_0 should be suppressed only on very rare occasions.

For example: $C_i = \beta_0 + \beta_1 Q_i + \epsilon_i$ where,

- Excluding β_0 in this context would preserve one degree of freedom and would supposedly make the estimate of β_1 more accurate.
- However such examples in real life are rare!
- The risks associated with supressing β_0 most times outweight the benefits of doing so.


Should we include the intercept β_0 ?

- Beginner researchers may want to supress the intercept when there is no logical interpretation for β_0 .
- This would be a mistake. β_0 should be suppressed only on very rare occasions.

For example: $C_i = \beta_0 + \beta_1 Q_i + \epsilon_i$ where.


- Excluding β_0 in this context would preserve one degree of freedom and would supposedly make the estimate of β_1 more accurate.
- However such examples in real life are rare!
- The risks associated with supressing β_0 most times outweight the benefits of doing so.

By omitting the constant term, we force the impact of the constant into the estimates of the other coefficients.




- The estimated regression will go through the origin
- $\hat{\beta}_1 \neq \beta_1$ biassed coefficient
- t-scores ↑ also biased

By omitting the constant term, we force the impact of the constant into the estimates of the other coefficients.




- The estimated regression will go through the origin
- $\hat{\beta}_1 \neq \beta_1$ biassed coefficient
- t-scores ↑ also biased

By omitting the constant term, we force the impact of the constant into the estimates of the other coefficients.



- The estimated regression will go through the origin
- $\hat{\beta}_1 \neq \beta_1$ biassed coefficient
- t-scores ↑ also biased

By omitting the constant term, we force the impact of the constant into the estimates of the other coefficients.



- The estimated regression will go through the origin
- $\hat{\beta}_1 \neq \beta_1$ biassed coefficient
- t-scores ↑ also biased


Polinomial Regressions

Sometimes the relationship between X and Y cannot be explained by a line.

Relationship between Earnings and Age.

- As a young worker gets older, his or her earnings will increase.
- Beyond some point an increase in age will not increase earnings, and around retirement earnings will start to fall abruptly with age.

Relationship between Earnings and Age




$$Earnings_i = \beta_0 + \beta_1 Age_i + \beta_2 Age_i^2 + \, \cdots \, + \varepsilon_i$$

$$\frac{\Delta Y}{\Delta X_1} = \, \beta_1 + 2\beta_2 X_1$$

General Advice: Rely of Theory when Defining the Functional Form of Your Model.

Relationship between Earnings and Age



$$Earnings_i = \beta_0 + \beta_1 Age_i + \beta_2 Age_i^2 + \, \cdots \, + \varepsilon_i$$

$$\frac{\Delta Y}{\Delta X_1} = \, \beta_1 + 2\beta_2 X_1$$


General Advice: Rely of Theory when Defining the Functional Form of Your Model.


- Multicollinearity
- ② Serial Correlation $Cov(u_i, u_j) \neq 0$
- **1** Heteroskedasticity $Var(u_i) \neq \sigma^2$

- Multicollinearity
- **2** Serial Correlation $Cov(u_i, u_j) \neq 0$
- **3** Heteroskedasticity $Var(u_i) \neq \sigma^2$

- Multicollinearity
- ② Serial Correlation $Cov(u_i, u_i) \neq 0$
- **3** Heteroskedasticity $Var(u_i) \neq \sigma^2$

- Multicollinearity
- ② Serial Correlation $Cov(u_i, u_i) \neq 0$
- **3** Heteroskedasticity $Var(u_i) \neq \sigma^2$





Let's get Started!

Access Google Colaboratory through your Gmail account