All−method bias distributions

BC marginal A−C RD | common | n = 100 | ESS ratio = 0.5

Adjustment: scores | fit weights: unit | BRM: linear_atanh | source: diffuse | EM: high | calibration: z1,z2,z3,z4,z5,z6 | complete 1000 / 1000



White points: full−data means. Red circles: display−clipped counts; statistics use all valid values.

Bias reference = 0; critical values +/−1; adaptive axis, no observations clipped.

All−method SE distributions

BC marginal A−C RD | common | n = 100 | ESS ratio = 0.5

Adjustment: scores | fit weights: unit | BRM: linear_atanh | source: diffuse | EM: high | calibration: z1,z2,z3,z4,z5,z6 | complete 1000 / 1000



White points: full−data means. Red circles: display−clipped counts; statistics use all valid values.

SE critical value = 1; adaptive axis, no observations clipped.

All−method SE calibration

BC marginal A−C RD | common | n = 100 | ESS ratio = 0.5

Adjustment: scores | fit weights: unit | BRM: linear_atanh | source: diffuse | EM: high | calibration: z1,z2,z3,z4,z5,z6 | complete 1000 / 1000



Target = 1. Matched SE−valid replicates; BC and MN_eff model SE not provided. Darker stems = larger deviation.

All−method coverage rate

BC marginal A−C RD | common | n = 100 | ESS ratio = 0.5

Adjustment: scores | fit weights: unit | BRM: linear_atanh | source: diffuse | EM: high | calibration: z1,z2,z3,z4,z5,z6 | complete 1000 / 1000



Target = 95%. Wald: blue; BC: orange; MN−ESS: purple. Darker stems = larger deviation. Valid returned intervals only.