

Reference excerpt 01-027

Source: arXiv:1604.05171v1 | License: CC BY 4.0 | Word window: 924-1183

is transmitted along the direction AR-IJV according to the following equation $p(z) = p(t) + \frac{1}{c} \frac{dp}{dt} z$

(2) For each cardiac cycle, the pressure in the RA reaches its maximum value between t_P and t_Q , where t_P indicates the ECG P wave and t_Q indicates the wave Q. In this note, for the sake of simplicity, it is assumed that the pressure in the RA is maximum in correspondence of the instant t_{PQ} which is the average between t_P and t_Q . The relationship between the pressure in the RA and the pressure in the point G at the same instant of time is given by: $p(t + l/c, z_G) = p(t, z_{RA})$ (3) Where z_G is the coordinate of the US scanning point and z_{RA} is the RA coordinate. The pressure is, at the same instant t_{PQ} , maximum in the RA ($p(t_{PQ}, z_{RA})$), and equal to $p(t_{PQ} - l/c, z_{RA})$ at the insonation point G. The qualitative JVP trace of 2 Figure 2: The figure a) shows the JVP trace of pressure in the IJV, measured in G, together with the ECG trace. The interval t_{aQP} is represented over the curves a and QP. In figure b) it is shown the pressure along the z-axes at instant t_{QP} . pressure in the IJV, measured in G, can be obtained simultaneously with the acquisition of the ECG trace [Sisini et al.(2016)]. On this trace, that reports both JVP and ECG waves, the time interval t_{aQP} between t_P and