

Reference excerpt 01-028

Source: arXiv:1604.05171v1 | License: CC BY 4.0 | Word window: 959-1218

and t_Q , where t_P indicates the ECG P wave and t_Q indicates the wave Q. In this note, for the sake of simplicity, it is assumed that the pressure in the RA is maximum in correspondence of the instant t_P Q which is the average between t_P and t_Q . The relationship between the pressure in the RA and the pressure in the point G at the same instant of time is given by: $p(t + lG/c, zG) = p(t, zRA)$ (3) Where zG is the coordinate of the US scanning point and zRA is the RA coordinate. The pressure is, at the same instant t_P Q, maximum in the RA ($p(t_P Q, zRA)$), and equal to $p(t_P Q - l/c, zRA)$ at the insonation point G. The qualitative JVP trace of 2 Figure 2: The figure a) shows the JVP trace of pressure in the IJV, measured in G, together with the ECG trace. The interval ta_{QP} is represented over the curves a and QP. In figure b) it is shown the pressure along the z-axes at instant t_{QP} . pressure in the IJV, measured in G, can be obtained simultaneously with the acquisition of the ECG trace [Sisini et al.(2016)]. On this trace, that reports both JVP and ECG waves, the time interval ta_{QP} between t_P Q and ta is given by: $ta_{QP} = ta - t_{QP} = lG/c$ (4) where ta is the instant corresponding the a wave. The parameter ta_{QP} is the time interval between the instant when the pressure