

Reference excerpt 01-029

Source: arXiv:1604.05171v1 | License: CC BY 4.0 | Word window: 995-1254

correspondence of the instant t_{PQ} which is the average between t_P and t_Q . The relationship between the pressure in the RA and the pressure in the point G at the same instant of time is given by: $p(t + lG/c, z_G) = p(t, z_{RA})$ (3) Where z_G is the coordinate of the US scanning point and z_{RA} is the RA coordinate. The pressure is, at the same instant t_{PQ} , maximum in the RA ($p(t_{PQ}, z_{RA})$), and equal to $p(t_{PQ} - l/c, z_{RA})$ at the insonation point G. The qualitative JVP trace of 2 Figure 2: The figure a) shows the JVP trace of pressure in the IJV, measured in G, together with the ECG trace. The interval t_{aQP} is represented over the curves a and QP. In figure b) it is shown the pressure along the z -axes at instant t_{QP} . pressure in the IJV, measured in G, can be obtained simultaneously with the acquisition of the ECG trace [Sisini et al.(2016)]. On this trace, that reports both JVP and ECG waves, the time interval t_{aQP} between t_{PQ} and t_a is given by: $t_{aQP} = t_a - t_{QP} = lG/c$ (4) where t_a is the instant corresponding the a wave. The parameter t_{aQP} is the time interval between the instant when the pressure is maximum at point G (instant t_a) and the instant when the pressure is maximum at point RA (instant t_{QP}). METHODS The applicability of the relationship expressed in Eq. 4, can be tested by comparing