

Reference excerpt 01-044

Source: arXiv:1604.05171v1 | License: CC BY 4.0 | Word window: 1527-1786

CSA x) during the cardiac cycle was 0.2 cm^2 and the compliance for unit of length was $9.8 \text{ } 10^{-3} \text{ cm}^2/\text{mmHg}$. Flow velocity calculation $w(t, z)$ The velocity was calculated following Eq. (1) and its trace is shown in Fig. 3 together with the experimental trace obtained by the Doppler examination. The two traces in agreement, nevertheless, since the velocity $w(t, G + l)$ was calculated up to an additive constant, only the wave amplitude has physical meaning whereas its shift along the velocity axis is not significant. ACKNOWLEDGEMENT The author wants to thank Eleuterio Toro for his critical comments to the paper and Giacomo Gadda, Valentina Tavoni and Valentina Sisini to have kindly reviewed the manuscript. REFERENCES [Applefeld(1990)] Applefeld MM. The Jugular Venous Pressure and Pulse Contour. In: Source Clinical Methods: The History, Physical, and Laboratory Examinations. Boston: Butterworths, 1990. chapter 19. [Beulen et al.(2011)] Beulen, B. et al. Toward noninvasive blood pressure assessment in arteries by using ultrasound. Ultrasound in Medicine; 2011: 37(5), 788-797. [Kalmanson et al. (1972)] Kalmanson D, Veyrat C, Derai C, Savier CH, Berkman M, Chiche P. Non-invasive technique for diagnosing atrial septal defect and assessing shunt volume using directional Doppler ultrasound. Correlations with phasic flow velocity patterns of the shunt. British Heart Journal. 1972;34:981-991. [Mackenzie(1902)] Mackenzie J. The Study of the Pulse, Arterial, Venous, and Hepatic and of the Movements of the Heart. Edinburgh: Young J Pentland, 1902. [Sisini et al.(2015)] Sisini F, Tessari M, Gadda G, Di Domenico G, Taibi A, Menegatti E, Gambaccini M, Zamboni P Sisini