

Reference excerpt 01-046

Source: arXiv:1604.05171v1 | License: CC BY 4.0 | Word window: 1599-1858

additive constant, only the wave amplitude has physical meaning whereas its shift along the velocity axis is not significant. ACKNOWLEDGEMENT The author wants to thank Eleuterio Toro for his critical comments to the paper and Giacomo Gadda, Valentina Tavoni and Valentina Sisini to have kindly reviewed the manuscript. REFERENCES [Applefeld(1990)] Applefeld MM. The Jugular Venous Pressure and Pulse Contour. In: SourceClinical Methods: The History, Physical, and Laboratory Examinations. Boston: Butterworths, 1990.chapter 19. [Beulen et al.(2011)] Beulen, B. et al. Toward noninvasive blood pressure assessment in arteries by using ultrasound. Ultrasound in Medicine;2011: 37(5), 788-797. [Kalmanson et al. (1972)] Kalmanson D, Veyrat C, Derai C, Savier CH, Berkman M, Chiche P. Non-invasive technique for diagnosing atrial septal defect and assessing shunt volume using directional Doppler ultrasound. Correlations with phasic flow velocity patterns of the shunt. British Heart Journal. 1972;34:981-991. [Mackenzie(1902)] Mackenzie J.The Study of the Pulse, Arterial, Venous, and Hepatic and of the Movements of the Heart. Edinburgh: Young J Pentland, 1902. [Sisini et al.(2015)] Sisini F, Tessari M, Gadda G, Di Domenico G, Taibi A, Menegatti E, Gambaccini M, Zamboni P Sisini F, Tessari M, Gadda G, Di Domenico G, Taibi A, Menegatti E, Gambaccini M, Zamboni P. An Ultrasonographic Technique to Assess the Jugular Venous Pulse: a Proof of Concept. Ultrasound Med Biol, 2015;41:1334-1341 [Sisini et al.(2015b)] Sisini F, Toro E, Gambaccini M, and Zamboni P, The Oscillating Component of the Internal Jugular Vein Flow: The Overlooked Element of Cerebral Circulation, Behavioural Neurology. 2015; Article ID 170756 [Sisini et al.(2016)]