

Reference excerpt 01-048

Source: arXiv:1604.05171v1 | License: CC BY 4.0 | Word window: 1669-1928

1990.chapter 19. [Beulen et al.(2011)] Beulen, B. et al. Toward noninvasive blood pressure assessment in arteries by using ultrasound. *Ultrasound in Medicine*;2011: 37(5), 788-797. [Kalmanson et al. (1972)] Kalmanson D, Veyrat C, Derai C, Savier CH, Berkman M, Chiche P. Non-invasive technique for diagnosing atrial septal defect and assessing shunt volume using directional Doppler ultrasound. Correlations with phasic flow velocity patterns of the shunt. *British Heart Journal*. 1972;34:981-991. [Mackenzie(1902)] Mackenzie J.The Study of the Pulse, Arterial, Venous, and Hepatic and of the Movements of the Heart. Edinburgh: Young & Pentland, 1902. [Sisini et al.(2015)] Sisini F, Tessari M, Gadda G, Di Domenico G, Taibi A, Menegatti E, Gambaccini M, Zamboni P. An Ultrasonographic Technique to Assess the Jugular Venous Pulse: a Proof of Concept. *Ultrasound Med Biol*, 2015;41:1334-1341 [Sisini et al.(2015b)] Sisini F, Toro E, Gambaccini M, and Zamboni P, The Oscillating Component of the Internal Jugular Vein Flow: The Overlooked Element of Cerebral Circulation, *Behavioural Neurology*. 2015; Article ID 170756 [Sisini et al.(2016)] Sisini F, Tessari M, Menegatti E, Vannini ME, Ganesini S, Tavoni V, Gadda G, Gambaccini G, Taibi A, Zamboni P. Clinical applicability of the assessment of the jugular flow over the individual cardiac cycle compared with current ultrasound methodology. *Ultrasound Med Biol*, In press. DOI: 10.1016/j.ultrasmedbio.2016.03.002 [Womersley(1955a)] Womersley, J. . Method for the calculation of velocity, rate of flow and viscous drag in arteries when the