

Reference excerpt 02-042

Source: arXiv:1604.05177v1 | License: CC BY 4.0 | Word window: 1556-1815

human expert. This paper has not discussed the clinical use of these parameters.

ACKNOWLEDGEMENT The author wants to thank Giacomo Gadda, Valentina Tavoni and Valentina Sisini to have kindly review the manuscript.

REFERENCES [Applefeld(1990)] Applefeld MM. The Jugular Venous Pres- sure and Pulse Contour. In: SourceClinical Methods: The History, Physical, and Laboratory Examinations. Boston: Butterworths, 1990: Chapter 19. [Beulen et al.(2011)] Beulen, B. et al. Toward noninvasive blood pressure assessment in arteries by using ultrasound. Ultrasound in Medicin, 2011:37(5), 788-797. [Beulen et al.(2010)] Beulen, B., Verkaik, A. C., Bijmens, N ., Rutten, M., Van De Vosse, F. (2010). Perpendicular ul- trasound velocity measurement by 2D cross correlation of RF data. Part B: Volume ow estimation in curved vessels. Experiments in Fluids, 2010:49, 1219-1229. [Chua Chiaco et al.(2013)] Chua Chiaco JM, Parikh NI, Fer- gusson DJ. The jugular venous pressure revisited. Cleve Clin J Med, 2013:80, 638-644. [Kalmanson et al. (1972)] Kalmanson D, Veyrat C, Derai C, Savier CH, Berkman M, Chiche P. Non-invasive technique for diagnosing atrial septal defect and assessing shunt vol - ume using directional Doppler ultrasound. Correlations with phasic ow velocity patterns of the shunt. British Heart Journal, 1972:34, 981-991. [Kohler(2002)] K ohler, B. U., Hennig, C., Orglmeister, R. (2002). The principles of software QRS detection. IEEE Engineering in Medicine and Biology Magazine, 2002:21(1), 42-57. doi:10.1109/51.993193 [Mackenzie(1902)] Mackenzie J.The Study of the Pulse, Arte - rial, Venous, and Hepatic and of the Movements of the Heart. Edinburgh: Young J Pentland, 1902. [Patel et al.(1999)] Patel, M. C., Berman, L. H., Moss, H. a, McPherson, S. J. (1999).