

## Reference excerpt 02-043

Source: arXiv:1604.05177v1 | License: CC BY 4.0 | Word window: 1594-1853

Venous Pressure and Pulse Contour. In: *SourceClinical Methods: The History, Physical, and Laboratory Examinations*. Boston: Butterworths, 1990: Chapter 19. [Beulen et al.(2011)] Beulen, B. et al. Toward noninvasive blood pressure assessment in arteries by using ultrasound. *Ultrasound in Medicine*, 2011:37(5), 788-797. [Beulen et al.(2010)] Beulen, B., Verkaik, A. C., Bijmens, N., Rutten, M., Van De Vosse, F. (2010). Perpendicular ultrasound velocity measurement by 2D cross correlation of RF data. Part B: Volume flow estimation in curved vessels. *Experiments in Fluids*, 2010:49, 1219-1229. [Chua Chiaco et al.(2013)] Chua Chiaco JM, Parikh NI, Ferguson DJ. The jugular venous pressure revisited. *Cleve Clin J Med*, 2013:80, 638-644. [Kalmanson et al. (1972)] Kalmanson D, Veyrat C, Derai C, Savier CH, Berkman M, Chiche P. Non-invasive technique for diagnosing atrial septal defect and assessing shunt volume using directional Doppler ultrasound. Correlations with phasic flow velocity patterns of the shunt. *British Heart Journal*, 1972:34, 981-991. [Kohler(2002)] Kohler, B. U., Hennig, C., Orglmeister, R. (2002). The principles of software QRS detection. *IEEE Engineering in Medicine and Biology Magazine*, 2002:21(1), 42-57. doi:10.1109/51.993193 [Mackenzie(1902)] Mackenzie J. *The Study of the Pulse, Arterial, Venous, and Hepatic and of the Movements of the Heart*. Edinburgh: Young & Pentland, 1902. [Patel et al.(1999)] Patel, M. C., Berman, L. H., Moss, H. a, McPherson, S. J. (1999). Subclavian and internal jugular veins at Doppler US: abnormal cardiac pulsatility and respiratory phasicity as a predictor of complete central occlusion. *Radiology*, 1999:211(2), 579-583. [Pyhel(1978)] H. JACK PYHEL, JANIE STEWART, A. M. E. T. Clinical assessment of