

Reference excerpt 02-044

Source: arXiv:1604.05177v1 | License: CC BY 4.0 | Word window: 1632-1891

in Medicine, 2011;37(5), 788-797. [Beulen et al.(2010)] Beulen, B., Verkaik, A. C., Bijnens, N., Rutten, M., Van De Vosse, F. (2010). Perpendicular ultrasound velocity measurement by 2D cross correlation of RF data. Part B: Volume flow estimation in curved vessels. Experiments in Fluids, 2010;49, 1219-1229. [Chua Chiacio et al.(2013)] Chua Chiacio JM, Parikh NI, Ferguson DJ. The jugular venous pressure revisited. Cleve Clin J Med, 2013;80, 638-644. [Kalmanson et al. (1972)] Kalmanson D, Veyrat C, Derai C, Savier CH, Berkman M, Chiche P. Non-invasive technique for diagnosing atrial septal defect and assessing shunt volume using directional Doppler ultrasound. Correlations with phasic flow velocity patterns of the shunt. British Heart Journal, 1972;34, 981-991. [Kohler(2002)] Kohler, B. U., Hennig, C., Orglmeister, R. (2002). The principles of software QRS detection. IEEE Engineering in Medicine and Biology Magazine, 2002;21(1), 42-57. doi:10.1109/51.993193 [Mackenzie(1902)] Mackenzie J. The Study of the Pulse, Arterial, Venous, and Hepatic and of the Movements of the Heart. Edinburgh: Young & Pentland, 1902. [Patel et al.(1999)] Patel, M. C., Berman, L. H., Moss, H. A., McPherson, S. J. (1999). Subclavian and internal jugular veins at Doppler US: abnormal cardiac pulsatility and respiratory phasicity as a predictor of complete central occlusion. Radiology, 1999;211(2), 579-583. [Pyhel(1978)] H. JACK PYHEL, JANIE STEWART, A. M. E. T. Clinical assessment of calibrated jugular pulse recording. British Heart Journal, 1978, 297-302. [Mackay(1947)] Mackay IF. An experimental analysis of the jugular pulse in man. J Physiol (Lond.), 1947;106, 113-118. [Rizkallah(2014)] Rizkallah, J., Jack, M., Saeed, M., Shaffer, L. A.,