

Reference excerpt 02-046

Source: arXiv:1604.05177v1 | License: CC BY 4.0 | Word window: 1708-1967

Veyrat C, Derai C, Savier CH, Berkman M, Chiche P. Non-invasive technique for diagnosing atrial septal defect and assessing shunt volume using directional Doppler ultrasound. Correlations with phasic flow velocity patterns of the shunt. *British Heart Journal*, 1972;34, 981-991. [Kohler(2002)] Kohler, B. U., Hennig, C., Orglmeister, R. (2002). The principles of software QRS detection. *IEEE Engineering in Medicine and Biology Magazine*, 2002;21(1), 42-57. doi:10.1109/51.993193 [Mackenzie(1902)] Mackenzie J. The Study of the Pulse, Arterial, Venous, and Hepatic and of the Movements of the Heart. Edinburgh: Young & Pentland, 1902. [Patel et al.(1999)] Patel, M. C., Berman, L. H., Moss, H. A., McPherson, S. J. (1999). Subclavian and internal jugular veins at Doppler US: abnormal cardiac pulsatility and respiratory phasicity as a predictor of complete central occlusion. *Radiology*, 1999;211(2), 579-583. [Pyhel(1978)] H. JACK PYHEL, JANIE STEWART, A. M. E. T. Clinical assessment of calibrated jugular pulse recording. *British Heart Journal*, 1978, 297-302. [Mackay(1947)] Mackay IF. An experimental analysis of the jugular pulse in man. *J Physiol (Lond.)*, 1947;106, 113- 118. [Rizkallah(2014)] Rizkallah, J., Jack, M., Saeed, M., Shaffer, L. A., Vo, M., Tam, J.. Non-invasive bedside assessment of central venous pressure: scanning into the future. *PloS One*, 2014;9. doi:10.1371/journal.pone.0109215 [Sahani et al.(2015)] Sahani, A K, Shah, M, Joseph, J, Sivaprakasam, M. Carotid and Jugular Classification in ARTSENS. *JBHI*, 2015;PP:1 [Schaller(2004)] Schaller B. Physiology of cerebral venous blood flow: from experimental data in animals to normal function in humans. *Brain research. Brain research reviews*, 2004;46. [Sisini et al.(2015)] Sisini F, Tessari M, Gadda G, Di Domenico