Asthma care plan for education and care services

CONFIDENTIAL: Staff are trained in asthma first aid (see overleaf) and can provide routine asthma medication as authorised in this care plan by the treating doctor. Please advise staff in writing of any changes to this plan.

To be completed by the treating doctor and parent/guardian, for supervising staff and emergency medical personnel.

PLEASE PRINT CLEARLY

Photo of child (optional)

Plan date /____/201_

				_	Review date //201	
Child's name			Date of birth			
Managing an as Staff are trained i	rthma attack n asthma first aid (see overl	eaf). Please write down an	ything different this child m	ight need if they l	nave an asthma attack:	
Daily asthma m	_					
This child's usual	asthma signs	Frequency and severion Daily/most days	ity	Known triggers for this child's asthma (eg exercise*, colds/flu, smoke) — please detail:		
☐ Wheeze			e than 5 x per year)			
 □ Difficulty bre	athing		s than 5 x per year)			
Other (please		Other (please de				
Does this child n Does this child u *Does this child Medication pla		dication? nedication before exercise	☐ Yes ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐	No No No	ind to staff	
	s asthma medication, please			er/mask are suppi		
Name of medi	cation and colour	Dose/nur	mber of puffs		Time required	
Doctor Name of doctor Address		attachments listed. I approve to and emergency medical perso there are any changes to these	I have read, understood and agreed with this care plan and any attachments listed. I approve the release of this information to staff and emergency medical personnel. I will notify the staff in writing if there are any changes to these instructions. I understand staff will seek emergency medical help as needed and that I am responsible		Emergency contact information Contact name	
		for payment of any emergency medical costs.		Phone		
	Phone	Signature Date		Mobile		
Signature	Date	Name		Email		









Asthma First Aid

- **1** Sit the person upright
 - Be calm and reassuring
 - Do not leave them alone



- Give 4 separate puffs of blue/grey reliever puffer
 - Shake puffer
 - Put **1 puff** into spacer
 - Take 4 breaths from spacer

Repeat until 4 puffs have been taken

Remember: Shake, 1 puff, 4 breaths

OR give 2 separate doses of a Bricanyl inhaler (age 6 & over) or a Symbicort inhaler (over 12)



- Wait 4 minutes
 - If there is no improvement, give <u>4 more separate puffs of blue/grey reliever</u> as above

OR give 1 more dose of Bricanyl or Symbicort inhaler



- If there is still no improvement call emergency assistance Dial Triple Zero (000)
 - Say 'ambulance' and that someone is having an asthma attack
 - Keep giving <u>4 separate puffs</u> every <u>4 minutes</u> until emergency assistance arrives

OR give 1 dose of a Bricanyl or Symbicort every 4 minutes - up to 3 more doses of Symbicort



Call emergency assistance immediately - Dial Triple Zero (000)

- If the person is not breathing
- If the person's asthma suddenly becomes worse or is not improving
- If the person is having an asthma attack and a reliever is not available
- If you are not sure if it's asthma
- If the person is known to have Anaphylaxis follow their Anaphylaxis Action Plan, then give Asthma First Aid

Blue/grey reliever medication is unlikely to harm, even if the person does not have asthma



Contact your local Asthma Foundation **1800 ASTHMA Helpline** (1800 278 462) **asthmaaustralia.org.au**

