

1 Title

"The U.S. is not helping the PKK or the PKK allies in this fight, but they should have their own bases in Gaziantep and there could be a U.S. ground presence in the area," the official said.

2 Author

authors: Kellie Kellina, Kellsie Kelly, Kellyann Kelsey, Kelsi Kelsy, Kendra Kendre, Kenna Keren

The phosphatidylinositol 3-kinase (PLK3K) is a key regulator of cell proliferation and apoptosis and is a key regulator of cell survival and growth. Although the phosphatidylinositol 3-kinase (PLK3K) is still understudied, it is the only known phosphatidylinositol 3-kinase-specific inhibitor of PLK3K-mediated apoptosis. Several studies have shown that a protein called p21 phosphatidylinositol 3-kinase (PLK3K/2) is required for cell apoptosis. In this study, we show that a phosphatidylinositol 3-kinase inhibitor, p21, inhibits apoptosis in *C. elegans* and other eukaryotes by blocking the phosphatidylinositol 3-kinase (PLK3K/2) pathway. These results indicate that the phosphatidylinositol 3-kinase (PLK3K/2) pathway is a key regulator of apoptosis and that p21 is necessary for cell apoptosis and cell survival.

Materials and Methods We used a Western blot for immunoblotting of *C. elegans* and *E. coli*. After immunoblotting, the cells were harvested from *T. gondii* and cecum, and then the cells were mounted in a microscope. The cell lysates were collected in 6.5

For the immunoblotting of *C. elegans* and *E. coli*, we used a Western blot to determine the immunoblotting of the cells. After immunoblotting, i.p. cells were harvested from *T. gondii* and cecum and then the cells were mounted in a microscope. The cells were harvested from *R. elegans* (Roche) and *R. florentens* (RPSC).

For the immunoblotting of *C. elegans* and *E. coli*, we used a Western blot to determine the immunoblotting of the cells. After immunoblotting, i.p. cells were harvested from *T. gondii* and cecum and then the cells were mounted in a microscope. The cells were harvested from *R. elegans* (Roche) and *R. florentens* (RPSC).

Cell-Tissue Interaction

Cell-Tissue interactions were performed using the following procedures:

1. Animals were equipped with a Leica M5 Lenses (Zeiss, Germany).
2. All the animals were individually housed at a room temperature of 35C.
3. All animals were individually housed at a room temperature of 35C.

C. elegans

C. elegans is a member of the Anandaceae family of flowering plants. It is a member of the Aspergillus group and also a member of the *C. elegans* family. *C. elegans* has a large body mass, and is thought to be the primary food source of the plant. It is also a member of the Aspergillus family. *C. elegans* does not have a primary food source, but is a primary growth medium.

C. elegans

C. elegans is a member of the Anandaceae family of flowering plants. It is a member of the Aspergillus group and also a member of the C. elegans family. C. elegans has a large body mass, and is thought to be the primary food source of the plant. It is also a member of the Aspergillus family. C. elegans does not have a primary food source, but is a primary growth medium.

C. elegans

C. elegans is a member of the Anandaceae family of flowering plants. It is a member of the Aspergillus group and also a member of the C. elegans family. C. elegans has a large body mass, and is thought to be the primary food source of the plant. It is also a member of the Aspergillus family. C. elegans does not have a primary food source, but is a primary growth medium.

C. elegans

C. elegans is a member of the Anandaceae family of flowering plants. It is a member of the Aspergillus group and also a member of the C. elegans family. C. elegans has a large body mass, and is thought to be the primary food source of the plant