

1 Title

A potential target for MMP-2, a dimerase involved in taurine biosynthesis and cell growth modulation

2 Author

authors: Nikoletta Nikolia, Nil Nina, Ninetta Ninette, Ninnetta Ninnette, Ninon Nisa, Nissa Nisse

P.Larsson, G.P. Scheifler, N.C. Norvig, O.M. Meier, and J.R. Bergstedt. 2013. The effect of monoclonal antibodies on the expression of a protein that is the basis for the protein tyrosine hydroxylase in human blood. *Blood*. 346: 13411352.

(C) 1996.

Abstract

The role of the cytoplasmic protein tyrosine hydroxylase in the structural stability of blood vessels.

The tyrosine hydroxylase is a pivotal component of the cyclic-functioning proteins that control the cyclic-functioning activity of blood vessels. The cytoplasmic protein tyrosine hydroxylase (Tyr) is essential for the cyclic-functioning activity of blood vessels.

The tyrosine hydroxylase (Tyr), which is a cytoplasmic protein, is essential for the cyclic-functioning activity of blood vessels. The tyrosine hydroxylase (Tyr) is also the backbone of the cytoplasmic protein, to which the cyclic-functioning activity of blood vessels can be transferred by the cytoplasmic protein tyrosine hydroxylase [31].

The tyrosine hydroxylase (Tyr) is required for the cyclic-functioning activity of blood vessels. The tyrosine hydroxylase (Tyr) is also the backbone of the cytoplasmic protein, to which the cyclic-functioning activity of blood vessels can be transferred by the cytoplasmic protein tyrosine hydroxylase (Tyr). The tyrosine hydroxylase (Tyr) is also a cytoplasmic protein, to which the cyclic-functioning activity of blood vessels can be transferred by the cytoplasmic protein tyrosine hydroxylase (Tyr) [31].

The cyclic-functioning activity of blood vessels can be transferred by the cytoplasmic protein tyrosine hydroxylase (Tyr). The cyclic-functioning activity of blood vessels is also the backbone. The cytoplasmic protein is essential for the cyclic-functioning activity of blood vessels.

The tyrosine hydroxylase (Tyr) is required for the cyclic-functioning activity of blood vessels. The tyrosine hydroxylase (Tyr) is also the backbone of the cytoplasmic protein, to which the cyclic-functioning activity of blood vessels can be transferred by the cytoplasmic protein tyrosine hydroxylase (Tyr). The tyrosine hydroxylase (Tyr) is also a cytoplasmic protein, to which the cyclic-functioning activity of blood vessels can be transferred by the cyclic-functioning protein tyrosine hydroxylase (Tyr).

The cytoplasmic protein tyrosine hydroxylase (Tyr) is essential for the cyclic-functioning activity of blood vessels. The cytoplasmic protein tyrosine hydroxylase (Tyr) is also the backbone of the cytoplasmic protein, to which the cyclic-functioning activity of blood vessels can be transferred by the cytoplasmic protein tyrosine hydroxylase (Tyr). The

cytoplasmic protein is also a cytoplasmic protein, to which the cyclic-functioning activity of blood vessels can be transferred by the cytoplasmic protein tyrosine hydroxylase (Tyr).

The tyrosine hydroxylase is required for the cyclic-functioning activity of blood vessels. The tyrosine hydroxylase (Tyr) is also the backbone of the cytoplasmic protein, to which the cyclic-functioning activity of blood vessels can be transferred by the cytoplasmic protein tyrosine hydroxylase (Tyr). The tyrosine hydroxylase (Tyr) is also a cytoplasmic protein, to which the cyclic-functioning activity of blood vessels can be transferred by the cytoplasmic protein tyrosine hydroxylase (Tyr). The tyrosine hydroxylase (Tyr) is also a cytoplasmic protein, to which the cyclic-functioning activity of blood vessels can be transferred by the cytoplasmic protein tyrosine hydroxylase (Tyr).

In the case of the cytoplasmic protein tyrosine hydroxylase (Tyr), the cytoplasmic protein tyrosine hydroxyl